


THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO









BUREAU OF FISHERIES

REPORT OF
THE COMMISSIONER OF FISHERIES
FOR THE FISCAL YEAR 1910
AND
SPECIAL PAPERS

GEORGE M. BOWERS

Commissioner



WASHINGTON
GOVERNMENT PRINTING OFFICE

1911



CONTENTS.

- REPORT OF THE COMMISSIONER OF FISHERIES FOR THE FISCAL YEAR ENDED JUNE 30, 1910.** Document 734, 40 p. (Issued November 25, 1910.)
- THE DISTRIBUTION OF FISH AND FISH EGGS DURING THE FISCAL YEAR 1910.** Document 740, 112 p. (Issued June 1, 1911.)
- DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES STEAMER ALBATROSS DURING THE PHILIPPINE EXPEDITION, 1907-1910.** Document 741, 98 p. (Issued November 29, 1910.)
- CONDITION AND EXTENT OF THE NATURAL OYSTER BEDS OF DELAWARE.** By H. F. Moore. Document 745, 30 p., 1 chart. (Issued February 10, 1911.)
- THE FISHERIES OF ALASKA IN 1910.** By Millard C. Marsh and John N. Cobb. Document 746, 72 p. (Issued April 19, 1911.)
- SPECIAL INVESTIGATION OF THE ALASKA FUR-SEAL ROOKERIES, 1910.** By Harold Heath. Document 748, 22 p. (Issued November 10, 1911.)
- THE FUR-SEAL FISHERIES OF ALASKA IN 1910.** By Walter I. Lembkey. Document 749, 40 p. (Issued November 8, 1911.)
- THE SALMON FISHERIES OF THE PACIFIC COAST.** By John N. Cobb. Document 751, 180 p. (Issued November 25, 1911.)

**REPORT OF THE COMMISSIONER OF FISHERIES
FOR THE FISCAL YEAR ENDED JUNE 30, 1910**

Bureau of Fisheries Document No. 734

CONTENTS.

	Page.
General considerations	5
Propagation of food fishes.....	6
Extent of work	6
Review of operations	7
New stations and improvements.....	11
Acclimatization and results of fish culture	12
Fish-cultural relations with States and foreign countries	13
Biological inquiries and experiments	15
Oyster investigations and surveys.....	15
Pearl-mussel investigations.....	16
Experiments in sponge culture	17
Study of fish diseases	17
Other inquiries and experiments	18
Marine biological laboratories	19
Alaska salmon service	19
Alaska fur-seal service.....	21
The fishery industries	22
Statistics and methods of the fisheries	22
Shad and alewife fisheries.....	27
Investigation of the mackerel fishery	28
Fisheries of Mississippi	31
Miscellaneous activities	33
Relations with other government bureaus.....	33
International fishery matters	33
Employment of vessels	34
Publications and library	35
Appropriations.....	35
Recommendations.....	36
Reorganization of personnel.....	36
Salaries and personnel.....	36
Sponge law	37
Extension of fish culture.....	38
Laboratory for the study of fish diseases.....	39
Fishery intelligence service	40
New building	40

REPORT
OF THE
COMMISSIONER OF FISHERIES.

DEPARTMENT OF COMMERCE AND LABOR,
BUREAU OF FISHERIES,
Washington, August 24, 1910.

SIR: I have the honor to submit herewith a report of the operations of the Bureau of Fisheries for the fiscal year ended June 30, 1910.

GENERAL CONSIDERATIONS.

This Bureau was organized as the United States Fish Commission in February, 1871, and on June 30, 1910, therefore, it completed the fortieth fiscal year of its existence. Originally clothed solely with functions of investigation and inquiry into the reputed or real decrease in the food fishes of the coastal and interior waters, it soon manifested that it could perform important service in actually increasing the supply of such fishes. In recognition of this fact acts of Congress from time to time have enlarged the functions of the Bureau until to-day the purely practical work of increasing and conserving aquatic food resources through cultural and experimental operations has become the dominant feature of the Bureau's activities.

For a long while wholly relieved of executive control of the fisheries by reason of the constitutional reservation of that right to the States, the Bureau recently has been invested with the administration of the important fisheries of Alaska, including the entire control of the Pribilof Islands and the fur-bearing animals of the Territory at large.

The steady increase in the volume and importance of the Bureau's work has been especially rapid in the past ten years, and the fiscal year just closed, which witnessed a drastic change in the control of the seal herd, has added considerably to the sum of the Bureau's duties. The probable adoption of joint international regulations in respect to the fisheries of the waters contiguous to our northern boundary presents the possibility of a great enlargement of the

Bureau's executive functions in the near future. Each year brings increasing demands from the several States for aid and advice in respect to the drafting of laws and regulations, the establishment of state fishery services, and the best measures for the conservation and development of fishery resources, and the Bureau feels that its influence for good in matters relating to the fisheries is yearly becoming more important. The salient features of the work during the fiscal year are exhibited in the following pages.

PROPAGATION OF FOOD FISHES.

EXTENT OF WORK.

It is gratifying to be able to record another successful year in fish-cultural work. Methods have not varied appreciably from those of former years, and attention has been directed principally to enlarging the output.

The widespread and increasing interest taken in the Bureau's work by people in all sections of the country and the growing conception of the benefits resulting from the stocking of public and private waters are manifested by the large number of applications for fish received during the year, the number being 10,635, an increase of 523 over 1909.

Work was conducted at 35 permanent stations and 86 field and collecting stations, located in 32 States. With reference to the fishes propagated, the regular hatcheries may be classified as follows: Marine species, 3; river fishes of the eastern seaboard, 5; fishes of the Pacific coast, 5; fishes of the Great Lakes, 7; fishes of the interior, 15.

The results of fish culture depend largely upon climatic conditions, the most elaborate and carefully executed plans ending in success or failure according to the state of the weather in the spawning season. In 1910 these conditions were generally unfavorable, resulting in the curtailment of egg collections of most of the important species, but owing to the superior quality of the majority of the eggs obtained, the Bureau was able to exceed its record year of 1909 by 126,800,000, or 4 per cent, the total output of fish and eggs being in excess of 3,233,000,000. This was accomplished without increased funds, the available appropriations being the same as in the preceding two years, and was made possible largely through the faithful and efficient service rendered by the Bureau's employees in their several lines of work.

The following is a table summarizing the distribution of fish and fish eggs for the year. Of these, 443,177,000 eggs and 7,425 fish were delivered to various state fish commissions, and 600,000 eggs of salmon and trout were shipped to foreign countries.

SUMMARY OF DISTRIBUTION OF FISH AND EGGS, FISCAL YEAR ENDED JUNE 30, 1910.

Species.	Eggs.	Fry.	Fingerlings, yearlings, and adults.	Total.
Catfish.....			544,350	544,350
Carp.....			22,710	22,710
Buffalofish.....			201,475	201,475
Shad.....	2,160,000	89,076,000		92,236,000
Whitefish.....	55,428,000	195,964,000		251,392,000
Lake herring.....	1,440,000	70,300,000		71,740,000
Silver salmon.....	375,000	10,918,025		11,293,025
Chinook salmon.....	37,531,417	16,342,556	67,525	53,941,498
Blueback salmon.....	100,000	121,136,995	21,719,000	142,956,995
Steelhead trout.....	250,000	3,570,287	179,718	3,900,005
Humpback salmon.....		1,308,000		1,308,000
Rainbow trout.....	536,494	552,716	1,771,128	2,860,338
Atlantic salmon.....	5,000	1,217,366	238,212	1,460,578
Landlocked salmon.....	115,000	985,040	304,364	1,404,404
Blackspotted trout.....	2,748,550	1,765,834	884,154	5,398,538
Loch Leven trout.....			68,248	68,248
Lake trout.....	10,210,000	33,649,622	4,286,150	48,145,772
Brook trout.....	516,000	7,405,545	4,228,461	12,150,006
Sunapee trout.....		171,029		171,029
Grayling.....	25,000	81,000	18	106,018
Pike.....			43,300	43,300
Pickereel.....			500	500
Crappie and strawberry bass.....			414,477	414,477
Rock bass.....			69,985	69,985
Warmouth bass.....			792	792
Smallmouth black bass.....		537,600	113,305	650,905
Largemouth black bass.....		36,600	673,482	730,082
Sunfish (bream).....			345,035	345,035
Pike perch.....	321,455,000	155,025,000	4,769	476,484,769
Yellow perch.....	5,200,000	326,885,000	109,245	332,194,245
Striped bass.....	4,566,000	2,784,000		7,350,000
White bass.....			6,050	6,050
White perch.....	16,500,000	338,450,000		354,950,000
Yellow bass.....			250	250
Sea bass.....		808,000		808,000
Smelt.....	4,500,000		9,060	4,509,060
Mackerel.....		764,000		764,000
Freshwater drum.....			11,950	11,950
Cod.....	9,854,000	210,354,000		220,208,000
Pollock.....		38,140,000		38,140,000
Haddock.....		712,000		712,000
Flatfish.....		930,755,000		930,755,000
Lobster.....	780,000	162,505,000	2,052	163,287,052
Total.....	474,295,461	2,722,310,215	36,326,896	3,233,332,572

REVIEW OF OPERATIONS.

The conspicuous increases in the output of fish and eggs over the year 1909 were in blueback, silver, and Atlantic salmons, lake trout, lake herring, yellow perch, shad, cod, flatfish, and steelhead trout, the production of the latter three species exceeding all previous records.

There was a slight decrease from last year in the number of chinook salmon liberated from the Pacific coast stations. Notwithstanding a normal run in the Sacramento, the season at the California stations was the poorest for thirteen years, due partly to such low water that the fish were unable to ascend the tributary streams on which the hatcheries are located, and, later, to freshets which carried away the racks and permitted the impounded fish to escape, with the loss of millions of eggs. Two causes are at present militating against the increase of salmon in these streams—the increasing numbers of black bass, which prey upon the young salmon after planting, and the

ascent of the fry by thousands into a recently constructed irrigating ditch, where they are left on the land to die. The only remedy that can be suggested is to plant the fry in the lower reaches of the rivers or establish a large hatchery at tide water, the latter method involving less expense. Unless some action is taken the number of salmon in these rivers will decline rapidly.

Taken as a whole, the work of the Oregon stations was satisfactory, although high water during the spawning of the chinook salmon shortened the season and reduced the collections to slightly below those of the previous year.

At the Washington stations, where attention is devoted chiefly to the sockeye, humpback, and silver salmons and the steelhead trout, the work was augmented by the opening of two new field stations. In Alaska, where the sockeye salmon is propagated, the yield of the two hatcheries was highly satisfactory, especially the Afognak station, operated for the first time this year.

The lake-trout, whitefish, and pike-perch work of the Great Lakes stations, while not equal to that of some seasons, gave better results than had been anticipated in view of the obstacles encountered. Potent factors in the shortage at the Michigan stations were the unusually early spawning season, followed by unfavorable weather, and the necessity of complying with recently enacted state legislation, which stipulates that the operations of the Bureau must be supervised by the state fish and game warden's department and that all eggs must be taken and fertilized by fishermen licensed by that department, thus placing the work in the hands of inexperienced men. Compliance with the provisions of this law curtailed the output of Northville and its substations fully one-fourth. The law also prohibited pike-perch collections on the St. Clair River, one of the Bureau's most productive fields in past years.

At the Duluth station the weather and other conditions were favorable, permitting increased lake-trout work, but whitefish and pike-perch operations on Lake Erie were materially interfered with by storms, although the poor collections of the latter species were offset to a great extent by the superior quality of the eggs secured.

The lobster output from the three marine stations was about equal to that of 1909. The impounded stock at the Boothbay Harbor station was stripped in April, and though the lobsters were in vigorous health the average yield of eggs was smaller than usual, due, it is believed, to their greater activity in the pound during the mild winter and the consequent shedding of many eggs. The construction of two substantial lobster pounds during the year places this station on a greatly improved basis. At the Gloucester and Woods Hole stations, which are not equipped with pounds, the lobsters collected during the fall are cared for in live cars through the winter.

The number of cod fry produced at these stations was nearly 100,000,000 greater than in 1909, the greatest gain being at the Gloucester station, where more eggs than could be handled were obtained from fishing grounds in the vicinity.

The collection of flatfish eggs was the largest ever made by the Bureau, numbering 1,195,911,000, from which 930,755,000 fry were hatched and distributed. At Boothbay Harbor, where this work has only recently been undertaken, the output was increased 100 per cent over that of the previous year.

Other marine species propagated included pollock at Gloucester, haddock at Boothbay Harbor, and mackerel and sea bass at Woods Hole.

In view of the steady decline in the shad fishery in rivers tributary to the Atlantic for the past fifteen years, it is gratifying to be able to record an increased egg collection of this species and a corresponding increase in the output of fry. The results are attributable partly to recently enacted legislation regulating the methods of fishing in the Albemarle Sound and partly to an exceedingly early spring, which started the run of fish in the Potomac River before the pound nets could be equipped, each factor permitting a larger number of fish than usual to ascend to the spawning grounds.

On the Susquehanna River, at one time the Bureau's most productive field, there was no improvement over recent years, emphasizing anew the destructive influences of unregulated fisheries and the necessity for concerted action by the States concerned if any practical results are to be obtained in the rehabilitation of this important fishery.

White and yellow perch were again produced in considerable numbers at the station on the Susquehanna River, and on the Potomac River the output of yellow perch exceeded all previous records, due to the enlargement of facilities for propagating the species.

Owing to the passage of a state law prohibiting the capture of striped bass by commercial fishermen during the spawning season, the Bureau was unable to secure eggs of this species at its California station in 1909, and as this law remains in force no attempt was made to conduct operations in 1910. The prospects are good for effective work with the striped bass in this field, and its propagation will be resumed in the event of a change in the law.

As in previous years, most of the brook-trout eggs handled at the fisheries stations are purchased from dealers, this course having proved more economical in most sections of the country than reliance upon collections from waters available for the purpose. At present only two stations—one in New England and one in Colorado—obtain their supplies of eggs from wild fish, and the fields heretofore

open to them are narrowing each year because of the encroachments of commercial fish culturists. In 1910 Wellington Lake and the Grand Mesa Lakes, heretofore the most productive sources of the Colorado station for eggs of the blackspotted, brook, and rainbow trout, had to be given up to private enterprise.

The Bureau having been requested to undertake the propagation of the blackspotted trout on the Truckee River with the view of replenishing the stock, depleted through excessive fishing, a field station was established at Derby Dam, Nevada, in the winter of 1909-10. In a normal season several millions of eggs might have been obtained, but owing to low water in the river and the destruction of large numbers of eggs by market fishermen the collections amounted to only 1,371,900. These were hatched without unusual losses and the fry deposited in the river. It seems advisable to continue operations here next season, as it is apparently a promising field for fish-cultural work.

Investigation of the streams in Yellowstone Park demonstrates the possibility of greatly extending operations with the black-spotted trout, and it is intended to increase the force of experienced men in this field with the view of making it a source of supply for the Leadville, Spearfish, and Bozeman stations. The work in the park during the past season was entirely satisfactory.

Taken as a whole, the output of the basses, sunfish, and catfish from stations in various parts of the country was good, the improved results being largely due to increased knowledge of the factors governing the successful propagation of these species. The production of pond stations was supplemented by the collections on the Mississippi and Illinois rivers, where, in addition to securing sufficient bass and allied species for restocking many depleted waters, large numbers of other fishes were seined from shallow sloughs formed by the floods and returned to the main streams. If not removed, the fish would perish from drought or cold, and their rescue conserves a valuable local food resource. A new station established at Helena, Ark., late in the summer rescued over half a million fish.

With the view of extending rescue operations over a larger territory, temporary collecting stations have been located at Caruthersville, Mo., and Rosedale, Miss., which will be made permanent auxiliary stations if experience proves favorable. It is believed that similar inexpensive stations can be advantageously established at various points on the Mississippi River from New Orleans to St. Paul, as the field for this work is extensive and the number of fish that can be economically reclaimed from the drying sloughs and lakes is governed only by the amount of money available for the purpose.

Although the propagation and general distribution of carp was discontinued many years ago, the Bureau constantly receives applica-

tions for this fish, and in instances where the waters described are unsuited to other species the requests are complied with by transferring carp from other waters. In this connection it may be interesting to quote from the census records that in 1903 the total catch of carp in the United States was 18,942,763 pounds, valued at \$442,255, and in 1908 the total catch was 42,763,100 pounds, valued at \$1,135,390.

NEW STATIONS AND IMPROVEMENTS.

Under authority of the act providing for two or more new fish-cultural stations on Puget Sound or its tributaries, a careful investigation has been made and two suitable sites decided on. As soon as title can be obtained construction will begin.

At Holden, Vt., 24.3 acres of land were acquired for an auxiliary to the station at St. Johnsbury, the facilities of which were too limited for the requirements of northern New England.

The opportunities for fish-cultural and biological work in the valley of the upper Mississippi prompted Congress to authorize a station auxiliary to that at Fairport, Iowa, but to be more particularly devoted to propagation and the saving of fishes from overflowed lands. A site of about 31 acres was purchased at Homer, Minn., about 5 miles from Winona, and a pumping plant and ponds are now nearly completed and other buildings begun. The station will be ready for operation at an early date.

Results in the past having warranted the extension of the station at Mammoth Spring, Ark., 10.5 additional acres have been purchased there for the construction of several large ponds, which will soon be ready for use.

At the Fairport, Iowa, biological station much work in grading, construction of roads, and laying out ponds was done. A building 20 by 50 feet, with pebble-dash finish, containing an office, storage room, and small laboratory equipped for experimental work in fresh-water mussel culture, was practically completed during the year. A pumping plant consisting of two gasoline engines and two centrifugal pumps was installed in a small frame building 20 by 30 feet constructed for that purpose. Eleven cement ponds (4 small ones, 6 of medium size, and 1 large one) were also constructed for practical experiments in mussel propagation.

Improvements provided for by special appropriations were made at many of the stations. At Bozeman, Mont., cement hatching troughs were installed in place of wooden ones, in accordance with modern practice, and are giving excellent results. At Boothbay Harbor, Me., a coal house was built, the wharf extended and altered, and the dams at the lobster pound completed. At Erwin, Tenn., a new hatchery was built on modern plans, the old one hav-

ing become badly dilapidated and beyond repair. The new building is a frame structure 32 by 72 feet on a concrete foundation, and contains besides the hatching room, equipped with cement troughs, an office and workrooms. The water-supply and drainage systems have also been improved and extended, and to a considerable degree built in concrete. At Duluth, Minn., a dwelling for the superintendent has been erected which is in harmony with the surrounding private structures of the city and adds to the efficiency and appearance of the reservation. It is a two-story frame structure 32 by 36 feet, containing 7 rooms and basement, with the necessary office facilities. At Greenlake, Me., the new road has been completed, facilitating the distribution of fish and eggs, shortening materially the distance over which it is necessary to haul supplies, and doing away in great part with unreliable boat transportation. At Neosho, Mo., the new pipe line providing an extra supply of water has been completed and connected with the hatchery and ponds in approved manner, and the woodwork about the ponds has been replaced by concrete. It is believed there will be no further trouble with the water supply at this point for many years to come.

At Leadville, Afognak, Yes Bay, and the Pribilof Islands no expenditures of importance have been made for account of special appropriations.

The plans and specifications for the constructions described have been prepared in the office of the Bureau's architect and engineer and the work planned and supervised by him. In addition, various surveys have been made and plotted, and maps and charts of a special nature prepared.

For fish-cultural work on Lake Erie, in connection with the Put-in-Bay station and to take the place of a boat obsolete and worn out, there was built a steel steamboat of the lake tug type 85 feet long, 16 feet beam, and 8 feet 6 inches in depth. The vessel is equipped for the special requirements, has machinery and appliances of approved design, and it is expected will be a valuable addition to the facilities of the Bureau.

ACCLIMATIZATION AND RESULTS OF FISH CULTURE.

After nearly forty years of endeavor to establish the chinook salmon of the Pacific coast in waters of the United States where it is not indigenous, conclusive evidence of success in one instance has come to hand. Within the past year it has been ascertained that the species has become established in Lake Sunapee, New Hampshire, where numerous specimens from 3 to 5 pounds in weight have been taken by anglers. This is undoubtedly the result of a plant made in 1904 by the New Hampshire fish commission, the eggs having been supplied from the Bureau's hatchery at Baird, Cal. Encouraged by the

outcome of this experiment, the Bureau made a plant of 40,000 fingerling chinook salmon in Lake Champlain in the spring of 1910.

There unquestionably has been an increase in Atlantic salmon in the Penobscot River, as evidenced by the results of the Bureau's operations in 1910 compared with 1908 and 1909. Though receiving the catch of a smaller number of weirs the past season, the collection of spawning fish was twice as great as in 1909 and 60 per cent greater than in 1908.

It is believed that owing to the planting of the species by the Bureau pike perch have become sufficiently abundant in the St. Lawrence River to warrant the collection of eggs at the Cape Vincent station, and plans will be made accordingly. The fishermen on Lake Ontario report that lake trout and whitefish, which have been planted extensively by the Bureau, are increasing rapidly, and that numbers of fishermen who were driven to other pursuits by the former depletion of the fishery are resuming operations. In 1908 the catch of these two species was 5,567 pounds, while in 1909 it increased to 12,532 pounds. A corresponding increase is shown in the take of pike perch in this lake.

The following statistics show the increasing catch of the striped-bass fishery in California, the species having first been introduced from the Atlantic coast into the waters of that State in 1879:

Year.	Pounds.	Value.	Year.	Pounds.	Value.
1889.....	16,296	\$4,073	1893.....	252,454	\$13,037
1890.....	20,119	4,021	1899.....	1,234,320	61,814
1891.....	30,674	4,602	1904.....	1,570,404	92,116
1892.....	56,209	6,488	1908.....	1,775,700	134,660

For a series of years it has been the custom at the Baird, Cal., station to select for spawning purposes large fish only, a practice which appears to be developing a larger breed of fish. Chinook salmon of the run of 1909 averaged 20 pounds in weight, an increase of about 3 pounds over the previous run. The possibilities of selective breeding are indicated by this experience.

FISH-CULTURAL RELATIONS WITH STATES AND FOREIGN COUNTRIES.

Several States still continue in force certain laws and regulations in respect to the fisheries which tend to curtail and hamper the activities of the Bureau. In some cases the States show a willingness to mitigate as far as possible the effects of laws which inadvertently interfere with the Bureau's work, but in one or two instances the legislative and executive attitude appears to be unreasonable if not hostile.

With the States in general the relations of the Bureau have always been harmonious, and a system of cooperation has developed which

has been mutually beneficial to the participants and advantageous to the public. Eggs taken and fertilized at the Bureau's stations are transferred to the state fish commissions, by which they are hatched and planted. The Bureau's expenses and difficulties in distribution are thereby reduced and simplified, and the superior local knowledge usually at the service of the state authorities is of value in indicating the most suitable localities in which to plant the fry. On Lake Erie the Ohio and Pennsylvania fishery authorities cooperated with the Bureau in the collection of eggs of the whitefish, lake cisco, and pike perch.

As shown in the following table, the fish eggs allotted to the state commissions during 1910 aggregated over 443,000,000 and were sent into 17 States:

ALLOTMENT OF FISH EGGS TO STATE FISH COMMISSIONS, FISCAL YEAR ENDED JUNE 30, 1910.^a

State and species.	Eggs.	State and species.	Eggs.
California:		New York:	
Chinook salmon.....	28,764,467	Blackspotted trout.....	50,000
Colorado:		Rainbow trout.....	41,500
Blackspotted trout.....	225,000	Landlocked salmon.....	15,000
Connecticut:		White perch.....	15,000,000
Yellow perch.....	5,200,000	North Dakota:	
Illinois:		Steelhead trout.....	100,000
Lake trout.....	500,000	Pike perch.....	10,000,000
Whitefish.....	4,000,000	Ohio:	
Pike perch.....	8,000,000	Whitefish.....	18,000,000
Rainbow trout.....	41,264	Pike perch.....	170,725,000
Michigan:		Oregon:	
Landlocked salmon.....	20,000	Chinook salmon.....	6,465,300
Lake trout.....	5,000,000	Blackspotted trout.....	175,000
Pike perch.....	34,280,000	Pennsylvania:	
Missouri:		Silver salmon.....	75,000
Brook trout.....	100,000	Blackspotted trout.....	50,000
Rainbow trout.....	25,000	Whitefish.....	31,428,000
Pike perch.....	2,000,000	Pike perch.....	96,000,000
Montana:		Washington:	
Blackspotted trout.....	550,000	Steelhead trout.....	50,000
Whitefish.....	500,000	Brook trout.....	100,000
Nevada:		Wisconsin:	
Blackspotted trout.....	422,000	Lake trout.....	4,500,000
New Hampshire:		Wyoming:	
Chinook salmon.....	100,000	Blackspotted trout.....	675,000
		Total.....	443,177,531

^a Also there were allotted to Michigan 3,500 lake trout, to Oregon 45 blackspotted trout, and to Wisconsin 3,880 lake trout, or a total of 7,425 fingerlings, yearlings, and adults.

In response to requests coming through diplomatic channels the Bureau furnished eggs to the governments of foreign countries as follows:

Country and species.	Eggs.	Country and species.	Eggs.
Argentina:		France:	
Chinook salmon.....	200,000	Blackspotted trout.....	10,000
Silver salmon.....	100,000	Japan:	
Sockeye salmon.....	100,000	Rainbow trout.....	110,000
Landlocked salmon.....	25,000	Brook trout.....	5,000
Lake trout.....	50,000	Total.....	130,000

BIOLOGICAL INQUIRIES AND EXPERIMENTS.

OYSTER INVESTIGATIONS AND SURVEYS.

The field work of the survey of the public oyster beds of James River, Virginia, which was undertaken at the request of the governor and the fish commissioner of Virginia, was brought to a conclusion on September 15, the charts and report were finished on November 30, and the printed report was issued about February 1. This survey was designed to furnish definite data concerning the location, extent, and condition of the public grounds in the James and Nansemond rivers above Newport News and to provide a foundation for needed legislation by the State. The present boundary lines are based on the survey of 1892-1894, and their justice has long been a matter of contention, the oystermen claiming that much productive bottom was omitted from the public grounds, and the planters contending that a large area of barren bottom was included. The present survey could not demonstrate the validity of the first claim, as such bottoms, if they existed, have been long since occupied for planting purposes, but it was shown that about 58 per cent of the present area of the grounds consists of barren bottom and an additional 15 per cent bears oysters too sparsely scattered to be commercially valuable. Of the 26,408.4 acres surveyed, but 7,153 acres can be regarded as actually productive. It was found also that in certain places oyster planters have encroached on the public rocks, and it was evident that in other places adjoining the planted beds the rocks had been depleted by illicit operations.

To release from the public grounds and throw open to rental a considerable area of the barren bottom and to rectify the boundary lines so as to permit adequate policing, the state fish commissioner had an enabling act introduced in the legislature at its latest session. To attain the ends sought, it unfortunately was necessary to exclude from the public grounds a small proportion of the productive bottom, and as the legislature held that this was in contravention of a constitutional provision relating to the oyster beds, the proposed law failed of passage.

At the request of the governor of Delaware, acting in his capacity as chairman of the Delaware Oyster Survey Commission, the Bureau, at the close of the fiscal year, was engaged in a survey of the natural oyster beds of Delaware, the State defraying part of the expenses for necessary temporary assistance. As in the case of the James River survey, the steamer *Fish Hawk* was detailed for the service, and a considerable part of the work was performed by her personnel.

The authorities of Alabama and Mississippi have also requested assistance and advice in connection with the management of oyster

bottoms, and a preliminary inquiry has been made to determine the most profitable and practicable assistance feasible with the resources available to the Bureau.

Cooperation with the Coast and Geodetic Survey and the Maryland Shell Fish Commission in the survey of the oyster beds of Maryland, pursuant to an act of Congress, has been continued, and the field work will be completed early in the next fiscal year. It is believed that the Bureau will have discharged all of its obligations in this connection prior to the end of the fiscal year 1911.

The experiments in the fattening of oysters at Lynnhaven Bay, Virginia, have produced better results than for several years past. During a period when practically no fat oysters could be obtained from the open waters of the bay the experimental claire was regularly producing oysters of very fine quality. In this connection the Bureau is conducting investigations of the food and feeding of oysters which have already developed some unexpected results, throwing light on practical problems confronting the oyster grower. Some minor modifications of the claire were made near the end of the fiscal year, and it is hoped that it will be possible to fatten oysters earlier in the season than has been possible heretofore.

PEARL-MUSSEL INVESTIGATIONS.

The Bureau has continued its investigations of the pearl-mussel beds of the Mississippi Valley, the material depletion of which has seriously threatened the prosperity of an important industry of that region. With the aid of persons connected with various educational institutions of the States principally interested, field parties were established for the examination of various streams in Virginia, West Virginia, Michigan, Indiana, Illinois, Kentucky, Tennessee, Arkansas, Missouri, and Oklahoma. The habits, distribution, abundance, and commercial availability of the mussels found in the several localities were studied with the view of opening new sources of supply for the manufacturers of pearl buttons and for the purpose of laying a foundation for the protection, conservation, and improvement of the existing beds.

Owing to the severity of the weather during the winter, progress in the erection of the biological station at Fairport, Iowa, authorized by Congress near the close of the preceding fiscal year, was less rapid than was desired, but on the improvement of conditions in the spring construction work went on more rapidly, and at the close of the fiscal year mussel-propagating operations were being conducted on a scale promising to yield some practical results. As was pointed out in the preceding report of the Bureau, this station is designed for the study of problems relating to the general fisheries and aquatic biology of

the Mississippi Valley, but particularly for the cultivation of the mussels employed as raw material in the pearl-button industry, a manufacturing interest giving employment to a large number of persons.

Progress has also been made in the construction of the substation at Homer, Minn., which recent investigations show can be employed for various economic purposes connected with the fisheries, in addition to mussel culture.

EXPERIMENTS IN SPONGE CULTURE.

Although the experiments in growing sponges from artificial cuttings have already developed what the Bureau regards as a practical system of sponge culture, work is still being carried on with the purpose of improving the methods and testing the effects of different environments on the rate and character of sponge growth.

The sponges grown in Cape Florida Channel, which, as reported last year, attained an average weight of 1.25 ounces each at the end of twenty-nine months, were found to average 2 ounces ten months later, some of the largest specimens weighing from 3 to 6 ounces each when thoroughly cleaned and dry. The same disparity in the rate of growth of different specimens observed in other localities was found to occur in this place, while at Soldier Key, about 7 miles distant, where the conditions appear to be equally favorable, growth was very slow.

STUDY OF FISH DISEASES.

During the fiscal year the Bureau has continued cooperation with the New York State Cancer Laboratory in the investigation of thyroid tumor or cancer in domesticated fishes. An aquarium with two independent systems of closed-water circulation, with proper means of refrigeration, has been established for the observation of salmon and trout and experiments in inoculation and treatment. Investigation at various stations of the Bureau and at other hatcheries have shown that the disease is even more widespread and general than was suspected. Considerable difficulty has been encountered in obtaining for purposes of experiment a sufficient number of fish above suspicion of infection, and it has been necessary in this effort to secure a quantity of wild trout from remote streams. Owing to the technical difficulties attending this work, which are equal to those retarding the advance of knowledge relating to the cause and nature of cancer in human beings, progress is made only by slow and painstaking steps and by the use of the most approved appliances and methods. For this reason it is highly important that the Bureau should be provided with a well-equipped laboratory

and experimental hatchery, not only for the purposes of the present investigation but for the study of the many other diseases affecting fishes, both under domestication and in a state of nature. The President, in a special message to Congress dated April 9, 1910, urgently recommended an appropriation for this purpose.

During the year the Bureau was called on to investigate epidemics among hatchery fish at Spruce Creek, Pa., and Roxbury, Vt. At the former place the mortality was due in part to the thyroid tumor or cancer before alluded to, but the majority of the deaths were apparently caused by a bacterial infection which the Bureau has found at other places, but which it has not the facilities to study at present. At Roxbury the disease is also infectious and annually causes large losses. The Bureau has likewise made investigations in Pennsylvania, Ohio, and West Virginia upon the kindred subject of the pollution of streams in its relation to fishes and the fisheries.

OTHER INQUIRIES AND EXPERIMENTS.

The investigations of the Pacific coast salmons have materially advanced knowledge of the subjects during the year, particularly in respect to parasitism and the changes in the tissues affecting the food value of the fish at and near the breeding season, and in regard to the relationship of the steelhead trout and rainbow trout.

In connection with the State Geological and Natural History Survey, the Bureau has continued examination of lakes in Wisconsin, with particular regard to the gaseous content of their waters. The relationship of this subject to practical fish culture is highly important, and the data so far obtained have thrown light on certain failures in the acclimatization of fishes, the causes of which have been obscure. The study of the physical environment and habits of the salmon, smelt, and other fishes of Sebago Lake, Maine, were continued, and in response to a request a somewhat similar line of research was undertaken in Sunapee Lake, New Hampshire. In the latter locality there is a considerable fishery for smelts as they ascend the streams to spawn, and it was learned that young chinook salmon planted in the brooks were taken with the smelts.

The survey of the fishing grounds and investigation of the aquatic resources of the Philippine Islands, in which the steamer *Albatross* has been employed since the autumn of 1907, was brought to a conclusion in October, 1909. The vessel returned to San Francisco on May 4, 1910. The Philippine expedition has yielded extensive collections and a large amount of information relating to the fisheries and fishery resources, and the material is now in course of study for the preparation of comprehensive reports on the scientific and economic results.

MARINE BIOLOGICAL LABORATORIES.

The marine biological laboratories maintained by the Bureau at Woods Hole, Mass., and Beaufort, N. C., were open as usual for several months during the summer and fall, and their facilities were availed of by the usual number of investigators. The researches carried on covered a considerable range of subjects and embraced investigations of a number of species of economic importance, including the diamond-back terrapin, fishes, stone crab, quahog or hard clam, oysters, mussels, and seaweeds. The year witnessed the completion of an elaborate report by the director of Woods Hole laboratory on the marine biology of the waters adjacent to the station, embodying the results of investigations carried on for many years.

ALASKA SALMON SERVICE.

The report of the agents at the salmon fisheries of Alaska, which was published in April, 1910, includes the data for the fishing season of 1909, practically all of which was embraced in the fiscal year 1910.

The number of salmon taken during the season was about equal to the catch of 1907, but fully 20 per cent less than the number caught in 1908. In 1909 there were taken 34,692,608 fish of a gross weight of 175,028,594 pounds, as compared with 43,304,979 fish weighing 213,378,570 pounds caught in 1908. The decrease was apparent in all species excepting the king salmon, which exhibited an increase of about 55 per cent. The catch of red salmon was 115,120,670 pounds, as compared with 124,713,630 pounds in 1908; of humpbacks, 37,965,928 pounds, as compared with 60,424,620 pounds; of dog salmon, 9,456,048 pounds, as compared with 18,066,576 pounds; of king salmon, 8,959,544 pounds, as compared with 5,757,246 pounds; and of cohos, 3,526,404 pounds, as compared with 4,416,498 pounds.

The total pack of canned salmon in 1909 was 2,403,669 cases, valued at \$9,439,152. There were 45 canneries in operation, a decrease of 5 since 1908, and the total investment in the industry, excluding cash capital, was \$8,631,345. In addition to the canned pack, the fishery produced pickled salmon to the value of \$208,758, mild-cured salmon valued at \$149,300, and some minor products.

The total yield of the salmon industry was valued at \$9,796,210, produced by an investment of \$9,007,037 and the labor of 11,439 persons.

Owing to the vigilant enforcement of the laws by the agents of the Bureau during the preceding year, there were comparatively few complaints of violations during 1909. Several convictions were obtained for fishing during the weekly close season, but those engaged in the fishery showed a general desire to comply with the laws and the regulations of the Department. The pernicious practice of

“jigging” for salmon, which results in the cruel mutilation of fish which afterwards escape and die, has been stopped, and prohibition has been placed on the tourists’ practice of catching in their hands the nutritively useless but reproductively valuable spawning fish struggling up the falls and rapids.

The effort to prevent the waste of edible portions of salmons, the choice parts of which have been pickled under former practices, has been successful, the salteries now pickling the entire fish or utilizing in other ways the edible parts formerly thrown away.

The statistics relating to the operations of the government and private fish hatcheries in Alaska will not be available until the return of the agents from the Territory.

The counting of the salmon passing into Wood River, which was begun in the preceding year, was continued during the run of 1909. The spawning fish numbered but 893,000, as compared with 2,600,000 in 1908, and the catch of fish in Nushagak Bay, to which Wood River is a tributary, was but 4,900,000, as compared with 6,400,000 in the year before. It is estimated that between 6,200,000 and 7,400,000 fish entered the Nushagak basin, and that between 20 and 35 per cent escaped to the spawning grounds, as compared with a total run of between 10,100,000 and 13,600,000 fish and an escape of between 37 and 53 per cent in 1908. From the valuable but still insufficient data so far obtained it appears that for every salmon reaching the spawning grounds from two to five return several years later, and that of these from one to four may be taken without impairing the fishery. These are highly probable extremes, and the present rate of reproductive increase is between the two.

In the minor fisheries of Alaska cod were taken to the value of \$118,821 and halibut worth \$195,529. There were employed in these fisheries fixed capital to the value of \$503,837 and 548 persons. In addition there is a fleet of vessels from California and Washington fishing in Alaskan waters, the data for which are not included in the above.

The Bureau is making an effort to stop the use of food fishes for fertilizer and to stimulate the utilization of scraps and waste fishes for that purpose. This is not only in the interest of economy of consumption, but to prevent the pollution of waters through the discharge of putrescent wastes. It therefore recommends the enactment of laws prohibiting the manufacture of fertilizer from food fishes and the extension of the antipollution act of March 3, 1899, in such manner as to protect the fisheries of Alaska.

Suitable vessels for the use of the salmon-inspection service are urgently required, and provision should be made by law for the regulation and limitation of the future establishment of plants for utilizing salmon.

Attention is again called to the fact that the personnel of the Alaska salmon service is entirely inadequate to a proper enforcement of the laws and regulations and the carrying on of investigations essential to a proper and intelligent administration of these important fisheries. Several additional scientific assistants are urgently needed in this service.

ALASKA FUR-SEAL SERVICE.

By an act of Congress approved April 21, 1910, that portion of the previous law requiring the Secretary of Commerce and Labor to lease the privilege of killing seals on the Pribilof Islands was repealed, and as the lease of the North American Commercial Company expired by limitation on April 30, 1910, the Bureau, under the direction of the Secretary of Commerce and Labor, assumed the entire administration of the islands, including the functions and obligations previously imposed on the lessees. The present duties of the Bureau on the islands therefore embrace all matters whatsoever relating to the seal herd and the care, education, and welfare of the native population.

Owing to the abuses connected with pelagic sealing mentioned in the preceding report of the Bureau, the condition of the seal herd is more precarious than at any previous period of its known history, and the utmost care must be exercised to save it from commercial extinction. In anticipation of the expiration of the lease recently in force and in view of the advisability of a change in the methods of administering the islands, the Bureau called a meeting of the advisory board mentioned in the last report, which, together with the employees of the Bureau, embraces practically all of the available naturalists and officials whose experience on the islands qualifies them to pass in judgment upon the present requirements of the seal herd. The Bureau has based its policy in respect to the islands upon the unanimous advice and recommendations of the parties to this conference.

The preponderance of the pelagic kill on the high seas, which is beyond the Bureau's control, consists of mature cow seals, and for reasons that are recognized by those having knowledge of the habits of the fur seal the killing of a limited number of the excess of immature males has been deemed advisable. No definite quota has been fixed, but the number is to be determined by the agents on the islands governed by certain rigid limitations as to age, sex, size, and the minimum number to be reserved for future breeding. The breeding reserve is to be selected, as far as possible, from the most vigorous and perfect individuals, with a view to the gradual improvement of the herd.

Under the provisions of the act of April 21, 1910, the Secretary of Commerce and Labor is charged with all matters pertaining to the care and preservation of all the fur-bearing animals of Alaska. Under this authority the Bureau has drawn regulations relating to the killing or capture in Alaska of certain fur-bearing animals other than seals, and said regulations, having been signed and promulgated by the Secretary of Commerce and Labor, are now effective in the Territory.

For the purpose of putting into effect the provisions of the act above referred to, the sum of \$150,000 was appropriated. The immediately necessary additional employees required by the enlargement of the Bureau's functions on the islands have been appointed. The Bureau, under authority of the law and by direction of the Secretary of Commerce and Labor, has entered into negotiations for the purchase of the buildings, boats, and other property of the North American Commercial Company on the islands. The company has placed an apparently reasonable valuation on its property, and the proposition is under consideration subject to the results of an inventory now being made by an agent of the Bureau on the islands.

The data relating to the killing and the condition of the seal herds to July 31, 1909, were published in the preceding report of the Bureau. Those for the season of 1910 are not available at the time of writing the present report, and in any event are more strictly germane to the succeeding fiscal year.

THE FISHERY INDUSTRIES.

STATISTICS AND METHODS OF THE FISHERIES.

The commercial fisheries of the United States, including the various fishery industries dependent upon them, represent an investment of about \$95,000,000, and the value of the products derived from the fisheries proper is about \$62,000,000. With the exception of the mackerel and some other fisheries that for a number of years have not been as extensive as formerly, all of the more important branches of the industry are in a prosperous condition. The catch of mackerel during the past year was smaller than in the previous year, amounting to 46,439 barrels fresh and 17,542 barrels salted in 1909, against 57,566 barrels fresh and 21,267 barrels salted in 1908. The spring fishery in 1910 was poorer than for a number of years past, the catch up to July 1 being only 16,410 barrels of fresh mackerel and only 2,490 barrels of salted mackerel. It was an exceptionally unfavorable season for the seiners, as they took only about 2,200 barrels of the total catch of fresh mackerel, the remainder being caught by the gill-net fishermen. The fish were larger than usual, many of them weighing from 3 to 4 pounds each, but the greater portion from 2 to 3 pounds each. The fleet numbered about 50 seiners and 125 netters.

Prices were good and some of the netters made large stocks. The first mackerel of the season were landed on April 8, at Fort Monroe, Va., the fare consisting of 1,200 fish weighing $2\frac{1}{2}$ pounds each. The seiners reported seeing a good body of fish off the southern coast, but they were wild and could not be caught with seines. Of the fresh mackerel landed, 1,000 barrels were caught on Nantucket Shoals and the remainder mostly off the coast of New Jersey and in the vicinity of Block Island. The salted mackerel were all from the Cape Shore, and were all large fish. The light catch so far during the season on the Cape Shore is attributed to the fact that the fish passed along the coast far offshore outside of the fleet.

The investigation of the fisheries of the Philippine Islands was completed before the close of the year, and the statistics and other information relating to the commercial fisheries are being compiled.

A canvass of the salmon fisheries of the Pacific coast has also been made and the returns will be published at an early date.

In the spring of 1910 a beginning was made in the collection of comprehensive statistics of the oyster fishery. This is the greatest single national fishery in the world, and of itself yields a more valuable product than that derived from the entire fisheries of many important maritime countries. The work is demanded in the interests of the trade and for enlightened legislative regulation of the fishery. A canvass of the shad fisheries of the South Atlantic States was begun at the same time, and both inquiries were in progress at the close of the year.

The usual information was collected by the local agents at Boston and Gloucester, Mass., as to the quantity and value of fishery products landed at those ports by American fishing vessels during the year. The investigation of the movements of mackerel was concluded, and an inquiry was made regarding the condition of the shad and alewife fisheries of Chesapeake Bay and tributaries, and the fisheries of Mississippi.

The statistics collected by the local agents at Boston and Gloucester, Mass., of the extensive vessel fisheries at those ports have been published as monthly bulletins and distributed to the trade in various parts of the country, and also as annual bulletins giving the quantity and value of fishery products landed by American fishing vessels by months and by fishing grounds for the calendar year. The number of trips landed at these ports in 1909 was 6,306, aggregating 173,102,224 pounds of fish, valued at \$4,616,444. Compared with the previous year the receipts have decreased 8,363,023 pounds in quantity and \$12,981 in value. There was a decrease in the catch of cusk, hake, and mackerel, but an increase in that of cod, pollock, and halibut. The statistics are given in detail on the following pages.

QUANTITIES AND VALUES OF CERTAIN FISHERY PRODUCTS LANDED AT BOSTON AND GLOUCESTER, MASS., BY AMERICAN FISHING VESSELS DURING 1909, BY MONTHS.

Month.	No. of trips.	Cod.		Cusk.		Haddock.	
		Fresh.	Salted.	Fresh.	Salted.	Fresh.	Salted.
		Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
LANDED AT BOSTON.							
January.....	287	1,211,700	\$31,170	121,900	\$2,265	3,394,000	\$84,561
February.....	310	1,373,000	50,359	169,700	3,766	4,801,100	127,821
March.....	327	2,408,300	70,673	289,000	2,749	5,210,500	132,470
April.....	518	1,689,700	47,942	289,000	4,288	4,043,800	91,698
May.....	357	1,984,400	44,892	310,000	3,609	1,715,600	44,588
June.....	305	2,696,800	75,888	90,500	1,398	2,284,100	45,400
July.....	317	3,082,000	76,982	67,900	1,143	2,388,200	42,791
August.....	362	2,931,800	75,275	43,700	2,573	2,000,800	68,714
September.....	330	2,843,800	76,922	144,500	2,941	3,800,450	66,096
October.....	464	2,440,900	79,378	195,000	2,941	3,584,600	96,513
November.....	475	1,931,850	62,177	134,700	2,038	80,795	2,295,100
December.....	410	1,245,250	50,010	233,500	3,982	2,306,400	85,237
Total.....	4,798	25,840,700	741,578	1,962,700	31,521	38,485,250	966,744
LANDED AT GLOUCESTER.							
January.....	77	337,373	5,982	30,395	478	327,080	6,120
February.....	43	131,311	3,464	58,170	850	571,686	11,626
March.....	76	782,729	13,131	21,510	327	8,342	553,864
April.....	65	383,267	5,827	245,890	3,082	624,560	3,748
May.....	121	164,685	2,461	185,236	3,657	19,460	118
June.....	202	783,700	17,569	131,585	1,973	159,073	955
July.....	202	1,128,505	20,920	227,736	3,611	47,000	2,810
August.....	143	1,727,503	29,753	86,743	1,413	468,668	1,142
September.....	147	3,117,323	50,052	83,990	1,348	203,350	1,320
October.....	153	2,781,693	42,899	29,942	4,749	227,540	2,078
November.....	197	1,284,533	20,040	78,359	1,215	284,185	4,273
December.....	82	86,719	2,371	11,182	1,172	172,815	5,348
Total.....	1,508	12,749,431	213,569	1,188,238	18,370	3,915,516	56,275
Grand total.....	6,306	38,590,131	955,147	3,147,938	49,891	42,400,766	1,023,019
Grounds E. of 66° W. long.....							
Grounds W. of 66° W. long.....	752	10,555,761	189,694	532,702	8,770	5,055,621	115,084
Landed at Boston in 1908.....	5,554	28,034,370	765,453	2,610,236	41,121	37,845,145	907,965
Landed at Gloucester in 1908.....	4,542	28,329,200	791,821	3,669,100	32,204	39,834,500	1,027,866
Landed at Gloucester in 1908.....	2,007	13,286,077	250,862	21,832,454	702,801	7,008,365	640,507
Total.....	12,555	80,195,308	2,297,632	11,144,792	330,997	102,723,775	2,772,216

Month.	Hake.				Follock.				Halibut.			
	Fresh.		Salted.		Fresh.		Salted.		Fresh.		Salted.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
LANDED AT BOSTON.												
January.....	376,000	\$7,611	183,700	\$4,450	69,500	\$7,250
February.....	362,100	13,181	172,600	6,183	74,200	6,435
March.....	393,500	10,300	264,000	7,933	180,200	12,899
April.....	508,100	10,620	273,200	4,926	117,300	7,976
May.....	1,049,800	13,755	276,900	3,770	163,300	11,825
June.....	875,400	9,013	283,600	3,379	215,900	11,988
July.....	777,600	10,854	663,000	8,293	141,900	9,222
August.....	715,100	12,028	821,500	13,252	27,800	2,280
September.....	1,076,700	15,936	845,300	14,495	138,100	12,597
October.....	2,536,900	28,967	1,707,700	16,929	21,650	2,887
November.....	1,965,900	26,343	1,402,950	14,511	40,400	4,577
December.....	832,300	23,445	1,004,400	20,633	14,700	2,772
Total.....	11,419,400	182,053	7,968,850	118,751	1,204,950	92,178
LANDED AT GLOUCESTER.												
January.....	10,585	89	4,747	\$72	22,820	196	43,860	\$659	158,316	15,798	4,685	\$328
February.....	19,448	223	625	8	11,820	168	8,796	106	246,827	24,545	510	36
March.....	1,190	9	820	10	12,546	117	3,882	39	306,329	24,724	5,115	345
April.....	49,836	300	6,910	69	45,028	317	3,592	36	395,156	26,161	9,759	652
May.....	299,207	1,796	665	7	1,125,580	7,426	69,570	696	412,121	20,304
June.....	359,649	2,158	5,700	58	933,193	5,598	116,755	1,169	261,205	14,462	21,301	1,067
July.....	314,908	1,892	10,305	113	344,258	2,069	484,614	6,342	248,464	16,652	34,436	1,921
August.....	60,432	1,392	10,282	103	58,504	351	147,911	1,478	135,807	11,098	22,159	1,226
September.....	73,425	511	38,594	386	102,070	844	154,039	1,537	72,799	6,735	575,421	45,887
October.....	272,655	2,151	7,026	71	355,047	2,061	155,166	1,552	62,443	7,366	175,715	14,066
November.....	218,348	1,697	24,605	246	1,459,862	8,088	96,955	971	59,549	6,932	9,394
December.....	15,007	277	2,985	30	63,013	1,077	93,945	956	24,664	3,051	1,618	130
Total.....	1,093,841	11,765	113,324	1,173	4,533,741	28,312	1,380,645	15,541	2,383,685	177,828	800,113	66,471
Grand total.....	13,163,241	193,818	113,324	1,173	12,502,591	147,063	1,380,645	15,541	3,588,635	270,006	800,113	66,471
Grounds E. of 66° W. long.....												
Grounds W. of 66° W. long.....	456,978	7,415	88,148	921	147,182	1,951	1,006,776	11,736	3,169,944	231,476	855,653	66,163
Landed at Boston in 1908.....	12,706,263	186,403	25,176	252	12,355,469	145,112	373,869	3,805	418,691	38,530	4,460	308
Landed at Gloucester in 1908.....	214,780	214,780	6,286,800	87,508	303,450	26,677
Landed at Gloucester in 1908.....	7,968,350	64,522	122,442	1,833	6,141,926	47,600	1,090,205	16,364	2,875,802	205,957	946,558	66,263

QUANTITIES AND VALUES OF CERTAIN FISHERY PRODUCTS LANDED AT BOSTON AND GLOUCESTER, MASS., BY AMERICAN FISHING VESSELS DURING 1909, BY MONTHS—Continued.

Month.	Mackerel.				Other fish. ^a				Total.				Grand total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
LANDED AT BOSTON.														
January.....
February.....
March.....
April.....
May.....
June.....
July.....
August.....
September.....
October.....
November.....
December.....
Total.....	3,521,300	196,826	490,900	25,400	1,631,000	157,917
LANDED AT GLOUCESTER.														
January.....
February.....
March.....
April.....
May.....
June.....
July.....
August.....
September.....
October.....
November.....
December.....
Total.....	599,760	27,111	2,967,000	186,751	5,485,106	122,303	9,304,664	165,428	32,546,318	655,533	47,980,256	1,447,943	80,526,374	2,103,476
Grand total.....	4,121,060	223,937	3,457,900	212,151	7,116,706	280,220	9,304,664	165,428	124,631,068	3,143,101	48,471,156	1,473,343	173,102,224	4,616,444
Grounds E. of 66° W.
Grounds W. of 66° W.
Grounds at Boston in 1908.
Landed at Gloucester in 1908.....	1,085,510	75,469	3,200,600	158,416	7,464,804	131,078	7,948,796	124,002	49,883,233	921,476	35,922,334	1,142,939	85,805,567	2,064,415

^a Includes herring from Newfoundland (4,296,250 pounds; frozen, \$113,635, and 9,029,756 pounds; salted, \$160,529).

More than 60 per cent of the quantity and nearly the same proportion of the value of the fishery products landed at Boston and Gloucester by the American fishing fleet during the year were caught on fishing grounds lying off the coast of the United States. A little over 28 per cent of the catch was from banks off the coast of the Canadian Provinces and 11.25 per cent from grounds off the coast of Newfoundland. The Newfoundland herring fishery furnished less than 8 per cent of the fishery products landed at these ports. The quantity and value of the catch from each of these fishing regions are given by species in the following table:

QUANTITY AND VALUE OF FISH LANDED BY AMERICAN FISHING VESSELS AT BOSTON AND GLOUCESTER, MASS., IN 1909, FROM GROUNDS OFF THE COASTS OF THE UNITED STATES, NEWFOUNDLAND, AND CANADIAN PROVINCES.

Species.	United States.		Newfoundland.		Canadian Provinces.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Cod:								
Fresh.....	28,031,010	\$765,402	88,810	\$1,492	10,470,311	\$188,253	38,590,131	\$955,147
Salted.....	4,158,127	137,120	3,828,665	113,087	24,757,580	753,446	32,744,372	1,003,653
Cusk:								
Fresh.....	2,608,026	41,022	7,660	123	531,652	8,746	3,147,938	49,891
Salted.....	105,627	2,637	7,690	191	72,218	1,809	185,535	4,637
Haddock:								
Fresh.....	37,345,145	907,965			5,055,621	115,054	42,400,766	1,023,019
Salted.....	186,428	1,885	11,235	113	226,940	2,291	424,603	4,289
Hake:								
Fresh.....	12,668,503	186,176	11,278	70	483,460	7,572	13,163,241	193,818
Salted.....	25,176	252	10,947	132	77,201	789	113,324	1,173
Pollock:								
Fresh.....	12,355,229	145,111	100	1	147,202	1,951	12,502,591	147,063
Salted.....	373,869	3,805	36,620	367	970,156	11,369	1,380,645	15,541
Halibut:								
Fresh.....	418,691	38,530	1,349,221	94,603	1,820,723	136,873	3,588,635	270,006
Salted.....	4,460	308	803,489	63,004	52,164	3,159	860,113	66,471
Mackerel:								
Fresh.....	2,461,000	132,707			1,660,060	91,230	4,121,060	223,937
Salted.....	794,400	55,250			2,663,500	156,901	3,457,900	212,151
Herring:								
Fresh.....	99,600	1,651	4,296,250	113,535	25,000	500	4,420,850	115,686
Salted.....	85,800	1,481	9,029,756	160,529	162,108	2,844	9,277,664	164,854
Swordfish:								
Fresh.....	1,626,520	157,185	394	47	10,242	1,297	1,637,156	158,529
Other fish:								
Fresh.....	1,058,700	6,005					1,058,700	6,005
Salted.....	27,000	574					27,000	574
Total.....	104,433,911	2,585,066	19,482,115	547,294	49,186,198	1,484,084	173,102,224	4,616,444

SHAD AND ALEWIFE FISHERIES.

The canvass relating to the methods, apparatus, extent, and condition of the shad and alewife fisheries of Chesapeake Bay and tributaries, which was begun in the spring of 1909, was completed before the close of that year, the work being done by the steamer *Fish Hawk* and field agents. The fishing apparatus used in the capture of shad and alewives was located on charts, and statistics of the catch for the season of 1909 were obtained. The fishing apparatus included 3,332 pound nets, 12,768 gill nets, and a considerable number of seines, fyke nets, and other appliances. The catch consisted of 2,924,018 shad, having a value to the fishermen of \$785,739, and 128,618,249 alewives, with a value of \$284,039. The shad were sold

fresh, and the alewives were disposed of in both a fresh and salted condition, the number salted being 16,827,000, valued at \$74,419. The shad catch has declined nearly 50 per cent in quantity since 1897, the number of shad taken that year in the Chesapeake and tributaries being 5,341,751. In 1901 the number had decreased to 3,000,544, and in 1904 to 2,950,492. A still further decrease of 26,474 occurred in the past year. This large falling off during these years is obviously due to overfishing and to the fact that the large number of pound nets and other apparatus operated prevent the anadromous species from reaching their spawning grounds, thus seriously interfering with both natural and artificial propagation. In Virginia in 1909 there were fished in these waters for shad and alewives 2,043 pound nets and 7,121 gill nets, and in Maryland 1,289 pound nets and 5,620 gill nets, the remainder of the gill nets being in Pennsylvania and Delaware. The catch apportioned by States in 1909 was as follows:

State.	Shad.		Alewives.	
	Number.	Value.	Number.	Value.
Virginia.....	1,855,446	\$488,336	69,469,949	\$128,375
Maryland.....	1,000,827	272,869	59,093,300	155,499
Pennsylvania.....	60,045	22,224	25,000	75
Delaware.....	7,700	2,310	30,000	90
Total.....	2,924,018	785,739	128,618,249	284,039

INVESTIGATION OF THE MACKEREL FISHERY.

The mackerel investigation, which was begun in April, 1909, at the request of the Board of Trade and Master Mariners' Association of Gloucester, Mass., representing many of the firms and vessel owners interested in the mackerel fishery, was concluded in October of that year, occupying a period of about six months. The schooner *Grampus* was detailed for the work, and Capt. Jerry E. Cook, an experienced mackerel fisherman of Gloucester, was in charge of the inquiry. The vessel was equipped with gill nets and lines for locating the fish and with tow nets for use in detecting the presence of the minute crustaceans which form the principal food of the mackerel. The object of the inquiry was chiefly to determine the movements of the mackerel, which usually make their first appearance on the American coast in the spring off Cape Hatteras and gradually move northward to the Gulf of St. Lawrence, to locate any bodies of mackerel that may frequent grounds remote from those cruised over by the fishermen, and also to assist the mackerel fishermen by furnishing them with information as to the schools of mackerel seen and their location and movements.

It is thought by some fishermen that the introduction of purse seines and gill nets in the fishery, replacing hooks and lines and a

plentiful supply of toll bait, has had a tendency to disperse the schools of mackerel and is partly responsible for the prevailing scarcity of that species during the past twenty or more years. This opinion, however, has not become sufficiently strong or general to lead to any concerted action on the part of the vessel owners with a view to abandoning the use of these forms of apparatus in the mackerel fishery and returning to the former methods.

The *Grampus* sailed from Gloucester April 7 and proceeded southward to Lewes, Del., where she joined the seining fleet. On May 2 the vessel sailed from that port to begin the work of investigating the movements of the mackerel. The first experiments were made on that date in latitude 38° N. and longitude $74^{\circ} 21'$ W. The work was continued along the coast from this locality to Georges Bank until the 1st of August, but chiefly on the southern grounds in order to ascertain whether the mackerel remain there after making their first appearance early in the spring or move northward. The fish were not located there, however, after the early run in the spring, nor were any of the usual signs of them, such as sea geese, red feed, whales, etc., observed. The vessel worked over Georges Bank and continued eastward over Browns Bank, and on August 5 anchored at Sandy Point, Shelbourne, Nova Scotia. She sailed from there on the 8th of August, and from Halifax on the 12th, reaching North Sydney, Cape Breton, on the 15th. For the remainder of August and during September the work was pursued in the Gulf of St. Lawrence and on the southerly part of the coast of Newfoundland. The *Grampus* left the Gulf of St. Lawrence early in October and arrived at Gloucester on the 16th of that month. At all times during the cruise a masthead lookout for mackerel was kept day and night when the weather was favorable for observation, and net trials for locating the fish were made at every opportunity.

During the first part of the trip the work was frequently interrupted by stormy weather, which also at times greatly interfered with the operations of the seining fleet. The mackerel were late in showing, and were unusually far offshore. Investigation showed that the latter condition was caused by the appearance of great schools of bonito, which came up the coast over the usual mackerel route and kept the schools of mackerel well offshore, and later, when the mackerel approached their regular course, caused them not to show, but to move along under water. This was indicated by the many large hauls made by the vessels of the mackerel netting fleet, which did unusually well, while the purse-seine fishermen, depending on the mackerel to school and show, had a poor season as a whole. There was a good catch on the Nova Scotia coast, or Cape Shore, and the vessels did well for a time on Nantucket Shoals, but otherwise the season was practically a failure. A few hauls were made on the southern edge

of Georges Bank, but the fish stayed there only a short time, disappearing as suddenly as they came. In Massachusetts Bay and on the Maine coast there was practically no mackerel fishing, a few small schools taken off Monhegan being all that showed on the latter shore. The season in the Gulf of St. Lawrence was also a poor one, the catch of the 22 American purse-seining vessels that went there being only 1,785 barrels. No fish were found schooling, and the catches made were secured by throwing toll bait and using line and jig, the seine being run around the vessel while the fish were attracted by the bait. Bad weather set in early and fishing off North Sydney was discontinued before the usual time.

The cruise did not result in ascertaining where the southern body of mackerel goes after coming as far north as Long Island nor in locating the great body of mackerel which goes into the Gulf of St. Lawrence, but that large schools entered the Gulf of St. Lawrence in June and, some of them at least, came out in the fall and went south is indicated by the fact that large catches were made off Halifax and La Have and westward as far as Cape Sable, and that great schools were reported off Halifax and other Cape Shore ports late in the season. It is evident from the experiments and observations made during the cruise that the food supply and spawning habits of the mackerel are not the only factors to be considered in the study of their migrations, but that the weather conditions and the presence of bonito and other predatory species have a decided influence on their movements. A plentiful supply of food was frequently found in localities where there were either no mackerel or only scattering individuals.

The *Grampus* kept in as close touch with the seining fleet as was consistent with the work, and furnished the vessels with all information obtained regarding the schools of mackerel seen.

Considerable information was also obtained during the cruise regarding the movements of menhaden. These fish were reported by the mackerel seiners early in April about 25 miles off Bodie Island, North Carolina, in 45 fathoms of water. They were in large schools and appeared to be working northward and keeping well offshore. About the 20th of the month large schools were observed in 36° 30' north latitude and extending a distance of about 90 miles. A number of menhaden were taken in the mackerel seines about this time, and were large and moderately fat fish. During April large schools of bonito were seen some distance inshore of the menhaden, which was apparently the reason why the latter remained so far offshore. One vessel reported sailing 25 miles with bonito constantly in sight, moving rapidly and at times breaking water, probably in pursuit of some small feed other than menhaden. No schools of bonito were seen north of the Virginia capes, and the menhaden appeared on the coast of New Jersey early in May after the bonito disappeared.

FISHERIES OF MISSISSIPPI.

At the request of citizens of Biloxi, Miss., through their Representative in Congress, an investigation was made of the condition of the fisheries at that place, and incidentally of the coast fisheries of the entire State, not including oysters.

The fisheries of Mississippi are chiefly carried on at Biloxi and Scranton, the former place having about 250 vessels and boats and the latter about 50 engaged in this industry. The principal species taken are shrimp (which ranks first in importance), bluefish, Spanish mackerel, pompano, mullet, flounders, trout or squeteague, and crabs. In former years the supply was generally equal to the demand, but in the last three or four years it is claimed by fishermen and others interested in the fisheries that there has been considerable falling off in the annual catch of most species, on account of overfishing.

While the coast fisheries of Mississippi are not as extensive as those of some other States, they are of great value to the State and should receive as careful attention in the way of protection as other States give to their fisheries. Many fishermen and dealers at Biloxi are of the opinion that artificial propagation is the only means by which the more important commercial species can be saved from extermination, but a judicious enforcement of laws that should be enacted to prevent the wholesale capture of fish during the spawning season, and making it a penal offense to capture fish by the use of dynamite, lime, or other explosives in rivers, lakes, bayous, or along the coast, would have a tendency to restore the fisheries to their former prosperous condition. This course of action would no doubt produce beneficial results in a comparatively short time. The first requisite in the present circumstances seems to be to take the necessary steps to save certain species by natural rather than by artificial means. The establishment of a state fish commission, with authority to recommend and enforce fishery legislation, would also be of great assistance in protecting and maintaining the fisheries. Without such an organization the fisheries are destined to decline more rapidly in the future than they have in the past.

The principal forms of fishing apparatus used in the fisheries of the State are drag seines, gill nets, and trammel nets. There is also a considerable quantity of fish taken with cast nets. In no part of the country is this apparatus used with greater skill than in this region, and according to some of the fishermen its extensive use is responsible for the great scarcity of crappie, black bass, and pike. When the water in the rivers and bays is low, many species of fish take refuge in pools and deep holes and are easily captured.

In Red and Black creeks it is said that fish were once abundant, but in recent years the use of dynamite has nearly exterminated them. According to reports, this method is not only employed by commer-

cial fishermen but also by people who wish to supply their own table. Jugs filled with lime and lowered to the bottom are also the means of destroying large quantities of fish. The water coming in contact with the lime causes the jugs to burst, scattering the lime, which either kills the fish or causes them to rise to the surface in a dazed condition, making their capture easy.

At Biloxi the harbor for vessels is at Back Bay. Six or seven years ago considerable fishing was carried on from 2 to 3 miles above the anchorage, toward the head of the bay. At the present time very few fish are taken in this vicinity. At times during a freshet buffalo-fish are caught in considerable quantities near the mouth of the bay. At other times this species is generally observed up the bay some 6 or 8 miles above Popp's Point, where commercial fishing is prohibited. During a heavy freshet it is said that the current runs 15 miles an hour.

In the upper part of the bay there are numerous small islands covered with tule grass; these islands afford excellent seining grounds. In the channels formed by the islands fishing is carried on with trammel nets. There are many snags in the channels, which prevent the use of drag seines.

Shrimp were quite scarce in 1909, but in the spring of 1910 they were plentiful, and the usual pack was made. It is estimated that in the vicinity of Gulfport and Biloxi 6,000 barrels of shrimp were caught during the season. It is stated that only about one-half the quantity of shrimp is now taken as compared to the catch ten years ago, although nearly double the number of men and boats is employed. In the last few years, however, there has been considerable increase in the catch, owing to an extension of the fishing grounds. Vessels now fish for shrimp 30 miles east and west of Biloxi and from 75 to 100 miles south.

Shrimp arrive from the south in the latter part of February and remain on the coast until May. In the latter part of July or the first of August a school of mixed sizes of shrimp appear, and in September another school of marketable shrimp strikes the coast.

In 1909, 14,000 pounds of mullet were taken in one haul of a seine and all were said to be spawn fish. Many fishermen are of the opinion that such wholesale slaughter of spawn fish should be stopped.

Redfish or channel bass, trout or squeteague, and sheepshead have not decreased as rapidly as some other species, being taken in deep water, and principally with hook and line.

The shipping facilities and method of handling fish at Biloxi compare favorably with those in other parts of the country. On account of the scarcity of many shore species, attention is being directed to the red-snapper fishery. To engage in this fishery would require deeper draft vessels and the building of plants for handling fish on the outlying islands, where vessels arriving from the banks could

land their fares and transship them in scows or other shallow boats to Biloxi. In this manner the red-snapper fishery might be established and successfully prosecuted.

MISCELLANEOUS ACTIVITIES.

RELATIONS WITH OTHER GOVERNMENT BUREAUS.

During the year the Bureau has cooperated with other branches of the Government, both giving and receiving assistance in the interests of an economical and efficient administration of the public business. The assistance rendered to the Bureau of the Census in the statistical canvass of the fisheries in the preceding fiscal year was supplemented by the detail of an agent of the Bureau to aid in certain technical matters connected with the compilation of the data. This assistance consisted principally in the identification and consideration of the involved and often dubious nomenclature of the fishes exhibited in the field schedules.

A large number of samples of fishery products have been identified and passed on at the request of the food and drug board of the Department of Agriculture, and other assistance has been rendered in connection with the functions of that board.

In March, 1910, on request of the Secretary of War preferred through the Department, an examination and appraisal was made of certain oyster bottoms adjoining the military reservation at Fort Monroe, Va., recently ceded by the State of Virginia to the Federal Government. A full report, accompanied by tracings, was transmitted to the War Department.

The Bureau expresses its appreciation of the services of the Bureau of Chemistry of the Department of Agriculture for analyses of water from various hatcheries and to the Coast and Geodetic Survey for various charts and projections and for other courtesies extended.

INTERNATIONAL FISHERY MATTERS.

In 1909, as in the four previous years, at the request of the Department of State, the Bureau detailed a representative to visit Newfoundland for the purpose of observing the operations of American fishing vessels engaged in the herring fisheries there under the provisions of the *modus vivendi*. The detail extended from October, 1909, to January, 1910. No vessel was assigned to the work this year. In June, 1910, two representatives from the Bureau's official staff were detailed to The Hague to assist the American counsel in the case before that tribunal for a settlement of the dispute as to the rights of our fishermen in Newfoundland and Canadian waters under the treaty of 1818.

The Bureau continued its cooperation with the State Department, through the International Fisheries Commission, in securing basic

data for the regulations required by the treaty between the United States and Great Britain, signed April 11, 1908, which provides for the joint control by the United States and Canada of the fisheries in the waters contiguous to the boundary between the two countries. Field work was conducted in Passamaquoddy Bay and eastern Maine and on Lake Erie and Lake Huron.

There is every reason to believe that both of these international questions, which have long been a source of irritation to the fishermen of the countries involved, will be satisfactorily adjusted during the present year.

EMPLOYMENT OF VESSELS.

The investigation concerning the aquatic resources of the Philippine Islands was continued by the steamer *Albatross* until February 12, when she went to Nagasaki for a general overhauling before undertaking the voyage to the United States. She arrived at San Francisco in excellent condition May 4, and was promptly made ready for immediate work in Alaskan waters. While the vessel underwent considerable repairs in Hongkong the year before, these were necessitated by work previous to the Philippine expedition and the fact that she returned to San Francisco in such good condition after a cruise beginning in 1907 reflects credit on the construction of the vessel and the care given by her commanding officers.

The steamer *Fish Hawk* was occupied from the beginning of the fiscal year until the middle of September in a comprehensive survey of the public oyster grounds of Virginia in the James River, and afterwards in collecting aquarium specimens. In October the ship went to Woods Hole, where her machinery was put in good order by the station force and the crew and the vessel made ready for further work. In the spring, shad hatching on the Delaware River was begun and continued until June, when a survey of the public oyster grounds of Delaware was commenced and at the close of the year was still in progress. Fuller references to the surveys mentioned are embodied elsewhere in this report.

The schooner *Grampus* was engaged in the mackerel investigation referred to elsewhere until October 10, 1909, her sphere of operations extending from Newport to Bay of Islands, Newfoundland, and the Gulf of St. Lawrence, and including the offshore fishing banks. During the late fall and winter the vessel was laid up and the crew utilized in connection with marine fish-cultural work on the New England coast until April, when she was made ready for sea and began the collection of lobster eggs and distribution of lobster fry for the hatchery at Boothbay Harbor, Me., and was so engaged the remainder of the year.

The smaller steamer *Phalarope* was used during the entire year in fish-cultural work on the New England coast and on the Potomac

River, and as a collecting vessel for the Woods Hole laboratory. The *Curlew* was employed on the Mississippi River, especially in collecting fishes from the overflowed lands.

PUBLICATIONS AND LIBRARY.

The collection of special books maintained by the Bureau for purposes of reference and technical investigation has received 260 accessions in Washington from gifts, purchases, and exchanges, and over 200 accessions at the laboratories and stations elsewhere. The intimate relations maintained with other libraries result in exchanges and transfers which are mutually profitable, and particularly advantageous to the Bureau in view of the limited funds available for the purchase of books and periodicals. The use of the library has been much facilitated by the progress made during the year on the systematic subject catalogue.

The continued interest of the public in the work of the Bureau is shown by the facts that during the year 2,916 bound volumes and 21,832 pamphlets of its publications were sent out on request, 45,890 were required for the regular mailing list, and 2,020 issued to authors. There were received from the Government Printing Office for distribution 87 new reports and bulletins published by the Bureau and 5 reprints of important documents the supply of which had been exhausted. The titles of the new issues (No. 646 to No. 732) may be found in the Bureau's list of publications available for distribution.

APPROPRIATIONS.

The total appropriations for the Bureau for the fiscal year amounted to \$823,490, or \$16,610 less than the aggregate for the previous year.

Salaries:

General	\$316, 860
Agents at Alaska salmon fisheries.....	4, 500
Agents at seal fisheries.....	11, 430

Miscellaneous expenses:

Administration	8, 000
Propagation of food fishes.....	275, 000
Inquiry respecting food fishes.....	30, 000
Statistical inquiry	7, 500
Maintenance of vessels.....	55, 000
Supplies for native inhabitants, seal islands.....	19, 500

Specials:

Establishment of fish-cultural stations on Puget Sound or its tributaries	50, 000
Establishment of a fish-cultural station in the upper Mississippi Valley	25, 000
Purchase of a steamboat, Put-in-Bay, Ohio.....	15, 000
Construction of roadway, Greenlake, Me.....	2, 700
Repairs to buildings, Pribilof Islands.....	3, 000

In addition to the above funds, the sum of \$150,000 was appropriated and made immediately available for the purpose of carrying out the provisions of the act of April 21, 1910, which placed under the Secretary of Commerce and Labor the administration of the fur-seal islands and the preservation of the fur-bearing animals of Alaska.

An itemized statement of expenditures authorized by the foregoing appropriations will be made as required by law.

RECOMMENDATIONS.

REORGANIZATION OF PERSONNEL.

The foregoing report exhibits briefly the rapid growth of the activities and responsibilities of the Bureau by natural accretion to lines of work long established and by the addition of functions not contemplated when the present organization was adopted. The assignment of new duties to the Bureau has made it necessary to impose them upon persons whose time and attention were already fully taxed by the natural development of their previous responsibilities, and it therefore appears to be essential to the continued efficiency of the Bureau that there should be a reorganization of the personnel. The Alaska salmon service and the fur-seal service, now assigned to the Bureau, both involving executive and police functions of an exacting character, are administered by the Division of Scientific Inquiry, from which it is desirable that they be separated. The original requirements of the division are incompatible with the added functions, and their continued administration by one person can only be at the sacrifice of the efficiency of both. It is therefore recommended that the present organization be augmented by the creation of a new division to be known as the Division of Alaska Fisheries, with sufficient additions to the present force to make its work effective.

The United States has entered into certain treaty obligations in respect to the waters adjacent to the Canadian boundary, whereby it is proposed to assume international control of the fisheries in the interest of their conservation and development. Regulations making this agreement effective were submitted to the Senate but were returned to the joint commissioners for further consideration. It is assumed that they will be reduced to a satisfactory basis in the near future, in which event the Bureau will find itself charged with enforcing them. Should this be the case, since under the present organization there is no provision for the discharge of this duty, it will be necessary to provide a Division of International Fisheries.

SALARIES AND PERSONNEL.

The recommendations of the preceding fiscal year in reference to the increase of the salaries attached to certain positions in its service are renewed. Congress at its recent session increased the pay of low-

grade clerks, firemen, and messengers, but did not authorize any advance in the salaries of those on whose work the efficiency of the Bureau is more directly dependent. The experience of another year has made more apparent the desirability of making remuneration more commensurate with duties and responsibilities.

The Bureau is in constant receipt of requests from Members of Congress and state authorities for special investigations and experiments in the interests of the public fisheries, and in many cases prompt compliance with these legitimate demands is difficult or impossible, because the personnel has not kept pace with either the growth of the work or the increase of general appropriations. There are certain fisheries to which, on account of their peculiar requirements, it has not been possible to render the service which those engaged in them have the right to expect. To the oyster industry, for instance, which yields \$16,000,000 annually, about 30 per cent of the value of the entire fisheries of the United States, the Bureau's assistance has been wholly inadequate. Proportionately to the value of the respective fisheries, sixty-five dollars are profitably expended in shad culture for every dollar spent for the benefit of the oyster industry. The inequality arises not from the inability to allot money from the appropriations, but to the lack of trained and experienced men. Fish-cultural methods can not be applied in oyster culture, and the only valuable aid which can be offered is through the medium of research and practical experiment, which experience has shown lead to profitable and lasting benefits from disproportionately small expenditures. For carrying on such work provision should be made for additional scientific assistants.

SPONGE LAW.

The act of June 20, 1906, to provide for the protection of the sponge fisheries of the United States on the high seas of the Gulf of Mexico and the Straits of Florida, has shown itself futile and impossible of enforcement. The purpose of this law was to prohibit the fishery by diving in depths of less than 50 feet, and during the period from May 1 to October 1 to prevent the taking, by whatever means, outside of the 3-mile limit, of sponges smaller than 4 inches in diameter.

The offenses aimed at are not specifically prohibited, but they were supposed to be prevented by the prohibition of certain subsidiary acts—the landing, curing, or offering for sale in the United States of sponges taken in contravention of the real purpose of the law. To secure a conviction it is therefore necessary to establish a connection between the act of taking under the objectionable circumstances and certain subsequent and secondary acts which per se are innocuous. A diving vessel operating during the close

season can not be interfered with until the sponges are landed, cured, or offered for sale in the United States. The sponges, therefore, must be followed or traced from their beds in the high seas to a point of territorial jurisdiction, a requirement that is usually impossible of enforcement.

Moreover, the law provided the Department with no machinery for its enforcement. It has been necessary to depend upon the courtesy of the Treasury Department for the personnel required, and no provision has been made for expenses.

In view of the circumstances narrated, and in the interest of the unimpaired maintenance of the sponge beds, it is recommended that the act of June 20, 1906, be amended to correct its defects and that the Bureau be provided with an inspector, a suitable boat, and funds for the proper enforcement of the law. It is further recommended that the minimum size of sponges which it shall be legitimate to take be established at 5 inches diameter, and if this be done that the close season be curtailed by not exceeding two months.

EXTENSION OF FISH CULTURE.

It is again urged that provision be made for the establishment of additional stations for the rescue of fishes from overflowed lands in the Mississippi Valley. Millions of fish now annually left by the receding waters to die of exposure can by this means be saved at small expense.

The Bureau is of the opinion that a highly important work of the near future will be the stocking of ponds and streams on the farms of the country with hardy species of fish requiring little care or attention and omnivorous as to diet. The several species of catfishes appear to fulfill the requirements more completely than any other fish. They will grow in sluggish and muddy water, they are very tenacious of life, their diet is of wide variety, and as food they are excelled by but few fresh-water fish. While some of the smaller species can be made important additions to the home food supplies of the farms, certain others, particularly the larger ones, are already the basis of important commercial fisheries. For the propagation of both kinds the establishment of a station at some point in the lower Mississippi Valley, preferably near Morgan City, La., is regarded as highly desirable.

The fish-cultural work in Yellowstone Park has been conducted heretofore with inadequate means as an adjunct to the operation of Spearfish Hatchery, but it is believed that the opportunities in the national park are such as to warrant an independent station. One of the chief difficulties encountered in the efforts to replenish the depleted fisheries of the United States arises from the lack of control

over the fishes after they are planted and the neglect of certain States to make provision for their protection. Yellowstone Park, being under federal jurisdiction, offers an exceptional opportunity to demonstrate the possibilities of fish culture under rational and consistent regulations.

The Bureau also recommends the establishment of one marine and one additional fresh-water hatchery on the Pacific coast, and an additional station in Texas for the supply of a demand for fish in the Southwest which it is at present impossible to satisfy.

LABORATORY FOR THE STUDY OF FISH DISEASES.

There is again urged the importance of a station for the study of fish diseases and experiments in the interests of fish culture. In some of the hatcheries of the Bureau and in similar establishments under state and private auspices certain fish diseases have become so prevalent as to make it a matter of grave consideration whether the propagation of certain species, especially the trouts, should not be abandoned. It frequently occurs that the fish and fry are decimated by epidemics for which there are no known remedies, in consequence of which there are annually entailed on fish culture large wastes of time and money. In addition to the financial loss, embarrassment arises at times in filling legitimate demands for fish for restocking depleted waters, and the effect on the morale of the employees of the Bureau who have to struggle hopelessly against an obscure disease is not unworthy of consideration. The gravest phase of the matter, however, is the possible relationship of some of these diseases to more or less kindred affections occurring in human beings. It has been determined that a type of cancerous affection is of widespread distribution among domesticated trout and their offspring planted in the streams. Whether this disease has a causal relation to cancer in human beings, or whether the two are to be even traced to the same source, is a matter of doubt, but the annually increasing mortality from cancer in man and certain remarkable coincidences in the geographical distribution of the disease in man and fish render it imperative that it should be made the subject of minute inquiry. The matter therefore has not only economic but humanitarian aspects, and the consideration of the serious character of the latter prompted the President to submit to Congress on April 9, 1910, a special message advocating an appropriation of \$50,000 for the construction and equipment of a laboratory adequate to enable the Bureau to discharge its plain obligations. The Bureau in the meantime is proceeding in the investigation to the limit of its powers, but it may be stated emphatically that it can make but little progress without the special facilities asked for.

FISHERY INTELLIGENCE SERVICE.

For many years the Bureau has maintained at Boston and Gloucester, Mass., a service making current statistical reports on the fisheries of those ports. This service has the strong support of the commercial interests, and a proposition for its abandonment would result in instant and vigorous protest. The large fishery interests of the Pacific coast are becoming insistent in their requests that a similar service be inaugurated in that region, and the Bureau regards the work of such importance as to impel it to recommend provision for a suitable personnel for the purpose. In view of the regard in which the reports at Boston and Gloucester are held by the fishery interests, it would appear desirable to gradually extend the service to other places on the Atlantic and Gulf coasts having extensive vessel fisheries.

NEW BUILDING.

As has been repeatedly indicated in these reports, the quarters of the Bureau are antiquated, crowded, unsafe, and inadequate in every respect. They impede the transaction of the public business and interfere with efficiency and development. It is again earnestly recommended that provision be made for a building which will furnish offices, laboratories, workrooms, and an aquarium national in scope and in keeping with necessitous requirements.

Respectfully,

GEO. M. BOWERS,
Commissioner.

To Hon. CHARLES NAGEL,
Secretary of Commerce and Labor.

THE DISTRIBUTION OF FISH AND FISH EGGS DURING
THE FISCAL YEAR 1910

Bureau of Fisheries Document No. 740



CONTENTS.

	Page.
Character of the work.....	5
Method of distribution.....	5
Size of fish when distributed.....	6
Size of allotments.....	7
Species cultivated in 1910.....	7
Output.....	9
Statement of output by stations.....	10
Allotments to State fish commissions.....	24
Shipments to foreign countries.....	24
Summarized statement of distributions.....	25
Details of the distributions.....	26

INDEX TO SPECIES.

	Page.		Page.
Atlantic salmon.....	39	Perch, yellow.....	107
Bass, large-mouth black.....	88	Pickereel.....	80
rock.....	83	Pike.....	80
sea.....	110	Pike perch.....	105
small-mouth black.....	86	Pollock.....	111
strawberry.....	81	Rainbow trout.....	31
striped.....	109	Rock bass.....	83
warmouth.....	85	Salmon, Atlantic.....	39
white.....	109	blueback.....	30
yellow.....	110	chinook.....	30
Blackspotted trout.....	40	humpback.....	30
Blueback salmon.....	30	landlocked.....	39
Bream.....	101	silver.....	29
Brook trout.....	44	Sea bass.....	110
Buffalofish.....	28	Shad.....	28
Carp.....	27	Silver salmon.....	29
Catfish.....	26	Small-mouth black bass.....	86
Chinook salmon.....	30	Smelt.....	80
Cisco.....	29	Steelhead trout.....	31
Cod.....	111	Strawberry bass.....	81
Crappie.....	81	Striped bass.....	109
Drum, fresh-water.....	111	Sunapee trout.....	80
Flatfish.....	112	Sunfish.....	101
Fresh-water drum.....	111	Trout, brook.....	44
Grayling.....	80	blackspotted.....	40
Haddock.....	111	lake.....	43
Humpback salmon.....	30	Loch Leven.....	43
Lake herring.....	29	rainbow.....	31
Lake trout.....	43	steelhead.....	31
Landlocked salmon.....	39	Sunapee.....	80
Large-mouth black bass.....	88	Warmouth bass.....	85
Lobsters.....	112	White bass.....	109
Loch Leven trout.....	43	White perch.....	110
Mackerel.....	110	Whitefish.....	29
Perch, pike.....	105	Yellow bass.....	110
white.....	110	Yellow perch.....	107



THE DISTRIBUTION OF FISH AND FISH EGGS DURING THE FISCAL YEAR 1910.

CHARACTER OF THE WORK.

More than 95 per cent of the output of the fish-cultural stations consists of important commercial species, notably the salmons, shad, whitefish, pike perch, yellow perch, white perch, lake trout, cod, pollock, flatfish, and lobsters. These are hatched in lots of many millions annually and planted by the Bureau, the fresh-water species principally in the large coastal streams and in the Great Lakes, the marine species upon the inshore fishing grounds of the Atlantic.

The cultivation of the fishes of the interior waters generally classed as game fishes, although a comparatively small factor in the total output, is a very important feature of the Bureau's work, supplying as it does various kinds of young fish for public streams, lakes and ponds, fishing preserves, private ponds, streams, etc., in all parts of the United States. Among the fishes most extensively cultivated for these purposes are the landlocked salmon, several species of trout, the grayling, the basses, crappie, bream, and catfish; various others also are handled. The trouts are artificially hatched from eggs taken from both wild and domesticated stock; the basses, catfish, and others are derived from mature fish held in ponds for breeding purposes, or (except the small-mouth black bass) they are rescued from the overflows of the Mississippi and Illinois Rivers. Collections from the latter sources include also pike and pickerel, which are not distributed to applicants but are returned immediately to the main streams.

METHOD OF DISTRIBUTION.

The first consideration in the Bureau's distribution of fishes is to make ample return to the waters from which eggs or fish have been collected. The remainder of the product is consigned to suitable public or private waters upon application indorsed by a United States Senator or Representative, the Bureau furnishing to persons interested an application blank for this purpose. The blank calls for a description of the waters to be stocked, and by this information is determined the species of fish that is suitable and the number that may be allotted to the water area in question. Certain predaceous species, such as the basses, perches, and pickerel, are not furnished

for waters inhabited by trout or other valuable fishes to which they would be destructive. Nor, of course, are species like trout and salmon furnished for waters already stocked with fish that would prey upon them.

The fish are carried to their destination in railroad cars equipped for the purpose, or by messengers who accompany the shipments in baggage cars, and are delivered to the applicant free of charge, at the railroad station nearest the point of deposit. The applicant is advised by telegraph when the shipment will arrive, and is expected to make due provision for care of the fish until planted. Definite instructions in this respect are furnished at the time of shipment.

During the past fiscal year (July 1, 1909, to June 30, 1910) the Bureau received 10,635 applications for fish, nearly all for the game species. The demand, especially for the basses, crappie, and the catfishes, has for some time been greater than could be met with available resources. The number of applications this year was 523 more than in 1909.

SIZE OF FISH WHEN DISTRIBUTED.

Fishes are distributed at various stages of development, according to the species, the numbers in the hatcheries, and the facilities for rearing. The commercial fishes—such as the shad, whitefish, lake trout, pike perch, cod, etc., hatched in lots of many millions—are necessarily planted as fry shortly after hatching. Atlantic salmon, landlocked salmon, and various species of trout are reared, in such numbers as the hatchery facilities permit, to fingerlings from 1 to 6 inches in length; the remainder are distributed as fry.^a

The basses, bream, and other sunfishes are distributed from some three weeks after they are hatched until they are several months of age. When the last lots are shipped the basses usually range from 4 to 6 inches and the sunfishes from 2 to 4 inches in length. The numerous fishes collected in overflowed lands—basses, crappie, sunfishes, catfishes, yellow perch, and others—are 2 to 6 inches in length when taken and distributed.

Eggs are distributed only to state hatcheries and, occasionally, to applicants who have hatchery facilities.

^a The varying usage in the classification of young fish as to size has caused such confusion and difficulty that the Bureau has adopted uniform definitions, as follows:

Fry—fish up to the time the yolk sac is absorbed and feeding begins.

Advanced fry—fish from the end of the fry period until they have reached a length of 1 inch.

Fingerlings—fish between the length of 1 inch and the yearling stage, the various sizes to be designated as follows: No. 1, a fish 1 inch in length and up to 2 inches; no. 2, a fish 2 inches in length and up to 3 inches; No. 3, a fish 3 inches in length and up to 4 inches, etc.

Yearlings—fish that are 1 year old, but less than 2 years old from the date of hatching; these may be designated No. 1, No. 2, No. 3, etc., after the plan prescribed for fingerlings.

SIZE OF ALLOTMENTS.

The Bureau does not attempt to furnish to any one applicant more than a brood stock of fish for a given private pond or stream, it being expected that these will be protected until they have had time to reproduce. The number of fish in an allotment is, however, a variable quantity, depending upon the species and the age at which distributed. Brook trout, which are distributed both as fry and fingerlings, are allotted in much larger numbers as fry than as fingerlings 3 or 4 inches long. Pike perch, which, owing to their excessive cannibalism, can not be reared and are consequently distributed as fry, may be supplied in lots of half a million, where an equal water area would receive only 200 or 300 young bass from 2 to 5 inches long. These latter larger fish have a much better chance of reaching maturity than have the fry, and the actual value for stocking purposes of a few hundred fingerling bass may therefore equal many thousand times this number of pike perch fry.

SPECIES CULTIVATED IN 1910.

The species cultivated by the Bureau in 1910 numbered some 50 fishes and the lobster. Of these the following were artificially propagated:

THE CATFISHES (SILURIDÆ):

- Horned pout, bullhead, yellow cat (*Ameiurus nebulosus*).
- Marbled cat (*Ameiurus nebulosus marmoratus*).

THE SHADS AND HERRINGS (CLUPEIDÆ):

- Shad (*Alosa sapidissima*).

THE SALMONS, TROUTS, WHITEFISHES, ETC. (SALMONIDÆ):

- Common whitefish (*Coregonus albus* and *C. clupeaformis*).
- Lake herring, cisco (*Leucichthys artedii*).
- Chinook salmon, king salmon, quinnat salmon (*Oncorhynchus tshawytscha*).
- Silver salmon, coho (*Oncorhynchus kisutch*).
- Blueback salmon, redfish, sockeye (*Oncorhynchus nerka*).
- Humpback salmon (*Oncorhynchus gorbuscha*).
- Steelhead trout, hardhead (*Salmo gairdneri*).
- Rainbow trout (*Salmo iridicus*).
- Atlantic salmon (*Salmo salar*).
- Landlocked salmon (*Salmo sebago*).
- Blackspotted trouts: Yellowstone Lake trout or cutthroat trout (*Salmo lewisi*); Colorado River trout (*Salmo pleuriticus*); Tahoe trout (*Salmo henshawi*).
- Loch Leven trout (*Salmo trutta levensis*). Introduced species, propagated in limited numbers for observation.
- Lake trout, Mackinaw trout, longe, togue (*Cristivomer namaycush*).
- Brook trout, speckled trout (*Salvelinus fontinalis*).
- Sunapee trout (*Salvelinus aureolus*).

THE GRAYLINGS (THYMALLIDÆ):

- Montana grayling (*Thymallus montanus*).

THE SMELTS (ARGENTINIDÆ):

- American smelt (*Osmerus mordax*).

THE BASSES, SUNFISHES, AND CRAPPIES (CENTRARCHIDÆ):

- Crappie (*Pomoxis annularis*).
- Strawberry bass, calico bass (*Pomoxis sparoides*).
- Rock bass, red-eye, goggle-eye (*Ambloplites rupestris*).
- Warmouth, goggle-eye (*Chænobryttus gulosus*).
- Small-mouth black bass (*Micropterus dolomieu*).
- Large-mouth black bass (*Micropterus salmoides*).
- Bluegill bream, bluegill sunfish (*Lepomis pallidus*).
- Other sunfishes, chiefly *Eupomotis gibbosus*.

THE PERCHES (PERCIDÆ):

- Pike perch, wall-eyed pike, yellow pike, blue pike (*Stizostedion vitreum*).
- Yellow perch, ring perch (*Perca flavescens*).

THE SEA BASSES (SERRANIDÆ):

- Sea bass (*Centropristes striatus*).
- Striped bass, rockfish (*Roccus lineatus*).
- White bass (*Roccus chrysops*).
- White perch (*Morone americana*).
- Yellow bass (*Morone interrupta*).

THE MACKERELS (SCOMBRIDÆ):

- Mackerel (*Seomber scombrus*).

THE CODS (GADIDÆ):

- Cod (*Gadus callarias*).
- Haddock (*Melanogrammus æglefinus*).
- Pollock (*Pollachius virens*).

THE FLOUNDERS (PLEURONECTIDÆ):

- Winter flounder, American flatfish (*Pseudopleuronectes americanus*).

CRUSTACEANS:

- American lobster (*Homarus americanus*).

After the annual seasons of high water in the Mississippi basin, great numbers of young fish are left in sloughs and pools when the waters have receded, and would eventually die by the drying up of these shallow places in summer or freezing in winter. Large collections are made from such sources, for return to the original stream and, of the most abundant species, also to supplement the hatchery stock for distribution. The fishes so collected in 1910 were as follows:

THE CATFISHES (SILURIDÆ):

- Spotted cat, blue cat, channel cat (*Ictalurus punctatus*). Only limited numbers obtainable.
- Horned pout, bullhead, yellow cat (*Ameiurus nebulosus*).

THE SUCKERS AND BUFFALOFISHES (CATOSTOMIDÆ):

- Small-mouth buffalofish (*Ictiobus bubalus*).

THE MINNOWS AND CARPS (CYPRINIDÆ):

- Carp (*Cyprinus carpio*). Distributed in rare instances, for waters unsuited to other species.

THE PIKES AND PICKERELS (ESOCIDÆ):

- Pike (*Esox lucius*). Restored to the streams; not distributed.
- Pickereel (*Esox reticulatus*). Restored to the streams; not distributed.

THE BASSES, SUNFISHES, AND CRAPPIES (CENTRARCHIDÆ):

- Crappie (*Pomoxis annularis*).
- Rock bass, red-eye, goggle-eye (*Ambloplites rupestris*).
- Warmouth, goggle-eye (*Chænobryttus gulosus*).
- Large-mouth black bass (*Micropterus salmoides*).

THE BASSES, SUNFISHES, AND CRAPPIES (CENTRARCHIDÆ)—Continued.

Small-mouth black-bass (*Micropterus dolomieu*).

Bluegill bream, bluegill sunfish (*Lepomis pallidus*).

Other sunfishes (chiefly *Eupomotis gibbosus*).

THE PERCHES (PERCIDÆ):

Yellow perch, ring perch (*Perca flavescens*).

THE CROAKERS (SCLENIDÆ):

Fresh-water drum, sheepshead, gaspergou (*Aplodinotus grunniens*). Only limited numbers obtainable. Not distributed.

Certain introduced species are propagated to a limited extent, as follows:

THE MINNOWS AND CARPS (CYPRINIDÆ):

Goldfish (*Carassius auratus*). Propagated for ornamental purposes; not distributed.

Ide (*Leuciscus idus*). Cultivated variety, golden ide. Propagated for ornamental purposes; not distributed.

OUTPUT.

Although unfavorable climatic conditions, in 1910, prevented the collection of as large numbers of eggs as usual, the superior quality obtained from the most important species made possible a 4 per cent increase over the previous record year of 1909. As appears in the Report of the Commissioner of Fisheries for 1910, this year's output of the stations was something over 473,000,000 eggs, 2,720,000,000 fry, and 36,000,000 fingerlings, yearlings, and adults, or more than 3,230,000,000 fish and eggs in all. The yield of the various species showed the usual fluctuations, there being notable increases in the blueback, silver, and Atlantic salmons, lake trout, lake herring, yellow perch, shad, cod, flatfish, and steelhead trout, offset to some extent by decreases in chinook salmon, whitefish, pike perch, and less important fishes.

The following table shows the work of the different stations in 1910, the period of operation, and the eggs and fish delivered by each station for distribution. It will be noted that transfers of eggs and fish from station to station are frequent, serving economy and convenience in transportation where the shipment consists of eggs, and giving advantageous distributing centers in the case of young fish. Transfers are in all cases credited to the receiving station in the column of totals, but for completeness of information are recorded opposite both shipping and receiving station in the columns headed "Transfers." The purpose of this table is to be distinguished from that of the summary of distributions on page 25 of this report, which is a statement of the number of eggs and fish actually delivered at their destination, all losses in shipment being deducted.

STATIONS OPERATED AND THE

NOTE.—The relative importance of the stations is in a degree indicated in the table by marginal indentations shifting in location from year to year. At all other substations eggs were both collected and hatched. Stations to which they are, for administration purposes, subordinate; but it is not always possible to show

Station and period of operation.	Species.	Eggs.		
		Dis-tributed.	Transfers to other stations.	Transfers from other stations.
Afognak, Alaska.....	Blueback salmon.....			
Entire year.	Humpback salmon.....			
Baird, Cal.....	Chinook salmon.....	7,331,217	Nashua, 100,000.....	
Entire year.			Central station, 15,000.	
	Rainbow trout.....	13,680		
	Brook trout.....			Leadville, 25,000.....
Battle Creek, Cal....	Chinook salmon.....	7,358,800		
Oct.-Jan.				
Derby, Nev.....	Blackspotted trout.....	438,550		
Jan.-May.	Rainbow trout.....	100,000		
Mill Creek, Cal.....	Chinook salmon.....	15,849,450		
Oct.-Jan.				
Baker Lake, Wash.....	Silver salmon.....	100,000		
Entire year.	Chinook salmon.....			
	Blueback salmon.....	100,000		
	Steelhead trout.....			
Birdsview, Wash....	Humpback salmon.....			
Entire year.	Silver salmon.....	275,000	Cape Vincent, 25,000.....	Day Creek, 769,000.....
	Steelhead trout.....	300,000	Spearfish, 25,000.....	
	Chinook salmon.....			Illabott Creek, 431,740.....
Day Creek, Wash....	Steelhead trout.....		Birdsview, 769,000.....	
Feb.-June.				
Illabott Creek, Wash.	Chinook salmon.....	439,990	Birdsview, 431,740.....	
July-Oct.				
Salmon Banks, San Juan Island, Wash.	Blueback salmon.....			
July-Oct.				
Battery, Havre de Gracc, Md.	Yellow perch.....	5,200,000		
Feb. 27-May 25.	White perch.....	16,500,000		
	Shad.....	800,000		
	Striped bass.....			
Boothbay Harbor, Me..	Lobster.....	780,000		
Entire year.	Flatfish.....			
	Cod.....			
	Haddock.....			
Portland, Me.....	Lobster.....			
July 1-Jan. 1.				
York, Me.....	do.....			
July 1-Oct. 31.				
Bozeman, Mont.....	Brook trout.....			
Entire year.	Blackspotted trout.....			Spearfish, 544,000.....
	Rainbow trout.....	85,000	Clackamas, 85,000.....	
	Grayling.....	25,000		
	Landlocked salmon.....			
	Lake trout.....			
Grayling, Mont.....	Steelhead trout.....			
Mar. 1-June 30.	Rainbow trout.....			
Soda Butte, National Park, Mont.	Grayling.....			
June 16-20.	Blackspotted trout.....			
Bryans Point, Md.....	Yellow perch.....	4,030,000	Central Station, 4,030,000.....	
Feb. 21-May 23.	Shad.....	1,077,000	Central Station, 1,077,000.....	
Cape Vincent, N. Y....	Steelhead trout.....			Birdsview, 25,000.....
Entire year.	Whitefish.....			Put-in-Bay, 25,000,000.....
	Brook trout.....			
	Lake trout.....			Duluth, 5,100,000.....
	Pike perch.....			Put-in Bay, 5,000,000.....
	Landlocked salmon.....			Grand Lake Stream, 15,000.....
	Rainbow trout.....			Wytheville, 50,000.....
	Yellow perch.....			

Station and period of operation.	Species.	Eggs.			
		Dis-tributed.	Transfers to other stations.	Transfers from other stations.	
Central Station, Wash- ington, D. C. Entire year.	Sunfish.....				
	Crappie.....				
	Catfish.....				
	Smelt.....				
	Warmouth bass.....				
	Rock bass.....				
	Small-mouth black bass.....				
	Large-mouth black bass.....				
	Rainbow trout.....			Wytheville, 15,000....	
	Steelhead trout.....			Baird, 15,000.....	
	Chinook salmon.....			Bryans Point, 4,030,000	
	Yellow perch.....			Put-in Bay, 6,000,000..	
	Pike perch.....			St. Johnsbury, 20,000..	
	Brook trout.....			Put-in Bay, 640,000...	
Whitefish.....			Detroit, 500,000.....		
Clackamas, Oregon City, Oreg. Entire year.	Shad.....			Bryans Point, 1,077,000	
	Rainbow trout.....			Bozeman, 85,000.....	
	Brook trout.....			Leadville, 100,000....	
	Steelhead trout.....			Eagle Creek, 75,000....	
	Blackspotted trout			Spearfish, 100,000....	
	Lake trout.....				
	Chinook salmon.....	150,000		Rogue River, 61,600..	
	do.....				
	Big White Salmon, Wash. Aug. 1-Feb. 28.				
	Cazadero, Oreg.....	Steelhead trout.....			Eagle Creek, 410,000..
Eagle Creek, Clacka- mas River, Oreg. Mar. 15-June 25.	Chinook salmon.....	2,452,000	Cazadero, 410,000..		
	do.....	485,000	Clackamas, 75,000..		
Eagle and Tanner Creeks, Oreg. Aug. 1-Oct. 1.	Chinook salmon.....	269,300			
<i>Illinois River, Oreg</i> Aug. 1-Apr. 30.	do.....	14,200	Rogue River, 14,200.		
Little White Sal- mon, Wash. Entire year.	do.....	3,805,000			
Rogue River, Oreg. Entire year.	do.....	484,000	Clackamas, 61,600....	Illinois River, 14,200..	
Wilamette, Oreg.....	Steelhead trout.....				
Jan. 1-July 15.	Shad.....				
<i>Bybee Bridge, Rogue</i> <i>River, Oreg.</i> Aug. 1-Nov. 1.	Chinook salmon.....				
Cold Springs, Bulloch- ville, Ga. Entire year.	Large-mouth black bass.....				
Craig Brook, East Or- land, Me. Entire year. Upper Penobscot, Me. Oct. 15-June 1.	Sunfish.....				
	Catfish.....				
	Warmouth bass.....				
	Rock bass.....				
	Brook trout.....			St. Johnsbury, 5,000..	
	Atlantic salmon.....	1,345,000	Upper Penobscot, Me., 1,340,000.		
	do.....			Craig Brook, 1,340,000.	
	Duluth, Minn.....	Landlocked sal- mon.....			Grand Lake Stream, 15,000.
	Entire year.	Brook trout.....			Detroit, 25,000,000...
		Whitefish.....			Put-in Bay, 15,000,000.
	Pike perch.....				
	Steelhead trout.....				
	Lake trout.....	5,425,000	Cape Vincent, 5,100,000 Green Lake, 125,000.	Northville, 5,000,000..	
<i>Grand Marais, Mich.</i> Oct. 16-Nov. 18.	do.....				
<i>Grand Marais, Minn.</i> Sept. 19-Nov. 26.	do.....				
<i>Grand Portage, Minn.</i> Sept. 24-Oct. 15.	do.....				
<i>Keeweenaw Point,</i> <i>Mich.</i> Oct. 4-Nov. 2.	do.....				
<i>Marquette, Mich.</i> Oct. 16-Nov. 11.	do.....				

OUTPUT OF EACH, 1910—Continued.

Fry.			Fingerlings, yearlings, and adults.			Total output.
Dis-tributed.	Transfers to other stations.	Transfers from other stations.	Dis-tributed.	Transfers to other stations.	Transfers from other stations.	
			5,600			5,600
			247			247
			450			450
			9,000			9,000
			752			752
			2,010			2,010
			1,000			1,000
			440			440
7,000						7,000
12,000						12,000
			10,000		Nashua, 10,000.	10,000
3,700,000						3,700,000
5,000,000						5,000,000
18,700						18,700
774,000						774,000
977,000						977,000
51,116						51,116
64,800						64,800
126,000						126,000
82,214			1,418			83,632
12,000						12,000
3,686,200			225			3,836,425
3,512,200						3,512,200
1,808,835						1,808,835
534,197						2,986,197
49,503						49,503
						269,300
4,808,000						8,613,000
660,292						1,082,692
89,850						89,850
1,678,000						1,678,000
			107,850			107,850
			7,080			7,080
			100			100
			40			40
			100			100
196,000			76,550	Nashua, 2,200.		272,550
155,799			82,413			243,212
1,217,366						1,217,366
			11,400			11,400
			370,000			370,000
25,000,000						25,000,000
13,800,000						13,800,000
			161,000			161,000
8,825,000			4,246,500			13,271,500

STATIONS OPERATED AND THE

Station and period of operation.	Species.	Eggs.		
		Dis-tributed.	Transfers to other stations.	Transfers from other stations.
Duluth, Minn.—Cont'd.				
<i>Munising, Mich.</i>	Lake trout.....			
Oct. 16—Nov. 12.	do.....			
<i>Ontonagon, Mich.</i>	do.....			
Oct. 16—Nov. 13.	do.....			
<i>Two Harbors, Minn.</i>	do.....			
Oct. 15—Nov. 1.	do.....			
Edenton, N. C.	Shad.....	1,360,000		
Jan. 2—June 30.	do.....			
Weldon, N. C.	Striped bass.....	4,566,000		
Apr. 1—May 30.	do.....			
Erwin, Tenn.	Small-mouth black bass.....			
Entire year.	Large-mouth black bass.....			
	Brook trout.....			
	Rainbow trout.....			Wytheville, 503,000.
	Catfish.....			
	Yellow perch.....			
	Sunfish.....			
	Rock bass.....			
Gloucester, Mass.	Lobster.....			
Entire year.	Pollock.....			
	Cod.....	34,689,000	Woods Hole, 24,835,000	
	Flatfish.....			
Green Lake, Me.	Landlocked salmon.....	55,000	St. Johnsbury, 5,000.	Grand Lake Stream, 704,799.
Entire year.	Brook trout.....	25,000		
	Smelt.....	4,500,000		
	Lake trout.....			Duluth, 100,000.
	do.....			Northville, 300,000.
<i>Branch Pond, Me.</i>	Landlocked salmon.....			
Sept. 13—Nov. 30.	do.....			
<i>Grand Lake Stream, Me.</i>	Landlocked salmon.....	824,799	Duluth, 15,000.	
Entire year.	do.....			
	do.....			Spearfish, 25,000.
	do.....			Cape Vincent, 15,000.
	do.....			Green Lake, 704,799.
	do.....			Baird, 25,000.
	do.....			Clackamas, 100,000.
	do.....			Manchester, 10,000.
Leadville, Colo.	Brook trout.....	605,000		
Entire year.	Rainbow trout.....	55,000		
	Lake trout.....			
	Blackspotted trout.....	235,000		
	Rainbow trout.....			
<i>Cheesman Lake, Colo.</i>	do.....			
Apr. 6—May 8.	do.....			
<i>Darrah Lake, Colo.</i>	Brook trout.....			
Nov. 11—Nov. 30.	do.....			
<i>Edith Lake, Colo.</i>	do.....			
Oct. 18—Nov. 28.	do.....			
<i>Engelbrecht Lake, Colo.</i>	do.....			
Oct. 16—Nov. 12.	do.....			
<i>Grand Mesa Lakes, Colo.</i>	Blackspotted trout.....			
July 1—Aug. 1.	Rainbow trout.....			
Oct. 25—Nov. 11.	Brook trout.....			
<i>Musgroves Lake, Colo.</i>	do.....			
Oct. 12—Dec. 6.	do.....			
<i>Woodbridge, Colo.</i>	do.....			
Nov. 27—Dec. 3.	do.....			
Mammoth Spring, Ark.	Large-mouth black bass.....			
Entire year.	Small-mouth black bass.....			
	Rainbow trout.....			
	Rock bass.....			
	White bass.....			
<i>Des Arc, Ark.</i>	do.....			
Mar. 4—May 7.	do.....			
<i>Helena, Ark.</i>	Catfish.....			
Aug. 24—Dec. 29.	Buffalo fish.....			
	Rock bass.....			
	Pike perch.....			
	Fresh-water drum.....			
	Sunfish.....			
	Crappie.....			
	Large-mouth black bass.....			
	Yellow bass.....			
	White bass.....			

OUTPUT OF EACH, 1910—Continued.

Fry.			Fingerlings, yearlings, and adults.			Total output.
Dis-tributed.	Transfers to other stations.	Transfers from other stations.	Dis-tributed.	Transfers to other stations.	Transfers from other stations.	
.....
.....
.....
48,262,000	49,622,000
2,669,000	7,235,000
.....	700	700
17,600	4,860	22,460
.....	233,600	233,600
.....	490,780	490,780
.....	230	230
.....	1,450	1,450
.....	18,535	18,535
.....	3,860	3,860
16,900,000	16,900,000
38,140,000	38,140,000
134,053,000	143,907,000
312,820,000	312,820,000
586,100	237,264	873,364
1,001,500	1,026,500
351,922	4,500,000
.....	351,922
.....	468,640
381,440	22,200
2,612,880	379,640	3,472,520
325,600	217,625	588,225
24,700	37,000	24,700
565,600	837,600
.....
.....
.....
.....
.....	1,400	1,400
.....	82,510	82,510
.....	200	200
.....	4,300	4,300
.....
.....	21,540	21,540
.....	178,675	178,675
.....	10,215	10,215
.....	800	800
.....	8,950	8,950
.....	85,365	85,365
.....	177,010	177,010
.....	18,230	Tupelo, 1,600	18,230
.....	250	250
.....	5,950	5,950

STATIONS OPERATED AND THE

Station and period of operation.	Species.	Eggs.		
		Dis-tributed.	Transfers to other stations.	Transfers from other stations.
Manchester, Iowa Entire year.	Rock bass.....			
	Pike perch.....			
	Brook trout.....			Put-in Bay, 3,500,000..
	Lake trout.....			
	Rainbow trout.....	125,650		Leadville, 10,000.....
<i>La Crosse, Wis.a</i> July 15-Oct. 19.	Small-mouth black bass.....			
	Sunfish.....			
	Yellow perch.....			
	Large-mouth black bass.....			
	Catfish.....			
	Pickereel.....			
	Crappie.....			
	Carp.....			
	Buffalofish.....			
	Pike.....			
	Pike perch.....			
	White bass.....			
<i>North McGregor, Iowa</i> July 15-Oct. 6.	Crappie.....			
	Sunfish.....			
	Large-mouth black bass.....			
	Catfish.....			
	Yellow perch.....			
	Carp.....			
	Pike.....			
	Fresh-water drum.....			
	Small-mouth black bass.....			
	Brook trout.....			
Nashua, N. H. Entire year.	Chinook salmon.....			Baird, 100,000.....
	Brook trout.....			Wytheville, 50,000.....
<i>Lake Sunapee, N. H.</i> Oct. 13-Nov. 22.	Rainbow trout.....			
	Sunapee trout.....			
	Landlocked salmon.....			
Neosho, Mo. Entire year.	Rainbow trout.....	41,264		
	Large-mouth black bass.....			
	Rock bass.....			
	Crappie.....			
	Carp.....			
	Yellow perch.....			
	Pike perch.....			Put-in Bay, 1,800,000..
	Small-mouth black bass.....			
	Brook trout.....			
	Lake trout.....	34,894,000	Duluth, 5,000,000..... Green Lake, 300,000. Sault Ste. Marie, 5,000,000. Alpena, 4,000,000. Charlevoix, 10,584,000.	Wytheville, 100,000..... Charlevoix, 3,066,560..
Alpena, Mich. Feb. 23-May 4.	Lake trout.....			Northville, 4,000,000.....
	Whitefish.....			Detroit, 15,000,000.....
<i>Bay City, Mich.</i> Apr. 1-Apr. 29.	Pike perch.....			
	Whitefish.....			
<i>Belle Isle, Mich.</i> Oct. 25-Dec. 12.	Whitefish.....			
	Whitefish.....			
Charlevoix, Mich. Oct. 20-Dec. 21.	Lake trout.....	3,066,560	Northville, 3,066,560..	Northville, 10,584,000..
	Whitefish.....			Detroit, 15,000,000.....
<i>Cheboygan, Mich.</i> Feb. 28-May 4.	Lake trout.....			
	Lake trout.....			
<i>Detour, Mich.</i> Oct. 18-Nov. 15.do.....			
do.....			
<i>Detour, Mich.</i> Oct. 15-Nov. 10.do.....			
do.....			

^a Station for the collection of fishes from overflowed lands.

OUTPUT OF EACH, 1910—Continued.

Fry.			Fingerlings, yearlings, and adults.			Total output.
Dis-tributed.	Transfers to other stations.	Transfers from other stations.	Dis-tributed.	Transfers to other stations.	Transfers from other stations.	
			8,300			8,300
3,300,000			866,500			3,300,000
			3,880			866,500
			85,700			3,880
			9,695			211,350
						9,695
			53,875			53,875
			10,320			10,320
			77,025			77,025
			111,500			111,500
			500			500
			102,820			102,820
			22,300			22,300
			22,800			22,800
			39,500			39,500
			4,460			4,460
			100			100
			95,125			95,125
			136,100			136,100
			162,025			162,025
			384,700			384,700
			84,700			84,700
			115			115
			3,800			3,800
			3,000			3,000
21,600						21,600
171,029						171,029
788,000						788,000
		St. Johnsbury, 104,000.	57,300		Craig Brook, 2,200. Central Station, 10,000.	57,300
168,500			52,855			262,619
			11,650			11,650
			30,025			30,025
			12,950			12,950
			115			115
			50			50
1,400,000						1,400,000
162,000			14,000			176,000
426,000			106,200			532,200
500			82,000			82,500
			3,500			10,013,500
4,000,000						4,000,000
15,000,000						15,000,000
10,584,000						10,584,000
15,000,000						15,000,000

STATIONS OPERATED AND THE

Station and period of operation.	Species.	Eggs.		
		Dis-tributed.	Transfers to other stations.	Transfers from other stations.
Northville, Mich.—Con- Detroit, Mich..... Entire year.	Whitefish.....	74,500,000	Central Station, 500,000 Duluth, 20,000,000. Sault Ste. Marie, 20,000,000. Alpena, 15,000,000. Charlevoix, 15,000,000.	
<i>Fairport, Mich.</i> Oct. 20—Nov. 23.	Pike perch.....	34,280,000		
<i>Grand Haven, Mich.</i> Nov. 6—Nov. 18.	Lake trout.....			
<i>Grassy Island, Mich.</i> Oct. 25—Dec. 12.	do.....			
<i>Naubinway, Mich.</i> Nov. 15—Nov. 24.	do.....			
<i>Northport, Mich.</i> Oct. 26—Nov. 18.	do.....			
<i>Port Huron, Mich.</i> May 1—May 20.	do.....			
<i>St. James, Mich.</i> Nov. 1—Nov. 24.	do.....			
Sault Ste. Marie, Mich. Feb. 20—May 21.	Whitefish.....			Detroit, 20,000,000....
<i>Manistique, Mich.</i> Oct. 15—Nov. 22.	Lake trout.....			Northville, 5,000,000....
Put-in Bay, Ohio..... Entire year.	do.....			
	Pike perch.....	324,475,000	Duluth, 15,000,000.... Central Station, 6,000,000. Neosho, 1,800,000. Meredosia, 5,000,000. Wytheville, 1,000,000. Manchester, 3,500,000. Cape Vincent, 5,000,000. Cape Vincent, 25,000,000. Central Station, 640,000.	
	Whitefish.....	77,068,000		
	Lake herring.....	1,440,000		
<i>Kelleys Island, Ohio.</i> Nov. 10—Nov. 23.	Whitefish.....			
<i>Middle Bass, Ohio.</i> Nov. 7—Dec. 3.	do.....			
<i>Monroe, Mich.</i> Nov. 1—Nov. 28. Apr. 1—Apr. 20.	do.....			
<i>North Bass Island, Ohio.</i> Nov. 5—Dec. 3. Apr. 16—28.	Pike perch.....			
<i>Port Clinton, Ohio.</i> Nov. 3—Dec. 2. Apr. 3—May 7.	Whitefish.....			
<i>Toledo, Ohio.</i> Apr. 1—May 11.	Pike perch.....			
Quincy, Ill..... Entire year.	do.....			
Meredosia, Ill. ^a July—Dec.	Crappie.....			
	Carp.....			
	Large-mouth black bass.....			
	Catfish.....			
	Yellow perch.....			
	Sunfish.....			
	Pike perch.....			Put-in Bay, 5,000,000....
St. Johnsbury, Vt..... Entire year.	Brook trout.....	35,000	Central Station, 20,000. Craig Brook, 5,000.	
	Small-mouth black bass.....			
	Landlocked sal- mon.....			Green Lake, 5,000....
	Yellow perch.....			
<i>Darling Pond, Vt.</i> Sept. 1—Dec. 21.	Brook trout.....			
<i>Hatch Pond, South Ryegate, Vt.</i> Aug. 9—Nov. 13.	do.....			
<i>Lake Mitchell, Vt.</i> Sept. 1—Dec. 17.	Brook trout.....			

^a Station for the collection of fishes from overflowed lands.

DISTRIBUTION OF FISH AND FISH EGGS, 1910.

19

OUTPUT OF EACH, 1910—Continued.

Fry.			Fingerlings, yearlings, and adults.			Total output.
Dis-tributed.	Transfers to other stations.	Transfers from other stations.	Dis-tributed.	Transfers to other stations.	Transfers from other stations.	
25,000,000						29,000,000
12,100,000						46,380,000
20,000,000						20,000,000
5,000,000						5,000,000
89,375,000						376,550,000
75,020,000						126,448,000
70,300,000						71,740,000
			20,100			20,100
			35			35
			108,045			108,045
			25,350			25,350
			9,055			9,055
			25,000			25,000
4,250,000						4,250,000
1,661,000	Holden,300,000.		346		Holden, 31,425.	1,267,346
	Nashua,104,000.					
140,000			2,550			142,550
4,800	Holden,1,800.				Holden,1,000.	3,000
			3,595			3,595

Station and period of operation.	Species.	Eggs.		
		Dis-tributed.	Transfers to other stations.	Transfers from other stations.
St. Johnsbury, Vt.—Con. Holden, Vt. July 1—Nov. 13. Apr. 12—June 30.	Brook trout.....			
	Landlocked salmon.....			
	Lake trout.....			
Swanton, Vt.....	Pike perch.....			
Mar. 15—June 2.	Yellow perch.....			
San Marcos, Tex.....	Sunfish.....			
Entire year.	Rock bass.....			
	Large-mouth black bass.....			
	Crappie.....			
	Carp.....			
Spearfish, S. Dak.....	Brook trout.....			
Entire year.	Landlocked salmon.....			Grand Lake Stream, 25,000.
	Loch Leven trout.....			
	Blackspotted trout.....	2,719,000	Clackamas, 100,000. Bozeman, 544,000.	
	Rainbow trout.....			Wytheville, 100,000.....
	Steelhead trout.....			Birdsview, 25,000.....
	Brook trout.....			
Sand Creek, Beulah, Wyo. Oct. 20—Jan. 15.do.....			
Schmidt Lake, S. Dak. Oct. 20—Dec. 31.	Blackspotted trout.....			
Thumb of Lake, Yellowstone National Park, Wyo. May 25—Aug. 1.do.....			
Clear Creek, Yellowstone National Park, Wyo. June 1—Aug. 10.do.....			
Columbine Creek, Yellowstone National Park, Wyo. June 1—Aug. 10.do.....			
Cub Creek, Yellowstone National Park, Wyo. June 1—Aug. 10.do.....			
Steamer Fish Hawk, Delaware River, Philadelphia, Pa. May 6—June 1.	Shad.....			
Tupelo, Miss.....	Sunfish.....			
Entire year.	Large-mouth black bass.....			
	Crappie.....			
	Catfish.....			
White Sulphur Springs, W. Va. Entire year.	Rainbow trout.....	100,900		
	Brook trout.....	1,000		
	Large-mouth black bass.....			
	Small-mouth black bass.....			
	Blackspotted trout.....			
Woods Hole, Mass.....	Lobster.....			
Entire year.	Cod.....			Gloucester, 24,835,000..
	Mackerel.....			
	Flatfish.....			
	Sea bass.....			
	Lobster.....			
Chilmark, Mass..... Oct. 1—Oct. 9.do.....			
East Greenwich, Mass. Mar. 1—Apr. 1.	Flatfish.....			
Gosnold, Mass..... Sept. 16—Oct. 9. May 23—June 23.	Lobster.....			
Newport, R. I..... Mar. 10—Apr. 1.	Flatfish.....			

OUTPUT OF EACH, 1910—Continued.

Fry.			Fingerlings, yearlings, and adults.			Total output.
Dis-tributed.	Transfers to other stations.	Transfers from other stations.	Dis-tributed.	Transfers to other stations.	Transfers from other stations.	
		St. Johnsbury, 300,000.	177,975	St. Johnsbury, 31,425.		177,975
		St. Johnsbury, 1,800.	3,500	St. Johnsbury, 1,000.		3,500
			3,370			3,370
20,000,000						20,000,000
1,000,000						1,000,000
			4,130			4,130
			3,335			3,335
			138,239			138,239
			9,675			9,675
			25			25
			684,000			684,000
			12,000			12,000
			68,248			68,248
	Bozeman, 400,000		514,750			2,989,750
			234,775			234,775
1,703,000						1,703,000
			9,950			9,950
			18,850	Helena, 1,600.		18,850
			1,550			1,550
			100			100
			262,275			363,175
59,000			821,870			881,870
			3,200			3,200
200,000			1,750			201,750
			2,480			2,480
17,499,000						17,499,000
61,413,000						61,413,000
764,000						764,000
215,770,000						215,770,000
808,000						808,000

STATIONS OPERATED AND THE

Station and period of operation.	Species.	Eggs.		
		Dis-tributed.	Transfers to other stations.	Transfers from other stations.
Woods Hole, Mass.— Continued.				
<i>Noank, Conn.</i>	Lobster.....			
Sept. 29–Oct. 21.				
<i>Plymouth, Mass.</i>	Cod.....			
Nov. 10–Mar. 22.				
<i>Sandwich, Mass.</i>	Lobster.....			
May 3–June 23.				
<i>Waquoit, Mass.</i>	Flatfish.....			
Jan. 20–Mar. 23.				
<i>Westport, Mass.</i>	Lobster.....			
May 3–June 23.				
<i>West Tisbury, Mass.</i> ..	do.....			
May 3–June 23.				
Oct. 1–Oct. 10.				
<i>Wickford, R. I.</i>	Flatfish.....			
Mar. 17–Apr. 1.				
Wytheville, Va.....	Large-mouth black bass.			
Entire year.	Small-mouth black bass.			
	Rock bass.....			
	Yellow perch.....			
	Rainbow trout....	948,000	Erwin, 503,000. Cape Vincent, 50,000. Nashua, 50,000. Spearfish, 100,000. Central Station, 15,000. Northville, 100,000.	
	Brook trout.....			
	Carp.....			
	Pike perch.....			
Yes Bay, Alaska.....	Blueback salmon..			Put-in Bay, 1,000,000.
Entire year.				
Total output of Bureau.				

ALLOTMENTS TO STATE FISH COMMISSIONS.

As usual, various state fish commissions were supplied from the Bureau's stock with eggs to be hatched and distributed under their respective auspices. Following is a record of such allotments in 1910:

ALLOTMENTS OF FISH AND EGGS TO STATE FISH COMMISSIONS, FISCAL YEAR 1910.

State and species.	Eggs.	Finger- lings, year- lings, and adults.	State and species.	Eggs.	Finger- lings, year- lings, and adults.
California:			New York:		
Chinook salmon.....	28,764,467		Blackspotted trout.....	50,000	
Colorado:			Rainbow trout.....	41,500	
Blackspotted trout.....	225,000		Landlocked salmon.....	15,000	
Connecticut:			White perch.....	15,000,000	
Yellow perch.....	5,200,000		North Dakota:		
Illinois:			Steelhead trout.....	100,000	
Lake trout.....	500,000		Pike perch.....	10,000,000	
Whitefish.....	4,000,000		Ohio:		
Pike perch.....	8,000,000		Whitefish.....	18,000,000	
Rainbow trout.....	41,264		Pike perch.....	170,725,000	
Michigan:			Oregon:		
Landlocked salmon.....	20,000		Chinook salmon.....	6,465,300	60
Lake trout.....	5,000,000	3,500	Blackspotted trout.....	175,000	45
Pike perch.....	34,280,000		Pennsylvania:		
Missouri:			Silver salmon.....	75,000	
Brook trout.....	100,000		Blackspotted trout.....	50,000	
Rainbow trout.....	25,000		Whitefish.....	31,428,000	
Pike perch.....	2,000,000		Pike perch.....	96,450,000	
Minnesota:			Washington:		
Large-mouth black bass..		18,250	Steelhead trout.....	50,000	
Montana:			Brook trout.....	100,000	
Blackspotted trout.....	550,000		Wisconsin:		
Whitefish.....	500,000		Lake trout.....	4,500,000	3,880
Nevada:			Wyoming:		
Blackspotted trout.....	422,100		Blackspotted trout.....	675,000	
New Hampshire:			Total.....	443,627,631	25,735
Chinook salmon.....	100,000				

SHIPMENTS TO FOREIGN COUNTRIES.

In response to requests reaching the Bureau through diplomatic channels, fish and fish eggs have been donated to foreign countries as follows:

SHIPMENTS OF FISH AND EGGS TO FOREIGN COUNTRIES, FISCAL YEAR 1910.

Country.	Species.	Eggs.	Finger- lings, year- lings, and adults.
Argentina.....	Chinook salmon.....	200,000	
	Silver salmon.....	100,000	
	Blueback salmon.....	100,000	
	Landlocked salmon.....	25,000	
France.....	Lake trout.....	50,000	
	Blackspotted trout.....	10,000	
Japan.....	Rainbow trout.....	110,000	
	Brook trout.....	5,000	
Mexico.....	Carp.....		25
Total.....		600,000	25

SUMMARIZED STATEMENT OF DISTRIBUTIONS.

The following table shows the numbers of eggs and fish actually distributed during the fiscal year 1910; or, in other words, the output of the hatcheries with all losses in transportation deducted. It thus does not agree with the tabulated summary in the Annual Report of the Commissioner for this year, compiled at an earlier date, which shows the numbers of eggs and fish delivered by the stations for distribution, the subsequent losses in transportation not being considered:

SUMMARY OF DISTRIBUTION OF FISH AND EGGS, FISCAL YEAR 1910.

Species.	Eggs.	Fry.	Fingerlings, yearlings, and adults.	Total.
Catfish.....			531,892	531,892
Carp.....			22,710	22,710
Buffalofish.....			201,475	201,475
Shad.....	2,160,000	89,076,000		91,236,000
Whitefish.....	55,428,000	195,719,000		251,147,000
Lake herring.....	1,440,000	70,300,000		71,740,000
Silver salmon.....	375,000	10,888,025		11,263,025
Chinook salmon.....	37,531,417	16,342,556	66,045	53,940,018
Blueback salmon.....	100,000	121,136,995	21,719,600	142,956,595
Humpback salmon.....		1,731,740		1,731,740
Steelhead trout.....	250,000	3,570,287	179,718	4,000,005
Rainbow trout.....	556,494	595,616	1,705,328	2,857,438
Atlantic salmon.....	5,000	1,217,366	238,212	1,460,578
Landlocked salmon.....	115,000	974,040	301,064	1,390,104
Blackspotted trout.....	2,748,550	1,756,094	906,654	5,411,298
Loch Leven trout.....			68,248	68,248
Lake trout.....	10,210,000	33,645,922	4,286,150	48,142,072
Brook trout.....	516,000	7,365,945	4,085,174	11,967,119
Sunapee trout.....		171,029		171,029
Grayling.....	25,000	81,000	18	106,018
Smelt.....	4,500,000		9,000	4,509,000
Pike.....			43,300	43,300
Pickereel.....			500	500
Crappie and strawberry bass.....			410,428	410,428
Rock bass.....			66,035	66,035
Warmouth bass.....			792	792
Small-mouth black bass.....		537,400	109,986	647,386
Large-mouth black bass.....		56,600	665,868	722,468
Sunfish (bream).....			342,825	342,825
Pike perch.....	321,455,000	154,480,000	5,260	475,940,260
Yellow perch.....	5,200,000	326,885,000	108,439	332,193,439
Striped bass.....	4,566,000	2,784,000		7,350,000
White bass.....			6,050	6,050
White perch.....	16,500,000	338,480,000		354,980,000
Yellow bass.....			250	250
Sea bass.....		808,000		808,000
Mackerel.....		764,000		764,000
Freshwater drum.....			11,950	11,950
Cod.....	9,854,000	210,354,000		220,208,000
Pollock.....		38,140,000		38,140,000
Haddock.....		712,000		712,000
Flatfish.....		930,755,000		930,755,000
Lobster.....		162,505,000	1,532	162,506,532
Total.....	473,535,461	2,721,832,615	36,094,503	3,231,462,579

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS, FISCAL YEAR 1910.

CATFISH.

Disposition.	Finger- lings, year- lings, and adults.	Disposition.	Finger- lings, year- lings, and adults.
Arizona:		Minnesota:	
Grand Canyon, Berry's pond.....	100	Brownsville, Mississippi River.....	43, 250
Summit Pond.....	100	Mahnomen, Mayzhuckegishig Lake.....	800
Holbrook, Becker's reservoir.....	100	Rochester, Zumbro River, South Fork.....	500
Pratt's pond.....	100	Mississippi:	
Prescott, American Ranch Lake.....	100	Guntown, Cochran's pond.....	100
Wilcox, Adling's pond.....	100	Missouri:	
Ditmar's pond.....	100	Brandsville, Niessen's pond.....	150
Arkansas:		Richland, Gasconade River.....	400
Boonville, Branch Pond.....	100	Seligman, Mountain Pond.....	200
Green Forest, Willow Pond.....	100	Springfield, Appleby's pond.....	200
Harrison, Estes's pond.....	125	New Jersey:	
Helena, Mississippi River.....	20, 640	Mullica Hill, Mullica Hill Pond.....	400
Hiawassee, Rucker's pond.....	100	Pompton Lakes, Pompton Lakes.....	400
McNeil, Stevens's pond.....	273	Washington, Fair Haven Pond.....	100
Mammoth Spring, Warm Fork Creek..	100	New Mexico:	
Stamps, Price Pond.....	150	Clovis, Laughing Water Pond.....	100
Colorado:		Columbus, Kennedy's pond.....	80
Pueblo, Skinner's reservoir.....	100	Corona, Ingram's pond.....	80
Rifle, White River.....	200	Deming, Burney's pond.....	100
Georgia:		Harris's pond.....	200
Chamblee, Jones's pond.....	100	Hon's pond.....	100
Idaho:		Jacobson's pond.....	100
Grangeville, Tolo Lake.....	300	Kelly's pond.....	100
Naples, Stampede Lake.....	300	Elida, Brown's pond.....	100
Illinois:		La Lande, McGill's reservoir.....	100
Avena, Sycamore Lake.....	400	Las Vegas, Asylum Lake.....	100
Chicago, Armour's pond.....	450	Pecos River.....	100
Otis's pond.....	450	Montoya, Paloma Springs.....	80
Galva, Mirror Pond.....	500	Portales, Humble's pond.....	100
Odell, Odell Pond.....	500	Twin Mill Ponds.....	100
Tremont, Pfederer's pond.....	500	Silver City, Central Creek Pond.....	100
Indiana:		Texico, Stafford's pond.....	100
Boonville, Hemenway's pond.....	500	Tucumcari, Buchanan's pond.....	80
Buckskin, Buck's pond.....	100	New York:	
Centerville, Townsend's pond.....	100	Cooperstown, Schuylers Lake.....	300
Evansville, Bockstede's pond.....	100	Greenport, Sills Pond.....	150
Heltonville, Ramsey's pond.....	100	Unadilla, Susquehanna River.....	300
Lewis, Freeze's pond.....	100	Walden, Walkkill River.....	152
Pleasant Lake, Pleasant Lake.....	200	Walkill, Dwaarskill Creek.....	155
Tilden, Hadley's pond.....	300	North Dakota:	
Iowa:		Devils Lake, Devils Lake.....	3, 000
Chester, Upper Iowa River.....	400	Glen Ullin, Burns's pond.....	100
Independence, Wapsipinicon River.....	400	Gwinner, Edmon's pond.....	150
Line Springs, Upper Iowa River.....	2, 500	Milnor, Stone Lake.....	100
Manchester, Maquoketa River.....	4, 000	Oakes, Christenson's pond.....	150
North McGregor, Mississippi River.....	187, 500	St. John, Bouvin Lake.....	400
Kansas:		Ohio:	
Goddard, Clear Creek Pond.....	65	Bethel, McCarty's pond.....	100
Kansas City, Hosps's pond.....	80	Bradford, Greenville Creek.....	250
Marquette, Sunny Pond.....	65	Upper Stillwater Creek.....	150
Pawnee, Payton's pond.....	65	Cincinnati, Lake Como.....	150
Kentucky:		Cridersville, Retreat Lake.....	100
Elizabethtown, Hagan's pond.....	200	Dola, Hively's pond.....	100
Hodgensville, Nolin Creek.....	400	Ironton, Kucker's pond.....	150
Nolin Creek, North Fork.....	300	Jackson, Long's pond.....	100
Tharpe's pond.....	200	Marion, Scioto River.....	250
Louisiana:		Orbiston, Orbiston Lakes.....	100
Grand Cane, Clear Springs Pond.....	100	Ravenna, Infirmary Pond.....	150
Maryland:		Ripley, Hauke's pond.....	150
Loch Raven, Harrison's pond.....	150	Rock Creek, Parks's pond.....	100
Mountain Lock, Potomac River.....	450	Stryker, Juillard's pond.....	100
Rocky Ridge, Owings Creek.....	150	Wapakoneta, Brown Pond.....	400
Sharon, Rogers Pond.....	150	Youngstown, Mahoning River.....	100
Massachusetts:		Wickliffe Lake.....	200
Westdale, Taunton River.....	500	Oklahoma:	
Michigan:		Aline, Elliott's pond.....	100
Collins, Grand River.....	480	Bison, Springdale Pond.....	100
Jackson, Big Portage Lake.....	480	Chilocco, Chilocco Lagoon.....	200
Grass Lake.....	480	Collinsville, Ellingswood Lake.....	200
Lakeview, Brimmer Lake.....	1, 000	Cushing, Prairie Lake.....	100
Tamarack Lake.....	1, 000	Twin Elm Lake.....	125
Town Line Lake.....	1, 000	Wild Horse Pond.....	150
Penn, Mud Lake.....	650	Enid, Spring Valley Creek.....	100
Portland, Grand River Pond.....	480	Erick, Garrett's pond.....	100

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

CATFISH—Continued.

Disposition.	Fingerlings, yearlings, and adults.	Disposition.	Fingerlings, yearlings, and adults.
Oklahoma—Continued.		South Carolina—Continued.	
Glencoe, Greenwood Lake.....	75	Starr, Pruitt's pond.....	125
South Side Pond.....	75	Walhalla, Carey's pond.....	125
Guymon, Jordan's pond.....	250	South Dakota:	
Hastings, Washash Pond.....	125	Fairfax, Manhalter's pond.....	150
Isabella, Wahl's pond.....	100	Phillip, Grindstone Pond.....	200
Lawton, Park Lake.....	175	Presho, Corkill's lake.....	200
Maramec, Maramec Lake.....	150	Scenic, Knutson's pond.....	200
Marshall, Crouch's pond.....	100	Warner, Papke's pond.....	200
Proctor's pond.....	100	Vermont:	
Perkins, Canon Pond.....	75	Bellows Falls, Connecticut River.....	400
Stigler, Hall's pond.....	200	Virginia:	
Stillwater, Boomer Creek.....	100	Covington, McAllister's pond.....	150
Kautz's ponds.....	75	Dillwyn, North River.....	300
Nash's pond.....	75	Slate River.....	300
Swartz's pond.....	75	Gainesville, Broad Run.....	300
Stratford, Davis's pond.....	100	Houston, Easley Mill Pond.....	230
Waynoka, Hancock's pond.....	275	Occoquan, Occoquan River.....	300
Yost, Newman's pond.....	75	Palmyra, Rivanna River.....	350
Yost Lake.....	75	Urbanna, Jackson's pond.....	550
Pennsylvania:		Washington:	
Birdsboro, Monocacy Creek.....	200	Addy, Blue Lake.....	75
Carbon Center, Carbon Center Pond.....	100	Spring Lake.....	75
Factoryville, Lake Carey.....	350	Anacortes, Lake Erie.....	150
Greensburg, Hacke Pond.....	100	Montesano, Silvia Lake.....	150
Kingston, Ryman's pond.....	400	Oroville, Lemonosky Lake.....	150
Rahns, Perkiomen Creek.....	400	West Virginia:	
Reading, Maiden Creek.....	400	Bedington, Emerson's pond.....	150
Rupert, Wide Water Canal.....	200	Benwood, Riedel's pond.....	250
Scottdale, Mill Race Pond.....	150	Grafton, Otter Creek Pond.....	250
Smiths Ferry, Woodlawn Pond.....	100	Nuttall, Chalybeate Spring Pond.....	250
Susquehanna, Churchill's lake.....	300	Romney, Potomac River, South Branch.....	550
Susquehanna River.....	300	Wisconsin:	
Troy, Cross Roads Creek.....	100	Brillion, Long Lake.....	300
Lillmary Creek.....	100	Round Lake.....	300
Mud Creek.....	100	Genoa, Mississippi River.....	4,166
Sugar Creek.....	450	La Crosse, Mississippi River.....	47,418
Wilkes Barre, Bear Lake.....	150	Mauston, Drainage Canal.....	300
Wolmesdorf, Tulpehocken Creek.....	400	Pelican, Little Mud Lake.....	300
Wagners Pond.....	400	Rice Lake.....	400
South Carolina:		Prairie du Chien, Mississippi River.....	172,500
Blackville, Rodgers Pond.....	175	Sheboygan Falls, Sheboygan River.....	500
Graycourt, North Rabun Creek.....	125	Victory, Mississippi River.....	1,666
Honea Path, Barkers Creek.....	150	Wyoming:	
Broad Creek.....	200	Lusk, "J. M." Company's pond.....	400
Haynie Pond.....	150	Moorcroft, Lone Tree Reservoir.....	200
Kays Pond.....	200	Newcastle, Lodge Pole Creek.....	250
Pickens, Bivers Lake.....	250	Sheridan, Big Horn Pond.....	150
Holders Lake.....	250	Total ^a	531,892
Spring Lake.....	300		
Thornley Pond.....	200		

CARP.

Kansas:		West Virginia:	
Pittsburg, North Lake.....	35	Moundsville, Jones's pond.....	15
Minnesota:		Wisconsin:	
Brownsville, Mississippi River.....	8,650	Genoa, Mississippi River.....	1,666
New York:		La Crosse, Mississippi River.....	10,318
Riverhead, Harrison's pond.....	100	Victory, Mississippi River.....	1,666
Oklahoma:		Mexico:	
Stillwater, Willow Pond.....	100	Sonora, Ysabel Lake.....	25
Vian, Allen's pond.....	15	Total.....	22,710
Virginia:			
Wytheville, Brownings Mill Pond.....	110		
Indian Creek.....	10		

^a Lost in transit, 12,078 fingerlings.

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BUFFALOFISH.

Disposition.	Finger- lings, year- lings, and adults.	Disposition.	Finger- lings, year- lings, and adults.
Arkansas:		Wisconsin:	
Helena, Mississippi River	178, 675	Genoa, Mississippi River	2, 666
Minnesota:		La Crosse, Mississippi River	11, 318
Brownsville, Mississippi River	8, 650	Victory, Mississippi River	166
		Total	201, 475

SHAD.

Disposition.	Eggs.	Fry.	Disposition.	Eggs.	Fry.
District of Columbia:			New Jersey—Continued.		
Washington, Anacostia River		295, 000	Riverton, Delaware River		80, 000
Potomac River		682, 000	Timber Creek, Delaware River		120, 000
Maryland:			New York:		
Accokeek Creek, Potomac River		980, 000	New York, New York Aquarium	800, 000	
Broad Creek, Potomac River		2, 504, 000	North Carolina:		
Carpenters Point, North East River		234, 000	Edenton, Albemarle Sound	1, 360, 000	47, 762, 000
Havre de Grace, Chesapeake Bay.		3, 485, 000	Tarboro, Tar River		500, 000
Susquehanna River		821, 000	Oregon:		
Swan Creek		396, 000	Willamette, Willamette River		1, 588, 000
Occoquan Bay, Potomac River		898, 000	Pennsylvania:		
Pamunkey Creek, Potomac River		5, 044, 000	Poquessing Creek, Delaware River		200, 000
Piscataway Creek, Potomac River		4, 621, 000	Virginia:		
Swan Creek, Chesapeake Bay		70, 000	Dogue Creek, Potomac River		2, 401, 000
Potomac River		3, 572, 000	Little Hunting Creek, Potomac River		2, 717, 000
Wild Duck Harbor, Susquehanna River		385, 000	Occoquan Creek, Potomac River		3, 391, 000
New Jersey:			Pamunkey Creek, Potomac River		600, 000
Camden, Delaware River		803, 000	Pohick Creek, Potomac River		4, 337, 000
Rancocas, Delaware River		500, 000	Washington:		
			Hamilton, Skagit River		90, 000
			Total	2, 160, 000	89, 076, 000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

WHITEFISH.

Disposition.	Eggs.	Fry.	Disposition.	Eggs.	Fry.
Illinois:			Montana:		
Havana, Illinois Fish Commission.....	4,000,000	Anaconda, Montana State Fishery.....	500,000
Michigan:			New York:		
Alpena, Lake Huron.....	1,000,000	Cape Vincent, Lake Ontario.....		1,500,000
Belle Isle, Lake St. Clair.....	9,000,000	Chaumont, Lake Ontario.....		2,000,000
Detour, Lake Huron.....	6,000,000	Cooperstown, Otsego Lake.....		387,000
Lake Michigan.....	3,000,000	Fox Island, Lake Ontario.....		3,500,000
Detroit, Detroit River.....	16,000,000	Fullers Bay, Lake Ontario.....		170,000
Escanaba, Lake Michigan.....	2,000,000	Grenadier Island, Lake Ontario.....		5,500,000
Fish Island, Lake Superior.....	490,000	Hayes Point, Lake Ontario.....		2,000,000
Isle Royale, Lake Superior.....	13,100,000	Mexico, Lake Ontario.....		4,000,000
McCargoes Cove, Lake Superior.....	210,000	New York, New York Aquarium.....	1,500,000
Manistique, Lake Michigan.....	2,000,000	Oneida Lake, Oneida Lake.....		387,000
Marquette, Lake Superior.....	4,655,000	Wilson Bay, Lake Ontario.....		1,500,000
North Point, Lake Huron.....	9,000,000	Ohio:		
Skulligallee Reef, Lake Michigan.....	5,000,000	Catawba Island, Lake Erie.....		10,000,000
St. Ignace, Lake Huron.....	2,000,000	Isle St. George, Lake Erie.....		10,000,000
Sand Bay Reef, Lake Michigan.....	5,000,000	Kelleys Island, Lake Erie.....		20,000,000
Scarecrow Island, Lake Huron.....	5,000,000	Lakeside, Lake Erie.....		20,000,000
Simmons Reef, Lake Michigan.....	5,000,000	Put-in Bay, Lake Erie.....		25,000,000
Whitefish Point, Lake Superior.....	5,000,000	Ohio State Fish Commission.....	18,000,000
Minnesota:			Toledo, Lake Erie.....		10,000,000
Duluth, Lake Superior.....	300,000	Pennsylvania:		
Grand Marais, Lake Superior.....	3,000,000	Erie, Pennsylvania Fish Commission.....	31,428,000
Susie Island, Lake Superior.....	3,000,000	Total ^a	55,428,000	195,719,000

LAKE HERRING, OR CISCO.

Ohio:			Ohio—Continued.		
Cleveland, Lake Erie.....	1,440,000	Port Clinton, Lake Erie.....		10,000,000
Isle St. George, Lake Erie.....	10,000,000	Put-in Bay, Lake Erie.....		10,000,000
Kelleys Island, Lake Erie.....	10,000,000	Toledo, Lake Erie.....		10,000,000
Lakeside, Lake Erie.....	300,000	Total	1,440,000	70,300,000
Middle Bass, Lake Erie.....	20,000,000			

SILVER SALMON.

California:			Washington:		
Brookdale, San Lorenzo River.....	100,000	Baker, Baker Lake.....		5,308,848
Santa Cruz County Hatchery.....	100,000	Lower Baker River.....		500,000
Pennsylvania:			Birdsview, Grandy Creek.....		5,079,177
Pleasant Mount, State Fish Commission.....	75,000	Argentina:		
			Buenos Aires, Argentine Government.....	100,000
			Total	375,000	10,888,025

^a Lost in transit, 245,000 fry.

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

CHINOOK SALMON.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
California:			
Baird, McCloud River.....		2, 286, 277	
Brookdale, Santa Cruz County Hatchery.....	1, 000, 000		
Eel River, California Fish Commission.....	1, 549, 500		
Point Reyes, applicant.....	300, 000		
Sisson, California Fish Commission.....	27, 214, 967		
New Hampshire:			
Edgemont, Lake Sunapee.....			7, 380
Laconia, New Hampshire Fish Commission.....	100, 000		
Newbury, Lake Sunapee.....			51, 200
New York:			
New York, New York Aquarium.....	5, 000		
Port Kent, Lake Champlain.....			3, 600
Tuxedo Park, applicant.....	25, 000		
Westport, Lake Champlain.....			3, 640
Oregon:			
Bonneville, Oregon Fish Commission.....	6, 465, 300		
Cazadero, Clackamas River.....		534, 197	
Clackamas, Clackamas River.....		3, 686, 200	70
Oregon Fish Commission.....			60
Rogue River, Elk Creek.....		100, 362	
Rogue River.....		499, 930	
Wedderburn, applicant.....	572, 400		
Washington:			
Baker, Baker Lake.....		349, 570	
Big White Salmon, Columbia River.....		2, 612, 200	
Spring Creek.....		900, 000	
Birdsview, Grandy Creek.....		705, 840	
Little White Salmon, Columbia River.....		1, 900, 000	
Little White Salmon River.....		2, 908, 000	
Seattle, Exposition Aquarium.....	99, 250		95
Argentina:			
Buenos Aires, Argentine Government.....	200, 000		
Total.....	37, 531, 417	16, 342, 556	66, 045

BLUEBACK SALMON.

Alaska:			
Afognak, Ahuyon Creek.....		34, 018, 060	
Letnik Lake.....		34, 404, 110	
Yes Bay, McDonald Lake.....			21, 719, 600
Yes River.....		48, 160, 000	
Washington:			
Baker, Baker Lake.....		4, 494, 825	
Lower Baker River, Lower Baker River.....		150, 000	
Argentina:			
Buenos Aires, Argentine Government.....	100, 000		
Total.....	100, 000	121, 136, 995	21, 719, 600

^a Lost in transit, 1,480 fingerlings.

HUMPBACK SALMON.

Disposition.	Fry.
Alaska:	
Afognak, Letnik Lake.....	363, 740
Washington:	
Birdsview, Grandy Creek.....	1, 368, 000
Total.....	1, 731, 740

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

STEELHEAD TROUT.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Maryland:			
Clear Spring, Tom Run Pond.....		12,000	
Michigan:			
Humboldt, Black River.....			10,000
Michiganme River.....			10,000
Spruce River.....			10,000
Munising, applicant.....	50,000		
Watersmeet, Duck Lake.....			14,000
Wetmore, Big Indian River.....			32,000
Minnesota:			
Duluth, Canosia Lake.....			12,000
Pike Lake.....			21,000
Knife River, Mic Mac Lake.....			12,000
Teteagonche Lake.....			16,000
Montana:			
Bozeman, Bridger Creek.....			8,300
Deer Lodge, Powell Lake.....			1,500
Libby, Kootenai River.....			400
Logging Creek, Belt Creek.....			2,500
Norris, Madison River Power Co. Lake.....			6,000
New York:			
Auburn, Owasco Lake.....		35,423	
Pulaski, Salmon River.....		11,338	
North Dakota:			
St. John, State fish commission.....	100,000		
Oregon:			
Cazadero, Clackamas River.....		1,934,835	
Eagle Creek, Eagle Creek.....		49,503	
Rogue River, Elk Creek.....		89,850	
Washington:			
Baker, Baker Lake.....		14,400	
Birdsview, Day Creek.....		40,300	
Grandy Creek.....		1,382,638	
Seattle, Exposition Aquarium.....			18
State Fish Commission.....	50,000		
Walla Walla, applicant.....	25,600		
Wisconsin:			
Hudson, applicant.....	25,000		
Lampson, Horse Shoe Lake.....			14,000
Spooer, Christie Lake.....			10,000
Total.....	250,000	3,570,287	179,718

RAINBOW TROUT.

Alabama:			
Tanner, Pecks Branch.....			2,400
Arizona:			
Flagstaff, Live Oak Creek.....			7,200
Rock Creek.....			7,200
Tucson, Sabino Creek.....			6,000
Winslow, Chevelon Creek.....			7,200
Arkansas:			
Bald Knob, Hart's pond.....			4,000
Berryville, Osage River.....		25,000	
Crickette, Yocum Creek.....			4,000
Decatur, Lakeside Pond.....		7,500	
Elkins, White River.....			800
Flippin, Goff's pond.....		7,500	
Greenwood, Vache Grass Creek.....		7,000	
Mammoth Spring, Spring River.....			200
Springdale, Lake Vaughan.....		7,000	
Sulphur Springs, Williams's pond.....		7,600	
California:			
Brookdale, Santa Cruz County hatchery.....	13,680		
Colorado:			
Buena Vista, Chalk Creek.....			6,000
Cottonwood Creek.....			6,000
Middle Cottonwood Creek.....			6,000
South Cottonwood Creek.....			6,000
Buffalo, Platte River.....			10,000
Cimarron, Little Cimarron Creek.....		2,000	
Colorado Springs, Frost's reservoir.....			25
Creede, applicant.....	100,000		
Eldora, Lake Eldora.....			8,535
Estabrook, Mendenhall Creek.....			3,750

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

RAINBOW TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Colorado—Continued.			
Grand Mesa Lakes, Ward Lake		55,000
Grant, Geneva Lake.....			2,500
Platte River.....			10,000
Ivanhoe, Frying Pan River.....		10,500
Jefferson, Platte River.....		25,000
Kline, Platte River.....		25,000
Leadville, Musgrove's pond.....			20,000
Malta, Big Thompson Stream.....	20,000	
Minturn, Eagle River.....			9,000
Moffat, Saguache Creek.....		25,000
Molina, Cottonwood Lakes.....		11,000
Montrose, East Dry Creek.....		4,000
New Castle, Divide Creek.....			9,000
Elk Creek.....			9,000
Pine Grove, Elk Creek.....			3,750
Wright's lake.....			2,500
Pueblo, Gunnison River.....		10,000
West Elk Creek.....		10,000
Salida, Little River.....		7,500
South Arkansas River.....		10,000
Shawnee, Price Creek.....			2,500
South Platte, South Platte River.....			10,000
South Platte River, South Fork.....			12,500
Telluride, Dolores River.....			10,000
Thomasville, applicant.....		25,000
Twin Lakes, Willow Lake.....		4,000
Webster, Beaver Creek.....			2,500
West Cliffe, Brush Creek Lake.....			6,000
Swift Creek.....			6,000
Georgia:			
Clayton, Hiwassee River.....			4,000
Mathias, Tallulah River.....			4,000
Oakman, Dry Creek.....			4,000
Rabun Gap, Charley Creek.....			3,200
Flat Branch.....			2,400
Mill Creek.....			3,200
Shook Creek.....			2,400
Tallulah River.....			4,000
Tate Creek.....			2,400
Ringgold, Murphy's pond.....			1,600
Idaho:			
Ashton, Eggbert Lake.....			1,000
Bliss, Far View Lakes.....			1,500
Cambridge, Little Weiser River.....			1,000
Hailey, applicant.....		5,000
Priest River, Skookum Pond.....			500
Troy, Pineview Pond.....			600
Illinois:			
Havana, Illinois Fish Commission.....	41,264	
Indiana:			
St. Paul, Mill Creek.....			2,000
South Bend, Beyer's lake.....			1,000
Leeper Pond.....			1,000
Iowa:			
Manchester, Maquoketa River.....			400
North McGregor, Bloody Run.....			3,000
Postville, Livingood Creek.....			600
Waukon, Silver Creek.....			1,000
Village Creek.....			1,500
Kansas:			
Erie, Canville Creek.....			200
Marion, Spring Creek.....			2,000
Maryland:			
Cumberland, Evitts Creek.....			3,000
Lakewood Lake.....			2,000
Minley Branch.....			2,500
Rocky Gap Creek.....			2,000
Mountain Lake Park, Broad Ford Creek.....			5,000
Little Youghiogheny River.....			480
Oakland, Browning Dam.....			5,000
Harvey's pond.....			320
Westminster, Fairview Pond.....			500
Michigan:			
Brentcreek, Gillett's pond.....			1,000
East Tawas, Cold Creek.....			5,000
Gaylord, Sturgeon River.....			15,000
Gladwin, Cedar River.....			1,250
Grayling, Tillula Lake.....			590

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

RAINBOW TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Michigan—Continued.			
Hillman, Thunder Bay River.....			10,000
Kalamazoo, applicant.....	10,000		
Halls Springs Brook.....			2,000
Portage Creek.....			2,000
Paris, Muskegon River.....			18,000
Petersburg, Crystal Pond.....			6,000
Plymouth, Millers Creek.....			500
Rose Center, Buckhorn Creek.....			12,000
West Branch, Chapman Creek.....			400
Tittabawassa River.....			1,250
Wingleton, Marquette River.....			18,750
Marquette River, South Branch.....			3,500
Minnesota:			
Duluth, Archer Creek.....			1,800
Silica, Little Swan Creek.....			3,000
Winona, Stockton Creek.....			2,500
Missouri:			
Aurora, Spring Creek.....			400
Turnback Creek.....			4,000
Wistman Creek.....			400
Bourbon, Blue Spring Branch.....			6,190
Brown Springs, Brown Springs Lake.....			400
Cabool, Flag Lake.....		12,500	
Clever, King's pond.....		7,500	
Lucas Branch.....			4,000
Silver Lake Branch.....		20,000	
Exeter, Roaring River.....		20,000	
Galena, Langley's pond.....			4,000
Marshfield, James River.....			6,000
Neosho, Hickory River.....			55
Newburg, Little Piney River.....			6,810
Mill Creek.....			4,000
Reeds Spring, Moose Springs.....		2,500	
St. James, Meramec Springs.....			6,000
St. Joseph, Missouri Fish Commission.....	25,000		
Springfield, Spring Creek.....		15,000	
Verona, Spring River.....		30,000	
Wheaton, Joys Creek.....			400
Pogues Creek.....			400
Shoal Creek.....			800
Montana:			
Armstead, McIntosh Creek.....			1,200
Spring Creek.....			1,200
Bozeman, Wild Horse Run.....			2,000
Chinook, Box Elder Creek.....			2,000
Columbia Falls, Fish Lake.....			2,000
Delphia, Half Moon Lake.....			1,000
Dillon, Ajax Creek.....			960
Blacktail Deer Creek.....			900
Carter Creek.....			2,400
Lake Creek.....			960
North Fork River.....			960
Stewart Gulch.....			960
Strowbridge's pond.....			960
Tent Lake.....			1,200
Van Camp Creek.....			1,200
Emigrant, Dailey's lake.....			2,000
Fortine, Fortine Creek.....			2,000
Lakeview, Cliff Lake.....		4,000	4,000
Elk Creek.....		10,000	
Elk Lake.....		5,000	
Hidden Lake.....		4,000	
Thompson, Clear Creek.....			1,500
Squaw Creek.....			1,500
Townsend, Duck Creek.....			2,000
Nebraska:			
Andrews, White River.....			10,000
Gretna, Chadron Creek.....			1,600
Nevada:			
Verdi, Boulder Riffles.....			4,000
Chalk Bluff Pools.....			4,000
Marble Works Pools.....			4,000
Truckee River.....			8,000
New Jersey:			
Jersey City, Witterman's pond.....			2,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

RAINBOW TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
New Mexico:			
Cimarron, Aqua Fria Creek.....			2,000
Canon Bonita Creek.....			1,000
Cimarroncita Creek.....			1,000
Cimarron River.....			1,000
Clear Creek.....			1,000
Fossil Creek.....			1,000
Rayado Creek.....			1,000
Las Vegas, Gallinas River, West Fork.....		2,400	
Raton, Myrtle Pond.....		1,200	
Sugarite Creek.....		6,000	
Roswell, Crystal Pond.....			2,000
New York:			
Adams, Big Sandy Creek.....		19,000	
Buffalo, New York State Cancer Laboratory.....		500	
Clifton, Wittenman Pond.....			2,000
Linthigo, Forest, Fish, and Game Commission.....	41,500		
New York, New York Aquarium.....	5,000		
Valhalla, Wygoda Pond.....			400
Willsboro, Warm Pond.....		19,000	
North Carolina:			
Addie, Buff Creek.....			1,600
Asheville, French Broad River.....			100
Midget Lake.....			1,600
Balsam, Scotts Creek.....			3,200
Barnard, Big Pine Creek.....			4,000
Black Mountain, Swanannoa River.....			75
Boonford, Big Crabtree Creek.....			2,100
South Toe River.....			125
Toe River.....			150
Brevard, Allison's lake.....			4,000
Bryson, Alarka Creek.....			3,200
Address Creek.....			2,400
Bald Creek.....			2,400
Bear Creek.....			2,400
Bear Meat Creek.....			2,400
Big Hurricane Creek.....			2,400
Bridge Creek.....			2,400
Cherry Creek.....			2,400
Clingman Creek.....			2,400
Cold Spring.....			2,400
Conley Creek.....			2,400
Cooper Creek.....			2,400
Cullasowah Creek.....			2,400
Deep Creek.....			3,200
Galbreath Creek.....			2,400
Grassy Branch.....			2,400
Indian Creek.....			2,400
Jenkins Creek.....			2,400
Jones Creek.....			2,400
Kirkland Creek.....			8,000
Lands Creek.....			2,400
Laurel Creek.....			2,400
Little Hurricane Creek.....			2,400
Long Creek.....			7,200
Middle Hurricane Creek.....			2,400
Mill Creek.....			2,400
Nettle Creek.....			2,400
Noland Creek.....			3,200
North Fork Creek.....			1,600
Peach Tree Creek.....			2,400
Pigeon Creek.....			2,400
Saw Mill Creek.....			2,400
Shepherd Creek.....			2,400
Silver Creek.....			2,400
Una Creek.....			2,400
Watkins Creek.....			2,400
West Fork Creek.....			1,600
Bushnell, Chambers Creek.....			3,200
Indian Camp Creek.....			2,400
Kirklin Creek.....			2,400
Little Laurel Creek.....			2,400
Stecoah Creek.....			2,400
Cherokee, Lufty Creek.....			2,400
Soco Creek.....			4,800
Cranberry, Blevin Creek.....			75
Cranberry Creek.....			75
Roaring Creek.....			3,200

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

RAINBOW TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
North Carolina—Continued.			
Dillsboro, Big Savannah Creek			2,400
Dick Creek			2,400
Savannah Creek, East Fork			2,400
Elk Park, Banners Elk Creek			3,200
Dutch Creek			75
Elk River			4,800
Flat Rock, Lake Anina			2,400
Forneys, Mill Creek			3,200
Franklin, Burningtown Creek			4,800
Ellijay Creek			3,200
Tesentee Creek			4,800
Goldsboro, Melton Pond			3,200
Hendersonville, Big Hungry Creek			4,000
Boylston Creek			4,000
Green River			4,800
Kanuga Lake			1,600
Laurel Creek			2,400
Kellerville, Buckeye Creek			50
Laurel Creek			75
Lake Toxaway, Lake Toxaway			64,800
Linville Falls, Caleb Creek			1,400
Cane Creek			1,400
Irish Creek			1,400
Katy Creek			1,400
Linville River			4,200
Magazine Creek			1,400
Marion, Allison Creek			1,400
Bill Creek			1,400
Bow Creek			4,000
Buffalo Creek			1,400
Burgin Creek			1,400
Camp Rock Creek			1,400
Cedar Creek			1,400
Cherry Creek			2,400
Chestnut Fork Creek			1,400
Cove Creek			1,400
Crooked Creek			1,400
Curtis Creek			1,400
Davidson Creek			1,400
Devils Fork Creek			1,400
Duncan Creek			2,100
Fall Branch			2,400
Ford Creek			2,400
Gladis Creek			1,400
Hall Creek			700
Harrar Creek			3,200
Harris Creek			1,400
Little Fork Creek			2,400
Little River			1,400
Mackey Creek			1,400
Maple Creek			1,400
Newberry Fork Creek			1,400
Paddy Fork Creek			1,400
Pigeon Roost Creek			1,400
Roaring Fork Creek			1,400
Sahadrec Creek			1,400
Singed Cat Creek			1,400
Six Mile Creek			2,400
Spring Creek			2,400
Stony Creek			3,200
Turkey Creek			1,400
Turkey Otter Creek			1,400
Vess Creek			2,400
Morrisville, Sorrell's pond			800
Sycamore Pond			3,200
Old Fort, Crooked Creek Pond			700
Otto, Tesenta Pond			3,200
Pineola, Linville River			450
Poplar, Poplar Creek			1,475
Relief, Lewis's pond			200
Sevier, Armstrong Creek			1,400
Back Creek			1,400
Ball Creek			2,800
Beaver Creek			1,400
Crib Creek			1,400

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

RAINBOW TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
North Carolina—Continued.			
Sevier, Dobson Creek.....			700
Dysart Creek.....			1,400
Indian Creek.....			1,400
Lime Kiln Creek.....			1,400
Nix Creek.....			1,400
North Fork Creek.....			1,400
Oil Mill Creek.....			1,400
Owens Creek.....			1,400
Rollins Creek.....			1,400
Steel Creek.....			1,400
Table Creek.....			1,400
York Creek.....			700
Swain, Oconalufly River.....			4,800
Sylva, Abs Creek.....			1,600
Chastain Creek.....			2,400
Cullowhee Creek.....			4,000
Johns Creek.....			2,400
Moses Creek.....			2,400
Mull Creek.....			3,200
Ruff Bitt Creek.....			3,200
Sugar Creek.....			2,400
Toecane, Big Rock Creek.....			75
Greasey Creek.....			75
Linn Creek.....			75
Tomotla, Peachtree Creek.....			3,200
Tryon, Poolet River.....			4,000
Vaughn Creek.....			4,000
Tuxedo, Green River.....			12,000
Pace Creek.....			3,200
Rock Creek.....			5,600
Vale, Cow Camp Creek.....			75
Willits, Scotts Creek.....			3,200
Winston-Salem, Nissen Park Pond.....			1,600
North Dakota:			
Braddock, Otter Creek.....			1,450
Edinburg, Park River, Middle Fork.....			2,000
Glen Ullin, Curlew Creek.....			1,000
Hebron, Knife River.....			500
Ohio:			
Akron, Adams's pond.....			2,000
Zanesville, Licking River.....			5,000
Oregon:			
Austin, Strawberry Lake.....		5,400	
Baker City, Burnt River, North Fork.....		6,000	
Deer Creek.....		3,000	
Downey Lake.....		3,000	
Eagle Creek.....		5,500	
Fish Lake.....		3,000	
Hilgard, Beaver Creek.....		3,000	
Five Points Creek.....		6,000	
Jordan Creek.....		3,000	
Meadow Brook.....		2,000	
Oregon City, Pine Creek.....		10,116	
Pennsylvania:			
Bainbridge, Engle Run.....			1,000
Hoffman Run.....			1,000
Stackstown Run.....			1,000
Benton, West Creek.....			5,000
Berlin, Blue Lick Creek.....			5,000
Brush Creek.....			4,000
Chambersburg, Birch Creek.....			6,000
Carbaugh Run.....			4,000
Hoosic Run.....			4,000
Cherry Tree, Cush Creek.....			4,000
Cherry Run, Penn Run.....			375
Clarendon, Arnots Run.....			3,000
Farensworth Creek.....			4,000
Four Mile Run.....			3,000
Tionesta Creek.....			3,000
Tionesta Creek, West Branch.....			5,000
Cresco, Bushkill River.....			2,400
Goose Run.....			1,500
Levis Branch.....			1,500
Ebensburg, Chest Creek.....			2,000
Frackville, Kaufman Dam.....			3,000
Glen Iron, Penns Run.....			1,125
Green Hill, Big Woods Pond.....			800

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

RAINBOW TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Pennsylvania—Continued.			
Lanesboro, Tunkhannock Creek.....			6,000
Lehighon, Wild Creek.....			3,000
Lenover, Weaver Run.....			2,000
Middleport, Morgan Dam.....			1,500
Millersburg, Forney Run.....			2,000
Little Wicanisco Creek.....			3,000
Norristown, Elmwood Park Lake.....			2,000
Paddy Mountain, Penns Run.....			2,250
Pardee, Penns Run.....			375
Ridgeway, Big Mill Creek.....			4,000
Rising Springs, Penns Creek.....			5,000
Somerfield, Youghiogheny Creek.....			7,000
Tunkhannock, Bowmans Creek.....			6,000
Welkert, Penns Run.....			375
South Carolina:			
Cleveland, Middle Saluda River.....			4,000
Greenville, South Saluda River.....			4,000
Rosman, Cane Creek.....			3,200
Estatoe Creek.....			4,000
South Dakota:			
Buffalo Gap, Beaver Creek.....			150
Cascade Springs, Cascade Springs.....			12,500
Custer, French Creek.....			5,775
Spring Creek.....			5,000
Deadwood, Polo Creek.....			8,000
Elmore, Spearfish Creek.....			5,325
Hermosa, Squaw Creek.....			150
Hill City, Newton Fork Creek.....			5,625
Palmer Creek.....			2,500
Slate Creek.....			2,500
Spring Creek.....			5,775
Sunday Creek.....			5,625
Hot Springs, Palmer Lake.....			12,500
Iron Creek, Spearfish Creek.....			4,500
Nahant, Tilson Creek.....			3,750
Pine Ridge Agency, Wolf Creek.....			12,500
Rapid City, Dark Canyon Pond.....			12,500
Rapid Creek.....			23,150
Slate Creek.....			150
Spring Creek.....			300
St. Onge, False Bottom Creek.....			600
Scenic, Conklin Lake.....			25,000
Snowina, Stearn's pond.....			500
Spearfish, Driskill's pond.....			500
Spearfish Creek.....			2,000
Sturgis, Deadmans Creek.....			10,000
Spring Creek.....			10,000
Tennessee:			
Bellevue, South Harpeth Creek.....			4,000
Blevins, Doe River.....			100
Bristol, Sinking Creek.....			125
Butler, Cable's pond.....			1,200
Lineback's pond.....			75
Spring Lake.....			800
Concord, Doughty's pond.....			800
Kirby's pond.....			800
Doyle Station, Sink Creek.....			1,600
Ducktown, Rough Creek.....			3,200
Dunn, Sugar Creek, West Fork.....			1,600
Elizabethtown, Hunter's Lake.....			100
Farner, Camp Creek.....			2,450
Fishery, North Indian Creek.....			2,185
Spring Branch.....			220
Fish Springs, Watauga River.....			4,800
Greenville, Camp Creek.....			4,000
Hampton, Laurel Creek.....			175
Hunter, Brush Creek.....			4,000
Johnson City, Brush Creek.....			3,200
Knoxville, Tennessee River.....			50
Marbleton, Garland's pond.....			1,000
Maryville, Mountain Pond.....			800
Oakdale, Emory Pond.....			125
Roan Mountain, Doe River.....			4,800
Hampton Creek.....			3,200
Heaton Creek.....			3,200
Rutledge, Manly's pond.....			1,600
Sievierville, Layman's pond.....			800

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

RAINBOW TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Tennessee—Continued.			
Sparta, Calf Killer Creek.....			4,000
Springfield, Red River.....			4,000
Telford, Bailey's pond.....			50
Tullahoma, Compton Creek.....			3,200
Walland, Hesser Creek.....			4,000
Little River.....			4,125
Wolf Creek, Wolf Creek.....			4,000
Utah:			
Charleston, applicant.....	125,000		
Murray, applicant.....	59,400		
Provo, Dry Creek Pond.....		6,000	
Provo River.....		48,000	
Virginia:			
Afton, Afton Pond.....			300
Ashland, South Anna River.....			8,000
Big Island, Hunting Creek.....			2,500
Cedar Bluff, Indian Creek.....			12,000
Cleveland, Bacon Creek.....			3,000
Big Cedar Creek.....			18,000
Burgess Creek.....			9,000
Gilmer Creek.....			3,000
Little Cedar Creek.....			12,000
Opossum Creek.....			6,000
Covington, Cedar Creek.....			4,000
Falling Springs Run.....			200
Culpeper, Hazel River.....			4,800
Miller Creek.....		7,000	
Faber, Cover Creek.....			6,400
Fairwood, Big Holton Creek.....			6,400
Marion, Holston River, South Fork.....			12,000
Mount Jackson, Garlic Hollow Run.....			800
Natural Bridge, Cedar Creek Dam.....			500
New Castle, Meadow Creek.....			2,000
Roanoke, Falling Creek Reservoir.....			3,200
Vinton Spring Lake.....			2,400
Rural Retreat, Buchanan's pond.....			2,400
Salem, Back Creek.....			8,000
Seven Mile Ford, Comer Creek.....			12,000
Holston River, South Fork.....			8,000
Somerset, Rapidan River.....			1,125
Springwood, Cummings's pond.....			300
Stanley, Henderson's pond.....			1,000
Sugar Grove, Holston River, South Fork.....			8,000
Waynesboro, Lithia Pond.....			300
West Point, Remick Hall Pond.....			3,000
Wytheville, Cove Creek.....			6,400
Washington:			
Colville, Black Lake.....			2,000
Colville River.....			3,000
Harrington, Crab Creek.....			4,000
Republic, Granite Creek.....			4,000
Seattle, Exposition Aquarium.....			18
Sumner, Salmon Creek Pond.....			1,000
Valley, Bond Lake.....			3,000
West Virginia:			
Blake, Loup Creek.....			1,500
Capon Springs, Trout Run.....			3,650
Yellow Stream Gap.....			3,650
Holly Junction, Elk River.....			750
Keyser, Patterson Creek.....			4,300
Marlinton, Elk River.....			2,500
Midvale, Middle Fork River.....			7,500
Rippon, Wiest's pond.....			1,000
Seebert, Cranberry Creek.....			38,500
Spring Creek, Sinking Creek.....			3,000
Stonewall, Piney Creek.....			21,000
Surveyor, Clay Pond.....			500
White Sulphur Springs, Howard Creek.....			3,000
Spring Branch.....			2,000
Wildell, Greenbrier River.....			5,000
Laurel Run.....			5,000
Wright, Piney Run.....			24,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

RAINBOW TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Wisconsin:			
Independence, Borst Valley Creek.....			3,000
Chimney Rock Creek.....			3,000
Cook Creek.....			1,200
Elk Creek.....			3,000
Fox Creek.....			1,500
Tamarack Creek.....			2,500
Traverse Valley Creek.....			3,000
Trempealeau River.....			3,000
Kendall, Lumsden Creek.....			1,200
Tunnell Creek.....			1,500
Sparta, Lower La Crosse River.....			3,000
Spring Valley, Eau Galle River.....			4,000
Wyoming:			
Beulah, Sand Creek.....			5,000
Cheyenne, Polaris Reservoir.....		12,000	
Lander, Glacier Lake.....			3,600
Lodge Pole Lake.....			2,000
Shoshone Lake.....			2,000
Laramie, Laramie River.....		7,000	
Moocroft, Riordan Lake.....			1,000
Sheridan, Patrick's reservoir.....			1,500
Wamsutter, Bens Lake.....			300
Wheatland, Development Company's reservoir.....			10,000
Yellowstone National Park, Rock Lake.....			10,000
Gibbon River.....			15,000
Japan:			
Tokio, Imperial Household Department.....	110,000		
Total	556,494	595,616	1,705,328

ATLANTIC SALMON.

District of Columbia:			
Washington, Central Station Aquarium.....			100
Maine:			
Brownville, Pleasant River.....			76,500
East Orland, Alamoosook Lake.....			5,139
Guilford, Piscataquis River.....			41,000
Milo, Pleasant River.....			33,000
Staceyville, Penobscot River.....	1,217,366		82,413
New York:			
Buffalo, New York State Cancer Laboratory.....			60
New York, New York Aquarium.....	5,000		
Total	5,000	1,217,366	288,212

LANDLOCKED SALMON.

Idaho:			
Hope, Lake Pend d'Oreille.....			4,000
Maine:			
Auburn, Lake Auburn.....			7,500
Taylor's pond.....	33,000		
Augusta, Cobboscocontee Lake.....			7,500
Baker, Baker's pond.....			2,000
Bingham, Rowe's pond.....			2,000
Brewer Junction Brewer Pond.....			2,751
Brownfield, Moose Pond.....	24,750		
Bryant Pond, Lake Christopher.....	16,500		
Twickell Pond.....	16,500		
Bucksport, Toddy Pond.....			6,000
Dedham, Branch Pond.....	30,000		5,000
Green Lake.....			15,000
Dover, Sebec Lake.....			10,500
East Orland, Alamoosook Lake.....			13
Ellsworth, Patten's pond.....	25,000		2,000
Ellsworth Falls, Alligator Lake.....			6,000
Beach Hill Pond.....	20,000		
Flood's pond.....	24,750		

α Lost in transit, 18,100 fry.

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LANDLOCKED SALMON—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Maine—Continued.			
Enfield, Cold Stream Pond.....			9,000
Farmington, Big Island Pond.....			4,500
Franklin, Donnell's pond.....		24,750	
George's pond.....		24,750	
Molasses Pond.....		24,750	
Green Lake, Arnold's pond.....			4,500
Grand Lake Stream, Dobsis Lake.....	65,000		4,500
Grand Lake.....	316,440		17,700
Holden, Fitz Pond.....	24,750		
Kennebunk, Kennebunk Pond.....	24,750		
Kinco Station, Moosehead Lake.....	32,000		10,500
Lincoln, Mattamawcook Lake.....	5,000		
Mosquito, Lake Moxie.....	18,000		
Newport, Lake Sebasticook.....			13,500
North Anson, Great Emden Lake.....	24,750		
Oquosoc, Rangeley Lakes.....			9,000
Otis, Green Lake.....	50,000		70,000
Peru, Worthley's pond.....	21,600		
Phillips Lake, Phillips Lake.....			6,000
Portage, Portage Lake.....	30,000		3,500
Sawyers Island, Campbell's pond.....			3,000
Sebago Lake, Sebago Lake.....	15,000		
Skowhegan, Lake George.....	24,750		
South Paris, Concord Pond.....			6,000
Strong, Sweet's pond.....			6,000
Thorndike, St. Georges Lake.....			7,500
Tunk Pond, Tunk Pond.....	24,750		6,000
Warren, Crawford's lake.....			6,000
Wescott, Little Ossipee Pond.....			5,700
Wilton, Wilson Lake.....		15,000	
Michigan:			
Munising, applicant.....	10,000		
Sault Ste. Marie, Michigan Fish Commission.....	20,000		
Montana:			
Gardner, Yellowstone Park waters.....			8,000
New York:			
Old Forge, applicant.....	15,000		
Forest, Fish, and Game Commission.....	15,000		
Pleasant Lake, Pleasant Lake.....		14,500	
Itaque Lake, Lake Kora.....	30,000		
Vermont:			
Averill, Averill Pond.....			1,000
Little Averill Lake.....		2,000	
Brandon, Lake Dunmore.....			2,500
Newport, Salem Pond.....		1,000	
Washington:			
Ephrata, Moses Lake.....			5,000
Wisconsin:			
Luck, McKenzie Lake.....			11,400
Wyoming:			
Lander, Christiana Lake.....			5,000
Grave Lake.....			5,000
Argentina:			
Buenos Aires, Argentine Government.....	25,000		
Total ^a	115,000	974,040	301,064

BLACKSPOTTED TROUT.

Arizona:			
Grand Canyon, Hull Pond.....			3,750
Little Hull Pond.....			3,750
Colorado:			
Antonito, Conejos River.....		19,440	
La Jara River.....		4,320	
Cardinal, Deyelin Lakes and Creek.....		9,500	
Cascade, Cascade Brook.....		10,000	
Cebolla, Elk Creek.....		10,000	
Gunnison River.....		25,796	
Red Creek.....		4,000	
Cimarron, Little Cimarron River.....		10,000	
Cliff, Platte River.....		4,800	
DeBeque, Bull Creek Lake.....		15,000	

^a Lost in transit, 11,000 fry and 2,300 fingerlings.

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BLACKSPOTTED TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Colorado—Continued.			
Denver, Colorado Fish Commission	225,000		
Dillon, Rock Creek		3,600	
Slate Creek		3,600	
Straight Creek		3,600	
Fort Collins, Cache la Poudre River		30,700	
Pine Creek		31,010	
Glenisle, Platte River		3,600	
Glenwood Springs, Mitchell Creek		10,000	
Grand Valley, Parachute Creek		10,000	
Gunnison, Bird Lakes		4,000	
Insmont, Rock Creek		2,400	
Loveland, Big Thompson		40,746	
Marshall, South Boulder Creek		14,400	
Molina, Cottonwood Creek		10,000	
Cottonwood Lakes		52,748	
East Bull Creek		10,000	
Monte Vista, Rock Creek, South Fork		6,000	
Montrose, Big Red Canyon Creek		8,000	
Spring Creek		6,000	
West Dry Creek		6,000	
Nast, Frying Pan River		10,500	
New Castle, Divide Creek		12,500	
Parlin, Quartz Creek		6,000	
Pine Grove, Elk Creek		4,800	
Ridgway, Cow Creek		12,000	
Dallas Creek		12,000	
Rifle, Williams River		22,000	
Salida, Arkansas River		22,500	
Little River		7,500	
Poncha Creek		10,000	
South Fork, Rio Grande River, South Fork		6,000	
Wheeler, West Tennesse Creek		8,400	
Idaho:			
Bonner County, Bonanza Lake			10,000
Darsey, Stevens Peak Lake			7,500
Greer, Wells Pond			2,500
McCammon, Mountainview Lake			3,000
Rupert, Lake Walcott			12,000
Soda Springs, Knollins Springs			3,000
Spirit Lake, Kit Carson Creek			5,000
Twin Falls, Blue Lake Creek	50,000		
Wallace, Lost Lake			7,500
Michigan:			
Detroit, Detroit Aquarium	10,000		
Montana:			
Anaconda, Montana Fish Commission	550,000		
Baker, Baker Lake			16,000
Ballantine, Arrow Creek			4,000
Belton, Lake McDonald			12,000
Big Timber, Big Boulder River			4,000
Bozeman, West Gallatin River, South Fork			5,000
Butte, Columbia Gardens Hatchery	440,000		
Chinook, Peoples Creek			8,000
Chinook, Snake Creek			8,000
Craig, Burke's reservoir			6,000
Darby, Tin Cup Lake			10,000
Dorsey, Cheekrboard Creek			6,000
Little Birch Creek			6,000
Woods Gulch Creek			6,000
Harlowton, Musselshell River			8,000
Havre, Clear Creek			6,000
Helena, Chessman Reservoir			8,000
Josephine, Sixteen Mile Creek			8,000
Kalispell, Corneilson's spring			6,000
Corneilson's lake			6,000
Howser's lake			6,000
Lewistown, Beaver Creek			6,000
Big Casino Creek			6,000
Big Spring Creek			14,000
Casino Creek			6,000
Cottonwood Creek			12,000
Surprenant's pond			12,000
Livingston, Fitzpatrick's pond			2,000
Trowbridge Creek			4,000
Martinsdale, Musselshell River, North Fork			8,000
Missoula, Bitter Root River			10,000
Monarch, Tillinghast Creek			6,000
Neihart, Belt Creek			6,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BLACKSPOTTED TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Montana—Continued.			
Red Lodge, Silver Run.....			2,000
Somers, Lake Alexander.....			6,000
Skag Lake.....			6,000
Townsend, Due Creek.....			6,000
Twodot, Haymaker Pond.....			6,000
Winston, Stanbach Reservoir.....			6,000
Nebraska:			
Chadron, Big Bordeaux Creek.....			12,000
Nevada:			
Derby, Nevada Fish Commission.....	298,300		
Truckee River.....		85,000	
Verdi, Bates's pond.....			3,000
Galena Creek.....			3,000
Nevada Fish Commission.....	123,800		
South Branch.....			3,000
Truckee River.....	16,450	633,020	6,000
Whites Creek.....			3,000
New Mexico:			
Cimarron, Cañon Bonito Creek.....			2,000
Cimarronciti Creek.....			2,000
Cimarron River.....			2,000
Clear Creek.....			2,000
Ponil Creek.....			2,000
Rayado Creek.....			2,000
Rayado River, West Fork.....			3,000
Ute Creek.....			4,000
Glorieta, Pecos River.....		14,400	
Las Vegas, Burro Branch.....		4,800	
Gallinas River.....		6,000	
Mountain Park, Frenal Creek.....			15,000
Sante Fe, Rio Testuque River.....		7,200	
New York:			
New York, New York Aquarium.....	25,000		
Saranac Inn, Forest, Fish, and Game Commission.....	50,000		
Oregon:			
Clackamas, Oregon fish commission.....			45
Medford, Four Bit Creek.....		12,000	
Rancharee Creek.....		12,000	
Rogue River.....		16,000	
Milwaukee, Lechler Lake.....		8,000	
Newberg, Walton's pond.....		14,214	
Oregon City, Clackamas River.....		20,000	
Portland, Oregon fish commission.....	175,000		
Pennsylvania:			
Pleasant Mount, Pennsylvania fish commission.....	50,000		
South Dakota:			
Aberdeen, Milwaukee Reservoir.....			10,000
Buffalo Gap, Beaver Creek.....			7,000
Custer, Flynn Creek.....			16,000
French Creek.....			6,000
Elmore, Spearfish Creek.....			35,000
Spearfish Creek, Southwest Branch.....			9,000
Englewood, White Wood Creek.....			30,000
Hermosa, Squaw Creek.....			5,000
Hill City, Castle Creek.....			30,000
Spring Creek.....			21,000
Hisega, Rapid Creek.....			35,000
Iron Creek, Spearfish River.....			8,000
Maitland, Fredbert Pond.....			5,000
Mystic, Rapid Creek.....			30,000
Rapid City, Electric Light Pond.....			12,500
North Side Park Pond.....			2,500
Price Pond.....			6,000
Rapid Creek.....			47,750
Slate Creek.....			5,000
Spring Creek.....			6,000
Saint Onge, False Bottom Creek.....			40,000
Spearfish, Spearfish Creek.....			25,000
Utah:			
Provo, applicant.....	50,000		
Provo River.....		20,000	
Virginia:			
Sweet Chalybeate, Sweet Springs Branch.....			2,480
Washington:			
Seattle, Exposition Aquarium.....	50,000		42
Spokane, Selheim Springs Pond.....			5,000
Walla Walla, Shelton's lake.....			737
Spring Creek.....			600
Winona, Palouse River.....			10,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BLACKSPOTTED TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Wyoming:			
Beulah, Crystal Springs.....			6,000
Crook County, Sand Creek.....			15,000
Yellowstone National Park, Cub Creek.....		400,000	
Lander, Grave Lake.....			11,200
Hobbs Lake.....			4,200
Raft Lake.....			5,600
Little Wind River, South Fork.....			5,600
Trail Lake.....			8,400
Laramie, Wyoming fish commission.....	175,000		
Moorcroft, Prairie Creek.....			21,250
Sheridan, Wyoming fish commission.....	500,000		
Shoshone, Big Wind Lake.....			15,000
Wamsutter, Stocks Lake.....			18,750
France:			
Bellefontaine, French Government.....	10,000		
Total^a.....	2,748,550	1,756,094	906,654

LOCH LEVEN TROUT.

South Dakota:			
Savoy, Little Spearfish Creek.....			68,248

LAKE TROUT.

Colorado:			
Twin Lakes, Upper Twin Lake.....		24,700	
Idaho:			
Hope, Lake Pend d'Oreille.....			18,000
Rathdrum, Twin Lake.....			4,000
Illinois:			
Havana, Illinois Fish Commission.....	500,000		
Maine:			
Bridgton, Highland Lake.....		11,000	
Cherryfield, Mopang Lake.....		11,000	
East Wilton, Pease Pond.....		11,000	
Green Lake, Green Lake.....		263,922	
North Anson, Great Emden Lake.....		11,000	
Readfield, Parker's pond.....		11,000	
Skowhegan, Lake George.....		10,000	
Unity, Unity Pond.....		11,000	
Massachusetts:			
Marlboro, Lake Williams.....		9,000	
Michigan:			
Big Rock Reef, Lake Michigan.....		756,000	
Cat Head Reef, Lake Michigan.....		756,000	
Charlevoix Reef, Lake Michigan.....		2,268,000	
Charlevoix, Pine Lake.....		756,000	
Detour, Lake Huron.....		2,000,000	
Detroit, Detroit Aquarium.....	10,000		
Escanaba, Lake Michigan.....		150,000	
Fishermans Island, Lake Michigan.....		1,512,000	
Fish Island, Lake Superior.....		600,000	
Grand Marais, Lake Superior.....		700,000	
Isle Royale, Lake Superior.....		1,975,000	2,052,500
Long Point, Lake Superior.....			600,000
McCargoes Cove, Lake Superior.....		275,000	
McLeods Channel, Lake Superior.....		1,025,000	
Mandan, Lake Medora.....			16,000
Manistique, Lake Michigan.....		150,000	
Marquette, Lake Superior.....		1,400,000	
Munising, Lake Superior.....		1,400,000	
North Point, Lake Huron.....		2,050,000	
North Point Reef, Lake Michigan.....		756,000	
Norwood Reef, Lake Michigan.....		756,000	
Ontonagon, Lake Superior.....		1,400,000	
Paris, Michigan Fish Commission.....	2,000,000		3,500
Petosky, Lake Michigan.....		756,000	
Point Iroquois, Whitefish Bay.....		700,000	
Sault Ste. Marie, Michigan Fish Commission.....	3,000,000		

^a Lost in transit, 9,740 fry.

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LAKE TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Michigan—Continued.			
Scarecrow Island, Lake Huron.....		1,950,000	
Seven Mile Point, Lake Michigan.....		756,000	
Skulligallee Reef, Lake Michigan.....		1,512,000	
Tobins Harbor, Lake Superior.....			780,000
Washington Harbor, Lake Superior.....			660,000
Whitefish Point, Lake Superior.....		2,000,000	
Minnesota:			
Grand Rapids, Pokegama Lake.....			20,000
Little Falls, Lake Alexander.....			20,000
Montana:			
Helena, Lake Sewell.....			6,900
New York:			
Auburn, Owasco Lake.....		40,000	
Charity Shoals, Lake Ontario.....		450,000	
Cooperstown, Otsego Lake.....		40,000	
Dutch Point, Lake Ontario.....		100,000	
Fox Island, Lake Ontario.....		1,000,000	
Fulton Chain, Little Moose and Panther Lakes.....		32,000	
Grenadier Island, Lake Ontario.....		1,627,000	
Hayes Point, Lake Ontario.....		750,000	
McKeever, Bisby Chain of Lakes.....		24,000	
Point Peninsula, Lake Ontario.....		450,000	
Raquette Lake, Lake Kora.....	150,000		
Riverside, Schroon Lake.....		40,000	
Wilson Bay, Lake Ontario.....		100,000	
North Dakota:			
St. John, Lake Lindeman.....			20,000
Oregon:			
Haines, Rock Creek Lake.....		11,300	
Pennsylvania:			
Waterford, Lake Leboeff.....		17,500	
Vermont:			
Averill, Big Averill Lake.....		30,000	
Barnet, Harvey's pond.....		35,000	
Barton, Silver Lake.....		17,500	
Stone Pond.....		17,500	
Brandon, Lake Dunmore.....			3,370
Hardwick, Elligo Pond.....		15,000	
Orleans, Willoughby Lake.....		35,000	
Readsboro, Howe's pond.....		14,000	
West Burke, Newark Pond.....		17,500	
Wisconsin:			
Brule, Twin Lakes.....			10,000
Crandon, Dry Lake.....			12,000
Metonga Lake.....			12,000
Stone Lake.....			12,000
Haugen, Monday Lake.....		16,000	
New Auburn, Wisconsin Fish Commission.....			3,880
Oshkosh, Wisconsin Fish Commission.....	4,500,000		
State Line, Black Oak Lake.....			32,000
Stone Lake, Little Stone Lake.....		10,000	
Sand Lake.....		12,000	
Stone Lake.....		12,000	
Argentina:			
Buenos Aires, Argentine Government.....	50,000		
Total.....	10,210,000	33,645,922	4,286,150

BROOK TROUT.

Arizona:			
Jerome, Beaver Creek.....			2,000
Dragoon Creek.....			2,000
Thompson Creek.....			2,000
West Fork Creek.....			2,000
Tucson, Sabino Creek.....			15,000
California:			
McCloud, Wheelers Creek.....		24,165	
Point Reyes, Paper Mill Creek.....	50,000		
Colorado:			
Antonito, Conejos River.....		20,000	
Basalt, Luna Creek.....		25,000	
Berrys Ranch, Eagle River.....			7,000
Black Hawk, Dory Lake.....		9,000	

^a Lost in transit, 4,000 fry.

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Colorado—Continued.			
Breckenridge, Crystal Lake.....		30,000	
Saw Mill Creek.....			4,500
Buena Vista, Cottonwood Creek.....		8,000	
Middle Cottonwood Creek.....		16,000	
South Cottonwood Creek.....		8,000	
Cebolla, Cebolla Creek.....			12,500
East Elk Creek.....			7,000
Cimarron, Cimarron River.....		35,000	
Silver Tip Lake.....		15,000	
Van Place Lake.....		15,000	
Colona, High Top Lake.....			5,100
Twin Lake.....			10,200
Wilson Lake.....			5,100
Colorado Springs, City Reservoir.....		30,000	
Glimmer Glass Lake.....		20,000	
Jimmy Camp Lake.....		27,500	
North Cheyenne Creek.....			6,000
Creede, Red Mountain Creek.....		10,000	
Rio Grande.....		10,000	
Sylvester's ponds.....		10,000	
Cripple Creek, Barnard Creek Pond.....			1,500
De Beque, Big Creek.....			8,500
West Bull Creek.....			6,800
Del Norte, Pinos River.....		10,000	
Delta, Alexander Lake.....		100,000	
Surface Creek.....		25,000	
Youngs Creek.....		100,000	
Denver, Crystal Springs Trout Hatchery.....		12,500	
Eldora, Lake Eldora.....		30,000	
Lake Kanawha.....		30,000	5,000
Frisco, Uneva Lake.....		40,000	
Georgetown, Green Lake.....		38,000	
Glenwood Springs, Hermitage Creek.....		25,000	
Mesa Creek.....		15,000	
Roaring Fork River.....		25,000	
Granby, East Inlet.....		12,000	
Grand Lake.....		24,000	
Grand River, North Fork.....		20,000	
Stillwater Creek.....		16,000	
Supply Creek.....		12,000	
Grand Junction, West Evacuation Creek.....		15,000	
Granger, Embargo Creek.....		12,500	
Graneros, Oak Lodge Ponds.....			3,000
Grant, Duck Lake.....		15,000	
Kirby Creek.....		15,000	
Gunnison, Bird Lakes.....			1,000
Hillside, Koch Branch.....		10,000	
Idaho Springs, Chinn Lake.....		15,000	
Edith Lake.....		50,000	
Saint Mary Lake.....		10,000	
Silver Lake.....		10,000	
Slater Lake.....		15,000	
Truesdale Creek.....		18,000	
Ivanhoe, Ivanhoe Creek.....		25,000	
Lyle Creek.....		15,000	
Jefferson, Rainbow Lake.....		15,000	
La Jara, Hamilton Ranch Pond.....		8,000	
La Jara River.....		19,950	
Pursley's pond.....		10,000	
Spring Creek.....		11,950	
Leadville, Arkansas River.....		39,000	
Austin's pond.....			40
Columbine Lake.....			2,000
Darrah's pond.....		20,000	
Half Moon Creek.....		24,000	
Lake Creek.....		24,000	
Laws Lake.....			20,000
Lower Twin Lakes.....		25,000	
Musgroves Pond.....		250,000	
Smith's ponds.....		20,000	
South Platte River.....		4,000	
Tennessee River.....		44,000	
Turquoise Lake.....		15,000	
Twin Lakes.....		25,000	
Upper Lake Creek.....		15,000	
Willow Creek.....		44,000	

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Colorado—Continued.			
Loveland, Big Thompson River, South Fork.....		30,000	
Big Thompson Pond.....		15,000	
Buckhorn Creek.....		15,000	
Lyons, Estes Park Hatchery.....	100,000		
Malta, Lake Creek.....		80,000	
Marshall, South Boulder Creek.....		30,000	
Minturn, Cross Creek.....			13,600
Eagle River.....			11,900
Gore Creek.....			10,200
Moffat, Artesia Pond.....			2,000
Monte Vista, Los Pinas Creek, Middle Fork.....		7,900	
Rock Creek.....		12,500	
South Fork Creek.....		16,000	
Montrose, Middle Spring Creek.....		10,000	
Spring Creek.....		15,000	
Nast, Frying Pan River.....		20,000	15,000
New Castle, Willow Creek.....			4,000
Norrie, Chapman Lake.....		15,000	
Olathe, Greys Creek.....		10,000	
Park Siding, South Platte River, North Fork.....		4,000	
Parlin, Quartz Creek.....			2,000
Parshall, Grand River.....		20,000	
Reads Lake.....		3,880	
Radium, Grand River.....		20,000	
Rico, Burnett Creek.....		10,000	
Ryman Creek.....		10,000	
Scotch Creek.....		15,000	
Ridgway, Dolores River.....		28,500	
Leopard Creek.....		15,000	
Rifle, Bear Creek.....			3,600
White River.....			1,800
Ruedi, Pond Creek.....		10,000	
Ruedi Lake.....		25,000	
Spearhead Lake.....			2,400
Salida, South Arkansas River.....		28,000	
Woodbridge Pond.....		40,000	
Sawpit, Sylvan Lake.....			6,700
South Fork, Beaver Creek.....		12,500	
Elk Creek.....		12,500	
Goupel Creek.....		12,500	
South Platte River.....			22,500
Trout Creek.....		12,500	
Steamboat Springs, Bear River.....		25,000	
Fish Creek.....		15,000	
Spring Creek.....		10,000	
Yampa River.....		15,000	
Texas Creek, Spruce Creek Reservoir.....			7,200
Thomasville, Spring Creek.....			2,400
Woods Lake.....		200,000	
Tolland, South Boulder Creek.....		23,000	
Trinidad, McWilliams Pond.....			4,000
South Lake.....			10,000
Twin Lakes, Lake Creek.....			5,000
Webster, Platte River.....			16,500
West Cliffe, De Weese Reservoir.....		98,000	
Venable Creek.....			10,800
Wheeler, Black Creek.....		15,000	
Wolcott, Eagle Creek.....			6,000
Wootton, Sugarite Creek.....			2,000
Connecticut:			
Botsford, Halfway River.....		12,000	
Danbury, Willow Brook.....			300
Greenwich, Byram River.....		8,000	
New Haven, Spring Glen Pond.....			300
Norwich, Billings Brook.....			400
Broad Brook.....			600
Choate Brook.....		7,500	
Pease Brook.....		7,500	
Stony Brook.....		7,500	
Saybrook Junction, Hart Brook.....			300
Stamford, Mill Creek.....		20,000	
Rippewan River.....		30,000	
Stratford, Brookdale Pond.....		12,000	
Tariffville, Three Cornered Pond.....		16,000	
Waterbury, Andrews Pond.....		32,000	
Hancock Pond.....		12,000	
Hop Brook.....		16,000	

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Connecticut—Continued.			
Waterbury, Long Hill Brook.....		16,000	
Osborne Brook.....		8,000	
Potatuck River.....		8,000	
Wilton, Norwalk River.....		23,000	
Delaware:			
Wilmington, Brandywine Creek.....			4,000
Georgia:			
Rabun Gap, Denton Creek.....			2,400
Young Harris, Brasstown Creek.....			4,000
Idaho:			
Bancroft, Eighteenmile Creek.....			1,800
Blackfoot, Tanner Spring Lakes.....			1,200
Bonnars, Spring Creek Pond.....			3,000
Buhl, Sand Spring Lake.....			1,000
Caldwell, Meyer Lake.....			900
Garner, Clifton Mill Pond.....			900
Hailey, Hartley Pond.....			900
Sheep Pond.....			900
Spring Creek.....			2,000
Hayden Lake, Hayden Lake.....			6,000
Jerome, Trail Springs.....			1,500
Kamiah, Little Duck Lake.....			2,000
Kingston, Pine Creek.....			6,000
Malad City, Waldon's pond.....			1,200
Montpelier, Mildred Pond.....			1,200
Naples, Fall Creek.....			4,500
Preston, Wilson Spring Pond.....			1,200
Rathdrum, Boeck Creek.....			1,500
Fish Lake Creek.....			2,000
Gilbert Creek.....			1,500
Lancaster Creek.....			2,000
Miller Creek.....			1,500
Rice Creek.....			1,500
Thorp Creek.....			2,000
Rexburg, Bell's pond.....			600
Illinois:			
Fox, Crystal Springs.....			300
Griggsville, Hatch Hollow Pond.....			300
Indiana:			
Angola, Clark Creek.....			1,950
Jackson Creek.....			2,000
Sauls Creek.....			1,950
Richmond, Henley Pond.....			1,000
St. Paul, Mill Creek.....			3,950
Iowa:			
McGregor, Bass Creek.....			6,000
Waukon, North Fork Creek.....			6,000
Patterson Creek.....			7,500
Kentucky:			
Compton Junction, Chimney Top Creek.....			10,000
Maine:			
Alfred, Nutter Brook.....			500
Annabessacook, Wilson Lake.....	30,000		
Belfast, Swan Lake.....	30,000		
Biddeford, Buzzell Brook.....	20,000		
Cold Spring Brook.....	15,000		
Runnells Brook.....	20,000		
Bingham, Pleasant Pond.....			1,800
Rowe Ponds.....		21,500	1,500
Bluehill, Woods Pond.....		25,000	
Brooks, Passachunkeag Pond.....		30,000	
Bryants Pond, Lake Christopher.....			1,500
Camden, Canaan Lake.....		30,000	1,500
Dedham, Green Lake.....		80,000	
Deering Junction, Bodge Brook.....		15,000	600
Machigonne Creek.....		15,000	750
Woodland Hatchery.....	25,000		
East Orland, Toddy Pond.....		21,000	
Ellsworth, Billings Pond.....		35,000	
Branch Pond.....		50,000	
Ellsworth Falls, Beach Hill Pond.....		20,000	
Floods Pond.....		25,000	
Long Pond.....		37,500	
Farmington, Beedy Brook.....			900
Big Island Pond.....			1,500
Cattle Brook.....			600
Chace Pond.....			1,500

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Maine—Continued.			
Farmington, Chain of Ponds.....			3,000
Dead River Pond.....			1,500
Grant Pond.....			1,500
Gull Pond.....			1,600
Lufkin Pond.....			1,500
Mt. Blue Pond.....			3,000
Redington Creek.....			1,500
Sandy River.....			1,500
Tufts Pond.....			1,400
Green Lake, Ducktail Pond.....	20,000		
Partridge Pond.....	25,000		
Snowshoe Pond.....	15,000		
Greenville Junction, Moosehead Lake.....			1,500
Harrington, Schoodic Lake.....	35,000		
Holeb, Little Pond.....			1,500
Jackman, Hatchery Brook.....	15,000		
Supply Pond.....	15,000		1,500
Thompson Brook.....	15,000		
Katahdin Iron Works, Big Houston Pond.....			1,500
Little Houston Pond.....	35,000		3,500
Kineo, Cany Creek.....	30,000		
Moosehead Lake.....	37,500		4,500
Lincoln, Long Pond.....	20,000		
Livermore Falls, Long Pond.....			1,500
Lowelltown, Bog Brook.....	12,500		
Deer Pond.....	12,500		
Lowell Pond.....	12,500		
Machias, Bog Lake.....	30,000		
Monmouth, Baker Pond.....			1,500
Jimmy Pond.....			1,500
Mosquito, Baker Pond.....	10,000		
Onawa, Upper Boarstone Pond.....	15,000		
Oquosoc, Rangeley Lakes.....			2,250
Otis, Green Lake.....	100,000		
Oxford, Hall Pond.....			1,200
Perry, Boyden Lake.....	40,000		
Phillips, Carlton Pond.....	37,500		
Phillips Lake, Phillips Lake.....	40,000		
Portage, Portage Lake.....			2,100
Rumford Falls, Howard Pond.....			1,500
Sedgwick, Thurston Brook.....			600
South Paris, Penneessevassee Lake.....	17,500		
Shagg Pond.....			1,500
Washburn Pond.....	15,000		
Tunk Pond, Tunk Pond.....			1,500
Unity, Sandy Creek.....	30,000		
West Ellsworth, Pattens Pond.....	25,000		
West Paris, Abbot Pond.....			1,200
Little Concord Pond.....			1,500
Washburn Pond.....			600
Wilton, Webb Pond.....	17,500		
York Beach, Otter Pond.....			450
Maryland:			
Annapolis, Alcorn Branch.....			1,000
Bel Air, Barnes Run.....			2,000
Cool Spring Run.....			1,000
Durham's brook.....			500
Elbow Brook.....			1,000
Flint Mill Brook.....			1,000
Graveyard Brook.....			1,000
Hollands Brook.....			1,000
Johnson's brook.....			1,000
Stoner Creek.....			1,000
Wysong Brook.....			500
Deer Park, Altamont Pond.....			500
Block Run.....			400
Pond Run.....			500
Trout Run.....			800
Elkridge, Stony Run.....			1,000
Fallston, South Fork Brook.....			1,000
Glyndon, Lake Jorosa.....			500
Hagerstown, Marsh Run.....			1,000
Mill Spring Run.....			500
Highland, Heaps Brook.....			500
Minefield Brook.....			1,000
Ramsey Brook.....			1,000
Hutton, Crystal Lake.....			2,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Maryland—Continued.			
Landover, Eccles Pond.....			500
Monkton, Curtis Brook.....			1,000
Matthews Branch.....			500
Patterson Brook.....			500
Phelps and Reynolds Branch.....			500
Mountain Lake Park, Pine Run.....			500
New Freedom, Ruhls Branch.....			1,000
Oakland, Cherry Creek.....			1,500
Deep Creek.....			2,200
Dunker Lick Creek.....			1,800
Hamill's lake.....			1,000
Harrington Creek.....			2,300
Harvey's pond.....			320
Millers Run.....			1,800
Wilson's Lake.....			500
Rockland Station, Green Springs Run.....			1,000
Ruxton, Rockland Creek.....			1,000
Sharon, Magnes Brook.....			500
Smithsburg, Oswald Run.....			500
Silver Falls Creek.....			500
Warner Gap Run.....			500
Stoyer, Sand Run.....			400
Thurmont, Hunting Creek.....			1,500
Westminster, Fairview Pond.....			500
Wilson, Laurel Run.....			1,500
Massachusetts:			
Athol, Swift River.....		20,000	
Clinton, Nashua River.....			600
Concord, Punkatasset Pond.....		16,000	
Fitchburg, Lord Brook.....			600
Mulpus Brook.....			900
Greenfield, Fisk Pond.....			500
Groton, Hunkerty Brook.....			600
Holyoke, Man Han River.....			700
Williamsett Brook.....			300
Lawrence, Schubert's pond.....		4,000	
North Adams, Hoosac River, North Branch.....			500
Hudson Brook.....			500
Northampton, Running Gutter Creek.....			700
South Hanson, Poors Creek.....		12,000	
Tolland, Slocum Brook.....			1,200
Waltham, Pequod Brook.....		8,000	
School House Brook.....		8,000	
Westfield, Big Powder Mill Brook.....			500
Farmington River, East Branch.....			1,400
Little River.....			700
Powder Mill Brook.....			500
Weston, Draper Brook.....			300
West Townsend, Allison's pond.....			180
Williamsburg, Clary Pond.....			300
Highland Brook.....			300
Michigan:			
Addison, Posy Creek.....			3,000
Alger, Bear Creek.....		5,000	
Wells Creek.....		10,000	
Alpena, Davis Creek.....		12,000	
Newton Creek.....		9,000	
Watson Creek.....		9,000	
Widner Creek.....		12,000	
Baldwin, Baldwin Creek.....		15,000	
Battle Creek, Sevenmile Brook.....			3,000
Bellaire, Shanty Creek.....			3,000
Biteley, Marquette River.....			3,000
Branch, Weldon Creek.....		10,000	
Brighton, Ore Creek.....		12,000	
Calumet, Eagle Creek.....			6,000
Mosquito Creek.....			4,000
Central Lake, Central Lake Brooks.....			3,000
Clare, Tobacco River, North Branch.....		18,000	
East Tawas, Vaughn Creek.....			1,000
Gladwin, Cedar River.....		15,000	
Smith Creek.....		10,000	
Grand Marais, Grand Marais Creek.....			10,000
Greenville, Berridges Creek.....			2,000
Hale, Hale Creek.....		9,000	
Smith Creek.....		9,000	

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Michigan—Continued.			
Hillsdale, Kirby Brook.....			3,000
Holland, Half Way Creek.....		6,000	
Interlochen, Betsie River.....			3,000
Kalamazoo, Haden Brook.....		15,000	
Silver Creek.....		12,000	
Kingsley, Boardman River.....			2,000
East Creek.....			2,000
Mayfield Brook.....			2,000
Little Manistee, Little Manistee River.....		20,000	
Lovells, Au Sable River, North Branch.....		25,000	
Big Creek.....		20,000	
Crapo Creek.....		10,000	
Mandan, Montreal River.....			6,000
Millersburg, Indian Creek.....		12,000	
Little Ocqueoc River.....		15,000	
Ocqueoc River.....		15,000	
Muskegon, Cedar Creek.....		12,000	
Silver Creek.....		9,000	
Newaygo, Bigton Creek.....		12,000	
Northville, Townsend Creek.....		10,000	
Peacock, Au Sable River.....			3,000
Manistee River.....			10,000
Petersburg, Crystal Pond.....			6,000
Phoenix, Gratiot River.....			6,000
Roscommon, Barnes Creek.....		5,000	
Beaver Creek.....		5,000	
Cedar Creek.....		5,000	
Cold Creek.....		15,000	
Durant Creek.....		10,000	
Willow Creek.....		5,000	
Standish, Lundy Creek.....			6,000
Sweetwater, Sweetwater Creek.....			4,000
White Cloud, White River.....			4,000
Wingleton, Bowman Creek.....			4,000
Cedar Creek.....			4,000
Danahar Creek.....		15,000	
Minnesota:			
Alborn, Ericsson Creek.....			600
Beaver Crossing, Beaver Creek.....			10,000
Budd Creek.....			4,000
Little Split Rock River.....			4,000
Split Rock River.....			9,200
Split Rock River, East Branch.....			6,000
Canton, Weisel Creek.....			5,300
Carlton, Otter Creek.....			10,000
Cloquet, Otter Creek.....			6,000
Squaw Creek.....			6,000
Deephaven, Jennison Creek.....			900
Kokesh Creek.....			2,000
Duluth, Endion Brook.....			12,000
Lester Creek, East Branch.....			6,000
Temperance River.....			1,200
Fond du Lac, Mission Creek.....			4,000
Fosston, Poplar Lake.....			10,000
Hibbing, O'Brien Brook.....			800
Hovland, Upper Brule River.....			7,500
Knife River, Miemac Lake.....			10,000
Mountain Brook.....			6,000
Nigadoo Brook.....			4,000
Lewiston, Enterprise Creek.....			2,000
Gunther Valley Creek.....			600
Hemmingway Creek.....			2,400
Laufenbergs Valley Creek.....			400
Pine Creek.....			2,000
Rush Creek.....			2,400
Stockton Valley Creek.....			2,000
Whitestone Creek, Middle Branch.....			600
Whitewater Creek, South Branch.....			2,800
Little Falls, Hillman Creek.....			10,000
Okesippi Creek.....			8,000
Skunk Creek.....			10,000
Minnesota City, Bear Creek.....			2,000
Rollingstone Creek, North Branch.....			2,000
Rollingstone Creek, Rupprecht Valley Branch.....			2,000
Preston, Bear Creek.....			2,000
Camp Creek.....			2,000
Forestville Creek, North Branch.....			1,000
Forestville Creek, South Branch.....			2,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Minnesota—Continued.			
Preston, Partridge Creek			1,000
Sugar Creek			1,500
Watson Creek			2,000
Redwood, Schmidts Creek			400
Rochester, Bear Creek			1,000
Rollins Siding, Bates Creek			4,000
Pine Creek			4,000
Rushford, Big Spring Creek			1,000
Camp Creek			1,000
Choice Creek			1,000
Coolidge Creek			1,000
Dalleys Creek			1,000
Diamond Creek			1,000
Emsend Creek			1,000
Enterprise Creek			1,000
Ferguson Creek			1,000
Gribbin Creek			1,000
Hemingway Creek			1,000
Iverson Creek			1,000
Jansens Creek			1,000
Johnson Creek			1,000
Meade Creek			1,000
Onstine Creek			1,000
Opheim Creek			1,000
Overland Creek			1,000
Paterson Creek			1,000
Pine Creek			1,000
Tangen Creek			1,000
Voagen Creek			1,000
Wilson Creek			1,000
Wisoy Creek			1,000
Saginaw, Demsey Creek			4,000
St. Charles, Campbells Spring Branch			1,000
Carters Run			1,500
Crows Creek			1,500
Drakes Creek			1,000
Fays Run			1,000
Logan Branch			400
Nichols Spring Branch			400
Pine Creek			2,000
Trout Run			2,000
Whitewater River			6,000
Savage, Nine Mile Creek			4,500
Two Harbors, Encampment River			800
Winona, Big Pickwick Creek			400
Cedar Creek			1,400
Corey Valley Creek			1,000
Dabelstein's ponds			800
East Burns Valley Creek			400
Ferguson Creek			400
Gilmore Valley Creek			1,000
Harvey Valley Creek			1,000
Hicks Valley Creek			1,600
Laufenberger Creek			1,000
Little Pickwick Creek			600
Marey Creek			1,000
Middle Valley Creek			400
Nunny Coulee Creek			600
Pine Creek			600
Pleasant Valley Creek			1,000
Rollingstone Creek			1,050
Rupprecht Valley Creek			600
Speltz Valley Creek			1,400
Straight Valley Creek			1,000
West Bruce Valley Creek			2,000
West Burns Valley Creek			400
Wisoy Creek			1,000
Missouri:			
St. Joseph, Missouri Fish Commission	100,000		
Montana:			
Alder, Moran Pond			1,200
Anaconda, Warm Springs Creek			2,800
Warm Springs Pond			1,600
Basin, Cataract Creek			22,500
Belt, Little Belt Creek			3,500
Belton, Fish Creek			2,000
Big Timber, Big Timber Creek			12,000
Boulder, Buffalo Creek			2,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Montana—Continued.			
Bozeman, Beaver Creek			4,000
Bridger Creek			36,000
Kelly Creek			2,000
Butte, Cauty's pond			2,000
Nez Perce Pond			2,000
White's lake			6,000
Chinook, Clear Creek			3,500
Columbus, Deep Creek			15,000
East Rosebud Creek			18,000
Fishtail Creek			1,500
Little Rosebud Creek			1,500
Skeleton Creek Pond			2,000
Spring Creek			2,000
Stillwater River			2,000
Crabtree, Spring Creek			1,500
Deer Lodge, Dog Creek			3,500
Dillon, Carter Creek			1,800
Landons Creek			600
Murray Spring Creek			4,000
Poindexter Creek			1,800
Dodson, Lodge Pole Creek			5,500
Emigrant, Dailey Lake			2,000
Helena, Papoose Creek			7,500
Hobson, Crescent Pond			2,000
Galbreath Coulee Lake			2,000
Lennep, Comb Creek			9,000
Lewistown, Arnell Creek			4,000
Box Elder Creek			4,000
Flat Willow Creek			3,000
Lima, Little Sheep Creek			1,500
Livingston, Holiday Spring Creek			9,000
Moore, Jones Spring			3,000
Sheridan, Branham Lake			2,800
Straw, East Buffalo Creek			9,000
Toston, Spring Creek Lake			7,500
Victor, Bear Creek			5,000
Big Creek			5,000
Sweathouse Creek			5,000
White Pine, Little Beaver Creek			2,000
Spring Lake			1,500
Winston, Staubach Creek			4,000
Nebraska:			
Chadron, Bordeaux Creek			15,000
Dead Horse Creek			30,000
Creighton, Bayile Creek			600
Nevada:			
Reno, Truckee River			3,600
New Hampshire:			
Ashland, Squam Lake		16,000	
Berlin, Chickwelnepy Creek		30,000	
Munn Pond		40,000	
Success Pond		40,000	
Bradford, Mountain Brook		12,000	
Campton, Bec Bee River		20,000	
Charlestown, Benware Brook			1,500
Hassom Brook			1,500
Mill Brook			1,500
Concord, Black Brook		8,000	
Bon Bog Brook		12,000	
Bow Brook Pond		8,000	
Bridge Brook		4,000	
Brown Brook		8,000	
Bumfogen Brook		16,000	
Deer Meadow Brook		8,000	
Monument Brook		8,000	
Pickard Brook		8,000	
Pine Island		12,000	
Trap Brook		12,000	
Enfield, Lovejoy Brook		12,000	
Epsom, Mountain Brook		8,000	
Exeter, Meadow Brook			180
Grafton, Wildmeadow Pond			220
Greenville, Shattuck Brook			180
Halcyon, Tilton Brook			
Keene, Alstead Brook		6,000	
Ashuelot River, East Branch		16,000	
		20,000	

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
New Hampshire—Continued.			
Laconia, Follett Brook		6,000	
Gilford Brook		12,000	
Lebanon, Cranberry Pond		8,000	
Lisbon, Star Crescent Pond		20,000	
Madison, Silver Lake		5,000	
Manchester, Dalton Brook			180
Manter Brook		12,000	
Nigger Creek		8,000	
Prescott Brook			180
Nashua, Budro Brook		6,000	
Chase Brook			180
Cider Mill Brook		8,000	
Gibson Brook		12,000	
Newbury, Lake Sunapee			6,000
New London, Barber Brook		6,000	
Newport, Cutts Brook			1,000
Penacook, Brickyard Brook		6,000	
Tannery Brook		6,000	
Peterboro, Nay Brook		12,000	
Pike, Eastman Brook			500
Plymouth, Little Glen Ponds		48,000	
Portsmouth, Marston Brook		8,000	
Peverly Brook			250
Potter Place, Fellows Meadow Brook		6,000	
Raymond, Fordway Brook			180
Jose Dudley Brook			180
Pine Hill Brook			180
Scribner Brook			180
Sanbornville, Pike Brook		16,000	
South Brookline, Rockwood Pond		8,000	
South Lyndeboro, Herrick Brook			180
Warner, Meadow Mills Creek		8,000	
Stevens Hill Creek		12,000	
Wentworth, Baker River		8,000	
Wilton, Miller Brook		12,000	
Purgatory Brook		16,000	
Stony Brook		12,000	
Winchester, Willard Pond			1,000
Wolfboro, Haith Brook		12,000	
New Jersey:			
Elberon, Whalepond Brook			1,000
Passaic, McDaniels Brook			1,000
Pattensburg, Manunselocwa Creek			1,000
Pompton Lakes, Haycock Brook			1,000
Princeton, applicant	1,000		
Salem, Collins Run			1,500
Cool Run			1,500
Easter Run			1,500
South Ogdenburg, Kinney Brook			500
Sparta, Pullis Stream			500
Sherman Mine Brook			500
New Mexico:			
Alamogordo, Spring Canon Pond			5,000
Glorieta, El Rcto de la Arrhaw			2,000
Las Vegas, Sapello River			4,000
Santa Fe, Rio Grande Live Stock Co.'s lake			2,000
Rio del Medio Creek			3,200
Santa Fe River			2,000
Tesuque Creek			2,000
Silver City, Glenwood Pond			4,000
Glenwood Springs			5,000
Wagon Mound, Tison Creek			1,600
New York:			
Adams, South Sandy Creek		24,000	
Afton, Cady Creek			500
Cornell Creek			500
North Afton Brook			1,000
Pixly Brook			1,000
Altmar, Beaver Dam Brook		12,000	
Potts Mill Brook		16,000	
Salmon River		24,000	
Apulia Station, Cascade Brook			1,500
Cold Brook			500
Conklin Brook			1,000
Dodge Brook			1,500
Gallinger Brook			600
Gleason Brook			1,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
New York—Continued.			
Apulia Station, Grady Brook.....			600
Johnson Brook.....			6,000
June Brook.....			1,500
Keeler Brook.....			1,000
Lee Brook.....			1,000
Newman Brook.....			1,000
Osborne Brook.....			1,000
Auburn, North Brook.....		20,000	
Salmon Brook.....		24,000	
Sennett Brook.....		20,000	
Barneveld, Big Drumlin Pond.....		12,000	
Beaver River, Beaver River.....		12,000	
Twitchell Creek.....		24,000	
Bellport, Osborne Creek.....			500
Berlin, Little Hoosick River.....		16,000	
Bliss, Wiscoy Creek.....		16,000	
Wiscoy Creek, North Branch.....		8,000	
Blossvale, Fish Creek.....		20,000	
Brainard, Black Brook.....		8,000	
Budlong Brook.....		6,000	
Buffalo, New York State Cancer Laboratory.....			250
Cambridge, Blair Brook.....		12,000	
Pammanook Creek.....		8,000	
Rice Brook.....		8,000	
Canton, Baldwin Brook.....		8,000	
Buck Brook.....		8,000	
Clark Brook.....		8,000	
Dean Brook.....		8,000	
Giffin Brook.....		8,000	
Granis Brook.....		6,000	
Howard Brook.....		8,000	
Leonard Brook.....		16,000	
Little River.....		16,000	
McFadden Brook.....		12,000	
Pleasant Brook.....		8,000	
Taylor Brook.....		12,000	
Cattaraugus, Cattaraugus Creek, West Branch.....		12,000	
Central Bridge, Grosvenor Pond.....			500
Cincinnati, Brakel Creek.....			1,500
Cooperstown, Iroquois Farm Ponds.....			600
Corinth, Sturdevan Brook.....		12,000	
Cornwell, Mineral Spring Creek.....			1,000
Dryden, Virgil Creek.....			1,500
Edmeston, Wharton Creek.....			2,000
Floodwood, Ledge Pond.....		24,000	
Georgetown Station, Gladding Brook.....			500
Mann Brook.....			1,000
Mariposa Creek.....			1,000
Middletown Creek.....			1,000
Plank Creek.....			600
Thompson Brook.....			600
Greene, Crandall Brook.....			1,000
Highland Falls, Queensboro Creek.....			1,500
Hoosick Falls, Case Brook.....		8,000	
Shingle Hollow Creek.....		12,000	
White Creek.....		16,000	
Iona Island, Doodletown Brook.....			1,000
Livingston Manor, Beaverkill River.....		10,000	
Elnore Lake.....		7,500	
Mahopac, Hillsboro Lake.....			2,500
Marathon, Hunts Creek.....			1,000
Merrills Creek.....			1,500
Newark, Military Brook Pond.....		8,000	
New Lebanon, Burnmead Brook.....		6,000	
Church Brook.....		6,000	
Cold Spring Brook.....		6,000	
Gillett Brook.....		8,000	
Hosmer Brook.....		8,000	
Hull Brook.....		8,000	150
Lost Brook.....		8,000	
Mahar Brook.....		6,000	
Meadow Brook.....		8,000	
Meander Brook.....		4,000	
Parker Brook.....		8,000	
Queechy Road Brook.....		8,000	
Shaker Mill Brook.....		16,000	150
Thomas Brook.....			150

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
New York—Continued.			
New Lebanon, Tilden Brook			150
West Meadow Brook		8,000	
Wyononock Creek			500
New York, New York Aquarium	10,000		
Northville, Barkers Stream		16,000	
Onativia, Hiscock Brook			1,000
Kennellys Brook			1,000
Morgan Brook			2,000
Montgomery Brook			1,000
Oneonta, Butternut Creek			2,500
Otsego Creek			1,500
Oulcous Creek			2,000
Otego, Otsdawa Creek			1,000
Paul Smiths, Lower St. Regis Lake		18,000	
Patterson, Croton River			2,500
Quaker Brook			2,500
Prospect, Big Rock Lake		24,000	
Randolph, Little Conewango Creek		16,000	
Rome, Canada Creek		16,000	
Point Rock Creek		16,000	
Roscoe, Abewood Brook		5,000	
Appley Brook		5,000	
Beaverkill River		7,500	
Berry Brook		6,000	
Darbee Brook		5,000	
Shin Brook		5,000	
Stewart Brook		5,000	
Tennanah Lake		10,000	
Willowemoc River		13,500	
Salamanca, Stoddards Pond		8,000	
Saugerties, Dwaskill Creek			2,000
Swartzwood, Jackson Hollow Creek			1,800
Syracuse, Carpenter Brook		16,000	
De Montforde Creek		8,000	
Thurman, Millington Brook		8,000	6,000
Veli Pond		20,000	
Valley Stream, Trout Lake			1,000
Watertown, French Creek		4,000	
Kings Creek		4,000	
Knapp Creek		6,000	
Waterville, Oriskany Creek		8,000	
Townsend Creek		10,000	
Williamstown, Carterville Pond		24,000	
Willsboro, Warm Pond		24,000	
North Carolina:			
Addie, Scotts Creek			3,200
Apalachia, Cane Creek			4,800
Sular Creek			4,000
Balsam, Dark Ridge Creek			1,600
Woodfin Creek			1,600
Black Mountain, Long Branch Creek			500
Middle Fork Creek			1,000
Montreat Lake			2,500
Silver Fork			2,000
Sugar Creek			1,000
Swannanoa River, North Fork			2,000
Boonford, Ayles Creek			1,500
Cane River, Elk Fork			1,000
Brevard, Middlesex Branch			4,000
Craggy, Wells's pond			800
Dillsboro, Brushyfork Creek			1,000
Elk Park, Elk River			2,400
Hickory Creek			2,400
Winkler Creek			6,400
Glenwood, Goose Creek			500
Mashburn Creek			500
Greenlee, Bear Creek			1,000
Bobs Fork Creek			1,000
Graybeard Creek			1,000
Greenlee Fork Creek			1,000
Haw Branch			1,000
Huskins Creek			1,000
Jarretts Creek			1,000
Little Shoals Creek			1,000
Logan Creek			500
Lone Fork Creek			500
Mountain Creek			1,000
Nahlets Creek			500

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
North Carolina—Continued.			
Greenlee, Pool Creek.....			500
Rock House Creek.....			500
She Bear Creek.....			1,000
Simmons Creek.....			1,000
Still House Creek.....			500
Teamster Creek.....			500
Thompson Fork Creek.....			1,000
Wild Cat Falls Creek.....			1,000
Wolf Creek.....			1,000
Hendersonville, Foley Creek.....			3,200
Kellerville, Beech Creek.....			14,000
Buckeye Creek.....			10,000
Linville Falls, Catawba River, North Fork.....			2,000
Green Mountain Branch.....			500
North Cove Creek.....			1,500
Pine Branch.....			500
Marion, Bee Rock Creek.....			1,000
Chalk Brook.....			500
Fourmile Creek.....			1,000
Garden Creek.....			1,000
Georges Creek.....			1,000
Greasey Creek.....			500
Honeycutte Creek.....			1,000
Jake Creek.....			1,000
Linnekin Creek.....			1,000
Little Buck Creek.....			1,000
Lost Cove Creek.....			1,000
Mill Creek.....			1,000
Osborne Creek.....			1,000
Paxton Creek.....			1,000
Rag Creek.....			1,000
Stott Creek.....			500
Minneapolis, Little Horse Creek.....			1,600
Montezuma, Deep Gap Branch.....			2,400
Emmonds Creek.....			2,400
Kawana Lake.....			4,000
Linville River.....			4,800
Stepup Branch.....			1,600
West Fork Creek.....			3,200
Penland, Brush Creek.....			1,000
Penrose, Brier Creek.....			2,400
Crab Creek.....			3,200
Grassy Creek.....			2,400
Laurel Creek.....			2,400
Little River.....			3,200
Reasonover Creek.....			2,400
Shoal Creek.....			2,400
Staghorn Creek.....			2,400
Racford, Pasture Branch.....			1,600
Toecane, Cane Creek.....			500
Club Creek.....			500
Hine Creek.....			500
Tomotla, Coloards Creek.....			3,200
Hayes Mill Creek.....			3,200
Waynesville, Balsam Spring Branch.....			1,600
Bennett Branch.....			1,600
Brindle Creek.....			1,600
Caldwell Fork Creek.....			1,600
Catatuchee Creek.....			3,200
Francis Branch.....			3,200
Hemlock Pond.....			3,200
Hyatts Branch.....			1,600
Indian Creek.....			1,600
Jaynes Branch.....			1,600
Jonathan Creek.....			3,200
Locust Grove Run.....			1,600
Long Branch.....			1,600
Loves Branch.....			1,600
Nick Creek.....			1,600
Pigeon River, Grassy Fork.....			1,600
Pigeon River, Middle Fork.....			1,600
Shelton Cove Creek.....			1,600
Ugly Creek.....			1,600
Whittier, Conley Creek.....			1,600

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Ohio:			
Bellefontaine, Macochee Creek.....			4,000
Spring Branch.....			3,000
Cleveland, Canyon Spring.....			2,000
Mansfield, Golf Spring Run.....			3,000
Mercer Creek.....			4,000
Mercer Lake.....			2,000
Niles Run.....			3,000
Reynolds Run.....			3,000
Ravena, Spring Creek.....			3,000
Urbana, Powells Brook.....			3,000
Oklahoma:			
Carrier, Spring Bark Creek.....			600
Weatherford, Deer Creek.....			400
Oregon:			
Baker City, Daly Creek.....		5,000	
Duncan, Meacham Creek.....		4,000	
Gibbon, Umatilla River.....		4,000	
Hilgard, Spring Creek.....		3,000	
Milwaukee, Crystal Lake.....		15,000	
Oregon City, Abernethy River.....		10,000	
Clear Creek.....		5,000	
Rock Creek Pond.....		9,000	
Woodcock River.....		9,800	
Pennsylvania:			
Allentown, Cedar Creek.....			3,000
Altoona, Big Laurel Run.....			500
Burgoon Run.....			500
Chondrius Run.....			500
Demmaree Run.....			500
Figarts Run.....			500
Green Springs Run.....			500
Juniata Gap Run.....			500
Laurel Run.....			500
Mill Run.....			500
Neb Run.....			500
Sandy Run.....			500
Arcadia, Powell's pond.....			500
Auburn, Bear Creek.....			1,200
Gold Mine Creek.....			1,000
Stony Creek.....			2,500
Austin, Bailey Run.....			1,000
Bark Shanty Run.....			500
Big Moores Run.....			1,000
Birch Run.....			1,000
Berg Run.....			1,000
Cowley Run.....			1,000
Darwin Run.....			1,000
East Fork Creek.....			1,000
Freeman Run.....			1,000
Hammersley Run.....			1,000
Jones Run.....			1,000
Little Nelson Run.....			1,000
Nelson Run.....			1,000
Portage Creek.....			1,000
Prouty Run.....			1,000
South Fork Run.....			1,000
South Woods Creek.....			1,000
Wild Boy Run.....			1,000
Bellefonte, Fulmers Run.....			1,500
Spring Run.....			3,000
Belleville, Kishacoquillas Creek.....			1,500
Kishacoquillas Creek, South Fork.....			2,000
Bellwood, Logan Spring Pond.....			500
Benton, Banks Run.....			1,000
Belles Run.....			1,500
Benjamin Run.....			500
Colley Brook.....			500
Fair Brook.....			500
Fishing Creek.....			1,000
Gallas Run.....			500
Hess Run.....			1,000
Hickory River.....			1,000
McHenry Run.....			500
Raven Creek.....			2,000
Wiles Run.....			500
Wynona Brook.....			1,000
Berlin, Laurel Run.....			2,000
Birdsboro, Molasses Pond.....			300

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Pennsylvania—Continued.			
Bloomsburg, Crouse Run.....			500
Brandonville, Torbert Run.....			600
Davis Run.....			1,200
Bridgeton, Wises Run.....			500
Bushkill, Bushkill Creek.....			2,000
Carrolltown Road, Ahles Run.....			500
Bash Run.....			500
Bearer Run.....			500
Boslet Run.....			500
Davis Run.....			500
Edwards Run.....			500
Farabaugh Run.....			500
Flemings Run.....			500
Flick Run.....			500
Griffith Run.....			500
Kane Run.....			500
Kirk Run.....			500
Lauer Run.....			500
Meisels Run.....			500
Mohler Run.....			500
Owens Run.....			500
Reese Run.....			500
Shettig Run.....			500
Snyder Run.....			500
Springer Run.....			500
Thomas Run.....			500
Tudor Run.....			500
Williams Run.....			500
Centerbridge, Rodgers's pond.....			500
Central, Beaver Run.....			500
Davis Brook.....			500
Jones Brook.....			500
Stony Brook.....			500
Chambersburg, Birch Run.....			4,000
Carbaugh Run.....			2,500
Hoosic Run.....			2,500
Cherry Run, Penns Run.....			375
Cherry Tree, Shryock Run, North Branch.....			500
Clarendon, Elk Run.....			1,000
Six Mile Creek.....			1,000
Wild Cat Creek.....			1,500
Clearfield, Cold Run.....			1,000
Lick Run.....			1,000
Moose Creek.....			1,000
Morgan Run.....			1,000
Stone Run.....			1,000
Trout Run.....			1,000
Coburn, Donners Deich Run.....			500
East Elk Creek.....			1,000
Elk Creek.....			1,000
Philips Creek.....			1,000
Rough Run.....			500
Spring Run.....			500
Turpentine Creek.....			1,000
West Elk Creek.....			1,000
Cold Springs, Pine Swamp Run.....			1,200
Coles Creek, Black Ash Run.....			500
Coudersport, Allegheny River.....			1,000
Big Morco Run.....			1,000
Lyman Run.....			1,000
Mill Creek.....			1,000
Pine Creek.....			1,000
Prouty Creek.....			1,000
Sinnamahoning Creek, South Branch.....			1,000
Crandalltown, Long Run.....			1,200
Cresco, Broadhead Creek.....			2,000
Buck Hill Creek.....			1,500
Honnet Hill Creek.....			1,000
Mill Creek.....			1,000
Rattlesnake Creek.....			1,000
Stony Run.....			1,500
Cresson, Clearfield Creek.....			1,000
Three Spring Run.....			500
Winterset Run.....			500
Daylesford, Darby Creek.....			500
Delta, Knell Run.....			1,000
Mine Run.....			1,000
Samples Run.....			500

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Pennsylvania—Continued.			
Downingtown, Dallin Run.....			2,000
Davis Run.....			2,000
Glen Isle Run.....			500
Rock Run.....			2,000
Dubois, Big Anderson Creek.....			2,000
Ebensburg, Abrams Run.....			500
Bash Run.....			500
Blacklick Creek.....			500
California Run.....			500
Clear Spring Run.....			500
David Evans Pond.....			500
Davis Creek.....			500
Factory Run.....			500
Farren Brook.....			500
James Run.....			500
Jones Creek.....			1,000
Kirschner Run.....			500
Laurel Branch.....			500
Lloyds Run.....			500
Noel Run.....			500
Roberts Run.....			500
St. James Run.....			500
Sakarak Run.....			500
Smith Run.....			500
Stewarts Run.....			500
Tudor Run.....			500
Williams Run.....			1,000
Ellenton, Rock Run.....			2,000
Emporium, Cooks Run.....			1,000
Crooked Run.....			500
East Cowley Creek.....			1,000
North Creek.....			1,000
Parker Creek.....			1,000
Salt Run.....			2,000
Sinnamaboning Creek.....			1,000
West Cowley Creek.....			1,000
Farrandsville, Lick Run.....			2,800
Fern Glen, Big Tomhicken Creek.....			600
Crooked Run.....			1,000
Roberts Run.....			1,000
Sand Spring Run.....			1,000
Fishing Creek, Fishing Creek.....			1,500
Martin Run.....			500
Forks, Huntingdon Creek.....			2,500
Little Pine Creek.....			1,000
Fort Washington, Kennedy's pond.....			400
Frackville, Crystal Creek.....			600
Little Mahanoy Creek.....			1,800
Tower Run.....			1,000
Frazer, Pigeon Run Pond.....			500
Glen Iron, Penns Run.....			1,125
Grays Run, Grays Run.....			1,800
Long Run.....			1,800
Yoder Run.....			1,000
Yoxtheimer Run.....			600
Greencastle, Willow Brook.....			1,000
Hawley, Wallen Paupac River.....			1,500
Hellam, Locust Run.....			1,000
High Rock, Livingston Run.....			500
Lockport Run.....			500
Tom Creek.....			1,000
Holidaysburg, Blairs Creek.....			1,000
Honesdale, Baker Brook.....			500
Bates Creek.....			500
Big Creek.....			1,000
Bramms Pond.....			500
Calkins Creek.....			500
Dyberry Creek.....			1,000
Fivemile Creek.....			500
Gageis Brook.....			500
Haines Brook.....			500
Kreglers Creek.....			500
Lackawaxen River.....			1,000
Lackawaxen River, North Branch.....			1,000
Middle Creek.....			1,000
Mitchell Creek.....			500
Old Log Cabin Creek.....			1,000
Paynter Brook.....			500

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Pennsylvania—Continued.			
Honesdale, Rattlesnake Creek.....			1,000
Rout Creek.....			500
West Branch.....			1,000
Hopewell, Beaver Creek.....			2,000
Otts Run.....			500
Three Spring Run.....			500
Yellow Creek.....			1,000
Howard, Lick Run.....			500
Hughesville, Muncy Creek.....			3,000
Huntingdon, Mill Creek.....			1,000
Stone Creek.....			1,500
Trough Creek.....			1,500
Jamison City, Bloody Run.....			1,000
Grassy Hollow Run.....			1,000
Haugh Run.....			500
Jersey Shore, Larrys Creek.....			2,400
Keating Summit, Brown Hollow Creek.....			500
Cowley Run.....			1,000
Indian Run.....			500
Portage Creek.....			1,000
Spring Creek.....			500
Knoxville, Troups Creek.....			1,500
Lancaster, Furnace Run.....			1,000
Middle Creek.....			1,000
Silver Run.....			1,000
Steinmans Run.....			1,000
Walnut Run.....			1,000
Landerberg, White Clay Creek, West Branch.....			500
Lanesboro, Brushville Creek.....			500
Canawacta Creek.....			1,000
Cascade Creek.....			1,000
Cold Spring Brook.....			1,000
Dodges Creek.....			500
Drinker Creek.....			1,000
Egypt Creek.....			1,500
Hemlock Creek.....			2,000
Roaring Brook.....			500
Wild Cat Brook.....			1,000
Laquin, Little Schrader Creek.....			1,800
Laubach Station, Hess Run.....			500
Longs Brook.....			500
Savage Brook.....			500
Laughlintown, McMullen Run.....			1,000
Lebanon, Tulpehocken Creek.....			1,000
Lehighton, Spring Brook.....			600
Lemont, Cedar Creek.....			500
Furnace Run.....			500
Hublers Gap Run.....			500
Laurel Run.....			1,000
Pine Swamp Run.....			500
Spring Creek.....			1,500
Lenover, Weavers Run.....			500
Lewisburg, Laurel Run.....			1,500
Rapid Run.....			1,500
White Deer Creek.....			1,500
Lilly, Bear Rock Creek.....			500
Dunn Creek.....			500
Hughes Spring Pond.....			500
Laurel Run.....			500
McTamany Run.....			1,000
Lock Haven, Bagley Run.....			500
Birds Run.....			1,200
Brewer Run.....			500
Castenea Run.....			1,200
Cherry Run.....			500
Chriss Faust Run.....			1,400
Clarks Run.....			500
Considines Run.....			700
Craig Run.....			500
Deise Run.....			1,200
Eady Run.....			500
Earon Run.....			500
Eckers Run.....			500
Ferney Run.....			500
Fogarty Run.....			700
Goulds Run.....			500
Grows Run.....			500

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Pennsylvania—Continued.			
Lock Haven,			
Halls Run.....			1,400
Hanna Run.....			500
Harlens Run.....			1,400
Harveys Run.....			1,200
Heaveners Run.....			700
Hurds Run.....			500
Jerry Run.....			500
Johnson Run.....			500
Kamp Run.....			1,200
Kirbys Run.....			1,200
Kissell Run.....			1,800
Little Bagley Run.....			500
Little Plum Run.....			500
Little Sugar Valley Run.....			1,200
Lloyds Run.....			700
Lucas Run.....			1,200
Lusk Run.....			500
McCloskey Run.....			500
McElhattan Creek.....			2,400
McKagnes Run.....			1,200
Martins Run.....			1,200
Mill Run.....			500
Mitchell Run.....			500
Moganhans Run.....			1,200
Muncher Run.....			700
Musters Run.....			1,200
North Fork Run.....			500
Packer Run.....			500
Pine Bottom Run.....			1,200
Plum Run.....			500
Queens Run.....			500
Quiggles Run.....			1,200
Ram Hollow Run.....			500
Reed Run.....			500
Rickers Run.....			1,400
Rock Run.....			700
Shadles Run.....			1,400
Shingle Hollow Run.....			500
Slab Run.....			500
South Fork Run.....			500
Spring Run.....			700
Sugar Run.....			500
Totanhorn Run.....			1,400
Tyler Run.....			500
Welsh Run.....			500
Wetzells Run.....			700
Widmans Run.....			500
Wiener Run.....			1,200
Wild Run.....			500
Winber Run.....			500
McElhattan,			
Bixler Run.....			700
Chathams Run.....			2,100
Comerdner Run.....			1,200
Jemersons Run.....			700
Little Chathams Run.....			1,200
Lucas Run.....			1,200
McElhattan Run.....			700
Motter Run.....			1,200
Nolans Run.....			700
Russells Run.....			700
Spring Run.....			700
Mahanoy City, Stony Run.....			600
Mansfield, Griffin Creek.....			3,000
Marienville, Bear Pen Run.....			500
Big Salmon Creek.....			1,500
Blue Jay Creek.....			1,000
Brush Creek.....			500
Centennial Run.....			500
Cherry Creek.....			500
Coleman Run.....			1,000
Crosman's pond.....			500
East Cherry Creek.....			500
East Millstone Creek.....			1,000
Guston Run.....			500
Hall's pond.....			500
Huling Run.....			500
Jakes Run.....			500

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Pennsylvania—Continued.			
Marienville, Maple Creek.....			1,000
North Salmon Creek.....			1,000
Six Mile Run.....			500
Truby Run.....			500
Warner Run.....			500
West Millstone Creek.....			2,000
Wild Cat Run.....			500
Marklesburg, Touse Run.....			500
Marsh Hill, Frozen Run.....			1,500
Maston, Pigeon Run.....			1,000
Pleasant Stream.....			2,000
Smith Run.....			1,000
Mauch Chunk, Bear Creek.....			600
Big Bear Creek.....			1,000
Drakes Creek.....			1,000
Glen Run.....			600
Heydst Run.....			600
Hickory Run.....			1,000
James Run.....			1,000
Keipers Run.....			600
Mauch Chunk Creek.....			1,000
Mud Run.....			1,500
Panther Creek.....			600
Pine Run.....			1,000
Robinsons Run.....			600
Ruddles Run.....			600
Sand Spring Run.....			500
Stony Creek.....			1,000
Wild Creek.....			1,000
Yellow Run.....			1,000
Mayport, Pine Run.....			2,000
Meadville, Berley Run.....			1,000
Brawley Run.....			500
Hamilton Run.....			1,000
Little Sugar Creek.....			1,000
Spring Run.....			1,000
Middleport, Cold Run.....			1,000
Mifflinburg, Brush Hollow Run.....			500
Buffalo Creek.....			1,500
First Gap Run.....			500
Fourth Gap Run.....			1,500
Halfway Gap Run.....			500
Hays Gap Run.....			500
Lukers Gap Run.....			500
Pine Swamp Creek.....			1,000
Rapid Run.....			1,500
Reeds Gap Run.....			500
Sand Run.....			500
Second Gap Run.....			500
Spruce Run.....			1,000
Third Gap Run.....			1,000
Yankee Run.....			500
Mifflintown, Big Run.....			1,500
East Lost Creek.....			1,500
Hornings Run.....			1,000
Sponhowers Run.....			1,000
Tennis Run.....			500
West Lost Creek.....			1,500
Millville, Bear Run.....			1,000
Milroy, Laurel Run.....			1,500
New Lancaster Stream.....			1,500
Mt. Joy, Big Spring Creek.....			500
Mt. Pocono, Wilson Spring Run.....			500
Mt. Union, Carters Run.....			500
Scrub Gap Run.....			1,000
Singers Gap Run.....			1,000
Muncy, Muncy Creek.....			2,500
New Freedom, Codorus Creek.....			1,000
Summitt Creek.....			500
New Holland, Goods Run.....			500
New Ringgold, Beaver Creek.....			600
Cold Run.....			600
Rausch Creek.....			1,000
Newton Hamilton, Licking Creek.....			1,000
Long Hollow Run.....			500
Nigger Creek.....			1,000
Orangeville, Achenbach Run.....			500

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Pennsylvania—Continued.			
Osceola Mills, Bear Run.....			2,000
California Run.....			2,000
Coal Creek.....			2,000
Flat Rock Creek.....			2,000
Mountain Creek.....			2,000
Trout Run.....			2,000
Paddy Mountain, Penns Run.....			2,625
Palm, Indian Creek.....			1,000
Parkersburg, Octorara Creek.....			1,500
Parsons, Bear Creek.....			1,500
Meadow Run.....			1,000
Pond Creek.....			1,000
Ten Mile Run.....			1,000
Patton, Carroll Creek.....			500
Shehan Run.....			500
Paxinos, Irish Creek.....			1,000
Petersburg, Garners Run.....			1,000
Globe Run.....			1,000
Irvin Run.....			500
Lick Run.....			1,000
Roaring Run.....			500
Philadelphia, Darbey Creek.....			1,000
Phillipsburg, Ardells Spring Run.....			500
Barker Run.....			500
Beaver Run.....			1,500
Bennens Run.....			1,000
Big Spring Run.....			500
Bilgers Run.....			1,000
Black Bear Run.....			1,500
Black Moshannon Creek.....			2,000
California Run.....			1,000
Clearwater Run.....			1,000
Clover Run.....			1,000
Cold Run.....			2,000
Dayton Run.....			1,000
Echo Glen Park Lakes.....			1,000
McCords Run.....			500
Morgan Run.....			1,000
Nooch Run.....			1,000
One Mile Run.....			500
Senser Run.....			500
Seven Springs Run.....			500
Shields Run.....			500
Six Mile Run.....			2,000
Smays Run.....			1,000
Tests Run.....			500
Tomtit Run.....			500
Upper Daugherty Run.....			600
Whetstone Run.....			500
Wolf Run.....			1,000
Pleasant Stream Junction, Potash Run.....			1,200
Pottstown, Powderdale Run.....			1,200
Pottsville, Big Creek.....			1,200
Black Creek.....			1,500
Breechlez Pond.....			600
Eichert Creek.....			600
Hells Creek.....			600
Neland's pond.....			600
Rattling Run.....			600
Seltzer Creek.....			600
Stony Creek.....			600
Strouser Creek.....			600
Powys, Cold Fork Run.....			500
Daugherty Run.....			600
Long Fork Run.....			600
Lower Daugherty Run.....			600
Wolf Run.....			600
Ralston, Rocky Run.....			2,000
Rattling Run, Rattling Run.....			500
Reading, Furnace Creek.....			600
Hartmens Creek.....			500
Hay Creek.....			1,000
Holdennan Creek.....			500
Laurel Creek.....			3,500
Limekiln Brook.....			600
Willow Creek.....			600
Wyomissing Creek.....			600

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Pennsylvania—Continued.			
Reedsville, Kishacoquillas Creek.....			2,000
Renovo, Bakers Run.....			2,400
Barneys Run.....			1,200
Benjamin Run.....			1,200
Boggs Run.....			1,200
Cranberry Run.....			1,200
Drurys Run.....			3,600
Fish Dam Run.....			1,400
Halls Run.....			1,800
Paddys Run.....			2,100
Shintown Run.....			1,400
Reynoldsville, Bear Pen Run.....			1,000
Black Run.....			1,000
Bollingers Run.....			1,000
Boyer Run.....			1,000
Britton Run.....			1,000
Bustop Run.....			1,000
Callen Run.....			1,000
Camp Run.....			1,500
Clover Run.....			1,000
Deans Run.....			1,000
Deemers Run.....			500
Degnan Run.....			1,000
Five Mile Run.....			1,000
Forest Run.....			1,000
Horn Run.....			1,000
Jenkins Run.....			1,000
Keys Run.....			500
Kyle Run.....			1,000
Laurel Run.....			1,000
McConnells Run.....			1,000
Manners Run.....			1,000
Mill Creek.....			1,000
Mitchells Run.....			1,000
Morrison Run.....			1,000
Mountain Run.....			1,000
Mowrey Run.....			1,000
O'Donnell Run.....			1,000
Panther Run.....			1,000
Pitch Pine Run.....			1,000
Rattlesnake Run.....			1,000
Schuckers Run.....			1,000
South Fork Creek.....			1,000
Stevenson Run.....			1,000
Toby Run.....			1,000
Trout Run.....			1,000
West Fork Creek.....			1,000
Whitstone Run.....			1,500
Windfall Run.....			1,000
Wolf Creek.....			1,000
Rising Springs, Laurel Run.....			2,000
Locust Run.....			1,000
Penns Creek.....			2,500
Rockport, Rapps Creek.....			1,000
Roulette, Bear Hollow Creek.....			1,000
Card Creek.....			500
Fishing Creek.....			2,000
Fish Hollow Creek.....			1,000
Laninger Creek.....			500
Reeds Run.....			1,500
Sartwell Creek.....			1,000
Trout Brook.....			1,000
Royer, McAllister Pond.....			500
Piney Creek.....			1,000
Sandy Run.....			500
Spring Run.....			1,000
Shade Gap, Scotts Run.....			500
Shenandoah, Fowler Pond.....			1,000
Knicker Hollow Run.....			1,000
Railroad Reservoir.....			1,000
Rattling Run.....			600
Shenandoah Reservoir.....			500
Thrashers Run.....			600
Trexler Run.....			1,000
Short Run Station, Short Run.....			1,000
Shrewsbury, Deer Creek.....			500
Smethport, Boyer Brook.....			1,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Pennsylvania—Continued.			
Somerfield, Youghiogheny River.....			400
Spruce Creek, Spruce Creek.....			2,000
Starrucca, Coxtown Creek.....			1,000
Farrell Creek.....			500
McKane Creek.....			1,000
Sampson Creek.....			500
Shadagee Creek.....			1,000
Shehawkin Creek.....			500
Starrucca Creek.....			2,000
Wild Cat Creek.....			500
Stewartstown, Grove Run.....			1,000
Stillwater, Myers Run.....			500
Roberts Run.....			500
Trout Run.....			500
Stroudsburg, Baker Run.....			500
Broadhead Creek.....			1,500
Brown Run.....			1,000
Cherry Creek.....			2,000
Deep Hollow Run.....			1,000
Kettle Run.....			1,000
McMichaels Creek.....			1,500
Mountain Creek.....			1,000
Pencil Creek.....			2,000
Pocono Creek.....			2,000
Sambo Creek.....			600
Wigwam Run.....			500
Tionesta, Bates Run.....			500
Bear Creek.....			500
Big Coon Creek.....			1,500
Chauncy Run.....			500
Council Run.....			500
Davis Run.....			500
Dawson Run.....			500
Hemlock Creek.....			1,500
Holeman Run.....			500
Indian Camp Creek.....			500
Jakes Run.....			500
Jamieson Run.....			500
Johns Run.....			500
Jug Handle Run.....			500
Korb Run.....			500
Lamentation Run.....			500
Little Coon Creek.....			1,000
Little Hickory Creek.....			1,000
Little Tionesta Creek.....			1,000
Pearson Run.....			500
Peters Run.....			500
Pigeon Run.....			500
Piney Run.....			500
Pit Hole Creek.....			1,500
Reck Run.....			500
Ross Run.....			1,000
Salmon Creek.....			1,500
Sandrock Run.....			500
Sibble Run.....			500
Stewarts Run.....			1,000
Sugar Run.....			500
Tubbs Run.....			1,000
Tower City, Clarks Creek.....			2,500
Rausch Creek.....			1,000
Troy, Becker Creek.....			600
Bullard Creek.....			600
Cleveland Creek.....			600
Covert Creek.....			600
Dry Run.....			600
Forbes Creek.....			600
Keith Creek.....			600
Kieff Creek.....			1,000
Kinar Creek.....			600
Morgan Creek.....			1,000
Palmer Creek.....			600
Rathborn Creek.....			1,200
Sherman Creek.....			600
Tamarack Creek.....			600
Webber Creek.....			500
Ulysses, Pine Creek.....			1,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Pennsylvania—Continued.			
Waynesboro, Antietam Spring, Branch			500
Weikert, Penns Run			375
West Chester, Broad Run			500
Wheeler'sville, Schrader Creek			3,000
Williamsburg, Brumbaugh's Run			500
Clover Creek			2,000
Marsh Run			500
Woodbine, Bells Hollow Branch			1,000
Boyd's Run			1,000
Kilgore Run			500
Rocky Run			1,000
Wade Hill Branch			500
York, Green Branch			1,000
South Carolina:			
Cleveland, Fall Creek			3,200
Headforemost Creek			3,200
Reeces Gap Creek			2,400
Pickens, Big Laurel Creek			1,600
Cane Creek			2,400
Dogwood Stump Creek			3,200
Laurel Ford Creek			2,400
Laurel Fork Creek			2,400
Lynchs Mill Creek			2,400
Mathers Creek			2,400
Siele Mountain Creek			2,400
Surveyors Camp Creek			2,400
Willis Creek			2,400
South Dakota:			
Custer, Willow Creek			7,500
Deadwood, Spruce Creek			8,000
Doyle, Big Elk Creek			20,000
Dumont, Spearfish Creek, East Fork			5,000
Elmore, Ice Box Canyon Creek			10,000
Spearfish Creek			15,000
Englewood, White Wood Creek			10,000
Hanna, Little Spearfish Creek, East Fork			10,000
Hermosa, Battle Creek			12,500
Hill City, Dismal Creek			7,500
Gibson Creek			10,000
Hutton Creek, South Branch			10,000
Palmer Creek			10,000
Spring Creek			7,500
Sunday Gulch Creek			7,500
Mystic, Prairie Creek			20,000
Tittles Springs Pond			20,000
Victoria Creek			20,000
Nemo, Box Elder Creek			12,000
Jim Creek			6,000
Knowlton's pond			6,000
South Box Elder Creek			6,000
Pine Ridge Agency, Bear Creek			12,500
Pringle, Beaver Creek			8,000
Cold Brook			8,000
Rapid City, Deer Creek			20,000
Pine Forest Lake			12,500
Rapid Creek			20,000
Spring Canyon Pond			30,000
Roubaix, Carroll Creek			6,000
Halls Pond			6,000
North Elk Creek			6,000
Rochford, Little Rapid Creek, North Fork			10,000
Sisseton, Long Hollow Creek			1,000
Spearfish, Cox Lake			2,500
False Bottom Creek			2,500
Hiltens Gulch Creek			8,000
Kingsley's lake			12,000
Lindley Spring Run			14,000
McGregor Spring Branch			10,000
Miller Creek			10,000
Normal Lake			12,000
Spearfish River			75,000
Todd's pond			6,000
Spring Gulch, McDonald Pond			12,500
Sturgis, Deadmans Creek			10,000
Walker, Rock Creek Pond			12,500
Tennessee:			
Blevins, Brushy Creek			2,400

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Tennessee—Continued.			
Butter, Greggs Branch.....			2,400
Greenville, Camp Creek.....			4,000
Knoxville, Fountain City Lake.....			4,000
Nashville, Lipscomb's pond.....			800
Newport, Ground Hog Creek.....			2,400
Pikeville, Bradens Creek.....			4,000
Cooper Branch.....			2,400
Glade Creek.....			5,600
Halls Creek.....			3,200
Skillern Creek.....			4,000
Shell City, Doll Branch.....			1,600
Shell Creek.....			5,600
Slocums, Farmer Branch.....			1,600
Shouns, McEwen Branch.....			1,600
Payne creek.....			1,600
Utah:			
Provo, applicant.....	100,000		
Applicant.....	25,000		
Grandview Pond.....			1,800
Provo River.....		18,600	
Robins Springs Pond.....			1,800
Spring Creek Pond.....			1,800
Springdale Pond.....			3,600
Upper Falls Ponds.....			1,800
Vineyard Ponds.....			3,600
Salt Lake, Spring Creek.....			3,000
Springville, Spring Creek.....			2,400
Vermont:			
Averill, Forest Lake.....		35,000	
Little Averill Lake.....		55,000	
Mild Brook.....			1,500
Bellows Falls, Morse Brook.....		25,000	
Bennington, Jackson Brook.....		12,000	
Brattleboro, Ames Brook.....			1,000
Brickyard Brook.....			1,000
Broad Brook.....			1,500
Houghton Brook.....			1,000
Johnson Brook.....			1,000
Weatherhead Hollow Brook.....			1,000
Whetstone Brook.....			2,000
Wilder Brook.....			1,000
Castleton, Castleton River.....			3,500
Chester, Fullerton Brook.....			1,000
Williams River.....			3,000
Cuttingsville, Shrewsbury Pond.....			4,800
Fair Haven, Eureka Pond.....			1,000
Fowler, Fowler Brook.....			1,500
Greensboro, Caspian Lake.....			6,000
Groton, Darling Pond.....		125,000	7,000
Holden, Furnace Brook.....			10,000
Pico Pond.....			3,000
Hydeville, Castleton River.....			4,000
Ferrin River.....		6,000	
Lyndonville, Vail's pond.....			850
Manchester, Batten Kill River.....		56,000	
Lye Brook.....			1,400
Mountain Brook.....		8,000	
Marshfield, Niggerhead Pond.....			3,000
Montpelier, Mallory Brook.....			2,500
North Bennington, Cold Springs Brook.....		12,000	1,250
Paran Creek.....		12,000	1,250
Northfield, Yatter Pond.....		16,000	
Pawlet, Pawlet River.....			5,000
Pittsford, Furnace Brook.....		10,000	
Sugar Hollow Brook.....			3,000
Plainfield, Laird's pond.....			4,000
Poultney, Poultney River.....			4,000
Pownel, Mattison Brook.....			2,000
Proctor, Fox Pond.....			4,000
Proctorsville, Williams River.....		20,000	
Putney, Sacketts Brook.....			1,500
Randolph, Ayers Brook.....		20,000	2,000
Bear Hill Brook.....		8,000	
Chandler Brook.....		16,000	
Clough Brook.....		8,000	
Eldredge Pond.....			500
Fisher Brook.....		8,000	

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Vermont—Continued.			
Randolph, Guilds Brook		8,000	
Halfway Brook		16,000	
Howard Hill Brook		8,000	
Meadow Brook		20,000	
Mud Pond		8,000	
Roods Brook		8,000	
Roxbury Brook		12,000	
Snow Brook		8,000	
White River, Middle Branch		24,000	
Readsboro, Lamb Brook			1,500
South Branch			1,500
Rutland, Atwood Brook			1,000
Beaver Meadow Brook		8,000	
Billings Brook			1,500
Brewer Brook		12,000	
Castleton River			5,500
Chittenden Reservoir			12,000
Cold River		16,000	1,350
Cold River, North Branch		12,000	
Cold River, South Branch		32,000	
Curtis Brook		12,000	
Deermont Creek		12,000	
East Brook			1,000
Eddy Brook		8,000	
Gleason Brook		12,000	
Ira Brook		8,000	
Ottaqueechee Brook		16,000	
Ripley Brook		8,000	
Sharon, Lake Mitchell		100,000	5,425
White River		8,000	
South Royalton, Pinehurst Lake		20,000	
South Ryegate, Hatch's pond		25,000	
South Wallingford, South Wallingford Branch		16,000	
St. Johnsbury, Blodgett Brook		15,000	
Fairbanks Ponds			648
Frog Pond		10,000	500
Green Mountain Brook		20,000	
Grouselands Pond			500
Joes Brook			7,500
Lawrence Ponds			500
Meadow Brook		20,000	
Sleeper River			1,677
Spaulding Brook			1,000
Stony Brook		20,000	
Water Endrick Creek			2,000
Waterman's pond			148
Springfield, Hazen's pond			500
Stockbridge, Tweed River		8,000	2,000
Taftsville, Beaver Brook			2,000
Townshend, Shanty Lot Brook			2,000
Walden, Haynesville Brook			1,500
Lyford Pond		40,000	
Meadow Brook		20,000	
Wells, Wells Brook		16,000	
West Hartford, Dimmick's ponds			1,000
Meadow Brook		8,000	
Northcote Brook			1,000
Rockland Brook			1,000
Whipple Brook			1,000
Woodland Brook			1,000
West Paulet, Indian River		20,000	
Windsor, Mill Brook			3,000
Woodstock, Lakota Lake			4,000
Moore Pond			1,500
Smith Brook		8,000	
Wyandale Brook		8,000	
Virginia:			
Alleghany Station, Cove Creek			500
Arcadia, North Creek			400
Arrington, Mountain Spring Pond			2,400
Basic City, Baker Springs			300
Jordan Pond			1,000
Bedford, North Otter River			2,400
Big Island, Hunting Creek			4,000
Reed Creek			4,000
Covington, Cast Steel Run			600
Laurel Run			3,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Virginia—Continued.			
Covington, Roaring Run			3,000
Craigsville, Campbell Run			500
Claytons Brook			1,500
Culpeper, Hazel River			4,800
Miller Creek		18,700	500
Ferrol, Trout Run			2,400
Glenvar, Callahan Brook			6,000
Goshen, Kelso Run			300
Grottoes, Big Run			300
Harrisonburg, Long Run			2,500
Hunters, Little Diffcult Run			600
Jenkins Ford, Cedar Creek			6,000
Maurertown, Cedar Creek			1,000
Mount Vernon, Washington Spring Branch			2,400
Pearch, Horsleys Creek			600
Richmond, Burke's pond			2,400
Rockfish, Goldmine Creek			6,400
Salem, Peters Creek			500
Spout Spring, Steele's pond			400
Stanley, Hendersons Mill Pond			50
Tates Run, Tates Run			2,400
Tye River, Cox Creek			
Washington:			
Addy, Stenger Creek			4,500
Bellingham, State Fish Commission	100,000		
Colville, Twin Lakes			4,500
Lake View, Clover Creek			5,000
Lamona, Crab Creek			5,500
Newport, Bead Lake			6,000
Mystic Lake			6,000
Seattle, Exposition Aquarium			18
Spangle, Spring Lake			3,000
Spokane, Newman Lake			6,000
Wenatchee, Spring Valley Pond			6,000
West Virginia:			
Berkeley, Cold Run			800
Beverly, Beaver Creek			1,000
Burner, Harper Run			1,500
Little River			2,000
Mountain Lick Run			2,000
Span Oak Run			2,000
Cairo, Lake Carrell			1,000
Capon Road, Laurel Lake			500
Capon Springs, Mutton Run			3,750
Davis, Blackwater River			2,500
Harman, Spruce Run			1,000
Harton, Candy Creek			3,000
Huttonsville, Elk River			500
Files Creek			1,500
Mill Creek			1,500
Riffles Creek			1,500
Keyser, Patterson Creek, North Fork			1,200
Marlinton, Cochrans Creek			750
Elk River, Crooked Fork			750
Indian Draft Creek			2,500
Mill Run			1,000
May, Greenbrier River			3,000
Orndorf Run			1,000
White Camp Run			1,000
Midvale, Cassity Fork Creek			3,000
Raleigh, Piney Creek			14,000
Renick, Spring Creek			500
Rippon, Bullskin Run			1,500
Seebert, Cranberry Creek			6,000
Terra Alta, Big Run			1,000
Big Wolf Creek			2,000
Buck Lick Creek			1,000
Dority Creek			2,500
Elsy Creek			2,000
Kinsinger Creek			800
Laurel Run			1,200
Little Wolf Creek			3,000
Muddy Creek			1,500
Roaring Creek			2,500
Salt Lick Creek			4,000
Snowy Creek			6,700
Spruce Run			1,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
West Virginia—Continued.			
Terra Alta, White Oak Creek.....			2,000
Webster Springs, Elk River, Buck Fork.....			600
White Sulphur Springs, Laurel Creek.....			1,000
Spring Branch.....		59,000	1,000
Turner Creek.....			1,000
Wildell, Elk Run.....			4,000
Mike Run.....			2,000
Snorting Lick Run.....			2,000
Wisconsin:			
Albertville, Little Elk Creek.....			3,000
Alma, Little Waumandee Creek.....			2,800
Alma Center, Pigeon Creek.....			1,200
Almena, Hay River.....			6,000
Arcadia, Bishop Creek.....			300
Eagle Valley Creek.....			300
French Creek.....			300
Gilman Creek.....			300
Haines Creek.....			300
Holcomb Coulee Creek.....			300
Hunters Creek.....			300
Kried Valley Creek.....			300
Lewis Valley Creek.....			300
Long Creek.....			300
Mineral Spring Brook.....			300
Montana Creek.....			300
Rocky Run Creek.....			300
Sandy Creek.....			300
Scharlow Valley Creek.....			300
Trout Run.....			300
Auburndale, Mohan Creek.....			4,000
Augusta, Beamans Creek.....			600
Bears Grass Creek.....			800
Beaver Creek.....			600
Bee Creek.....			300
Beef River.....			400
Bridge Creek.....			600
Browns Creek.....			300
Chaney Creek.....			300
Coon Gut Creek.....			300
Diamond Creek.....			300
Hathaway Creek.....			400
Hay Creek.....			400
Horse Creek.....			400
Muskrat Creek.....			400
Otter Creek.....			800
Sand Creek.....			300
Thompson Creek.....			400
Travis Creek.....			300
Bangor, Adams Creek.....			400
Big Creek.....			300
Burns Creek.....			900
Kalburan Creek.....			300
Sand Creek.....			600
Swamp Creek.....			300
Barneveld, Clavaln Stream.....			4,000
Four Mile Creek.....			800
Beldenville, Trimbelle Creek.....			900
Birchwood, Fullerton Pond.....			2,700
Black River Falls, Roaring Creek.....			6,000
Blair, Bear Creek.....			300
Beaver Creek.....			300
Lake Coulee Creek.....			300
Strum Creek.....			300
Tappan Creek.....			300
Tennison Creek.....			300
Vasse Coulee Creek.....			300
Welsh Coulee Creek.....			300
Bluff Siding, Bohlies Valley Creek.....			600
Bohn Valley Creek.....			1,000
Eagle Valley Creek.....			1,600
Fox Coulee Creek.....			1,000
French Creek.....			1,000
Holcomb Coulee Creek.....			1,400
Little Tamarack Creek.....			400
Norwegian Creek.....			400
Norway Coulee Creek.....			1,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Wisconsin—Continued.			
Bluff Siding, Pine Creek.....			1,600
Brule, Carlson Creek.....			2,000
Shade Creek.....			4,000
Stony Brook.....			3,000
Cable, Big Run.....			4,000
Cable Lake Brook.....			2,000
Caps Creek.....			2,000
Five Mile Creek.....			2,000
Garrison Brook.....			4,000
Lynch Creek.....			4,000
Namekagon River.....			8,000
Neffs Brook.....			2,000
Ole Lake Brook.....			4,000
Spring Brook.....			2,000
Twin Brooks.....			4,000
Cadott, Big Drywood Creek.....			3,000
Paint Creek.....			4,000
Willow Creek.....			1,000
Camp Douglas, Little Lemonweir River.....			1,000
Cashton, Coon Creek.....			1,000
Fremstead Creek.....			1,000
Hanson Creek.....			1,000
Heiser Creek.....			1,000
Jersey Creek.....			1,000
Lyons Creek.....			1,000
Meissner Creek.....			1,000
Timber Coulee Creek.....			1,000
Witelman Creek.....			1,000
Cassville, Furnace Branch.....			600
Chippewa Falls, Big Beaver Creek.....			1,000
Clear Creek.....			1,000
Drywood Creek.....			1,000
Duncan Creek.....			1,000
Eighteen Mile Creek.....			1,000
Elk Creek.....			1,000
Hay Creek.....			1,000
Jims Falls Creek.....			1,000
Little Beaver Creek.....			1,000
Little Drywood Creek.....			1,000
Little Hay Creek.....			1,000
McCann Creek.....			1,000
Murphy Creek.....			1,000
Nicoli Creek.....			1,000
Paint Creek.....			1,000
Seth Creek.....			1,000
Tenmile Creek.....			1,000
Trout Creek.....			1,000
Cochrane, Breams Valley Brook.....			300
Bulls Valley Brook.....			300
Dannser Valley Brook.....			300
Esbach Brook.....			300
Florin Valley Brook.....			300
Irish Valley Brook.....			300
Johns Valley Creek.....			600
Mill Creek.....			300
Montane Brook.....			300
Oak Valley Brook.....			300
Rebhahu Valley Brook.....			300
Rose Valley Brook.....			300
Rutsehous Brook.....			300
Schaub Brook.....			300
Schoepps Valley Brook.....			300
Schultz Brook.....			300
Weisenberger Brook.....			300
Wolf Valley Brook.....			300
Yaeger Brook.....			300
Crandon, Andrews Pond.....			1,000
Drake Creek.....			1,000
Mud Lake.....			4,000
Rice Creek.....			3,000
Swamp Creek.....			2,000
Wolf River.....			1,000
Cumberland, Miller Creek.....			4,500
Dodgeville, Bremker Creek.....			500
Edmunds Branch.....			3,000
Hoskins Branch.....			3,000
Middleberry Creek.....			800

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Wisconsin—Continued.			
Dodgeville, Smith Creek.....			3,000
Williams Stream.....			1,500
Drummond, Jaders Creek.....			4,500
Johnson Creek.....			1,500
Long Lake Branch.....			4,500
Durand, Averill Creek.....			300
Bear Creek.....			1,200
Big Arkansas Creek.....			2,000
Big Coulee Creek.....			1,000
Drier Creek.....			1,000
Fall Creek.....			1,200
Fox Creek.....			300
Gray Creek.....			300
Heron Creek.....			1,000
Little Arkansas Creek.....			2,000
Porcupine Creek.....			2,000
Spring Creek.....			600
Eau Claire, Beaver Creek.....			1,600
Clear Creek.....			1,600
Coon Creek.....			1,500
Craft Creek.....			1,000
Cranberry Creek.....			800
Deer Creek.....			500
Eight Mile Creek.....			1,000
Eighteen Mile Creek.....			1,000
Elk Creek.....			2,600
Five Mile Creek.....			1,500
Grace Creek.....			400
Hansen Creek.....			2,000
Little Niagara Creek.....			300
Little Rock Creek.....			500
Lowes Creek.....			1,800
Nine Mile Creek.....			1,800
North Creek.....			300
Otter Creek.....			1,000
Pine Creek.....			500
Rock Creek.....			1,600
Sandy Creek.....			500
Seven Mile Creek.....			1,500
Sherman Creek.....			1,600
Spring Creek.....			1,300
Trout Creek.....			1,800
Twelve Mile Creek.....			1,000
West Creek.....			1,000
Wrights Creek.....			800
Edgewater, Arfin Creek.....			1,000
Beaver Creek.....			1,000
Billikin Springs Creek.....			2,000
Casey Creek.....			1,000
Derosier Creek.....			1,000
Hay Creek.....			2,000
Laughing Water Creek.....			1,000
Mallard Creek.....			1,000
Moose Creek.....			2,000
Nelson Creek.....			1,000
Pigeon Creek.....			2,000
Plum Creek.....			1,000
Sissebagama Creek.....			1,000
Trout Creek.....			2,000
Yarnell Creek.....			2,000
Elcho, Hunting River.....			9,000
Eleva, Big Creek.....			1,000
Trout Creek.....			1,000
Ellsworth, Brush Creek.....			3,000
Cave Creek.....			3,000
Isabelle Creek.....			3,000
Lost Creek.....			3,000
Elmwood, Big Mosourie River.....			4,000
Cady Creek.....			3,000
Cave Creek.....			3,000
Eau Galle River.....			4,000
Plum Creek.....			4,000
Fairchild, Black Creek.....			300
Boatman Creek.....			300
Coon Fork Creek.....			600
Coon Gut Creek.....			300
Flick Creek.....			600

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Wisconsin—Continued.			
Fairchild, Johnson Creek			400
McLaren Creek			300
Marrin Creek			300
Pitts Creek			300
Toals Creek			300
Travis Creek			300
Fennimore, Legged Creek			4,500
Fond du Lac, Parson Brook			500
Foxboro, Big Balsam Creek			6,000
Empire Creek			4,000
Little Balsam Creek			4,000
State Line Creek			6,300
Galesville, Beaver Creek			300
Beaver Creek, North Branch			300
Beaver Creek, South Branch			900
Bean Creek			300
Corrigan Creek			300
Coulee Creek			300
Crystal Valley Creek			300
Dutch Creek			300
French Creek			300
Grant Creek			300
Hardy Creek			600
Silver Creek			300
Tamarack Creek			300
Gleason, Eight Mile Creek			2,000
Hay Meadow Creek			2,000
North Branch River			2,000
Pine River			2,000
Silver Creek			2,000
Glenwood, Balons Creek			400
Behrens Creek			400
Beleans Creek			300
Blakely Creek			300
Bolan Creek			400
Browns Creek			300
Camp Nine Creek			300
Conners Creek			300
DeSmith Creek			600
Eldridge Creek			300
Jacobson Creek			300
Johns Creek			300
Little Beaver Creek			400
Morgan Creek			300
Sachse Creek			300
Sand Creek			700
Sullivan Creek			300
Torgeson Creek			300
Vance Creek			400
Grand Rapids, Five Mile Creek			1,000
Green Bay, De Greef's pond			500
Greenwood, Alder Creek			300
Black Creek			2,400
Cawley Creek			500
Colby Creek			2,000
Dickerson Creek			400
Giler Creek			300
Hay Creek			300
Kawley Creek			2,000
Nichol Creek			300
Norwegian Creek			2,400
Rock Creek			500
Rocky Run			2,400
Wedges Creek			500
Hackley, Hackley Creek			3,000
Harshaw, Bearskin Creek			5,000
Little Bear Creek			1,000
Rice Creek			3,000
Hejneman, Prairie River			900
Hixton, Amo Creek			1,000
Curran Creek			1,000
Gaulster Creek			1,000
Holmes Creek			1,000
Judkins Creek			1,000
Larson Creek			1,000
Lowe Creek			1,000
Mortiboy Creek			1,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Wisconsin—Continued.			
Hixton, Nettleton Creek			1,000
North Branch			1,000
Pine Creek			2,000
Schmerhorn Creek			1,000
Simpson Creek			1,000
Tank Creek			1,000
Timber Creek			1,000
Hudson, Willow River			3,000
Independence, Bennett Valley Creek			300
Borst Valley Creek			1,300
Bruce Valley Creek			1,300
Burt Valley Creek			1,000
Chimney Rock Creek			1,300
Cookes Creek			1,000
Dubil Valley Creek			1,000
Elk Creek			1,300
Elk Creek Pond			300
Engum Creek			1,000
Finright Creek			1,000
Gunderson Creek			1,000
Hawkinson Creek			1,000
Husselgard Creek			1,300
Ignatz Lyga Creek			1,000
Kilniss Creek			1,000
Kurth Valley Creek			1,000
Lyga Creek			1,000
Maloney Creek			1,000
Nelson Valley Creek			1,000
North Branch Creek			1,300
Olson Creek			1,000
Plumb Creek			1,300
Poppies Creek			1,000
Roskos Creek			1,000
Russell Valley Creek			1,000
Rusts Creek			1,000
Schaffners Creek			1,000
Simonson Valley Creek			1,000
Skogstad Creek			1,300
Slanton Creek			1,000
Solfest Creek			1,300
Traverse Valley Creek			1,300
Uetz Creek			1,000
Ulbug Valley Creek			1,000
Vennis Creek			1,000
Zimmer Creek			1,300
Iron River, Iron River			4,000
Muskeg Creek			8,000
Pine Lake			4,000
Kendall, Brainard Creek			300
Davis Creek			300
Oborn Creek			300
Smiths Creek			300
La Crosse, Fays Creek			400
Sand Creek			300
La Farge, Dalton Spring Branch			300
Indian Creek			2,000
North Bar Creek			4,000
Spring Creek			4,000
Lancaster, Austin Branch			3,000
Beatham Branch			3,000
Borah Branch			3,000
Club Branch			3,000
Day Branch			1,500
McKenzie Branch			1,500
McPherson Branch			1,500
Millner Branch			3,000
Pollock Branch			1,500
Raines Branch			3,000
Trollope Branch			3,000
Walker Branch			3,000
Williams Branch			3,000
Lavalle, McGloy Creek			300
Long Lake, Coldwater Creek			4,000
Manitowoc, Francis Creek			3,000
Hermans Creek			3,000
Mellen, Bad River			10,000
Tyler Forks River			1,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Wisconsin—Continued.			
Menomonie, Anderson Creek.....			800
Annis Creek.....			800
Asylum Springs Creek.....			800
Balsbaugh Creek.....			800
Beaver Creek.....			800
Big Elk Creek.....			800
Big Hay Creek.....			800
Big Meadow Creek.....			800
Bishop Creek.....			800
Biss Creek.....			800
Blairs Creek.....			800
Boland Creek.....			1,600
Browns Creek.....			800
Clarks Creek.....			800
Coon Creek.....			800
Cowan Creek.....			800
Cranberry Creek.....			800
Dashone Creek.....			800
Denning Creek.....			800
Drowleys Spring Creek.....			800
Eau Galle River.....			800
Eddy Creek.....			800
Eighteen Mile Creek.....			800
Fall Creek.....			800
Foss Creek.....			800
Galloway Creek.....			800
Gilbert Creek.....			800
Hay River.....			1,600
Home Farm Creek.....			800
Iron Creek.....			800
Irvin Creek.....			800
John Creek.....			800
Kings Creek.....			800
Knights Creek.....			800
Kripple Creek.....			800
La Forge Creek.....			800
Lambs Creek.....			1,600
Little Beaver Creek.....			800
Little Elk Creek.....			800
Little Hay Creek.....			800
Little Missoni River.....			800
Little Otter Creek.....			800
Little Sand Creek.....			800
Losby Run.....			800
Louis Creek.....			800
Lower Pine Creek.....			800
Lynch Creek.....			800
McCarthy Creek.....			800
Missoni River.....			800
Mud Creek.....			800
Otter Creek.....			800
Palmers Run.....			800
Paradise Creek.....			800
Parker Springs Creek.....			800
Popple Creek.....			800
Roach Creek.....			800
Roek Creek.....			800
Rush Creek.....			800
Sand Creek.....			800
Shofer Creek.....			800
Simonson Creek.....			800
Sinking Creek.....			800
Sly Creek.....			800
Smith Creek.....			800
Spring Creek.....			800
Stoner Creek.....			800
Thum Creek.....			800
Tiffany Creek.....			800
Torgerson Creek.....			800
Trout Creek.....			800
Upper Pine Creek.....			800
Varney Creek.....			800
Weber Creek.....			800
White Creek.....			800
Wileox Creek.....			800
Wilson Creek.....			1,600
Wolf Run.....			800

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Wisconsin—Continued.			
Merrillan, Arnold Creek.....			2,000
Cesna Creek.....			2,700
Clark Creek.....			300
Farka Creek.....			1,300
Flood Creek.....			400
Gearing Creek.....			1,300
Hall Creek.....			2,000
Hayden Creek.....			500
Houghton Creek.....			300
Hunters Creek.....			300
Jones Creek.....			300
Pine River.....			1,200
Prairie Creek.....			2,000
Stocknell Creek.....			600
Van Hersey Creek.....			1,000
Visneau Creek.....			1,500
Millston, Alvord Creek.....			1,000
Clear Creek.....			1,000
Dunham Creek.....			1,000
Gebhardt Creek.....			1,000
Glen Creek.....			1,000
Hauser Creek.....			2,000
Indian Creek.....			1,000
Ketchum Creek.....			1,000
King Creek.....			1,000
Lamb Creek.....			1,000
Lambert Creek.....			1,000
Merritt Creek.....			1,000
Mill Creek.....			1,000
Patterson Creek.....			1,000
Pigeon Creek.....			1,000
Pongartz Creek.....			1,000
Pulling Creek.....			1,000
Pump Creek.....			1,000
Robinson Creek.....			1,000
Rudd Creek.....			2,000
South Wyman Creek.....			1,000
Spring Creek.....			1,000
Stanton Creek.....			2,000
Wyman Creek.....			1,000
Mondovi, Adams Creek.....			300
Bennett Valley Creek.....			400
Big Creek.....			300
Brown Creek.....			300
Carroll Creek.....			500
Cranberry Creek.....			500
Dutch Creek.....			300
Elk Creek.....			300
Fifteen Creek.....			300
Ford Creek.....			600
Gilman Valley Creek.....			300
Hoovey Creek.....			500
Merritt Creek.....			300
Neal Creek.....			300
Rock Creek.....			500
Rosman Creek.....			500
Silver Creek.....			300
Whelan Creek.....			300
Muscoda, Booth Hollow Creek.....			300
Byrds Creek.....			300
Indian Creek.....			600
Ludvick Branch.....			300
Sixmile Branch.....			300
Wall Branch.....			300
Nashville, Clear Lake.....			6,000
Spring Creek.....			3,000
New Auburn, Duncan Creek.....			1,000
Sand Creek.....			600
New Lisbon, White Creek.....			900
Oregon, Bodfish Creek.....			4,500
Pepin, Big Plum Creek.....			600
Bogus Creek.....			300
Elk Creek.....			300
Little Plum Creek.....			300
Lost Creek.....			300
Porcupine Creek.....			300
Roaring River.....			600

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Wisconsin—Continued.			
Phipps, McDermott Brook.....			1,500
Nemokagon River.....			6,000
Rogers Creek.....			4,500
Plymouth, Mullet Creek.....			10,000
Union River.....			3,000
Rice Lake, Angler Creek.....			300
Barker Creek.....			300
Big Bear Creek.....			1,600
Big Kettle Creek.....			1,000
Browns Creek.....			1,000
Cannon Creek.....			1,000
Cobb Creek.....			1,000
Cranberry Creek.....			300
Desair Creek.....			300
German Creek.....			1,000
Hay River.....			1,000
Heger Creek.....			300
Hemlock Creek.....			300
Kegamo Creek.....			1,000
Little Bear Creek.....			300
Little Spring Creek.....			300
Long Lake Stream.....			1,000
Meadow Creek.....			1,000
Miller Creek.....			300
Moosier Creek.....			1,000
Mud Creek.....			1,000
Olson Creek.....			1,000
Overby Creek.....			1,300
Pekegamo Creek.....			300
Prairie Creek.....			300
Renville Creek.....			1,000
Rice Creek.....			1,000
Savage Creek.....			1,000
Silver Creek.....			300
South Creek.....			300
Spoon Creek.....			1,000
Spring Creek.....			300
Spur Nine Brook.....			300
Sucker Creek.....			1,000
Weiss Creek.....			1,000
West Branch.....			300
Yellow River.....			300
Richland Center, Ash Creek.....			600
Fancy Creek.....			600
Little Willow Creek.....			600
Melancthon Creek.....			600
Pine River.....			200
Ridgeway, Mill Creek.....			1,000
River Falls, Kinnickinnic Creek.....			900
Nye Creek.....			600
South Fork River.....			900
Rosendale, Silver Creek.....			600
Solon Springs, Ox Creek.....			1,500
Sparta, Beaver Creek.....			300
Big Creek.....			1,100
La Crosse River.....			400
Little La Crosse River.....			400
Sargent Creek.....			300
Silver Creek.....			300
Soper Creek.....			300
Sparta Creek.....			300
Squaw Creek.....			300
Tarr Creek.....			300
Tuttles Creek.....			300
Walworth Creek.....			300
Spring Valley, Bahrs Creek.....			300
Burghardt Creek.....			300
Cady Creek.....			300
Cave Creek.....			600
Eagle Springs.....			300
French Creek.....			300
Gilbert Creek.....			1,200
Jacobson Creek.....			300
Johnson Creek.....			300
Lohns Creek.....			300
Lousy Creek.....			600
Mines Creek.....			300

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Wisconsin—Continued.			
Spring Valley, Rush River.....			1,200
Stanley, Hay Creek.....			4,000
Swim Creek.....			3,000
Stitzer, Bald Branch.....			4,500
Leggett Branch.....			4,500
Superior, Wisconsin Creek.....			6,000
Thorp, Bolin Creek.....			2,000
Lost Creek.....			2,000
Sterling Creek.....			3,000
Tomah, Brandy Creek.....			2,000
Council Creek.....			2,000
Deer Creek.....			1,000
Dodgeville Creek.....			2,000
Jennings Creek.....			2,000
Mill Creek.....			2,000
Mud Creek.....			1,000
Sand Creek.....			2,000
Silver Creek.....			1,000
Sparta Creek.....			2,000
Spring Bank Pond.....			1,000
Viola, Church Creek.....			3,000
Viroqua, Bishop Branch.....			4,000
Cotter Creek.....			2,000
Duck Eggs Branch.....			1,000
Pine Hollow Creek.....			1,000
See Branch.....			1,000
Warrens, Bettz Creek.....			300
Brandy Creek.....			300
Castle Rock Creek.....			300
Dampka Creek.....			300
Fish Creek.....			300
Harp Creek.....			300
Lowrie Creek.....			300
Matchett Creek.....			300
Sand Creek.....			300
Second Creek.....			300
Wausau, Jim More Creek.....			4,000
Wautoma, Beebe Creek.....			3,800
Birch Creek.....			600
Bird Creek.....			2,000
Chafee Creek.....			3,000
Lunch Creek.....			3,000
Pine Creek.....			2,000
Wedde Creek.....			2,000
White River.....			3,000
Westby, Bad Axe River.....			1,200
Clear Branch.....			300
Clockmakers Creek.....			2,300
Coon Creek.....			2,300
Crumo Spring Creek.....			2,000
Dauve Spring.....			300
Jown Vele Creek.....			1,000
Kickapoo Creek.....			1,200
Knapp Creek.....			1,300
North Bad Axe River.....			900
Paulsrud Creek.....			300
Paulson Creek.....			600
Sanding Creek.....			1,300
Seas Branch.....			2,000
Sherve Creek.....			1,000
Spring Coulee Creek.....			2,000
Spring Valley Creek.....			300
Sveen Creek.....			2,000
Timber Coulee Creek.....			2,000
Timber Valley Creek.....			300
Van Ruden Creek.....			2,300
West Salem, Adams Valley Creek.....			400
Bostwicks Valley Creek.....			400
Burns Creek.....			400
Cliff McClentock Creek.....			300
Gilles Coulee Creek.....			300
Green Creek.....			300
Holberg Creek.....			300
Johnson Creek.....			300
Jones Creek.....			600
Kincade Creek.....			300
Larson Creek.....			300

DETAILS OF DISTRIBUTION* OF FISH AND FISH EGGS—Continued.

BROOK TROUT—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Wisconsin—Continued.			
West Salem, Louis Valley Creek.....			400
Luce Creek.....			300
McEldowney Creek.....			600
Martin Creek.....			300
Memkings Creek.....			300
Rackley Creek.....			300
Raum Creek.....			300
Ruland Creek.....			300
Thronson Creek.....			400
Tousche Creek.....			300
Young Creek.....			300
Wheeler, Big Beaver Creek.....			1,000
Big Otter Creek.....			1,000
Blank Creek.....			1,000
Carey Creek.....			1,000
Hay River.....			1,000
La Forge Creek.....			1,000
Lambs Creek.....			1,000
Little Beaver Creek.....			1,000
Little Otter Creek.....			1,000
Page Creek.....			1,000
Whitehall, Barlow Valley Creek.....			300
Beaver Creek.....			300
Bruce Valley Creek.....			300
Elk Creek.....			600
Fly Creek.....			300
Hay Creek.....			300
Irvine Creek.....			300
North Valley Creek.....			300
Pigeon Creek.....			300
Wild Rose, Willow Creek.....			800
Willard, Cameron Creek.....			600
Tinning Horn Creek.....			500
Wilton, Adrian Creek.....			300
Beacher Creek.....			1,500
Hibbard Creek.....			300
Kinney Creek.....			300
Sinks Creek.....			3,300
Slaten Creek.....			300
Waage Creek.....			1,500
Winneboujou, Big Lake.....			2,000
Black Hoof Creek.....			4,000
Brule River.....			23,000
Cutler Creek.....			2,000
Govan Springs Pond.....			2,000
Lake Florence.....			4,000
Little Brule River.....			2,000
Lucius Lake.....			2,000
Miles Creek.....			2,000
Rock Creek.....			2,000
Sandy Run.....			2,000
Stones Creek.....			2,000
Wheatons Creek.....			2,000
Wyoming:			
Clark, Clarks Fork River.....			2,000
Green River, Green River.....			6,000
Kemmerer, Rock Creek.....			3,000
Lander, Cabin Lake.....			2,500
Hobbs's lake.....			2,500
Popo Agie River.....			2,500
Upper Lake.....			2,500
Laramie, Laramie River.....			5,500
Willow Creek.....			10,000
Manhattan, Spotted Tail Creek.....			12,500
Yellowstone National Park, Glen Creek.....			5,000
Indian Creek.....			15,000
Willow Creek.....			20,000
Japan:			
Tokyo, Japanese Government.....	5,000		
Total <i>a</i>	516,000	7,365,945	4,085,174

a Lost in transit, 23,600 fry and 158,687 fingerlings.

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

SUNAPEE TROUT.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
New Hampshire:			
Lake Sunapee, Lake Sunapee.....		115,029	
Newbury, Lake Sunapee.....		56,000	
Total.....		171,029	

GRAYLING.

Montana:			
Lakeview, Elk Creek.....		16,000	
Elk Lake.....		65,000	
Washington:			
Seattle, Exposition Aquarium.....			18
Wyoming:			
Sheridan, Bear Creek.....	25,000		
Total.....	25,000	81,000	18

SMELT.

Maryland:			
Mountain Lock, Potomac River.....			9,000
New York:			
Raquette Lake, Lake Kora.....	4,500,000		
Total.....	4,500,000		9,000

PIKE.

Iowa:			
Lime Springs, Upper Iowa River.....			700
North McGregor, Mississippi River.....			1,900
Minnesota:			
Brownsville, Mississippi River.....			18,650
Wisconsin:			
Genoa, Mississippi River.....			500
La Crosse, Mississippi River.....			19,650
Prairie du Chien, Mississippi River.....			1,900
Total.....			43,300

PICKEREL.

Wisconsin:			
Genoa, Mississippi River.....			166
La Crosse, Mississippi River.....			168
Victory, Mississippi River.....			166
Total.....			500

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

CRAPPIE AND STRAWBERRY BASS.

Disposition.	Finger- lings, year- lings, and adults.	Disposition.	Finger- lings, year- lings, and adults.
Arkansas:		Mississippi:	
Harrell, Spring Dale Pond.....	70	Booneville, Beach Bluff Lake.....	100
Helena, Blue Hole.....	7,000	Hollaway Lake.....	100
Long Lake.....	22,200	Red Elm Lake.....	200
Mississippi River.....	145,610	Columbus, Mullins Lake.....	100
Junction, Spring Lake.....	70	Corinth, Lake Billsville.....	250
Nashville, Mine Creek.....	250	Macon, Poplar Lake.....	100
Patmos, Mental Pond.....	100	Willow Glen Pond.....	100
Stamps, Mucille Lake.....	60	Noxapater, Estes's pond.....	100
Price Pond.....	185	Philadelphia, Spring Pond.....	100
Washington, Allen's pond.....	100	Tupelo, Sterns's pond.....	100
Connecticut:		West Point, Fortson Lake.....	100
Danbury, Kellogg's pond.....	250	Missouri:	
Wolf Pond.....	250	Aurora, Crane Creek.....	300
New Haven, Granniss Lake.....	200	Butler, Lake Katherine.....	100
Illinois:		Higginsville, Railroad Pond.....	275
Avena, Willow Lake.....	200	Mount Vernon, Honey Creek.....	300
Belleville, Club Pond.....	150	Hoshaw Lake.....	200
Heinemann's lake.....	400	Jaggerman Lake.....	200
Carbondale, Club Lake.....	200	Johnson's lake.....	200
Simons Lake.....	200	Spring River.....	400
Carterville, Peyton's pond.....	250	Nevada, Katy Allen Lake.....	100
Donnellson, Clover Leaf Lake.....	150	Springfield, Walnut Spring Lake.....	200
East Hannibal, Sni E'Carte River.....	750	Warrensburg, Meily's lake.....	100
Herrin, Manning Pond.....	500	West Plains, Carter's pond.....	100
Mine Pond.....	500	Willow Springs, Maple Pond.....	100
Hillsboro, Seymour Club Lakes.....	300	New York:	
Lake Forest, Whitehall Pond.....	320	Albany, Stevens's pond.....	100
Indiana:		Newark, Asylum Reservoir.....	100
Haubstadt, Oak Summit Pond.....	100	North Carolina:	
Lebanon, Bramble Gravel Pit.....	100	Hendersonville, Jane Mill Pond.....	200
Paoli, Willow Lake.....	100	Lake Osceola.....	300
Richmond, Crystal Lake.....	200	Rainbow Lake.....	150
Shell Brook Pond.....	200	North Dakota:	
Iowa:		Berlin, Rush Pond.....	55
Algona, Upper Des Moines River, East Branch.....	400	Fullerton, Appelquist Pond.....	100
Fort Madison, Green Bay.....	125	Glen Ullin, Sprecher's pond.....	100
Independence, Wapsipinicon River.....	400	Hankinson, Lake Elsie.....	200
North McGregor, Mississippi River.....	46,000	Lisbon, Prairie Farm Lake.....	100
Stockport, Silver's pond.....	100	Ohio:	
Kansas:		Bradford, Greenville Creek.....	350
Caldwell, Bluff Creek.....	1,000	Covington, Stillwater River.....	250
Farlington, Mitchell's pond.....	25	Gettysburg, Greenville Creek.....	250
Kentucky:		Winton Place, Hollywood Lake.....	200
Bradford, Locust Brook Pond.....	100	Oklahoma:	
Meadow Brook Pond.....	100	Alva, Harbaugh Lake.....	175
Campbellsburg, Sanford Pond.....	100	Apache, Morgan's ponds.....	100
Cropper, Willow Pond.....	100	Spring Pond.....	50
Emmons, Breezy Heights Pond.....	100	Sturman's pond.....	50
Lebanon, Graham's pond.....	150	Wogan's pond.....	50
Rogers's pond.....	150	Ardmore, Camp Brown Creek.....	400
Louisville, Cemetery Lake.....	100	Edward's pond.....	300
Lake Lansdowne.....	300	Hickory Creek.....	200
St. Mary, Forester Lake.....	200	Love's lake.....	300
Louisiana:		Silver Lake.....	400
Athens, Gandy's pond.....	100	Barron Fork, Yonah Pond.....	100
Marsalis Pond.....	100	Bliss, Arkansas River.....	200
Bernice, Chalybeate Spring Pond.....	70	Canute, Turkey Pond.....	150
Heard's pond.....	70	Chouteau, Bledsoe Pool.....	100
Keatchie, China Grove Lake.....	100	Cleveland, Silver Lake.....	100
Mansfield, Bickerstaff Lake.....	150	Cushing, Willow Pond.....	100
Brick Company's pond.....	100	Elgin, South Side Farm Pond.....	50
Many, Hoagland's pond.....	130	El Reno, Nettie Ruth Lake.....	300
Quitman, Harvey's pond.....	70	Fletcher, Cox Reservoir.....	150
Spring Lake.....	70	Gracemont, Walnut Grove Pond.....	100
Ruston, Hancock's pond.....	100	Marietta, Black Lake.....	50
Maryland:		McKinney's pond.....	65
Mountain Lock, Potomac River.....	247	Smith's pond.....	50
Prince George County, Goodloe's pond.....	100	Washington Lake.....	50
Minnesota:		Noble, Appleby's pond.....	50
Brownsville, Mississippi River.....	43,250	Oklahoma City, Deepwater Lake.....	150
Rochester, Zumbro Mill Pond.....	20	Fields's pond.....	175
Wheaton, Lake Traverse.....	200	Gaylord's pond.....	200

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

CRAPPIE AND STRAWBERRY BASS—Continued.

Disposition.	Finger- lings, year- lings, and adults.	Disposition.	Finger- lings, year- lings, and adults.
Oklahoma—Continued.		Texas—Continued.	
Oklahoma City, Gum's lake.....	200	De Kalb, Crump's pond	30
Turner's lake.....	100	Detroit, Oil Mill Pond	30
Oologah, Sunday's pond.....	100	Elgin, Elgin Lake.....	20
Pawhuska, Clear Creek.....	100	Elkhart, Elkhart Lake.....	100
Snyder, Deep Pond.....	150	Farwell, Hamlin Pond.....	31
Terral, Ewing's lake.....	100	Fort Worth, Lake Homewood.....	140
Yukon, Kralick Run.....	100	Garrison, Cedar Lake.....	50
Pennsylvania:		Giddings, Fisher's pond.....	65
Falls Station, Lake Winola.....	200	Jaehne's pond.....	30
York, Codorus Creek, South Branch..	150	Raube's lake.....	30
South Carolina:		Sumf's pond.....	25
Aiken, Black Poplar Pond.....	100	Symm's pond.....	30
Belton, Belton Mills Pond.....	100	Thonig Pond.....	30
Bishopville, Kelley's lake.....	125	Toepper's pond.....	25
Central, Arnold's pond.....	100	Volkers's pond.....	30
Chappells, Mills Pond.....	100	Graham, Norris's lake.....	106
Scurry Pond.....	100	Oak Grove Pond.....	50
Webb's pond.....	100	Worthington Knox Lake.....	50
Clover, Campbell's pond.....	75	Grand Saline, Malone Pond.....	20
Darlington, Creek Pond.....	150	Grapeland, Tyers Lake.....	50
Fair Forest, Fair Forest Creek.....	100	Willow Lake.....	30
Fountain Inn, Durbin Creek Pond.....	100	Groveton, Friday's pond.....	30
Greenville, Saluda Silver Lake.....	200	Nelms's lakes.....	80
Rembert, Evans's pond.....	125	Hamlin, Red Lake.....	20
Waterce, Griffin Creek Pond.....	100	Haysland, Lake Shelby.....	75
Yorkville, Turkey Creek Pond.....	100	Honey Grove, Fin and Feather Club Lake.....	100
Tennessee:		Jacksboro, Cooper Lake.....	50
Somerville, Allbright's lake.....	200	Mays Lake.....	20
Texas:		Jacksonville, Hillside Lake.....	75
Albany, Kellum's pond.....	30	Jordan Lake.....	75
Broyle's pond.....	30	Park Lake.....	75
Waterworks Pond.....	100	Sampson Lake.....	100
Annona, Capital Lake.....	40	Shearn Lake.....	75
Arlington, Jones's pond.....	30	Jonesville, Lake Sand Hill.....	100
Artesia, McWhorter's reservoir.....	20	Kaufman, Bond's pond.....	10
Athens, Gauntt's lake.....	15	Gilmore Lake.....	100
Koon Kreek Klub Lake.....	100	Hatch Pond.....	20
Prater's lake.....	20	Hindman's pond.....	20
Atlanta, Warren's lake.....	40	Sand Lake.....	20
Austin, Austin Lake.....	50	Taylor's pond.....	50
Slaughter Lake.....	100	Warenskjold Lave Lake.....	20
Windy Crest Lake.....	30	Kemp, Long Lake.....	100
Bay City, Austin's pond.....	20	Kent, Tatam's pond.....	25
Water Works Pond.....	20	Kerrville, Turtle Creek Pond.....	30
Beckville, Parker's lake.....	100	Lampasas, Collins's pond.....	20
Big Sandy, Big Sandy Lake.....	50	Lillian, Reese Branch Pond.....	40
Lake Everman.....	100	Llano, Llano Lake.....	315
Robinson's lake.....	50	Longview, Beale Lake.....	75
Blossom, Patton's pond.....	26	Fisher Lake.....	75
Brazoria, State Farm Lake.....	50	Lake Lomond.....	100
Canyon City, Paloduro Creek.....	100	Lovelady, Duck Lake.....	100
Spring Creek Lake.....	100	Kelley Pond.....	20
Carmona, Carmona Pond.....	40	McDade, Milton's pond.....	20
Carthage, Davis's lake.....	50	Manchaca, Bear Creek.....	50
Prior's pond.....	100	Marshall, Bentley Lake.....	30
Center Point, Guadalupe River.....	100	Bonita Lake.....	100
Verde Creek.....	100	Lake Ferns.....	100
Childers, Lake Scott.....	200	Thelma Lake.....	60
Clarksville, Clear Lake.....	50	Mart, Club Lake.....	150
Grassy Lake.....	100	Midlothian, Cooper's lake.....	40
Round Lake.....	100	Mineola, Goldsmith's pond.....	30
Coleman, Coleman Lake.....	100	Mineral Wells, Kearby Tank.....	25
Lost Creek.....	100	Mount Calm, Nelson Pond.....	10
Santa Anna Branch.....	100	Stovall Pond.....	40
Sunnyside Lake.....	75	Mount Selman, Brock's lake.....	30
Coolidge, Karner Lake.....	30	Mayfield's pond.....	20
Copperas Cove, Dewald's pond.....	20	Naples, Naples Club Lake.....	15
Corsicana, Corsicana Fish Association Pond.....	50	Walker's pond.....	30
Water Works Lake.....	40	Nash, Earnest's lake.....	100
Cotulla, Cartwright's reservoir.....	40	Normanna, Blackburn's pond.....	28
Counter Switche, Country Club Lake.....	175	Paige, Gropp Pond.....	30
Crockett, Daniel's lake.....	30	Horn's pond.....	20
Dallas, Munger's pond.....	20	Palestine, Bear Lake.....	200

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

CRAPPIE AND STRAWBERRY BASS—Continued.

Disposition.	Fingerlings, yearlings, and adults.	Disposition.	Fingerlings, yearlings, and adults.
Texas—Continued.		Texas—Continued.	
Palestine, Cartmell's lake	20	Tyler, Pine Hill Lake	30
East Side Park Pond	30	Tyler Pin Club Lake	100
Wallace Lakes	40	Waco, Katy Club Lake	100
Paris, Stannard's pond	20	Wills Point, Imperial Lake	100
Willow Lake	30	Virginia:	
Petty, Fielding Lake	100	Culpeper, Englands Mill Pond	200
Queen City, Prator's pond	20	Dillwyn, Fitzgerald Pond	125
Rockdale, Clear Lake	50	Fredericksburg, Boscobel Pond	500
Rotan, Willingham Pond	100	Leesburg, Goose Creek	300
Royston, Brooks's pond	30	Lynchburg, Murrell Pond	100
California Creek Lake	50	Midlothian, Midlothian Pond	100
Henry's tank	50	Natural Bridge, Cedar Creek	400
Stephens's tank	100	Petersburg, Belschers Pond	150
Saginaw, Kane's pond	35	Hauslik Pond	325
San Angelo, Concho River, Middle and		Spicer Pond	200
South Forks	133	Richmond, Crittenden Pond	200
Dove Creek	60	Darbytown Pond	200
Kickapoo Creek	133	Fulton Fishing Club Pond	200
Water Valley Country		Selden's pond	200
Club Lakes	74	Rockfish, Rockfish Lake	200
San Antonio, Lamm's tank	30	Scottsville, Chester Pond	100
Mitchell Lake	100	Soudan, Grass Creek	200
San Marcos, Blue Hole Pond	25	Suffolk, Lake Savage	2
Saron, William Lake	30	Sweet Briar, Sweet Briar Lake	200
Sulphur Springs, Elberta Lake	100	Winterpock, Indian Spring Pond	150
Picnic Lake	50	Zuni, Joyner's pond	200
Thomas Lake	50	Richardson's pond	200
Taylor, Roberts's lake	20	West Virginia:	
Temple, Lake Polk	75	Blueton, Holley's pond	150
Terrell, Bass Lake	20	Phillippi, Middle Fork River	400
County Club Lake	75	Salisbury, Salisbury's pond	200
Elm Pond	50	Wisconsin:	
Green Lake	20	Genoa, Mississippi River	5,832
Grinnan Pond	20	Independence, New City Pond	250
High Point Creek	75	Kewaskum, Beachwood Lake	200
Martin Pond	30	La Crosse, Mississippi River	49,086
Muckleroy Pond	40	Millston, Poley Creek	200
Sargent Pond	20	Mosinee, Half Moon Lake	350
Timpson, Bussey's pond	20	Prairie du Chien, Mississippi River	46,000
McWilliams's pond	20	State Line, Pickerel Lake	200
Tye, Crawford Lakes	25	Victory, Mississippi River	3,332
Tyler, Clear Spring Lake	50	Wausau, Lake Wausau	400
DeLay's lake	100	O'Day Lake	250
Lake Park Lake	100	Silver Creek Bay	400
Lakewood Country Club Lake	100		
Murphy's pond	30	Total a	410,428

ROCK BASS.

Alabama:		Illinois:	
Fivepoints, Poplar Springs	100	Belleville, Club Pond	100
Arizona:		Carbondale, Thompson's lake	300
Wilcox, McComb Ranch Pond	100	Donnellson, Cherry Grove Pond	100
Arkansas:		Wilson's pond	200
DeQueen, Gantlon's pond	500	DuQuoin, Egyptian Pond	100
Gravette, Dow's pond	250	McLeansboro, Goehring's pond	100
Harrison, Estes's pond	400	Indiana:	
Helena, Mississippi River	9,915	Bloomfield, Richland Creek	550
Mena, Irons Fork River	500	Boonville, Hemenway's pond	500
Mountain Fork River	500	Carlisle, Wellington Pond	150
Ouachita River	500	Cory, Prairie Lake	200
Prairie Creek	500	Woodland Lake	200
Rock Creek	500	Danville, Soper's pond	200
Twomile Creek	500	Evansville, Clear Pond	150
Pine Bluff, Trigg's pond	200	Stringtown Springs Pond	150
Connecticut:		Fairmont, Brookshire's pond	100
New Haven, Hubinger's lake	500	Fort Branch, Symond's pond	100
Georgia:		Greencastle, Lake Woodland	200
Etowah, Hill's pond	100	Greentown, Ayres's pond	100
Ringgold, Tiger Creek	300	Macy, Baker's pond	100

a Lost in transit, 9,049 fingerlings.

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

ROCK BASS—Continued.

Disposition.	Finger- lings, year- lings, and adults.	Disposition.	Finger- lings, year- lings, and adults.
Indiana—Continued.		Missouri—Continued.	
Plainfield, Spright's pond.....	500	Merwin, Corbin's ponds.....	3,000
Seymour, Bars Pond.....	200	Mount Vernon, Gillingham's pond.....	250
Kasting's pond.....	200	Skinner's pond.....	250
Summitville, McLain's pond.....	100	Tillotson's spring.....	200
Wawaka, Fountain View Pond.....	100	Truitts Creek.....	750
Winchester, Gravel Pit Pond.....	100	Williams Creek.....	500
Iowa:		Neosho, Twin Springs.....	500
Manchester, Maquoketa River.....	325	New Mexico:	
Kansas:		Ancho, Cooper's lake.....	200
Chanute, Durey Pond.....	100	Carlsbad, Dark Canon Creek.....	225
Cherokee, Allen Pond.....	150	Deming, Knowles's pond.....	100
Farlington, Mitchell's pond.....	150	Peterson's pond.....	100
Leavenworth, Park Lake.....	200	Ramsey's pond.....	100
Marion, Bruno Creek.....	50	Texico, Crescent Pond.....	100
East Creek.....	50	Tularosa, Silver Lake.....	200
French Creek.....	50	Vernigo Park, Adams Lake.....	300
Lyons Creek.....	50	New York:	
Medicine Lodge, Houchin's pond.....	100	Dover Plains, Lake Ellis.....	100
Kauffman's pond.....	100	Great River, Timber Point Pond.....	100
Peabody, Calbeck's pond.....	50	Middletown, Walkill Creek.....	400
Kentucky:		New Windsor, Walker's lake.....	100
Beaver Creek, Hindman Pond.....	125	North Carolina:	
May's pond.....	125	Carthage, Hannon's pond.....	150
Buechel, Blankenbaker's pond.....	200	Durham, Ellis's pond.....	75
Campbellsville, Creel's pond.....	175	Fayetteville, Cross Creek.....	50
Cropper, Turnpike Pond.....	100	Mollett Pond.....	50
Dover, Jennings Pond.....	150	Hendersonville, Lily Pond.....	150
Lebanon, McElroy's pond.....	175	Mebane, Lake Weda.....	75
Lexington, Lake Callahan.....	400	White Pond.....	75
Louisville, Parkview Club Lake.....	200	Salisbury, Josey's pond.....	75
Schroerluেকে's pond.....	200	Star, Hursey Spring Pond.....	150
Paris, Brannon's pond.....	100	Wake Forest, Walthonia Fish Club Pond.....	385
Clarke's pond.....	100	Weldon, Gooch's pond.....	75
Clay Pond.....	100	Ohio:	
Edwards Pond.....	100	Bidwell, Jones's pond.....	100
Frazier Pond.....	100	Blanchester, Reeves's pond.....	100
Grayson Pond.....	150	Chardon, Charlotte Pond.....	150
Hedge Pond.....	100	East Palestine, Freed's pond.....	200
Jackson's pond.....	100	Fronton, Sandusky River.....	500
Maher's pond.....	100	Fronton, Howell's pond.....	100
Paynes Pond.....	100	Kansas, Feasel Quarry Pond.....	100
Purnell's pond.....	100	Marion, Whetstone River.....	250
Vimont's pond.....	100	Springfield, Little Miami River.....	300
Watson Pond.....	100	Summit, Summit Lake.....	200
Wiggins Pond.....	100	Wickliffe, Morris Reservoir.....	100
Shawhan, Estes's pond.....	100	Oklahoma:	
Ewalt's pond.....	100	Chickasha, Harness Pond.....	150
Winchester, Twomile Creek.....	200	Crescent, Crescent Lake.....	50
Louisiana:		Osborn's pond.....	50
Arcadia, Boone's springs.....	100	Elgin, Glenn Pond.....	150
Grand Cane, Grand Cane Creek Pond.....	100	Guthrie, Hawley's pond.....	142
Homer, Gandy's pond.....	100	Highland Lake.....	141
Maryland:		Red Lake.....	142
Painsville, Quynn's pond.....	200	Hillsdale, Coldwater Creek.....	50
Monrovia, Cashour's pond.....	2,000	Lawton, Markeson's pond.....	250
Mountain Lock, Potomac River.....	2,010	Marlow, Jorgeson Pond.....	100
Thurmont, Hemler's pond.....	230	Newkirk, Lake Vanderpool.....	100
Michigan:		Santa Fe Lake.....	50
Bath, Park Lake.....	200	Okeene, Seigfried's pond.....	50
Minnesota:		Perry, Clear Lake.....	50
Rochester, Zumbro River, South Branch.....	200	Watson's pond.....	50
Willet's pond.....		Ponca, Bell Lake.....	50
Mississippi:		South Coon Creek.....	50
Guntown, Cochran's pond.....	100	Purcell, Brewer's lake.....	30
Pontotoc, Gardner's pond.....	125	Tryon, Bermuda Lakes.....	50
Highland Fish Co. Lake.....	100	Wanette, Laughlin's pond.....	100
Patterson's pond.....	125	Weatherford, Bear Creek Pond.....	30
Ripley, Keenin's pond.....	100	Pennsylvania:	
Missouri:		Birdsboro, Hay Creek.....	200
Butler, Lake Catherine.....	5,000	Bushkill, Delaware River.....	1,200
Glasgow, Steinmetz Pond.....	100	Indiana, Yellow Creek.....	300
Holmes, Dunlap's lake.....	100	Marion, Back Creek.....	400
Joplin, Wild Cat Spring.....	200		
Marshall, Stedem Pond.....	100		

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

ROCK BASS—Continued.

Disposition.	Finger- lings, year- lings, and adults.	Disposition.	Finger- lings, year- lings, and adults.
Pennsylvania—Continued.		Texas—Continued.	
Marion, Conococheague Creek.....	400	Grapevine, Hicks's pond.....	50
Rowlands, Lackawaxen River.....	600	Greenville, Birdsong Lake.....	100
Weissport, Big Creek.....	300	Swan Pond.....	40
Rhode Island:		Haskell, Cunningham's pond.....	100
Barrington Center, Wood's pond.....	200	Shook's pond.....	50
South Carolina:		Hico, Gilmore Creek.....	50
Blacksburg, Bear Creek.....	100	Joshua, Stephen's pond.....	200
Clover, Camp Run.....	100	Linden City, Dean's pond.....	50
Greenwood, Curtail Creek.....	100	Lufkin, Melville Delta Pond.....	100
Little Curtail Creek.....	100	Marfa, Barrel Springs Pond.....	50
Ridgeway, Hobby Lake.....	100	Mineola, Conger's pond.....	100
Rock Hill, Spring Ponds.....	200	Mount Vernon, Gardner's pond.....	50
Spartanburg, Moore's pond.....	100	Smith's pond.....	75
Starr, Branch Pond.....	100	Palestine, Spring Lake.....	100
Westminster, Branch Lake.....	100	Park Springs, Plum Pond.....	30
Woodruff, James Creek Pond.....	100	Rotan, Lake Cottonwood.....	100
Yorkville, Hart's pond.....	200	Tuxedo, Davis Lake.....	25
McNeil's pond.....	200	Waco, Fleming's pond.....	75
Tennessee:		Winchell, Hogland's pond.....	150
Chattanooga, Chickamauga Creek.....	200	Wimmsboro, Beggs's pond.....	200
Concord, Pepper's pond.....	200	Wolf City, Jones's pond.....	50
Gibson, Estes's pond.....	100	Utah:	
James's pond.....	100	Lund, Bur Oak Spring Pond.....	100
McMinnville, Sink Creek.....	400	Virginia:	
Murfreesboro, Stones River.....	400	Bumpass, Hill's pond.....	150
Paris, Russell's lake.....	100	Danville, McGuire's pond.....	300
Sparta, Cave Spring Pond.....	100	Hewlett, Duke's pond.....	200
Watauga Point, Buffalo Creek.....	500	Hurt, Dawson's pond.....	200
Texas:		Nace, Brugh's pond.....	150
Alpine, Jackson's pond.....	100	Natural Bridge, Cedar Creek.....	1,000
Anson, Hendrick's lake.....	40	Orange, Macon Spring.....	100
Aquilla, Vaughan's lake.....	75	Salem, Roanoke River.....	600
Blum, Mirror Lake.....	50	Scottsville, Moon's pond.....	600
Bowie, Waggoner Pond.....	50	Totier Creek Pond.....	600
Celina, Gearhart's pond.....	60	South Hill, Ferguson's pond.....	200
Chico, Largent's lake.....	40	Spout Springs, Webbs Pond.....	150
Comanche, Highland Lake.....	150	Stuart, Mothers Home Pond.....	200
Cooledge, Hardeman's pond.....	50	Tazewell, Wimmer's pond.....	250
Trinity and Brazos Valley Lake.....	100	Walkers Station, Vaidens Mill Pond.....	600
Crawford, Railroad Lake.....	100	Winchester, Back Creek.....	250
Cushing, Kinney's pond.....	50	Opequon River.....	250
Datura, Pritchard's pond.....	50	Woods Cross Roads, Valley Front Pond.....	150
Dublin, Johnson's pond.....	50	West Virginia:	
Easterly, Easterly's pond.....	75	Bruceon Mills, Kelley's pond.....	450
Edgewood, Brier Springs.....	75	Fairmont, Little Lakes.....	650
Fairlie, Martingon Pond.....	50	Fort Gay, Sweet Lake Pond.....	200
Franklin, Cedar Creek, West Fork.....	150	Wellsburg, Cross Creek.....	500
Duncan's pond.....	75	Wyoming:	
Love's pond.....	100	Sheridan, Cut Off Pond.....	300
Grapevine, Crowley's pond.....	100	Total ^a	66,035

WARMOUTH BASS.

Georgia:		Maryland:	
Chamblee, Jones's pond.....	40	Mountain Lock, Potomac River.....	752
		Total.....	792

^a Lost in transit, 7,360 fingerlings.

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

SMALL-MOUTH BLACK BASS.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Arkansas:			Maryland—Continued.		
Newport, Gamble Lake.....		2,000	Phoenix, Great Gunpowder River.....	2,000	
Watson Lake.....		2,000	Pinesburg, Potomac River.....	12,000	
Warren, Eagle Creek.....		2,500	Turnpike, Red Run.....	1,000	
Saline River.....		2,500	Massachusetts:		
Connecticut:			Congamond, Congamond Pond.....	750	
Wauregan, Moosup Pond.....	1,500		Halifax, Stetson Pond.....	900	
Quinebaug River.....	1,500		Kingston, Big Indian Pond.....	900	
Illinois:			Northampton, Highland House Lake.....	750	
Anna, Fairground Lake.....		1,000	Onset Junction, Flax Pond.....	900	
Bloomington, Heafer Lake.....		150	Webster, Peter Pond.....		300
Momence, Kankakee River.....		500	Webster Lake.....		300
Naperville, DuPage River, West Branch.....		200	Woods Hole, Watcha Pond.....	900	
Wilmington, Kanakee River.....		10,000	Michigan:		
Indiana:			Alpena, Long Lake.....	6,000	
Angola, Bass Lake.....	1,000		Au Sabie, Cedar Lake.....		400
Big Center Lake.....	1,000		Burr Oak, Hog Creek Lake.....	3,000	
Buck Lake.....	1,000		Clare, Bass Lake.....	1,500	
Clear Lake.....	1,000		Geroux Lake.....	1,500	
Elston Lake.....	1,000		Lake Dewey.....	1,500	
Falling Lake.....	1,000		Lily Lake.....	1,500	
Fox Lake.....	1,000		South Lake.....	1,500	
Hog Lake.....	1,000		Stevenson Lake.....	1,500	
Lake James.....	1,000		West Lake.....	1,500	
Lake Jimmerson.....	1,000		Clarion, Walloon Lake.....	10,000	
Little Silver Lake.....	1,000		Clyde, Fish Lake.....	1,500	200
Marsh Lake.....	1,000		Comins, Churchill Lake.....		400
Middle Center Lake.....	1,000		Dryden, Seven Ponds.....	1,500	
Pigeon Lake.....	1,000		Youngs Lake.....	1,500	
Silver Lake.....	1,000		East Tawas, Bass Lake.....	3,000	1,000
Snow Lake.....	1,000		Empire, Glen Lake.....		400
Batesville, Little Laughery Creek.....		180	Lake Florence.....		400
Bedford, Quarry Pool.....		500	Ewart, Garvison Pond.....		400
Bloomfield, Richland Creek.....		250	Fowlerville, School Lot Lake.....		400
Columbia City, Round Lake.....		300	Gaylord, Otsego Lake.....		1,000
Corydon, Big Indian Creek.....	1,500		Gogebic, Gogebic Lake.....		300
Fort Wayne, Cedar Creek.....	1,000		Harrisville, Cedar Lake.....	3,000	
Dunton Lake.....	1,000		Hubbard Lake.....	3,000	400
Lake James.....	1,000		Hart, Round Lake.....		400
Maumet River.....	1,000		Silver Lake.....		400
St. Joseph River.....	1,000		Hastings, Clear Lake.....	3,000	
St. Marys River.....	700		Leach Lake.....	3,000	
Viberg Lake.....	1,000		Long Lake.....	3,000	
Georgetown, Big Indian Creek.....	2,000		Middle Lake.....	3,000	
Goshen, Goshen Mill Pond.....		300	Pine Lake.....	3,000	
Greencastle, Big Walnut River.....	1,000		Hillman, Valentine Lake.....	3,000	
Deer Creek.....	1,000		Hillsdale, Bav Bees Lake.....	3,000	
Little Walnut River.....	1,000		Holly, Dickson Lake.....	1,500	
Indianapolis, Eagle Creek.....	2,540		Fish Lake.....	1,500	
Fall Creek.....	2,540		Ironwood, Beatons Lake.....		600
School Creek Pond.....	300		Langsford Lake.....		800
White River.....	3,240		North Lake.....		400
Lagrange, Royer River.....	225		Rowe Lake.....		600
Laporte, Pine Lake.....	300		Triplet Lake.....		400
Monticello, Monon River.....	345		Wolf Lake.....		800
Tippecanoe River.....	375		Kingsley, Hogsback Lake.....	2,000	
New Albany, Silver Creek.....	300		Munsey Lake.....	2,000	
Pendleton, Fall Creek.....	150		Rennie Lake.....	2,000	
Ray, Clear Lake.....	300		Spider Lake.....	2,000	
Rome City, Sylvan Lake.....	375		Lake George, Lake George.....	1,500	
Shelbyville, Big Blue River.....	2,000		Shingle Lake.....	1,500	
Kentucky:			La Rocque, Lake May.....	3,000	
Cadiz, Caney Creek.....	2,000		Lewiston, Twin Lake.....		1,000
Little River.....	2,000		Lincoln, McNally Lake.....	3,000	
Muddy Fork Creek.....	2,000		Trask Lake.....	3,000	
East View, Nolin River.....	2,500		Lupton, Sage Lake.....		400
Franklin, Sharps Creek.....	900		Mears, Silver Lake.....		400
Winchester, Goff's Lake.....	1,500		Middleville, Thornapple River.....	3,000	
Maine:			Millersburg, Barnhart Lake.....	3,000	
Fryeburg, Kezar Pond.....	1,600		Montague, Big Blue Lake.....		400
Winthrop, Lake Annabessacook.....	1,500		Muskegon, Big Black Creek.....		400
Lake Maranocook.....	4,500		Newaygo, Sylvan Lake.....		400
Maryland:			Oden, Crooked Lake.....	5,000	
Cropley, Potomac River.....	4,000		Omena, Dougherty Lake.....		400
Hagerstown, Potomac River.....	12,000		Orchard Lake, Cooley Lake.....	3,000	
Mountain Lock, Potomac River.....		1,000	Long Lake.....	3,000	
			Orion, Lake Orion.....	3,000	

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

SMALL-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Michigan—Continued.			Ohio—Continued:		
Pentecost, Sand Lake.....	2,800		West Milton, Stillwater River.....	1,500	
Pentwater, Pentwater Lake.....		400	Zanesville, Muskingum River.....		400
Rose Center, Mungers Lake.....		200	Oklahoma:		
North Buckhorn			Wyandotte, Sycamore Creek.....		200
Lake.....	1,500		Pennsylvania:		
South Buckhorn			Arcola, Perkiomen Creek.....		68
Lake.....	1,500		Carbondale, Crystal Lake.....		40
St. James, Barney Lake.....		400	Newton Lake.....		40
Shelbyville, Gun Lake.....	4,500		Water Company		
Topinabee, Mullet Lake.....		1,000	Dam.....		40
Twin Lake, Wallace Lake.....		400	Collegeville, Skipack Creek.....		50
Watersmeet, Bass Lake.....		300	Greenville, Shenango River.....		40
Crooked Lake.....		300	Harrisburg, Conodoguinet Creek.....		70
Witch Lake, Long Lake.....		300	Kratz, Perkiomen Creek.....		50
New Hampshire:			Lebanon, Big Swatara Creek.....		70
Claremont, Rocky Bound Pond.....	750		Farlings Creek.....		45
Peterboro, Cunningham Pond.....	1,500		Indian Town Creek.....		70
Pittsfield, Jeness Pond.....	1,500		Klimes Dam.....		70
New Jersey:			Lake Conewago.....		70
Blackwood, Blackwood Lake.....		200	Little Swatara Creek.....		70
Branchville, Culver Lake.....		200	Lowdermilk Dam.....		70
Lambertville, Lambertville			Mish Mill Dam.....		70
Reservoir.....		100	Mount Gretna Lake.....		70
Sewell, Chestnut Branch.....		125	Raccoon Creek.....		45
Sunset Lake.....		125	Stavers Mill Pond.....		70
Sterling Forest, Greenwood			Stover Lake.....		70
Lake.....	150		Weidman Dam.....		70
Sussex County, Lake Grinnell.....		100	Woomers Mill Pond.....		70
New York:			Lenape, Brandywine Creek.....		68
Auburn, Owasco Lake.....		500	Oil City, Allegheny River.....		50
Batavia, Godfrey Pond.....		40	Palm, Hosenack Lake.....		68
Horseshoe Pond.....		40	Leiberts Dam.....		50
Tonawanda Creek.....		200	Pottstown, Manatawany Creek.....		50
Binghamton, Susquehanna			Seranton, Cobbs Pond.....		50
River.....		40	Moosic Lake.....		50
Broadalbin, Kennyette Creek.....	5,000		Susquehanna, Butler Lake.....		40
Cambridge, Crystal Lake.....	5,000		Comfort Lake.....		40
Dead Pond.....	5,000		Susquehanna		
Lake Lauderdale.....	5,000		River.....		40
School House Pond.....	5,000		Troy, Sugar Creek.....		40
Fort Edward, Glen Lake.....	5,000		West Chester, Sharples' Lake.....		50
Highland, Long Pond.....	2,000		Wheelerville, Elk Lake.....		40
Johnstown, Caroga Lake.....	5,000		Rhode Island:		
East Caroga Lake.....	5,000		Kingston, Long Pond.....	1,500	
Mud Lake.....	5,000		Tucker Pond.....	1,500	
Kingston, Mohonk Lake.....	2,000		White Pond.....	1,400	
Mohonk Reservoir.....	2,000		Tennessee:		
Middletown, Walkill Creek.....		34	Denver, Trace Creek.....		6,000
Pelham, Hutchins Pond.....		150	McEwen, Hurricane Creek.....		3,000
Schenectady, Mariaville Lake.....	5,000		Waverly, Hurricane Creek.....		7,000
State Line, Queechee Lake.....	2,000		Vermont:		
Troy, Hudson River.....		400	Barnet, Martins Pond.....	6,000	
Water Mill, Howedona Lake.....		300	Wardens Pond.....	6,000	
West Point, Brooks Pond.....	2,000		Concord, Halls Pond.....	4,000	
Yonkers, Grassy Sprain Lake.....		300	Danville, Keaser Pond.....	5,000	
North Carolina:			Groton, Lake Groton.....	6,000	
Hendersonville, Lake Wajaw.....	200		Lunds Pond.....	6,000	
Mortimer, Johns River.....	150		Lyndonville, Bean Pond.....	5,000	
Mulberry Creek.....	200		Institute Pond.....	10,000	
Wilson Creek.....	150		Miles Pond, Miles Pond.....	4,000	
Ohio:			North Troy, Upper Missisquoi		
Columbus, Alum Creek.....	1,500	200	River.....		300
Black Lick Creek.....	1,500		Passumpsic, Passumpsic River.....	750	
Deer Creek.....	1,500		Poultney, Lake St. Catherine.....	5,000	
Hayden Run.....	1,500		Rutland, Lake Bomoseen.....	10,000	
Little Darby Creek.....	1,500		West Danville, Joe's pond.....	5,000	
Olentangy River.....	1,500	400	Wolcott, Wolcott Pond.....	5,000	
Rocky Fork Creek.....	1,500		Virginia:		
Scioto River.....	1,500		Ashby, Shenandoah River.....		350
Dayton, Mad River.....	3,000		Covington, Potts Creek.....		200
Miami River.....	3,000		Danville, Clarks Pond.....	3,000	
Stillwater River.....	3,000		McGuire's Ponds.....	8,000	
Delphos, Auglaize River.....	1,500		Drewrys Bluff, Falling Creek.....	9,000	
Germanatown, Big Twin Creek.....	3,000		Loudoun County, Potomac		
Marietta, Muskingum River.....		400	River.....		24,000
Newark, Raccoon Creek.....		600	Millboro, Cow Pasture River.....		12,000
Pleasant Hill, Stillwater River.....	1,500		Nathalie, Brown Pond.....		3,000
Portsmouth, Millbrook Park			Orange, Sharon Lake.....		1,000
Lake.....		400	Petersburg, Club Pond.....		3,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

SMALL-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Virginia—Continued.			West Virginia—Continued.		
Petersburg, Woody Pond.....	6,000	Roncevert, Greenbrier River.....		800
Providence Forge, Mirror Lake.	12,000	Sistersville, Middle Island Creek.		100
Relee, Relee Lake.....	1,000	Springfield, Potomac River,		
Remington, Rappahannock			South Branch.....	15,000
River.....	2,000	Wisconsin:		
Richmond, Falling Creek Pond.	3,000	Armstrong Creek, Lake Gordon.		300
Rockfish, Rockfish Pond.....	1,000	Lake Hilbert.....		300
Rockfish River.....	2,000	Cisco, Lake Tenderfoot.....		300
Warrenton, Cedar Run.....		300	Harshaw, Hancock Lake.....		300
Washington:			Hayward, Round Lake.....		800
Bellingham, Lake Wildwood.....	100	Hurley, Bear Lake.....		400
Loon Lake, Loon Lake.....	100	Okauchee, Okauchee Lake.....		488
Spokane, Liberty Lake.....	100	Princeton, Fox River.....		300
Tacoma, American Lake.....	100	Sobieski, Bass Lake.....		300
West Virginia:			Soperton, Otter Lake.....		300
Capon Springs, Great Cacapon			Spider, Spider Lake.....		500
River.....	45,000	State Line, Little Bass Lake.....		300
Harpers Ferry, Potomac River.		1,200			
Renick, Cuberson Creek.....	9,000	Total ^a	537,400	109,986
Greenbrier River.....	24,000			

LARGE-MOUTH BLACK BASS.

Alabama:			Colorado—Continued.		
Montgomery, Brick Yard Lake.....	1,000	Lamar, Neenoshe Lake.....		320
Whetstone Lake.....	2,000	Neeskah Lake.....		320
Seale, Evans's pond.....	2,000	Neesopah Lake.....		320
Arizona:			Parrish's lake.....		320
Flagstaff, Lake Mary.....	300	Thurston Lake.....		320
Tucson, Cienga Creek.....	300	Thurston Reservoir.....		320
Arkansas:			Littleton, Springer's pond.....		300
Bearden, Crystal Lake.....	150	Manzanola, Lewis' reservoir.....		450
Bentonville, Sugar Creek.....	500	Pueblo, Squirrel Creek Reser- voir.....		100
England, Clear Lake.....	400	Riffe, Bear River.....		150
Fairfield, Atkins Lake.....	350	Grand River.....		300
Helena, Blue Hole.....	1,000	Connecticut:		
Long Lake.....	1,800	Coscob, Pipestave Lake.....		250
Mississippi River.....	7,323	Danbury, Bradley's pond.....		225
Hope, Moses's lake.....	100	Weekapeeka Lake.....		300
Sandy Bois d'Arc River.....	250	East Hampton, Pocotopaug Lake.....		400
Lake Village, Lake Chicot.....	1,150	Goodspeeds, Bashan Lake.....		390
Lancaster, Frog Bayou.....	500	Higganum, Higganum Reser- voir.....		300
Little Rock, Asylum Pond.....	100	New Canaan, Lake Waccobuc.....		200
Mammoth Spring, Strawberry Creek.....	500	North Stonington, Wyassup Lake.....		390
Warm Fork.....	300	Waterbury, White Oak Pond.....		260
Mena, Big Brushy Creek.....	300	Weathersfield, Goff Pond.....		250
Big Fork Creek.....	350	Delaware:		
Carter Creek.....	300	Milton, Parkers Pond.....		300
Clear Creek.....	350	Parker Run.....		100
Cossatot River.....	300	Tea! Mill Pond.....		200
Dallas Creek.....	300	District of Columbia:		
Irons Fork River.....	300	Washington, Central Station Aquarium.....		150
Jansen Lake.....	300	Florida:		
Little Brushy Creek.....	300	Ehren, Muller's pond.....		500
Little Missouri River.....	350	Lake Como, Lake Como.....		500
Little Rock Creek.....	300	Ocala, Fry Lake.....		2,000
Mountain Fork River.....	300	Orlando, Smith's lake.....		500
Ouachita River.....	600	Sanford, Lake Bertha.....		500
Prairie Creek.....	300	Santos, Lake Madonna.....		500
Two Mile Creek.....	300	Sorrento, Lake Lucy.....		1,000
Paris, College Lake.....	500	Georgia:		
Rosboro, Caddo Pond.....	100	Douglas, Peterson's ponds.....		1,750
Scott, Old River.....	540	Greenville, Powers Hill Pond.....		1,500
Thornton, Pine Lake.....	125	Groveland, Cannoche River.....		1,000
Upland, Brazeal's pond.....	30	Lake Park, Long Pond.....		1,000
Colorado:			Ocean Pond.....		500
Boulder, Pitts' pond.....	150	Marietta, McKenzie's pond.....		500
Denver, Holliday's lakes.....	150	Mayfield, Cason's pond.....		100
La Jara, Laguna Escondida.....	480			
La Junta, Holbrook Reservoir.....	1,000			
Lamar, King Lake.....	320			
Neegrando Lake.....	320			

^aLost in transit, 3,319 fingerlings.

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Georgia—Continued.			Indiana:		
Millen, Buck Head Creek.....		1,000	Anderson, Bayview Pond.....		400
Ogeechee River.....		1,000	West Brook Pond.....		300
Oglethorpe, Buck Creek.....		750	Claypool, Caldwell Lake.....		250
Talbotton, Williams Pond.....		250	Simon Shultz Lake.....		200
Tifton, Hale's pond.....		500	Yellow Creek Lake.....		250
Valdosta, Loch Laurel.....		500	Corydon, Big Indian Creek.....		300
Vienna, Heard's pond.....		250	Elbert's lake.....		100
Idaho:			DeLong, Tippecanoe River.....		700
Nampa, Lake Lowell.....		250	Eaton, Hamilton's pond.....		125
Priest River, Lees Pond.....		100	Fort Wayne, Lake Emily.....		300
Illinois:			Indianapolis, Eagle Creek.....		100
Antioch, Lake Marie.....		1,200	Fall Creek.....		200
Aurora, Fox River.....		900	Nesom's pond.....		75
Belleville, Biebel's pond.....		250	White River.....		200
Fourmile Club Lakes.....		200	Jasper, Calumet Lake.....		300
Beech Ridge, Cache River.....		550	Schmitt's pond.....		100
Brighton, Kelsey's pond.....		300	Kendallville, Bixler Lake.....		200
Montgomery Lake.....		250	Liberty, White Water River,		
Cairo, Cache River.....		550	East Fork.....		375
Campus, Factory Pond.....		500	Macy, North Mud Lake.....		400
Carbondale, Cox's lake.....		200	South Mud Lake.....		400
Manning Pond.....		800	Monticello, Big Metamonong		
Mine Pond.....		800	Creek.....		300
Spillers Lake.....		300	Tippecanoe River.....		40
Thompsons Lake.....		400	New Albany, Silver Lake.....		300
Carter, Wellman's lake.....		300	North Liberty, Rupel Lake.....		100
Cartersville, Brandon Pond.....		100	Owensville, Stone's pond.....		100
Carroll's pond.....		300	Paoli, Brookside Reservoir.....		200
Carter Pond.....		150	Pierceton, Webster Lake.....		50
Coleman Pond.....		175	Richmond, Rettig Lake.....		45
Colp and Arnold			Rockville, Little Raccoon Creek,		
Lakes.....		150	Rome City, Lower Lake.....		400
Ferrell Pond.....		150	Sylvan Lake.....		200
Hofer Lake.....		100	Stewartsville, Footes Lake.....		200
Zimmerman's lake.....		350	Summitville, Roseboom's pond.....		400
Chester, Crisler's pond.....		350	Warren, Salamonie River.....		400
Fishing Club Lake.....		300	Iowa:		
Clay City, Doherty's pond.....		100	Bentley, Walnut Hill Pond.....		125
Crainville, Norton's pond.....		100	Cedar Falls, Cedar River.....		400
Crystal Lake, Crystal Lake.....		800	Hacketts Lake.....		400
Dallas City, Mississippi River.....		900	Chariton, McCoy's pond.....		100
Decatur, Club Lake.....		80	Rice Lake.....		600
Franklin, Burlington Reservoir.....		300	Charles City, Cedar River.....		400
Freeburg, Freeburg Lake.....		200	Chester, Upper Iowa River.....		800
Walnut Grove Pond.....		500	Clarion, Elm Lake.....		400
Glenwood, Glenwood Pond.....		1,000	Coggon, Buffalo Creek.....		400
Grays Lake, Gages Lake.....		1,000	Corning, Lake Vernon.....		100
Herrin, Cambon Pond.....		300	Decorah, Upper Iowa River.....		400
Egyptian Pond.....		450	DeWitt, Crystal Lake.....		400
Homewood, Calumet River.....		500	Silver Creek.....		200
Kankakee, Iroquois River.....		1,200	Edgewood, Funk's pond.....		150
Kankakee River.....		1,200	Forest City, Imogene Lake.....		150
Kansas, Hallock's lake.....		200	Glenwood, Glenwood Park		
Kewanee, Sans Souci Lake.....		750	Lake.....		700
Makanda, Roberts's pond.....		100	Hampton, Reed Lake.....		400
Marion, Hart's pond.....		175	Independence, Wapsipipicon		
Keystone Pond.....		150	River.....		400
Schwerdt's lake.....		400	Lime Springs, Upper Iowa		
Water Works Lake.....		150	River.....		2,250
Modoc, Bersche's pond.....		200	Manchester, Maquoketa River.....		7,100
Mulberry Grove, Hudson Pond.....		300	Marble Rock, Shell Rock River.....		400
Murphysboro, Stacher Lake.....		200	Maynard, Little Volga Creek.....		300
Naperville, Du Page River,			North McGregor, Mississippi		
East Branch.....		1,000	River.....		5,250
Stone Quarry Lake.....		450	Tuskego, Robertson's pond.....		100
O'Fallon, Henrys Lake.....		400	Kansas:		
Olney, Olney City Reservoir.....		400	Belmont, Bentley's pond.....		100
Richmond, Lake Elizabeth.....		800	Blue Rapids, Big and Little		
Riverside, Des Plaines River.....		900	Blue River.....		300
Sandusky, Round Pond.....		100	Bronson, Second Lake.....		100
Shepherd, Sni E'Carte River.....		600	Caldwell, Fall Creek.....		500
Sterling, Sinsippi Lake.....		900	Chanute, Valley View Pond.....		100
Ulin, Cache River.....		550	Cherryvale, City Lake.....		300
Utica, Fourth Quarry Pond.....		450	Colony, Clark's pond.....		125
Vandalia, Kaskaskia River.....		500	Conway Springs, Slate Creek.....		300
Virden, Maple Avenue Lake.....		300	Farlington, Mitchell's pond.....		125
Waterloo, Bissell Lake.....		600	Huron, Anthony's pond.....		225
Woodberry, Woodberry Lake.....		500	Isabel, Gibson's pond.....		100

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Kansas—Continued.			Kentucky—Continued.		
Kansas City, Idlewild Lake.....		100	Stephensburg, Blue Lake.....		200
Kingman, Harris Springs Pond.....		200	Stephensburg		
Reed's pond.....		100	Lake.....		200
Leavenworth, Fairgrounds Lake.....		150	Williamsburg, Jellico Creek.....		300
Marion, Catlin Creek.....		250	Louisiana:		
Clear Creek.....		250	Athens, Dullon Pond.....		100
Cottonwood River,			Benton, Sunnyside Pond.....		100
South Fork.....		250	Bogalusa, Bogalusa Pond.....		250
Middle Creek.....		250	Bowie, Hill Pond.....		50
Mud Creek.....		250	Broussard, Hazard Pond.....		25
Willowbrook Pond.....		100	Clinton, Gallent's pond.....		100
Medicine Lodge, Chapin Ponds.....		125	Edgerly, Chesson's pond.....		50
Currie Lake.....		125	Jeauerette, Albania Pond.....		75
Read Lake.....		250	Lake Charles, Brickyard Pond.....		25
Silver Springs			Laurel Hill, Rose Mound Lake.....		150
Lake.....		250	Lillie, Pin Oak Pond.....		30
Melvorn, Long Creek.....		250	Marthaville, Huff's pond.....		100
Peabody, Cotton Creek.....		100	Rustin, Lyles's pond.....		125
Country Club Lake.....		250	Maine:		
Crisfield Pond.....		250	Boothbay Harbor, Pine Lake.....		450
Doyle Creek.....		350	Redfield, Parker Pond.....		155
Gray's pond.....		250	Maryland:		
Henry Creek.....		250	Abell's Wharf, Forbes Pond.....		150
Johnson's pond.....		100	Alesia, Big Gunpowder River.....		320
Rock Island Lake.....		250	Gunpowder Falls.....		180
Spring Creek.....		350	Baltimore, Severn River.....		280
Townsend's pond.....		250	Brunswick, Potomac River.....		130
Pittsburg, Sporting Club Ponds.....		125	Cumberland, Potomac River.....		160
St. Francis, Spring Creek.....		75	Wills Creek.....		80
Selden, Prairie Dog Creek.....		325	Easton, Peach Blossom Creek.....		150
Tyro, Brick Company's Lake.....		150	Freeland, Rock Dale Ponds.....		280
Waverly, Rock Creek.....		200	Gwynnbrook, Gwynn Brook.....		100
Wilder, Woodson's pond.....		100	Hagerstown, Antietam Creek.....		150
Yates Center, Waterworks Reser- voir.....		250	Conococheague		
Kentucky:			Creek.....		200
Anchorage, Cox Lake.....		150	Potomac River.....		400
Pryor's pond.....		75	Hampstead, Patapsco River,		
Augusta, Licking River, North			North Branch.....		340
Fork.....		80	Hoods Mill, Patapsco River.....		270
Bonnieville, Riggs's pond.....		75	Lambson, Sassafras River.....		300
Campbellsburg, Little Ken- tucky River.....		200	Massey, Swan Branch.....		150
Ekron, Horse Lot Pond.....		100	Mountain Lock, Potomac River.....		440
Woods Pond.....		100	Phoenix, Gunpowder River.....		140
Yellow Lake.....		100	Riverdale, Anacostia River.....		100
Elizabethtown, Cedar Creek.....		80	Rocky Ridge, Monocacy River.....		300
Nolin River.....		80	Salisbury, Wicomico River.....		200
Rauboldt Pond.....		40	Taneytown, Goulden's pond.....		80
Valley Creek.....		80	Woodstock, Patapsco River.....		360
Youngers Creek.....		80	Massachusetts:		
Eminence, Thorne's pond.....		75	East Dedham, Mather Brook		
Glasgow, Beard Pond.....		75	Pond.....		250
Boyd's Creek.....		75	Fall River, Laurel Lake.....		390
Fallen Timber Creek.....		200	Falmouth, Morse Pond.....		250
Peters Creek.....		200	Greenfield, Deerfield River.....		1,500
Richardson Pond.....		150	Wareham, Big Sandy Pond.....		400
Skeggs Creek.....		200	Little Sandy Pond.....		400
South Fork Creek.....		75	West Gloucester, Haskell's pond.....		390
Glendale, Nolin Creek.....		150	Michigan:		
Hodgensville, Nolin Creek.....		80	Alpena, Grand Lake.....		350
La Grange, Highland Lake.....		150	Crystal Falls, Fortune Lake.....		200
Lebanon, Big Pond.....		75	Lake Mary.....		200
Cheyls Creek.....		75	Mud Lake.....		200
Indian Creek.....		75	Edwardsburg, Morn Creek.....		350
Peeps Creek.....		150	Greenville, Flat River.....		1,000
Rolling Fork Creek.....		150	Tufk Lake.....		200
Rolling Fork Creek,			Hanover, Crispell Lake.....		150
North Branch.....		150	Farewell Lake.....		150
Rolling Fork Creek,			Fox Lake.....		100
South Branch.....		150	Hart, Juniper Pond.....		175
Louisville, Green's pond.....		80	Ironwood, Long Lake.....		400
Lake Lansdowne.....		195	Mosquito Lake.....		400
Parkview Club Lake.....		80	North Lake.....		200
South Park Lake.....		160	Pomeroy Lake.....		400
Wagner's pond.....		80	Round Lake.....		200
			Silver Lake.....		400
			Sutherland Lake.....		400

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Michigan—Continued.			Mississippi—Continued.		
Ironwood, Tamarack Lake.....		400	Corinth, Lambert's lake.....		150
Taylor Lake.....		400	Long Pond.....		150
Ishpeming, Silver Lake.....		400	Marlows Mill Pond.....		100
Kingsley, Hogsback Lake.....		375	Parmithie Creek.....		200
Rennie Lake.....		375	Romine and Ward		
Oakley, Shiawassee River.....		200	Pond.....		150
Oden, Crooked Lake.....		700	Santa Fe Lake.....		150
Schoolcraft, Weed Lake.....		175	Seven Mile Creek.....		200
Sylvania, Katherine Lake.....		200	Tuscumbia River.....		300
West Bear Lake.....		200	Utley Mill Pond.....		150
Turtle, African Lake.....		400	Waukomis Lake.....		300
Clover Leaf Lake.....		200	Wilson's pond.....		100
Eel Lake.....		200	Dancy, Barefoot's pond.....		100
Emiline Lake.....		200	McCarter's pond.....		150
Gaylord Lake.....		200	Smith's pond.....		150
Hawk Lake.....		400	Walker's pond.....		100
Honey Moon Lake.....		400	White's pond.....		150
Independence Lake.....		400	Wilson's pond.....		100
Line Lake.....		400	Durant, Smith's pond.....		200
Mint Lake.....		200	Friars Point, Moon Lake.....		250
Moose Lake.....		400	Houlka, Reed's pond.....		25
Orms Lake.....		400	Houston, Busby's pond.....		150
Rowes Lake.....		400	Knox Pond.....		50
Toe Lake.....		200	Howells Switch, Rankin Pond.....		400
Minnesota:			Jackson, Curry's pond.....		300
Alexandria, Darling Lake.....		200	Farish Pond.....		150
Lake Agnes.....		150	Lewis's pond.....		100
Lake Carlos.....		700	Lynch's pond.....		150
L'Home die u			Morrison's pond.....		150
Lake.....		300	Richmond Lake.....		300
Brownsville, Mississippi River..		3,000	Spring Lake.....		300
Duluth, White Lake.....		400	Tapley's pond.....		150
Kelsey, Lake Rauppe.....		450	Lee County, King Creek.....		650
Mankato, Lake Washington.....		400	McCool, Fancher's pond.....		300
Minneapolis, Burnett's lake.....		300	Lily Pond.....		150
Pengilly, Swan Lake.....		900	Sweet Gum Lake.....		150
Preston, Root River.....		600	McDonald, Majure's pond.....		25
Root River, Middle			Ogletree's pond.....		25
Branch.....		600	Smith's pond.....		25
Rochester, Zumbro River, Mid-			Maben, Butler's pond.....		150
dle Branch.....		200	Macon, Eiland Pond.....		150
Zumbro River,			Howards Lake.....		200
South Branch.....		200	Madison Station, Glenarchen		
St. Paul, State Fish Commis-			Pond.....		100
son.....		18,250	Mantee, Lofton's pond.....		150
South Haven, Augusta Lake.....		400	Moseley Pond.....		200
Betsy Lake.....		400	Taylor's pond.....		150
Lake Caroline.....		400	Meridian, College Lakes.....		300
Stewartsville, Lake Florence.....		500	Pleasant Springs.....		45
Root River.....		600	Queen City Club		
Mississippi:			Pond.....		200
Aberdeen, Dead Lake.....		25	New Albany, Conner's pond.....		200
McNiece Lake.....		300	New Houlka, Chuquatonchee		
Medor Lake.....		600	Creek.....		150
Tombigbee River.....		275	DeLashmet Lake.....		150
Ackerman, Willow Pond.....		200	Houlka Creek.....		300
Agricultural College, McKell's			Reed Pond.....		200
pond.....		100	Okolona, Elliott Pond.....		200
Bexley, Leatherberry Mill Pond..		75	Mill Pond.....		200
Mill Pond.....		75	Okolona Lake.....		200
Biloxi, Howell Pond.....		75	Red Bud Creek.....		150
Lorenzo Pond.....		75	Sansom's lakes.....		600
Brandon, Raymond Pond.....		100	Osborn, Montgomery's pond.....		100
Canton, Factory Pond.....		100	Oak Grove Pond.....		100
McBride Pond.....		100	Pearson, Sweetwater Lake.....		200
Round Lake.....		100	Philadelphia, Wilson's pond.....		50
Columbus, Lake Katherine.....		300	Pickayune, Tate's lake.....		75
Corinth, Bridge Creek.....		200	Ripley, Morgan's pond.....		25
Cane Creek.....		300	Sallis, Temple's pond.....		200
Chambers Creek.....		300	Sessums, Ash Creek Pond.....		100
Clear Creek.....		200	Gay's pond.....		100
Clear Lake.....		150	Rush's pond.....		100
Conway Lake.....		200	Wild's pond.....		100
Coon Creek Pond.....		200	Shuqualak, Belle Pond.....		200
Derryberry Lake.....		100	Dugan Pond.....		100
Elams Creek.....		150	Hamilton's pond.....		100
Griffins Pond.....		150	Jenkins' pond.....		75
Gum Pond.....		200	Woodlawn Pond.....		200

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Mississippi—Continued.			Nevada:		
Starkville, Harmon Lake.....		200	Ely, Argus Lake.....		250
Johnson's pond.....		100	Cleveland Lake.....		250
McPherson Lake.....		100	New Jersey:		
Reynolds Lake.....		100	Ashland, Newtown Lake.....		300
Richey's pond.....		100	Boonton, Deckers Pond.....		400
Wade's pond.....		150	Collingswood, Newton Lake.....		600
Washington's pond.....		100	Denville, Sanitarium Lake.....		300
Strongs, Cox Branch.....		100	Dover, Longwood Lake.....		500
Lake Artecore.....		100	Shongum Lake.....		400
Tofulla Creek.....		200	Lakewood, Lanes Mill Pond.....		200
Sturgis, Hutchinson Pond.....		100	Lambertville, Lower Reservoir.....		250
Summit, Godbold's lake.....		250	Mullica Hill, Mullica Hill Pond.....		400
Toomsba, Live Oak Lake.....		35	Netcong, Bear Pond.....		200
Tupelo, Mill Pond.....		100	Ogdensburg, Hawthorne Lake.....		400
Park Lake.....		300	Paterson, Squaw Lake.....		400
Wiygul's lake.....		150	Penns Grove, Du Pont Pond.....		100
Union, Johnson's pond.....		25	Layton Pond.....		350
West Point, Evans Pond.....		150	Pompton Lakes, Pompton Lakes		800
Tibbee Lake.....		300	Princeton Junction, Carnegie		
Tipton's pond.....		150	Lake.....		500
Yazoo City Cedar Grove Pond.....		150	Rahway, Water Company's		
Missouri:			reservoir.....		500
Asbury, Blackberry Creek.....		200	Riverside, Beck's pond.....		250
Aurora, Flat Creek.....		400	Sicklerville, Brooklyn Lake.....		600
Bolivar, Pomme de Terre River.....		300	Sewell, Bethel Lake.....		300
Brandsville, Lake of the Four			South Vineland, Buckshietem		
Cantons.....		100	Mill Pond.....		400
Butler, Lake Katherine.....		400	Waterloo, Jefferson Lake.....		200
Cabool, Piney River.....		200	Wenonah, Pyle's lake.....		250
Clever, Bailey's lake.....		200	Westwood, Musquapsink Lake.....		400
Estes's pond.....		400	New Mexico:		
Clinton, Clinton Lake.....		300	Artesia, Clark's lake.....		250
Cole Camp, Cole Camp Creek.....		300	Carlsbad, Pecos River.....		500
Corkney, Niangua River.....		150	Rocky Arroyo Creek.....		150
Creve Coeur, Creve Coeur Lake.....		225	Cofax, Adams Lake.....		254
Dedwick, Livingston's pond.....		100	Dexter, Bishop's lake.....		150
Deepwater, Dickey Lake.....		150	Gallup, Ramah Reservoir.....		300
Fredericktown, St. Francis			Hagerman, Ware's reservoir.....		150
River.....		200	Las Vegas, Buena Vista Lake.....		280
Grand View, Spring Lake.....		100	Santa Fe, Miller's pond.....		100
Higginsville, Railroad Pond.....		475	Springer, Farmers Reservoir.....		320
Kansas City, Fairmount Lake.....		400	Jaritas Lake.....		320
Koshkonong, Lake Rowland.....		100	Wagon Mound, Santa Clara		
Knoblick, Little St. Francis			Creek Reservoir.....		195
Creek.....		140	New York:		
Langdon, Langdon Lake.....		300	Arcade, Crystal Lake.....		400
Maysville, Dieter's lake.....		150	Cambridge, Second Pond.....		400
Mexico, Railroad Lake.....		200	Craryville, Copake Lake.....		400
Water Works Reservoir.....		200	Dover Plains, Lake Ellis.....		400
Mount Vernon, Truitt Creek.....		300	East Worcester, Hudson Lake.....		400
Neosho, Crescent Pond.....		200	Gloversville, Mountain Lake.....		400
Nevada, Railroad Reservoir.....		200	Greatkills, Shore Acres Pond.....		150
Noel, Perry's ponds.....		200	Greene, Chenango River.....		400
Pleasant Hill, Leonards Lake.....		500	Greenport, Sills Pond.....		100
Richards, Richardson's pond.....		100	Highland Falls, Roe Park Lake.....		200
Rolla, Big Beaver Creek.....		80	Hudson, Hasbrouck Pond.....		400
Big Dry Fork Creek.....		150	Huntington, Koster's pond.....		100
Little Beaver Creek.....		100	Johnstown, Canada Lakes.....		400
Little Dry Fork Creek.....		100	Lockport, Red Creek.....		300
Love Creek.....		100	Middletown, Walkkill River.....		300
McBride Spring Branch.....		40	Monticello, Anawana Lake.....		200
Waltz Spring Branch.....		40	Brown Pond.....		200
Rosedale, Lewis's pond.....		40	Highland Lake.....		200
Springfield, Doling Lake.....		300	Klamesha Lake.....		200
Swope Station, Lagoon Lake.....		200	Metock Pond.....		200
Wooded Lake.....		200	Sacket Lake.....		200
Thayer, Warm Fork Creek.....		200	Sand Pond.....		200
Wayne, Woodruff Springs.....		300	White Lake.....		200
Waynesville, Gasconade River.....		150	Narrowsburg, Half Moon Lake.....		400
West Plains, Woolworth's			Nunda, Genesee River.....		400
bayou.....		200	Raquette Lake, Blue Mountain		
White River,			Lake.....		400
North Fork.....		200	Eagle Lake.....		400
Willow Springs, Willow Springs			Utawana Lake.....		400
Reservoir.....		200	Riverside, Big Pond.....		500
Nebraska:			Bullett Pond.....		500
Stuart, Clear Lake.....		200	Paradox Lake.....		500

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
New York—Continued.			Ohio—Continued.		
Riverside Schron Lake.....		500	Newton Falls, Mahoning River.....		250
Roscoe, Florence Lake.....		400	Nova, Railroad Reservoir.....		200
Thurman, Echo Lake.....		400	Paulding, Maumee River.....		500
Ticonderoga, Eagle Lake.....		400	Portsmouth, Millbrook Park Lake.....		500
Walden, Walkkill River.....		300	Rarden, Scioto Brush Creek.....		225
Walkkill, Schawangunk River.....		400	Ravenna, Lake Brady.....		300
Warwick, Wickham Lake.....		400	Ripley, Gardner's pond.....		75
Williamstown, Panther Lake.....		400	Rock Creek, Grand River.....		125
North Carolina:			St. Marys, Mercer County Res- ervoir.....		600
Charlotte, Catawba River, North Fork.....		96	Salem, Crumrine Dam.....		125
Franklin, Cartoogaja Creek.....		405	Springfield, Buck Creek.....		125
Cullasagee Creek.....		405	Warren, Youngs Run.....		150
Tennessee River.....		300	Wauseon, Miller and Becker Pond.....		295
North Dakota:			Woodsfield, Woodsfield Dam.....		200
Ambrose, Skjermo Lake.....		300	Youngstown, Lake Cohasset.....		125
Annamossee, Round Lake.....		400	Lake Katrine.....		50
Berlin, Cottonwood Creek.....		400	Mahoning River.....		175
Cottonwood Pond.....		100	Oklahoma:		
Bottineau, Lake Dana.....		300	Ada, Boggy Lake.....		250
Lake McArthur.....		300	City Lake.....		325
Burnstad, Beaver Lake.....		400	Lawrence Lake.....		175
Buttzville, Buttz's pond.....		300	Radka Creek.....		250
Cathay, Rocky Run Lake.....		300	Ames, Garden Lake.....		100
Cayuga, Anderson's lake.....		100	Jones's lake.....		100
Crystal Springs, Crystal Springs Lake.....		500	Apache, Cache Creek.....		300
Dawson, Lake Isabel.....		400	Chandler Creek.....		400
Devils Lake, Devils Lake.....		3,500	Gassoway's lake.....		250
Elliott, Lake Elliot.....		200	Mission Creek.....		300
Glen Ullin, Antelope Creek.....		100	Newcomb Pond.....		100
Burns Pond.....		100	Sturman's pond.....		100
Granville, Buffalo Lodge Lake.....		600	Ta-La Creek.....		250
Gwinner, Aliceton Lake.....		200	Toney Creek.....		300
Denning's lake.....		100	Ardmore, Ardmore Club Lake.....		250
Johnson's pond.....		100	Caddo Creek.....		200
Harvey, Sheyenne Lake.....		400	Club Lake.....		175
Jamestown, James River.....		10,500	Twin Lake.....		100
Kenmare, Des Laes Lake.....		300	Atoka, City Reservoir.....		300
Thompson Lake.....		300	Barron Fork, Owl Lake.....		300
Lisbon, Bale's pond.....		200	Bernardi, Bogardus Pond.....		100
Sheyenne River.....		600	Blanchard, Bridge Creek.....		150
Milnor, Storm Lake.....		600	Spring Lake.....		150
Nicholson, Jackson Hill Pond.....		150	Bliss, Lake 101.....		150
Nome, Carlson's pond.....		100	Ranch Lake.....		100
Pingree, James Lake.....		400	Broken Arrow, Prairie Lake.....		100
Pipestem River.....		1,100	Calumet, Mae Lake.....		125
Ray, Beaver Creek.....		150	Carney, Carney Lake.....		100
St. John, Cameron's lake.....		300	Chattanooga, Sunnyside Lake.....		100
Jarvis Lake.....		300	Checotah, Spring Lake.....		140
Strium, Medd's pond.....		100	Chickasha, Lanier Pond.....		150
Ohio:			Chilocco, Chilocco Lagoon.....		100
Alexandria, Raccoon Creek.....		50	Crescent, Kelly's pond.....		100
Aurora Station, Harmon Pond.....		100	Devol, Suter's pond.....		125
Bradford, Greenville Creek.....		575	Duncan, Bumpass's lake.....		100
Celina, Mercer County Reservoir.....		500	Norvell's pond.....		125
Cleveland, Swimming Pond.....		100	Elk City, Chambers's lake.....		100
Cloverdale, Myers's pond.....		100	Lake Coleman.....		200
Covington, Factory Pond.....		175	El Reno, Club Lake.....		200
Greenville Falls Dam.....		250	Enid, Clear Lake.....		250
Mohlers Eddy.....		300	Gross's pond.....		250
Stillwater River.....		500	Spring Lake.....		250
Defiance, Auglaize River.....		150	Eufaula, Lake Buford.....		140
Maumee River.....		150	Faxon, Cuddy Lake.....		100
Findley, Auglaize River.....		150	Fort Sill, Medicine Bluff Creek.....		250
Fremont, Sandusky River.....		400	Frederick, Ater Lake.....		125
Georgetown, Sunny Side Lake.....		50	Glencoe, Lake Louisa.....		100
Hebron, Buckeye Lake.....		775	North Side Pond.....		100
Kent, Twin Lakes.....		300	Granton, Alfalfa Pond.....		100
West Twins Lake.....		300	Prairie Pond.....		100
Lisbon, Furnace Run.....		100	Willow Pond.....		100
Furnace Run Reservoir.....		100	Guthrie, Ellison Lake.....		250
Nelsonville, Hocking River.....		200	Johnson's pond.....		250
Newark, Buckeye Lake.....		425	Martin Lake.....		250
Newcomerstown, Tuscarawas River.....		250	Reddington Lake.....		250
New Paris, White River, East Fork.....		125	Twin Lakes.....		250
			Walker Lake.....		250
			Hallett, Mirror Lake.....		200

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Oklahoma—Continued.			Oklahoma—Continued.		
Haskell, Oputtuna Pond.....		100	Poteau, Long Lake.....		300
Willows Pond.....		300	Purcell, Club Lake.....		300
Hennessey, Jarvis's pond.....		100	Ripley, Crain's pond.....		100
Hobart, Elk Lake.....		150	Sallisaw, Sallisaw River.....		300
Hydro, Deer Creek.....		250	Sentinel, Big Elk River.....		350
Jet, Saline Valley Pond.....		100	Stillwater, Carpenter's lake.....		100
Kelsey, Illinois River.....		300	McKinnon's pond.....		100
Lawton, Medicine Creek.....		650	Stroud, Loeh Kathrine.....		125
McAlester, Cole's lake.....		150	Sulphur, Lowrance Lake.....		200
Madill, McMillan Lake.....		125	Tahlequah, Wolfe Lake.....		100
Marietta, Bills Creek.....		125	Terral, Rock Island Lake.....		100
Cochron Creek.....		250	Tishomingo, Big Sandy River.....		200
Coreoran Creek.....		175	City Lake.....		100
George William Creek.....		125	Foley Lake.....		200
Haynes's lake.....		125	Little Sandy River.....		250
Hickory Creek.....		250	Trousdale, Livvix's lake.....		100
Kirkpatrick Lake.....		125	Tuttle, Davis's pond.....		100
Marietta Club Lake.....		200	Vinita, Electric Park Lake.....		100
Oil Creek.....		200	Hall's lake.....		100
Rock Creek.....		150	Walter, Johnson's pond.....		100
Shegan Creek.....		125	Watonga, Cunningham's lake.....		300
Simon Lake.....		150	Waukomis, McClennahan's pond.....		100
Marlow, Adkins Pond.....		100	Woodward, Reilly's springs.....		100
Boone Pond.....		100	Yukon, Maixner's pond.....		100
Cooper's pond.....		100	Pennsylvania:		
Findley's pond.....		150	Bath, Spring Reservoir.....		100
Marlow Park Lake.....		150	Big Bend, Conewago Creek.....		300
Marlow Pond.....		150	Fleuent Pond.....		300
Martin's pond.....		125	Birdsboro, Hay Creek.....		350
Murray's pond.....		150	Brillharts, Cadorus Creek, South Branch.....		420
Quinn's lake.....		150	Bushkill, Deer Lake.....		300
Sand Hill Pond.....		200	Forest Lake.....		300
Shaws Pond.....		100	Lake Taminant.....		300
Waldbridge Lake.....		125	Mud Pond.....		300
Mill Creek, Mill Creek.....		300	Chester Springs, Pickering Creek.....		300
Muskogee, Country Club Lake.....		300	Collegeville, Willow Hurst Dam.....		100
Newkirk, Santa Fe Lake.....		250	Connellsville, Indian Creek.....		80
Ninnekah, Nelson Lake.....		125	Danville, Susquehanna River.....		175
Noble, Clear Brook.....		100	Susquehanna River, North Branch.....		150
Wadley's pond.....		125	Denver, Cocalico Creek.....		250
Norman, Sunnybrook Lake.....		150	East Berlin, Conewago Creek.....		350
Oehelata, Water Works Reser- voir.....		200	Factoryville, Lake Carey.....		300
Okeene, Schallmo Pond.....		100	Lake Kewanna.....		300
Oklahoma City, Belle Isle Lake.....		300	Lake Manataka.....		300
Club Lake.....		200	Falls Station, Susquehanna River.....		350
Coleord's lake.....		250	Fort Washington, Sandy Run.....		200
Elm Lake.....		125	Gettysburg, Marsh Creek.....		250
Hogan's pond.....		200	Rock Creek.....		250
Kingkade's lake.....		250	Goldsboro, Susquehanna River.....		280
Lakeview Lake.....		300	Graftesford, Perkiomen Creek.....		300
Shepherd's lake.....		375	Greenville, Shenango River.....		300
Spring Creek.....		125	Hanover, Conewago Creek.....		200
Osage, Osage Lake.....		150	Little Conewago Creek.....		200
Pawhuska, Clear Creek.....		350	Hatboro, Little Neshaminy Dam.....		200
Pawnee, Walenciak's lake.....		100	Hickory, Allegheny River.....		350
Perkins, Jennings Pond.....		100	Huntingdon, Raystown Branch.....		180
Perry, Beers's lake.....		200	Indiana, Twolick Creek.....		150
Bostiek's pond.....		200	Kimberton, French Creek.....		300
Brown's pond.....		200	Lancaster, Conestoga River.....		300
Casey's pond.....		200	Mount Morris, Dunkard Creek.....		1,000
City Lake.....		500	New Oxford, Little Conewago Creek.....		250
Hansen's pond.....		200	Newtown, Neshaminy Creek.....		600
Hansing's lake.....		100	Oaks, Perkiomen Creek.....		200
Keaton's pond.....		200	Skippack Creek.....		200
McCune's pond.....		250	Oxford, Octoraro Creek, East Branch.....		500
Moore's pond.....		175	Palm, Gehard Dam.....		100
Tucker's pond.....		125	Hosensaek Creek.....		100
Ponea, Cottonwood Lake.....		200	Perkiomen Creek.....		100
Evans Lake.....		200	Phillipsburg, Lehigh River.....		200
Roekbound Lake.....		200			
Turkey Creek.....		325			
Willow Pond.....		150			
Pond Creek, Fairview Lake.....		250			
Guernsey's lake.....		250			

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Pennsylvania—Continued.			South Carolina—Continued.		
Pittsburg, Griffin Reservoir.....		180	Clover, Allison Creek.....		1,000
Wildwood Reservoir.....		270	Beaver Dam Creek.....		1,000
Pocono, Naomi Lake.....		300	Bigger's pond.....		500
Pocono Lake.....		350	Catawba Creek.....		1,000
Port Royal, Tuscarora Creek.....		180	Catawba River.....		2,000
Pottstown, Manatawney Creek.....		150	Crowders Creek.....		4,000
Rahns, Perkiomen Creek.....		300	Crowders Mill Pond.....		1,000
Reading, Angelica Creek.....		200	Lower Beaver Dam		
Jordan Creek.....		200	Creek.....		1,000
Maiden Creek.....		250	Mill Creek.....		1,000
Schuylkill River.....		350	Upper Beaver Dam		
Stony Creek.....		200	Creek.....		1,000
Tulpehocken Creek.....		1,000	Columbia, Cedar Creek.....		36
Schwenksville, Perkiomen			Congaree Creek.....		36
Creek.....		300	Cotton Mills Reser-		
Susquehanna, Susquehanna			voir.....		48
River.....		400	Dents Pond.....		96
Telford, Perkiomen Creek,			Gin Pond.....		48
Northeast Branch.....		200	Poplar Branch Pond.....		36
Temple, Ontelaunie Creek.....		300	Rodgers Spring.....		36
Troy, Cross Roads Creek.....		250	Croft, Bridge Pond.....		500
North Branch.....		300	Darlington, Charles Mill Pond.....		500
Trunkysville, Alleghany River.....		300	Easley, Silver Pond.....		1,000
Tunkhannock, Lake Carey.....		800	Eastover, Colonels Creek.....		1,000
Union City, Lake Pleasant.....		350	Edgefield, Beaverdam Creek.....		1,000
Warren, Jackson Creek.....		300	Edmund, Thresher Pond.....		500
Weissport, Poho Poco Creek.....		250	Eureka, Seizers Mill Pond.....		500
West Chester, Park's pond.....		200	Everett, Hilliard Pond.....		500
Wrightsville, Cabin Creek.....		140	Old Mill Pond.....		500
Fishing Creek.....		200	Fort Lawn, Abernathy's pond.....		500
Krentz Creek.....		200	Catawba River.....		1,500
Susquehanna			Crawfords Pond.....		500
River.....		300	Fishing Creek.....		1,000
York, Beaver Creek.....		140	Gilbert, Hamburg Branch.....		48
Big Conewago Creek.....		560	Great Falls, Catawba River.....		1,000
Codorus Creek, South			Catawba River		
Fork.....		140	Pond.....		2,000
Codorus Creek, West			Rocky Creek.....		1,000
Fork.....		280	Southern Power		
Fishing Creek.....		140	Co.'s pond.....		1,000
Fox Creek.....		280	Greenville, Saluda Lake.....		4,000
Keesey Dam.....		140	Greenwood, Bag Creek.....		75
Kreutz Creek.....		140	Curl Tail Creek		
Kreutz Pond.....		140	Pond.....		120
Little Badams Creek.....		280	Cutler Branch		
Little Conewago Creek.....		140	Pond.....		75
Susquehanna River.....		280	Davis's pond.....		75
York Haven, Big Conewago			Garys Pond.....		25
Creek.....		280	Harrison Creek.....		75
Conewago Creek.....		560	Johns Creek.....		75
Susquehanna			Little Curl Tail		
River.....		280	Creek.....		135
Zieglersville, Perkiomen Creek.....		300	Rays Pond.....		75
Rhode Island:			Wardlaws Pond.....		1,000
Kingston, Hundred Acre Pond.....		520	Hartsville, Ox Pen Branch.....		500
Westerly, Park Lake.....		390	Hickory Grove, Bullock Creek.....		1,000
South Carolina:			Honea Path, Broad Mouth		
Aiken, Branch Pond.....		75	Creek.....		150
Shaws Creek.....		500	Little Creek.....		75
Anderson, Branch Water Pond.....		48	Little River.....		75
Brown Pond.....		48	Mattison Mill		
Silver Lake.....		48	Pond.....		75
Angelus, Middleton's pond.....		500	Saluda River.....		75
Belton, Saluda River.....		96	Turkey Creek.....		75
Bethune, Estridge's pond.....		500	Hopkins, Chappelle Creek.....		1,000
Mill Branch Pond.....		500	Mill Creek.....		1,500
Mill Creek Pond.....		500	Tub Mill Creek.....		1,000
Blacksburg, Broad River.....		1,000	Inman, Ray's pond.....		500
Blaney, Black Lake.....		1,000	Lamar, Harrell Mill Pond.....		500
Borden, Pollard Mill Pond.....		70	Lancaster, Mosier's pond.....		500
Bowling Green, Crowders Creek.....		500	Langley, Power House Pond.....		150
Crowders Creek,			Laurens, Reedy River.....		48
South Fork.....		1,000	Leesville, Lightwood Creek		
Calhoun, Twenty-three Mile			Pond.....		75
Creek.....		2,000	Lightwood Pond.....		500
Camden, Savage's pond.....		25	Lexington, Gable's pond.....		500
Chester, Sandy River.....		48	Marietta, Middle Saluda River.....		1,500
Clinton, Enoree River.....		500	North Saluda River.....		2,500

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposit on.	Fry.	Finger- lings.
South Carolina—Continued.			South Dakota—Continued.		
Marietta, South Saluda River...		1,000	Phillips, Harding Grove Dam...		125
Montmorenci, Runtz Creek.....		75	Plankinton, James's lake.....		150
Mullins, Buck Swamp.....		1,000	Saunders's lake.....		300
North Augusta, Walkers Mill Pond.....		150	Redfield, Twin Lakes.....		300
Oakvale, Oakvale Lake.....		25	Sisseton, Lake Traverse.....		400
Orangeburg, Spring Lakes.....		1,000	Minnesota River.....		300
Pageland, Black Pond.....		500	One Road Lake.....		300
Little's pond.....		500	Strand Creek.....		150
Spring Pond.....		500	White Stone Lake.....		300
Thompson's pond.....		500	Wilcox Creek.....		150
Patrick, Big Juniper Creek.....		1,000	Springfield, Emanuel Creek.....		300
Pellon, Black Creek.....		500	Tabor, Rezac Lake.....		300
Beaver Pond.....		500	Tripp, Herr's lake.....		200
Pickens, Saluda River, South Fork.....		1,500	Van Metre, Inland Lake.....		125
Twelvemile River.....		1,000	Sun Flower Dam.....		125
Ridge Springs, Flatrock Creek.....		500	Watertown, Lake Pelican.....		400
Gunter's pond.....		500	Tennessee:		
Rock Hill, Catawba River.....		2,000	Austral, Childress Creek.....		150
Little Allison Creek.....		48	Towey Creek.....		150
St. Matthews, Milwood Pond.....		500	Chatanooga, Chickamauga Creek, East Branch.....		200
Zeigler's pond.....		75	Chickamauga Creek, North Branch.....		200
Santuck, Broad River.....		500	Jetts Pond.....		200
Selvern, Indian Branch.....		500	Lookout Creek.....		200
Sharon, Bullock Creek.....		48	Spring Creek.....		200
Silverstreet, Beaverdam Creek.....		1,000	Cleveland, Candas Creek.....	2,400	
Springfield, Goodland Creek Pond.....		150	Greater Wildwood Lake.....		200
Steadman, Barr Pond.....		500	Hall's pond.....	800	
Gantt's pond.....		500	Wildwood Lake.....	2,400	
Sunter, Cains Mill Pond.....		150	Clinton, Clinch River.....		200
Pocalla Springs Pond.....		500	Moore's pond.....	800	
Trenton, Bottis's pond.....		500	Coal Creek, Coal Creek.....	1,600	
Chevis Creek Pond.....		500	Cool Creek.....		150
Pace Run.....		1,000	Conasauga, Jack River.....		
Shaws Pond.....		1,000	Curryhee, Little River, East Fork.....	2,065	
Walkers Pond.....		500	Knoxville, Little Pigeon River, East Fork.....	2,055	
Troy, Clinkscales's pond.....		25	Pigeon River, East Fork.....	2,055	
Cane Creek.....		1,000	Loudon, Little Tennessee River.....		200
Cuffy Town Creek.....		1,000	McGhee, Eagle Lake.....		150
Dowtin's pond.....		25	Memphis, Toney Pool.....		105
Hardlabor River.....		1,000	Newcomb, Elk Fork Creek.....		200
Leard's pond.....		25	Oakdale, Emory River.....		300
Long Cane Creek.....		3,000	Tenna, Conasauga River.....		150
Talbert's ponds.....		50	Townsend, Little River.....	3,425	
Young's pond.....		500	Texas:		
Union, Buffalo Reservoir.....		48	Albany, Honeycutt's pond.....		100
Yorkville, Brown's pond.....		1,000	Roseland Lake.....		100
Catawba River.....		1,000	Alto, Bailey Pond.....		100
Clarks Fork Pond.....		1,000	Alvarado, Rentfro's pond.....		100
Inman's pond.....		500	Amarillo, Paladora Pond.....		900
Langdon Branch Pond.....		1,000	Angus, Stewart's pond.....		400
Turkey Creek Pond.....		1,000	Annona, Hill's pond.....		200
Woodruff, Chumley's pond.....		500	Kickapoo River.....		500
South Dakota:			Arp, Hughes's pond.....		200
Astoria, Oak Lake.....		300	Athens, Shelton Mill Pond.....		400
Bonesteel, Flurans Lake.....		250	Austin, Barton Creek.....		500
Canton, Big Sioux River.....		800	Avoca, Martin's pond.....		200
Carthage, Lake Magnuson.....		175	Axtell, Biggerstaff's pond.....		100
Clark, Antelope Lake.....		300	Bellevue, Ford Lake.....		400
Round Lake.....		300	Bennetts, Cat Tail Lake.....		150
Dell Rapids, Big Sioux River.....		400	Bettie, Sewell's pond.....		100
Forestburg, Watch Lake.....		125	Bland Lake, Bland's pond.....		800
Kimball, Pleasant Lake.....		300	Blossom, Blossom Club Pond.....		150
Lane, Flowing Wells Lake.....		175	Boerne, Cibola Pond.....		150
Lennox, Lake Thorsen.....		300	Bowie, Black Pond.....		200
Madison, Lake Herman.....		500	Waggoner Pond.....		100
Lake Madison.....		600	Brady, Live Oak Creek.....		200
Marion, Center Lake.....		300	Brandon, Giles's lake.....		400
Silver Lake.....		300	Bronson, Clear Lake.....		300
Vermillion River, West Branch.....		300	Travis Branch.....		950
Midland, Stafford's pond.....		125			
Oakton, Stang's pond.....		200			
Parker, Dorow's pond.....		100			

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Texas—Continued.			Texas—Continued.		
Brookesmith, Buena Vista Lake		100	Elgin, Keeble's lake		100
Brownsville, Horseshoe Resaca Lake		1,000	Elkhart, Elkhart Lake		2,000
Resaca de la Guerra Lake		1,000	Pate's pond		50
Brownwood, Allison's pond		200	Encinal, Johnson Lake		500
Camp's pond		150	Fluvanna, Little Bull Pond		300
Collins's pond		200	Fort Worth, Concrete Pond		32
McGaugh Pond		200	Davie Burns Lake		50
Snyder's pond		1,000	Happy Lake		300
Bryan, Nall's lake		150	Hush Lake		213
Buckholtz, Helmcamp Pond		25	Lake View		300
Calallen, Casa Morado Reservoir		160	Lake Wandry		213
Calvert, Calvert Country Club Lake		500	Tandy's lake		200
Canyon City, Canyon Lake		600	Franklin, Cavitt's pond		200
Paladora Creek		725	Frisco, Stewarts Creek Lake		500
Pritchard's pond		600	Gainesville, Gainesville Club Lake		600
Terra Blanco Creek		725	Garrison, Brickyard Reservoir		100
Carlos, Lake Carlos		800	Fishing Club Lake		500
Caro, Lower Saner Pond		150	Giddings, Braesel's pond		100
Celina, English Lake		329	Carmean's pond		200
Moore's lake		300	Dunk's pond		200
Smith's lake		350	Gily Lake		50
Stelzer's pool		125	Mitschkes Pond		100
Center, Wood Lake		300	Namkin's pond		100
Center Point, Medina River		1,500	Quarry Lake		150
Childress, Lake Keeler		1,150	Raube's pond		200
Lake Scott		500	Schautschick's pond		100
Clarendon, Allan Creek		300	Schkades Pond		150
Clarksville, Clarksville Country Club Lake		500	Sumf's pond		100
Cleburne, Cleburne Country Club Lake		200	Unger's pond		150
Willow Pond		100	Wilson's pond		100
Clifton, Christenson's lake		112	Gladwater, Phillips Spring Lake		150
Reeder's pond		112	Goldthwaite, Cain's pond		200
Clyde, Deadman Pond		150	Gordon, Lake Creek		1,000
Colmesneil, Lively's lake		150	Goree, Goode's lake		300
Colorado, McCreless's lake		200	Granbury, Lake Add-Ran		200
Plasted's pond		600	Roberson Creek		185
Spring Creek Pond		300	Grand Saline, Dunn Mill Pond		100
White Elephant Lake		300	Grandview, Country Club Lake		1,200
Cooledge, Cottonwood Lake		100	Sturges's pond		150
Long Branch Lake		200	Grapeland, Hodge's lakes		600
McReynolds's reservoir		50	Keen Crystal Pond		500
Valley Lake		201	Grapevine, Willey Lake		150
Corsicana, Burks Lake		1,000	Yancy Lake		200
Morse's lake		200	Greenbrier, Beckham Pond		400
Woodley Pond		500	Butler Pond		400
Cotulla, Chapman Lake		400	Country Club Lake		400
Poteet Lake		400	Indian Creek		400
Crowell, Burress's pond		300	Leek Creek		400
Campbell's pond		150	Mud Creek		400
Railroad Pond		400	South Side Lake		500
Cuero, Hickory Lake		1,500	Hamlin, Country Club Lake		800
Cushing, Becton Lake		50	Harry Wynn Pond		200
Dale, Eppright Pond		200	Harlingen, Dilworth Lake		500
Dalhart, Rita Blanca Lake		200	Harlingen, Harlingen Lake		500
Dallas, Bachman Pond		375	Harrold, Ayers's pond		360
Coombs Creek		775	Haskell, Bevers's lake		300
Tenison Lake		300	Hico, Fairview Lake		150
Decatur, Halsell Lake		100	Gilmore Creek		200
DeKalb, Hathcocks's pond		300	Higgins, First Creek		300
Del Rio, Devils River		500	Poor Farm Lake		150
Denison, Lake Denison		800	High Island, Smith's lake		1,000
Denton, Country Club Lake		300	Hillsboro, Park Lake		400
Detroit, Clarksville Club Lake		400	Hubbard, Jones's lake		100
Detroit Club Lake		150	Leftwich Lake		200
Sample's pond		100	Jacksboro, Spring Pond		100
D'Hanis, Clay Hill Pond		300	Sunny Brook Lake		100
Doucette, Pope's pond		400	Joaquin, Garrett's pond		150
Stewart's lake		200	Kaufman, Clark Lake		200
Eagle Pass, Rosita Creek		1,000	Pyle's lake		816
Eastland, Kinnebrew Pool		300	Sapp's pool		200
Edgewood, Davis Pond		150	Taylor's pond		634
Elgin, Christian Lake		150	Willow Springs		200
Egleston Lake		134	Kemp, Berry Lake		300
			Moorehead Lake		300
			Porters Bluff Lake		400
			Kingsville, Christenson's reser- voir		100

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Texas—Continued.			Texas—Continued.		
Kyle, Goforth Pond.....		150	Oakwoods, Glaze Lake.....		800
Ladonia, Burton's pond.....		300	Palestine, Huff Lake.....		900
Elliott's pond.....		300	Spring Park Lakes.....		600
Water Works Pond.....		300	Panhandle, West Dippon Creek.....		500
LaGrange, Crownover Lake.....	1,500		Paris, Bankhead Lake.....		400
LaMarque, Irrigation Reservoir.....	1,500		Gordon Country Club Lake.....		1,000
Laredo, Bulls Eye Lake.....		500	Oak Grove Lake.....		150
Davis's pond.....		300	Silver Lake.....		100
Moritas Lake.....		500	Pecos, Edward's pond.....		175
Perren's pond.....		400	Pawket's pond.....		150
Lillian, Ball's pond.....		150	Penelope, Sealy Pond.....		200
Lillian Lake.....		150	Pilot Point, Lake Feeley.....		100
Lindale, Roberts's pond.....		150	Pittsburg, Adair Pond.....		100
Llano Grande, Llano Grande Lake.....		1,000	City Lake.....		200
Llano, Llano River.....		5,000	Davis Club Lake.....		150
Shumake's pond.....		50	Ferndale Lake.....		1,000
Longview, Harris's lake.....		400	Flag Pond.....		300
Melton's lake.....		200	Flannagan Pond.....		1,000
Taylor's pond.....		300	Hargrove Pond.....		150
Lovelady, Patterson Lake.....	1,000		Holt Pond.....		300
Lyford, Bamboo Lake.....		100	Hopkins's lake.....		30
McGregor, Leon River.....		500	Knights Mill Pond.....		1,000
South Bosque Creek.....		400	Lilly Pond.....		200
Mabank, Caruthers's pond.....		200	Music Pond.....		200
Cockerell's pond.....		54	Pilk Lake.....		200
Grubb's pond.....		150	Reves Lake.....		1,000
Hebel's pond.....		200	Reynolds Lake.....		1,000
McCoy's pond.....		200	Star Lake.....		200
Pepper's pond.....		200	Titule Lake.....		300
Robertson's pond.....		250	Willow Lake.....		305
Wind Mill Pond.....		200	Plano, City Reservoir.....		1,000
Madison, Donaho's pond.....		50	Queen City, Griffin's pond.....		30
Mahl, Pleasant Hill Lake.....		75	Randolph, Randolph Pool.....		300
Watkins's pond.....		50	Ranger, Water Works Lake.....		1,000
Malakoff, Bartlett's pond.....		100	Ravenna, Eubanks's pond.....		150
Brickyard Pond.....		200	Seals's pond.....		150
Flag's lake.....		400	Ricardo, Bertelson's reservoir.....		100
Manchaca, Labenski Creek.....		400	Ringgold, Woolsey's pond.....		200
Onion Creek.....		500	Rockdale, Clear Lake.....		300
Marfa, Barker's pond.....		100	Rogers, Rogers Lake.....		200
Marshall, Fern Lake.....		500	Rosebud, Ocker's pond.....		100
McClaran's lake.....		250	Williams Creek.....		400
Maxwell, Schawe Lake.....	1,000		Rotan, Cave Pond.....		150
Memphis, Brice's lake.....		100	Royston, Lake View.....		150
Cottonwood Creek.....		500	Saginaw, Canes Pond.....		200
Jones Creek.....		400	Salesville, Herring's lake.....		800
Noel's lake.....		100	San Angelo, Bismark Lake.....		500
Parker Creek.....		500	Concho River.....		500
Salt Creek.....		900	Cunningham Lake.....		500
Spring Creek.....		500	Doorkey Lake.....		500
Spring Lake.....		100	Gardners Lake.....		500
Mercedes, Davis Lake.....	1,000		Mires Lake.....		500
Meridan, Johnson's lake.....		200	North Concho River.....		500
Merkel, Martin's lake.....		650	Pecan Creek.....		500
Miller's lake.....		400	Scines Lake.....		500
Valley Farm Lake.....		300	Spring Creek.....		500
Miles, Lipan Creek.....		410	Twin Mountains Lake.....		500
Milford, Katy Pond.....		300	San Antonio, Anderson Club Pond.....		600
Mineola, Conger Pond.....		28	Billy Lake.....		900
Lake Park Pond.....		100	Guinn's lake.....		60
Willow Pond.....		150	Lake Toft.....		400
Mingus, Nine Lake.....		300	Sanger, Duck Creek.....		400
Thurber Lake.....	1,000		Hughes's pond.....		50
Mount Calm, Herring Lake.....		100	Sarber, Sarber Lake.....		500
Mount Pleasant, Lake Dellwood.....		150	Schulenburg, Running Spring.....		50
Mount Selman, Phialpha Lake.....		250	Seguin, Duck Lake.....		50
Mount Vernon, Devall's pond.....		150	Sherman, O'Hanlon's pond.....		100
Holbrook Lake.....		150	Stamford, Boulevard Pond.....		500
Nacogdoches, Fern Lake.....	1,000		Park Pond.....		300
Stone Lake.....		800	Swenson Pond.....		500
Navasota, Shell Lake.....	1,000		Tank Lake.....		300
Yarboro Lake.....	1,000		University Park Lake.....		500
New Braunfels, Comal Creek.....		600	Wedington Pond.....		200
G u a d a l u p e River.....		300	Sulphur Springs, Booker's pond.....		200
Rebecca Creek.....	1,000		Byrd's pond.....		50
North Zulch, Railroad Reser- voir.....		600			

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Texas—Continued.			Virginia—Continued.		
Sulphur Springs, Higdon Pond.....		10	Clarkton, Staunton River Lake.....		500
Pound Lake.....		10	Cobham, Cobham Park Pond.....		100
Reiley Lake.....		20	Cohoke, Cohoke Club Pond.....		75
Thompson Pond.....		20	Cologne, Bland's pond.....		75
Woodland Lake.....		150	Craigsville, Campbell Pond.....		75
Taylor, Taylor Lake.....		150	Culpeper, Smith Run Pond.....		350
Temple, Lake Polk.....		300	Danville, Dan River.....	1,000	
Terrell, Arnolds Lake.....		100	Drakes Branch, Twitty Creek.....		350
Cooper Lake.....		200	Drewryville, Drewry Mill Pond.....		250
Country Club Lake.....		900	Pope's pond.....		250
Garrett's pond.....		100	East Lexington, North River		
Gordon Lake.....		500	Pond.....		200
Griffith League Lake.....		100	Elmont, Chickahominy Mill		
Landos Lake.....		400	Pond.....		75
Martin's lake.....		100	Evington, Haden Branch.....		200
Oleander Lake.....		100	Farmville, Boiling's pond.....		250
Sand Branch Lake.....		100	Richardson's pond.....		200
Walton Lake.....		100	Fishers Hill, Shenandoah River.....		200
White Rock Lake.....		150	Fredericksburg, Corenty Pond.....		40
Timpson, Green's lake.....		100	Rappahannock		
Wedgeworth's lake.....		300	River.....		80
Troup, Gourley Lake.....		200	Gordonsville, Atkinson's pond.....		300
Waco, Holloway Lake.....		300	Harrisonburg, Dry River.....		100
Oak Lake.....		300	Linville Cree k		
Turner's lake.....		100	Lake.....		100
Waller, Ellis Pond.....		400	North River.....		100
Walnut Springs, Smitham's lake.....		50	Hollins, Carvins Creek.....	2,000	
Waxahachie, Bell Branch Lake.....		800	Hot Springs, Jackson River.....		400
Bullard's lake.....		200	Hunters, Little Hunting Creek.....		150
Davis's lake.....		200	Heswick, Christian's pond.....		75
Katy Fishing Club			La Crosse, Meherrin River.....	3,000	
Lake.....		500	Lawrenceville, Great Creek.....		300
Spalding Lake.....		475	Meherrin River.....		300
West End Lake.....		485	Rose Creek.....		250
Weatherford, Briten Branch.....		50	Lawyers, Leech's pond.....	1,000	
Hammond Lake.....	2,300		Leesburg, Goose Creek.....		50
Webbs, La Zeta Pond.....		400	Potomac River.....		200
Weinert, Edwards Lake.....		150	Limeton, Shenandoah River.		
Lake Creek Tank.....		750	South Branch.....		300
West, McClellan Lake.....		400	Louisa, Kent Mill Pond.....		100
Wetmore, Classen's pond.....		50	Lynchburg, Odd Fellows Home		
Wichita Falls, Woodall's pond.....		300	Lake.....		700
Wills Point, McKinney Lake.....		100	Martinsville, Smith River.....	3,000	
Winsboro, Harris's pond.....		20	Moseley Junction, Oak Hill Pond.....		100
Worham, Hardy Gin Lake.....		150	Mt. Jackson, Mill Creek.....		100
Yoakum, Mergenthal Pond.....		100	Shenandoah River.....		100
Shampaign's lake.....		200	Shenandoah River.		
Zulch, Zulch Lake.....		150	North Branch.....		100
Utah:			Smith Creek.....		100
Centerville, Perkins' pond.....		100	Natural Bridge, Buffalo Creek.....		250
Ogden, Brigham Pond.....		200	Nelson, Arons Creek.....		125
Virginia:			New Castle, Craig Creek.....	5,000	
Alleghany, Dunlap Creek.....		300	Johns Creek.....		250
Ashland, Ashland Park Pond.....		75	Newport News, Jordan's lake.....		125
King Pond.....		75	Norfolk, Lake Madoo.....		200
Atlee, Cross Creek Pond.....		100	North River, North River.....		100
Blackstone, Webb's pond.....		150	Nottaway, Robertson's pond.....		125
Bristol, Columbian Paper Co.'s			Oak Ridge, Oak Ridge Pond.....		500
reservoir.....		200	Occoquan, Metzger's pond.....		40
Broad Run, Broad Run.....		75	Occoquan River.....		80
Brookneal, Falling Creek.....		250	Wells Pond.....		40
Buffalo Junction, Arons Creek.....	3,000		Overall, Shenandoah River.....		300
Hites Pond.....		250	Oyster Point, Oyster P o i n t		
Pools Pond.....		250	Pond.....		200
Watkins Mill			Youngs Mill Pond.....		200
Pond.....	1,000	250	Pamplin City, Bakers Mill Pond.....		250
Callaghan, Dunlop Creek.....		400	Calhoun Pond.....		250
Potts Creek.....		400	Rossers Mill		
Chatham, Crystal Lake.....	1,000		Pond.....		250
Hedrick's pond.....	1,000		Penola, Mataponi Pond.....		80
Church Road, Burnt Quarter			Petersburg, Brandon Pond.....		200
Pond.....		200	Cains Mill Pond.....		200
Claremont, Snyder's pond.....			Daniels Pond.....		75
Clarksville, Grassy Creek.....	2,000		Kutchan Pond.....		75
Island Creek.....	1,000		Lake Ferndale Park		
Lewis's lake.....	2,000		West End Park		
			Lake.....		150
			Rapidan, Taliaferro Lake.....		150

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Virginia—Continued.			West Virginia—Continued.		
Richmond, Broad Lock Pond.....		1,000	Felton, Tygarts Valley River....		400
Bryan Pond.....		1,000	Glouanum, Tug River.....		150
Dead Creek Pond.....		1,000	Grafton, Tygarts Valley River....		400
Falling Creek.....		1,500	Harpers Ferry, Potomac River....		1,150
Flat Rock Pond.....		1,000	Little Falls, Monongahela River...		400
Fulton Club Pond.....		1,500	Morgantown, Deckers Creek....		200
Garlick Pond.....		1,000	Monongahela River.....		640
Lakeside Pond.....		100	Orleans Road, Potomac River....		1,000
Licking Creek Pond.....		1,000	Paw Paw, Great Cacapon River....		200
MacGregor Hall Pond.....		1,000	Phillippi, Middle Fork River....		400
Newman Pond.....		1,000	Tygarts Valley River....		400
Powhite Pond.....		1,000	Ripley, Mill Creek.....		80
Providence Forge Pond.....		1,000	Romney, Potomac River, South Branch.....		560
Reservoir Lake.....		100	St. Albans, Coal River.....		400
Spring Pond.....		100	Springfield, Potomac River, South Branch.....		300
Rockfish, Hardwick Lake.....		75	Star City, Donkard Creek.....		400
Plainview Pond.....		100	Sutton, Elk River.....		1,400
Shawen's pond.....		100	Weston, Monongahela River, West Fork.....		600
Rocky Mound, Furnace Creek.....	1,000		Woodland, Fish Creek.....		400
Big River.....	2,000	200	Wisconsin:		
Roxbury, Etna Mill Pond.....		250	Albany, Sugar River.....		500
Rural Retreat, Scott's pond.....		2,000	Butternut, Butternut Lake.....		600
Salem, Roanoke River.....		200	Cable, Cable Lake.....		400
Saxe, Charlotte Pond.....		225	Henry Lake.....		150
Shipman, Oak Ridge Pond.....		100	Cisco, Palmer Lake.....		400
Soudan, Grass Creek.....	3,000		Cumberland, Beaver Dam Lake....		400
South Boston, Butram Creek.....		200	Durand, Bear Lake.....		200
Dan River.....		300	Plummer Lake.....		200
Strasburg, Shenandoah River, Shenandoah River, West Fork.....		75	Thompson Lake.....		200
Stuart, Mayo River.....		350	Elcho, Bass Lake.....		300
Swords Creek, Clinch River.....		200	Enterprise Lake.....		500
Syeamore, Hunt Mill Pond.....		250	Otter Lake.....		250
Tappahannock, Mornington Lake.....		200	Elkhart, Crystal Lake.....		300
Timber Ridge, North River.....		250	Elmwood, Eau Galle Mill Pond....		300
Urbanna, Jackson Mill Pond.....		200	Elroy, Mill Pond.....		250
Victoria, Abilene Reservoir.....	2,000		Fairchild, Eau Claire River, North Fork.....		400
Victoria Reservoir.....	2,000		Fox Lake, Fox Lake.....		800
Village, Smithers Mill Pond.....		100	Genoa, Mississippi River.....		1,668
Virginia Beach, Lake Christine....		300	Gordon, Bass Lake.....		400
Wadesville, Opequan Creek.....		200	Blue Gill Lake.....		200
Wakefield, Brittle's pond.....		100	Hartford, Pike Lake.....		400
Walkerford, James River.....		400	Hatfield, Lake Arbutus.....		600
Walkers Station, Vaidens Mill Pond.....		400	Haugen, Bear Lake.....		600
Warren, Ballinger Creek.....		100	Devils Lake.....		500
Waterlick, Shenandoah River....		75	Hawkins, Shamrock Lake.....		250
Weems, Carter Creek.....		200	Hayward, Grindstone Lake.....		300
Winchester, Back Creek.....		200	Lake Court O'Reilles.....		400
Hogue Creek.....		200	Tripp's lake.....		200
Woodstock, Shenandoah River, North Branch.....		100	Whitefish Lake.....		300
Wytheville, Reed Creek.....	3,000	350	Hillsboro, Baraboo River, South Fork.....		250
Washington:			Hurley, Island Lake.....		400
Anacortes, Lake Campbell.....		400	Independence, Bugle Lake.....		400
Paso Lake.....		300	Trempealeau River.....		500
Medical Lake, Clear Lake.....		400	La Crosse, Mississippi River....		4,666
Silver Lake.....		400	Lake Beulah, Lake Beulah.....		1,200
Montesano, Lake Neuwatzel.....		300	Lake Geneva, Lake Como.....		1,000
Newport, Casey Lake.....		250	Lavalle, Duren.....		200
Tacoma, Madrona Lake.....		200	Little Baraboo Pond.....		200
West Virginia:			Long Lake, Fay Lake.....		400
Belva, Peters Creek.....		150	Long Lake.....		400
Bretz, Deckers Creek.....	4,000		Lublin, Lublin Lake.....		300
Caddell, Cheat River.....	4,000		Medford, Lake Esadore.....		200
Capon Springs, Great Cacapon River.....		900	Lake Murat.....		200
Chapmansville, Guyandotte River.....		240	Lake Perkins.....		400
Charleston, Elk River.....		200	Powell Lake.....		200
Elm Grove, Big Wheeling Creek.....		400	Richter Lake.....		200
Fairmont, Monongahela River....		400	Sackett Lake.....		200
Tygarts Valley River.....		400	Twin Lakes.....		200
			Mellon, Beaver Lake.....		200
			Carrot Lake.....		400
			Herbert Lake.....		200

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

- LARGE-MOUTH BLACK BASS—Continued.

Disposition.	Fry.	Finger- lings.	Disposition.	Fry.	Finger- lings.
Wisconsin—Continued.			Wisconsin—Continued.		
Mellon, Island Lake		400	Solon Springs, Twin Lakes		450
Lake Caroline		400	Sparta, La Crosse River		800
Menomonie, Cub Lake		200	Perch Lake		300
Lake Menomonie		400	State Line, Bass Lake		200
Red Cedar		500	Black Oak Lake		400
Stumps Slough		300	Tomah, Water Mill Pond		300
Youngs Lake		400	Tomahawk Lake, Little New- man Lake		250
Merton, Lake Keesar		400	Turtle, Long Lake		400
Muscoda, Mill Creek Pond		400	Victory, Mississippi River		165
New Auburn, Jenstow Lake		200	Wonewoc, Baraboo River		500
Shatiek Lake		250	Baraboo River, North Branch		500
Okauchee, Okauchee Lake		600	Wyoming:		
Pelican, Pelican Lake		800	Basin, Red Canyon Reservoir		125
Post Lake		500	Shoshoni, Big Horn River		400
Prairie du Chien, Mississippi River	4,250		Total a	56,600	665,868
Richfield, Lake Amy Belle	400				
Sheboygan Falls, Sheboygan River		250			

a Lost in transit, 25,135 fingerlings.

SUNFISH (BREAM).

Disposition.	Finger- lings, year- lings, and adults.	Disposition.	Finger- lings, year- lings, and adults.
Alabama:		Georgia—Continued.	
Gordo, Hannah's pond	100	Cuthbert, Nochaway Creek	100
Haleyville, Haleyville Pond	125	Wade's pond	50
Hodges, Strifel's pond	100	Ellaville, Murray's pond	100
Kennedy, Savage's pond	100	Ellabelle, Tony Branch	200
Reform, Harper's pond	100	Flint, Stegall's lake	400
Sulligent, Maddox's pond	100	Forsythe, Bessie Tift Lake	50
Tuscumbia, Tuscumbia Spring	100	Jackson's pond	50
Vance, Lawrence's pond	125	Garfield, Oglesby's pond	100
Arkansas:		Glennville, De Loach's pond	50
Greenwood, Saling's pond	150	Lewis's pond	50
Harrison, Bates's pond	150	Graymont, Cowert's pond	100
Helena, Mississippi River	83,665	Wetherford's pond	100
Hope, Brandon's pond	100	Halcyondale, Simmons's pond	50
Johnson's pond	100	Junction City, Carlisle's pond	50
Little Rock, Asylum Pond	100	Moore's pond	50
Mammoth Springs, Mammoth Springs	200	Leesburg, Kinchatoonee Creek	100
Marshall, Horton's pond	150	Macon, Biarly Lodge Pond	150
Nashville, Mine Creek	250	Recreation Club Lake	100
Reese's pond	100	Manchester, Manchester Pond	100
Whelen, Edmond's pond	100	Marshallville, Grisold Spring Pond	100
Connecticut:		Outing Club Pond	100
Leonard Bridge, Hop River	600	Rumple's pond	100
Seymour, Beecher's pond	300	Mayfield, Long's pond	200
Florida:		Millen, Buckhead Creek	200
Ehren, Floral Lake	100	Ogeechee River	100
Tampa, Saddle Bag Lake	100	Munnerlyn, Rosemary Creek	100
Georgia:		Rupert, Bodford's pond	50
Adel, Beaver Dam Bay	50	Scarboro, Ogeechee River	100
Saddlebag Pond	50	Smithville, Kinchatoonee Creek	100
Americus, Mountain Creek Pond	50	Muckalee Creek	100
Ashburn, Clear Pond	50	Stillmore, Cannochee Pond	100
Fitzgerald's pond	50	Stinson, Lake Betson	225
Atlanta, Moccasin Lake	110	Summit, Bowie's pond	100
Blue Ridge, Carter's pond	120	Brown's pond	100
Chamblee, Jones's pond	25	Cowart's pond	50
Charing, Branch Pond	50	Spring Branch Pond	50
Clarkesville, Edward's pond	100	Turner's pond	50
Hazel Creek	150	Sylvester, Pope's pond	50
Clayton, Justus's pond	125	Talbotton, Maxwell's pond	50
Collins, Jarriel's pond	50	Parker's pond	50
Wilson's pond	50	Silver Lake	50
Wrenn's pond	50	Wilson's pond	50

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

SUNFISH (BREAM)—Continued.

Disposition.	Finger- lings, year- lings, and adults.	Disposition.	Finger- lings, year- lings, and adults.
Georgia—Continued.		Mississippi—Continued.	
Tennille, Boatright's pond.....	100	Corinth, Pound's pond.....	100
The Rock, Stafford's pond.....	50	Rilla Pond.....	100
Tifton, Purdy's pond.....	50	Waukomis Lake.....	100
Ty Ty, Parks's pond.....	50	Crenshaw, Berk's pond.....	100
Vienna, Lane's pond.....	50	Mitchell's pond.....	100
Wade, Brinson's pond.....	50	Durant, McDonald's pond.....	100
Illinois:		Enterprise, Kamper's pond.....	100
Belleville, Gauss's lake.....	200	Gandsi, Spring Pond.....	100
Rheins's lake.....	100	Hazelhurst, Harrison's pond.....	100
Olney, Olney Reservoir.....	100	Heidelberg, Vernon's pond.....	100
Indiana:		Hickory, White Oak Pond.....	100
Borden, Koerber's pond.....	100	Houston, Knox's pond.....	100
Spring Pond.....	100	Jackson, Spring Lake.....	100
Bristol, Newman's pond.....	100	Willow Pond.....	100
Carbon, Harrold's pond.....	300	Laurel, Park Lake.....	150
Chrisney, Oak Hill Pond.....	100	Liberty, Ball's pond.....	100
Dubois, Silver Pond.....	100	Lockhart, Harbour's pond.....	100
Edinburg, Spring Lake.....	300	McDonald, Ingrani's pond.....	100
Fairmount, Little's pond.....	100	Macon, Boswell's pond.....	150
Manzanita Lake.....	100	Eliland's pond.....	150
Farmersburg, Lewis's pond.....	200	Howard's lake.....	125
Kewanna, Bruce Lake.....	400	Stuart's pond.....	100
Lima, Still Lake.....	200	Meridian, Bailey's pond.....	100
Madison, Big Creek.....	350	College Lake.....	100
New Albany, Old Cave Pond.....	100	Miller's pond.....	200
Ossian, Willow Pond.....	100	Suttle's pond.....	350
Silver Lake, Silver Lake.....	100	New Albany, Strond's pond.....	100
Veedersburg, Coal Creek.....	800	Nicholson, Gentry's pond.....	100
Iowa:		Okolona, Colburn's pond.....	200
Casey, Spring Lake.....	200	Quitman, Lake Ruth.....	100
Cumberland, Hawthorn Lake.....	100	McNair's pond.....	100
Fort Madison, Green Bay.....	1,100	Sessums, Perkins' pond.....	100
Lime Springs, Upper Iowa River.....	4,500	Sherwood, Norris' pond.....	100
North McGregor, Mississippi River.....	73,250	Shuqualak, Adams' lake.....	150
Underwood, Geise's pond.....	100	Wigwam Lake.....	150
Kansas:		Strongs, Lake Bolivar.....	100
Grenola, Cana River.....	200	Spring Creek.....	100
Kentucky:		Williams' pond.....	100
Beard, Cypress Pond.....	100	Summit, Hillside Pond.....	200
Elizabethtown, Heady's pond.....	100	Willow Pond.....	150
Eminence, Boyne's pond.....	100	Taylor'sville, Robinson's pond.....	100
Helburn's pond.....	100	Tishomingo, Holley's lake.....	150
Glasgow, Fallen Timber Creek.....	150	Tupelo, Hill's pond.....	200
Grays, Lynn Camp Pond.....	400	Van Vleet, Arnett Place Pond.....	250
Louisville, Lake Lansdowne.....	300	Hickory Grove Pond.....	100
Saxton, Beech's pond.....	400	Waynesboro, Dyess Mill Pond.....	100
Sonora, Ireland's pond.....	150	Oakland Pond.....	100
Louisiana:		Patten's creek.....	100
Bogalusa, Bogalusa Pond.....	300	Taylor's lake.....	100
Homer, Gladney's pond.....	100	Wilkins Mill Pond.....	100
Spring Lake.....	100	West Point, Dunlap's lake.....	400
Ruston, Pugh's pond.....	100	Ivy's pond.....	400
Scotland, Scotland Plantation Lake.....	200	Trout Lake.....	100
Maryland:		Whittaker, Whittaker's pond.....	150
Bel Air, Barnes Run.....	150	Yazoo City, Hicks' pond.....	100
Chevy Chase, Locust Lake.....	400	Missouri:	
Landover, Oak Hill Pond.....	250	Arlington, Lukrofka's pond.....	400
Mountain Lock, Potomac River.....	5,600	Conway, Thomas' pond.....	200
Massachusetts:		Marquand, Clubb's pond.....	200
Plymouth, King's pond.....	300	Nebraska:	
West Pond.....	300	Cheney, Variety Grove Farm Pond.....	100
Minnesota:		Nevada:	
Brownsville, Mississippi River.....	17,300	Ely, Olsen's lake.....	150
Smiley, Pelican Lake.....	500	New Mexico:	
Mississippi:		Deming, Harris's pond.....	150
Blue Mountain, Simmons' pond.....	100	Elida, Mesa Lake.....	100
Booneville, Gin Pond.....	100	North Carolina:	
Brookhaven, Applewhite's pond.....	100	Aberdeen, Bonnie Brier Pond.....	75
Brooks ville, Dixie Pond.....	100	Sand Hill Branch Pond.....	300
Peterson's pond.....	150	Angier, Matthews' pond.....	150
Centreville, Dick's pond.....	150	Concord, Clark Creek.....	225
Willow Lake.....	150	Springville Pond.....	150
Collins, Mayfield's pond.....	125	Fayetteville, Pine Lake.....	450
Columbus, Fig Pond.....	300	Franklinton, Dickerson Mill Pond.....	75

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

SUNFISH (BREAM)—Continued.

Disposition.	Finger- lings, year- lings, and adults.	Disposition.	Finger- lings, year- lings, and adults.
North Carolina—Continued.		Oklahoma:	
Franklinton, Green Hill Pond.....	75	Ardmore, City Lake.....	300
Spring Branch.....	75	Dyer Lake.....	200
Whiteside Pond.....	75	Reed's lake.....	100
Williams's ponds.....	150	Santa Fe Lake.....	300
Gastonia, Crawford's pond.....	75	Asher, Merrill's pond.....	100
Lake Giles.....	225	Salt Creek Ponds.....	125
Payes Lake.....	150	Doxey, Topper's pond.....	100
Spencer's lake.....	300	Elk City, Hughes's lake.....	100
Glen Alpine, Silver Creek Pond.....	75	Hugo, Wright's pond.....	100
Gold Hill, Second Creek.....	150	Pryor, Miller's pond.....	100
Graham, Graham Country Club Pond.....	225	Stuart, Coal Creek.....	100
Guilford College, Ash Pond.....	75	Tyrone, Crites's pond.....	100
Hendersonville, Lake Osceola.....	600	Pennsylvania:	
Lake West.....	300	Canonsburg, Neill's pond.....	300
Rhett's pond.....	150	Danville, Susquehanna River.....	1,250
High Point, Willard's pond.....	75	Hanover, Little Onewago Creek.....	150
Landis, Codle Creek Pond.....	75	Huntingdon, Raystown Branch.....	200
Landrum, Greenway's pond.....	75	Icedale, Brandywine Creek.....	300
Hughes' pond.....	150	New Bethlehem, Leatherwood Creek.....	500
Lexington, Bock's pond.....	175	Palm, Hosenack Creek Lake.....	200
Hargrave's pond.....	75	Perkiomen Creek.....	200
Liberty, Cane Creek Pond.....	75	Reading, Maiden Creek.....	300
Thompson's pond.....	75	Tulpehocken Creek.....	300
Lillesville, Dockery's pond.....	225	Shoemakersville, Dreiblebis Creek.....	200
Island Creek.....	225	Moyer Creek.....	200
Mill Brook, Pineridge Pond.....	210	Temple, Ahren's pond.....	200
Morgantown, McDowell's pond.....	100	Bernhart's lake.....	200
Morven, Hamville Pond.....	75	Weissport, Big Creek.....	300
Mill Pond.....	150	Windber, Ice Company Pond.....	200
Spring Pond.....	150	York, Spring Lake.....	100
Pinnacle, Culler's pond.....	75	South Carolina:	
Pittsboro, Four Springs Pond.....	225	Aiken, Bridge Creek Pond.....	150
Hallborne Pond.....	75	Johnson's pond.....	100
Petty's pond.....	150	Shaw's pond.....	100
Raleigh, Country Club Lake.....	300	Thorpe's pond.....	100
Lynn's pond.....	75	Belton, Williams's pond.....	100
Rockingham, Dog Branch Pond.....	75	Bethune, Bell Branch Pond.....	100
Ronda, Bugaboo Pond.....	150	Blacksburg, Parris's pond.....	75
Little Elkin Pond.....	150	Blaney, Crystal Lake.....	100
Rutherfordton, Broad River Pond.....	75	Borden, Pollard Mill Pond.....	100
Salisbury, Cauble's pond.....	175	Camden, Boykin's pond.....	100
Glover's pond.....	125	McLeod's pond.....	200
Smithfield, Pou's pond.....	150	Central, Arnold's pond.....	50
Southside, Rhyme's pond.....	75	Holcomb's pond.....	50
Wake Forest, Allen's pond.....	75	Chester, Dry Fork Pond.....	75
Bobbitt's pond.....	100	Columbia, Cayce's pond.....	200
Dickson's pond.....	100	Gill Creek.....	200
Harrison's pond.....	100	Messer's pond.....	200
Maltonia Club Pond.....	150	Mill Creek Pond.....	200
Moore's ponds.....	300	Poplar Branch Pond.....	100
Spring Pond.....	100	Cope, Fogle's pond.....	100
Wilbon, Neill's pond.....	225	Cordova, Smoak's pond.....	75
Wilkesboro, Roberson's pond.....	75	Fort Mill, Spring Pond.....	75
Winston-Salem, Holton's pond.....	75	Gaffney, Parker's pond.....	75
Youngsville, Alexander's pond.....	75	Turner's pond.....	75
North Dakota:		Graniteville, Power House Pond.....	75
Devils Lake, Devils Lake.....	1,000	Greenville, Houston's pond.....	150
Granville, Buffalo Lodge Lake.....	300	Maple Creek Pond.....	75
Oriska, Beyer's pond.....	70	Greenwood, Logan Branch.....	75
St. John, Clear Lake.....	300	Moore Branch Pond.....	75
Hooker's lake.....	300	Spring Pond.....	75
Lake Alexander.....	300	Hartsville, Beaver Dam Pond.....	100
Lake Nemo.....	300	Prestwood Pond.....	100
Waukippa Lake.....	300	Honea Path, Big Spring Pond.....	100
Ohio:		Broadmouth Creek.....	175
Ada, Hubbell's pond.....	100	Kay's pond.....	150
Gallipolis, Safford's pond.....	100	Knight's pond.....	75
Hebron, Buckeye Lake.....	600	Little River.....	100
Orbiston, Orbiston Pond.....	100	Johnston, Brinson's pond.....	100
Perry, Shady Nook Pond.....	400	Butler's pond.....	75
Rarden, Taylor's pond.....	100	Calloun's pond.....	75
Rossmoyne, Taylor's pond.....	100	Hilliard's pond.....	100
Sharonville, Schatzle's pond.....	100	Hollingsworth's pond.....	75
Tippecanoe City, Kessler's pond.....	100	Lott's pond.....	75

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

SUNFISH (BREAM)—Continued.

Disposition.	Finger- lings, year- lings, and adults.	Disposition.	Finger- lings, year- lings, and adults.
South Carolina—Continued.		Texas—Continued.	75
Johnston, Spring Branch.....	75	Brady, Lime Oak Creek.....	
Ward Creek Pond.....	100	Brazos, Blucher's pond.....	30
Kershaw, Horton's pond.....	75	Carbon, Pierce's pond.....	30
Kinards, Oxner's pond.....	75	Carthage, Hill's lakes.....	50
Lancaster, Steele's pond.....	75	Cisco, Lake Borine.....	100
Wildcat Pond.....	75	Clifton, Manske's pond.....	50
Laney, Robeson's pond.....	100	Comanche, Highland Lake.....	25
Langley, Little Horse Creek Pond.....	250	De Leon, Spring Pond.....	100
McCormick, Britt's pond.....	75	Elkhart, Elkhart Lake.....	200
Spring House Pond.....	75	Eskota, Kurth's pond.....	50
Spring Pond.....	100	Fort Worth, Davie Burns Lake.....	50
Macedon, Bogy Pond.....	100	Friona, Mayflower Pond.....	30
Newberry, Kings Creek.....	100	Gorman, King's pond.....	60
North Augusta, Big Branch Pond.....	100	Lusk's pond.....	30
North, White's pond.....	100	Gordon, Chenault's pond.....	50
Orangeburg, Gue's pond.....	50	Horlin's pond.....	30
Pageland, Hicks's pond.....	100	Graham, Oak Grove Pond.....	20
Perry, Piney Branch Pond.....	125	Grand Saline, Brown's pond.....	30
Pickens, Colony Pond.....	50	Jacksonville, Belva Lake.....	200
Oolong Pond.....	75	Kaufman, Hofer Pond.....	30
Rock Hill, Mill Pond.....	100	Kemp, Trinity Lake.....	100
Ruby, Oliver's pond.....	100	Lindale, Mill Creek Pond.....	100
Salley, Branch Pond.....	150	Llano, Doel's pond.....	40
Seneca, Langston's pond.....	50	Lytle, Carter's pond.....	30
Shoals Junction, Dunn's pond.....	100	Mabank, Grubb's pond.....	30
Simpsonville, Rocky Creek Pond.....	75	Manor, Johnson's reservoir.....	20
Strother, McMahan's pond.....	50	Marlin, Clark's pond.....	50
Trenton, Hughes's pond.....	100	Marshall, Lake Ferns.....	300
Horn Creek.....	75	Lake Katrine.....	50
Hunt Creek Pond.....	100	Walker's lake.....	50
Marsh's pond.....	100	Merkel, Count's pond.....	30
Raus's pond.....	75	Nacogdoches, Mamie Ross Lake.....	300
Shaws Creek Pond.....	75	Rockdale, Coffield's pond.....	100
Webb's pond.....	75	Randle's lake.....	40
Union, Buffalo Mill Pond.....	100	Rotan, Hunter's pond.....	130
Municipal Reservoir.....	100	Saginaw, Beall's pond.....	50
Wagner, Dean Swamp Pond.....	75	Santo, Miller's pond.....	50
Walhalla, Bauknight's pond.....	75	Terrell, McCord's pond.....	25
Burley's pond.....	75	Renfro Creek Lake.....	25
Oconee pond.....	75	Toyah, Humphries's pond.....	50
Todd's pond.....	75	Tuxedo, Davis's lake.....	100
Verner's pond.....	75	Tye, Worthington Lake.....	40
Willington, Arial's pond.....	75	Tyler, Country Club Lake.....	200
Covin's pond.....	75	Lake Park.....	200
Gilbert's pond.....	75	Lake Wood.....	200
Le Roy's pond.....	75	Walnut Springs, Smitham's lake.....	100
Little River.....	100	Wichita, Railroad Pond.....	60
Winnsboro, Creight's pond.....	50	Winnsboro, Baker's pond.....	20
Haynes's pond.....	75	Spring Lake.....	150
Woodruff, Chumley's pond.....	75	Virginia:	
Ferguson Creek.....	50	Beaeton, Old Gum Spring Pond.....	150
Watson's pond.....	75	Beaver Dam, Thompson's pond.....	125
Yorkville, Smith's pond.....	75	Belmont Park, Goose Creek.....	350
South Dakota:		Charlottesville, New Reservoir.....	225
Hitecock, Cramer's pond.....	100	Cumberland, Burleighhall Pond.....	125
Scenic, Knutson's pond.....	425	Dillwyn, Fitzgerald's pond.....	125
Tennessee:		Disputanta, Belscher's pond.....	125
Builer, Cable's pond.....	175	Drewrys Bluff, Spring Lake.....	125
Concord, Turkey Creek Lake.....	200	Duugamon, Kilgore's pond.....	450
Cookeville, Clause's pond.....	225	East Lexington, North River Pond.....	400
Cumberland Gap, Holly Hill Pond.....	200	Evington, Irvine's pond.....	200
Lambert's pond.....	200	Farmville, Agee's pond.....	200
Johnson City, Aspen Bower Lake.....	500	Glady's, Maple Pond.....	250
Knoxville, Little River.....	75	Gordonville, Oak Hill Pond.....	400
Maryville, Housholder's pond.....	200	Orange, Mill Creek Pond.....	125
Tate Springs, Kirkham's pond.....	75	Pennington Gap, Hickory Flats Pond.....	200
Tate Springs Reservoir.....	150	Petersburg, Belscher's pond.....	400
Wautauga Point, Buffalo Creek.....	500	Shipman, Mountain Pond.....	125
Whitesburg, Shields's pond.....	75	Spout Springs, Poplar Pond.....	150
Texas:		Staunton, Gypsy Hill Lake.....	125
Amarillo, Famous Heights Park Lake.....	50	Troutville, Alderson's pond.....	150
Big Springs, Davis's pond.....	35	Troy, Poplar Grove Pond.....	125
Fisher's pond.....	35	Winton, Brown's pond.....	200
Blum, Klondike Lake.....	100	Warrenton, Cedar Run.....	100

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

SUNFISH (BREAM)—Continued.

Disposition.	Finger- lings, year- lings, and adults.	Disposition.	Finger- lings, year- lings, and adults.
Virginia—Continued.		Wisconsin—Continued.	
Warrenton, Forest Branch Pond.....	150	Independence, New City Pond.....	200
Washington:		La Crosse, Mississippi River.....	21,468
Oroville, Lemonosky Lake.....	300	Muscoda, Mill Creek Pond.....	300
West Virginia:		Prairie du Chien, Mississippi River.....	58,250
Bedington, Emerson's pond.....	500	Victory, Mississippi River.....	1,666
Weston, Walnut Fork Pond.....	200	Wyoming:	
Wisconsin:		Sheridan, Cut-Off Pond.....	150
Genoa, Mississippi River.....	4,166	Total ^a	342,825
Independence, Bugle Lake.....	300		

^a Lost in transit, 2,810 fingerlings.

PIKE PERCH.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Arkansas:			
Des Arc, Caloutchie Bay.....		50,000	
Elkins, White River.....		400,000	
Helena, Mississippi River.....			800
Connecticut:			
Wallingford, Lake Quonnipaug.....		500,000	
Illinois:			
Havana, Illinois State Fish Commission.....	8,000,000		
Meredosia, Illinois River.....		930,000	
Momence, Kankakee River.....		1,260,000	
Wilmington, Kankakee River.....		1,260,000	
Indiana:			
Angola, Buck Lake.....		800,000	
Fox Lake.....		1,000,000	
Columbia City, Shriner Lake.....		1,200,000	
Leesburg, Shoe Lake.....		1,000,000	
Monticello, Tippecanoe River.....		1,500,000	
Rome City, Sylvan Lake.....		1,500,000	
Iowa:			
Clear Lake, Clear Lake.....		750,000	
Estherville, Des Moines River, West Branch.....		600,000	
Manchester, Maquoketa River.....		300,000	
Orleans, East Okeboji Lake.....		400,000	
Spirit Lake.....		400,000	
Ruthven, Lost Island Lake.....		400,000	
Waterloo, Cedar River.....		250,000	
West Liberty, Cedar River.....		200,000	
Kansas:			
Marion, Cottonwood River.....		400,000	
Kentucky:			
Hopkinsville, Waterworks Lake.....		800,000	
Lebanon, Beech Fork River.....		1,500,000	
Cartwright Creek.....		800,000	
Lloyds Creek.....		800,000	
North Fork Creek.....		1,000,000	
Pitman Creek.....		1,000,000	
Popes Creek.....		800,000	
Rolling Fork River.....		1,500,000	
South Fork Creek.....		1,000,000	
Massachusetts:			
Palmouth, Shivericks Pond.....		400,000	
Greenfield, Connecticut River.....		1,000,000	
Deerfield River.....		800,000	
Pittsfield, Pontosuc Lake.....		500,000	
Shelburne Falls, Deerfield River.....		600,000	
Waltham, Nonsuch Pond.....		500,000	
Michigan:			
Alpena, Long Lake.....		1,200,000	
Bay City, Saginaw Bay.....		4,500,000	
Birmingham, Wing Lake.....		500,000	
Crystal Falls, Mary Lake.....		540,000	
Detroit, Michigan Fish Commission.....	34,280,000		
Edwardsburg, Eagle Lake.....		975,000	

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

PIKE PERCH—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Michigan—Continued.			
Hale Lake, Hale Lake.....		500,000	
Loon Lake.....		800,000	
Lincoln, Brownlee Lake.....		600,000	
Millersburg, Barnhart Lakes.....		800,000	
Paw Paw, Maple Lake.....		1,000,000	
St. Joseph, Lake Chapin.....		1,200,000	
Witch Lake, Horse Shoe Lake.....		360,000	
Minnesota:			
Alexandria, Lake Geneva.....		540,000	
Big Lake, Big Lake.....		500,000	
Brownsville, Mississippi River.....			1,730
Chub Lake, Chub Lake.....		400,000	
Hanging Horn Lake, Hanging Horn Lake.....		600,000	
Mankato, Lake Washington.....		720,000	
Missouri:			
Crocker, Gasconade River.....		400,000	
Roubidoux Creek.....		400,000	
St. Joseph, Missouri Fish Commission.....	2,000,000		
New Hampshire:			
Mountaintview, Ossipee Lake.....		1,000,000	
Winchester, Forest Lake.....		500,000	
New Jersey:			
Boonton, Rockaway River.....		700,000	
New York:			
Addison, Canister River.....		600,000	
Bliss, Eagle Lake.....		600,000	
Lisle, Tioughnioga River.....		400,000	
North Dakota:			
Cando, State Fish Commission.....	10,000,000		
Ohio:			
Columbus, Scioto River.....		1,000,000	
Fremont, Sandusky River.....		1,000,000	
Holliers Beach, Lake Erie.....		16,000,000	
Isle St. George, Lake Erie.....		16,000,000	
Lima, Lima Lake.....		1,000,000	
Port Clinton, Lake Erie.....		475,000	
Put-in Bay, Lake Erie.....		20,000,000	
Ohio State Commission.....	170,725,000		
Toledo, Lake Erie.....		10,000,000	
Upper Sandusky, Upper Sandusky River.....		1,500,000	
Oklahoma:			
Tahlequah, Illinois River.....		400,000	
Pennsylvania:			
Bushkill, Delaware River.....		800,000	
Coolbaugh, Echo Lake.....		600,000	
Erie, Pennsylvania Fish Commission.....	96,450,000		
Factoryville, Lake Kewanna.....		700,000	
Goldsboro, Susquehanna River.....		500,000	
Huntingdon, Raystown Branch.....		700,000	
New Freedom, Clipper Dam.....		300,000	
New Milford, Upper Lake.....		700,000	
Spruce Hill, Tuscarora Creek.....		500,000	
Susquehanna, Page Pond.....		800,000	
Susquehanna River.....		800,000	
Vicksburg, Armstrong Run.....		200,000	
Wilkes-Barre, Nuangola Lake.....		1,000,000	
Wrightsville, Susquehanna River.....		500,000	
York Haven, Susquehanna River.....		500,000	
South Dakota:			
Langford, Ninemile Lake.....		800,000	
Sixmile Lake.....		800,000	
Tennessee:			
Springfield, Milldale Pond.....		800,000	
Vermont:			
Boltonville, Tickle Necked Pond.....		600,000	
Ludlow, Plymouth Pond.....		800,000	
Miles Pond, Miles Pond.....		800,000	
Swanton, Lake Champlain.....		11,000,000	
West Danville, Joe's pond.....		1,000,000	
Virginia:			
Wytheville, Reed Creek.....		1,000,000	
West Virginia:			
Fairmont, Tygarts Valley River.....		500,000	
Morgantown, Cheat River.....		800,000	
Wisconsin:			
Antigo, Edith Lake.....		400,000	
Barronette, Deep Lake.....		500,000	

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

PIKE PERCH—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Wisconsin—Continued.			
Cable, Namekagon Lake.....		800,000	
Colgate, Lake Five.....		500,000	
Crandon, Oak Lake.....		800,000	
Genoa, Mississippi River.....			416
Gordon, Bass Lake.....		400,000	
Clear Lake.....		400,000	
Wagner Lake.....		400,000	
Greenwood, Popple River.....		600,000	
Hancock, Fish Lake.....		400,000	
Haugen, Pokagama Lake.....		500,000	
Iron River, Lower Pike Lake.....		720,000	
Kewaunee, Kewaunee River.....		450,000	
La Crosse, Mississippi River.....			2,148
Nashville, Dry Lake.....		600,000	
Okauchee, Okauchee Lake.....		2,500,000	
Stone Lake, Whitefish Lake.....		400,000	
Victory, Mississippi River.....			166
Wonewoc, Baraboo River.....		800,000	
Total	321,455,000	154,480,000	5,260

YELLOW PERCH.

Colorado:			
La Jara, Laguna Escondida.....			200
Connecticut:			
Hadlyme, State Fish Commission.....	5,200,000		
Delaware:			
Wilmington, Brandywine Creek.....		800,000	
Illinois:			
Carbondale, Horse Shoe Lake.....			200
Chicago, Armour's pond.....			900
Otis's pond.....			900
Eckerts, Deich's pond.....			100
Irving, Funk's lake.....			500
Millstadt, Grossman's pond.....			300
Shipman, Olmsted's pond.....			400
Indiana:			
Angola, Walled Lake.....			200
Centerville, Kitterman's pond.....			90
Edinburg, White River, East Fork.....			200
Lake Cicott, Lake Cicott.....			300
Lebanon, Saltmarsh Pond.....			75
Silver Lake, Silver Lake.....			200
Winchester, Summers's pond.....			100
Iowa:			
Lime Springs, Upper Iowa River.....			20
McGregor, Lake Como.....			900
North McGregor, Mississippi River.....			42,750
Kansas:			
Pittsburg, Gibson Pond.....			100
Kentucky:			
Cropper, Dunavent's pond.....			100
Pollard's pond.....			100
Louisville, Lake Lansdowne.....			300
Park View Lake.....			100
Woodbine, Lake Placid.....			300
Maryland:			
Acookeek Creek, Potomac River.....	66,117,500		
Baltimore, Patapsco River Pond.....			150
Bryans Point, Potomac River.....	10,945,000		
Bush River, Bush River.....	2,400,000		
Cecil, Chesapeake Bay.....	23,600,000		
Chase, Dundee River.....	600,000		
Freeland, South Lake.....			300
Gunpowder, Gunpowder River.....	2,200,000		
Harford, Swan Creek.....	9,500,000		
Harmony Grove, Richfield Pond.....	200,000		
Havre de Grace, Chesapeake Bay.....	12,600,000		
Pamunkey Creek, Potomac River.....	10,985,000		
Piscataway Creek, Potomac River.....	64,887,500		
Principio, Chesapeake Bay.....	15,000,000		

^a Lost in transit, 545,000 fry.

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

YELLOW PERCH—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
Maryland—Continued.			
Swan Creek, Potomac River.....		5,915,000	
Town Point, Elk River.....		41,000,000	
Waterbury, Old Place Creek.....		400,000	
Massachusetts:			
Merrimac, Sargent's pond.....		400,000	
Michigan:			
Alpena, Lake Esau.....			400
Minnesota:			
Brownsville, Mississippi River.....			4,000
Rochester, Zumbro Mill Pond.....			200
Missouri:			
St. Charles, Crystal Lake.....			100
New Hampshire:			
Meredith, Long Pond.....		400,000	
New Jersey:			
Hammonton, Hammonton Lake.....		800,000	
Netcong, Bear Pond.....			200
Pompton Lakes, Pompton Lakes.....		1,000,000	
Pompton River.....		1,000,000	
Red Bank, Shrewsbury Pond.....		200,000	
New Mexico:			
Colfax, Adams Lake.....			219
New York:			
Auburn, Owasco Lake.....			2,000
Fallsburg, Ruddick Pond.....		600,000	
Flushing, Iron Spring Lake.....			150
Middleton, Ketchens Pond.....		600,000	
Summit Lake.....		400,000	
Millers Place, Hopkins Pond.....		200,000	
Mohonk Lake, Mohonk Reservoir.....		200,000	
Monroe, Monebasha Lake.....		600,000	
Round Island Lake.....		600,000	
Walton Lake.....		600,000	
North Carolina:			
Hendersonville, Tulip Pond.....			60
Lexington, Hankins' pond.....			100
Nokomis Mill Pond.....			100
Sandy Creek Pond.....			100
Salisbury, Coolemece Pond.....			100
Miller's pond.....			100
Second Creek.....			100
Statesville, Buffalo Shoal Pond.....			100
North Dakota:			
Devils Lake, Devils Lake.....			1,000
Lisbon, Mulinex's pond.....			200
Milnor, Star Pond.....			175
Ohio:			
Marion, Scioto River.....			70
Oklahoma:			
Devol, Suter's pond.....			70
El Reno, Carter's pond.....			75
McAlester, Cole's lake.....			50
Marietta, Love's lake.....			100
Ochelata, Upper Pond.....			150
Oklahoma City, Lake View Lake.....			250
Pennsylvania:			
Bedford, Dunning Creek.....			120
Raystown Branch.....			120
Bunkney, Susquehanna River.....		600,000	
Danville, Susquehanna River.....			425
Devon, Eldonridge Pond.....		200,000	
Dushore, Headley Pond.....		600,000	
Housingers Pond.....		400,000	
Mill Pond.....			125
Factoryville, Gardners Pond.....		600,000	
Freeport, Briar Patch Pond.....			100
Greenville, Shenango River.....			325
Honey Brook, Mackelduff Pond.....		400,000	
Indiana, Crooked Creek.....			150
Ledys, Big Pond.....			150
Lenape, Brandywine River.....		1,000,000	
New Freedom, Smith Ponds.....			300
Sheridan, Tulpehoeken Creek.....		600,000	
Waltersburg, Big Redstone Pond.....			100
South Carolina:			
Calhoun, Twenty-three Mile Creek.....			120
Denmark, Savannah Lake.....			180

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

YELLOW PERCH—Continued.

Disposition.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
South Carolina—Continued.			
Gaffney, Sarrat's pond.....			60
Greenville, Greenville Lake.....			120
Trenton, Horse Creek Pond.....			60
Troy, Spring Branch.....			60
South Dakota:			
Madison, Lake Herman.....			600
Vermont:			
Brattleboro, West River.....		300,000	
Lyndonville, Chandler Pond.....		600,000	
Bean Pond.....		400,000	
Poultney, Lake St. Catherine.....		500,000	
St. Johnsbury, Passumpsic River.....			1,445
Walden, Coles Pond.....		800,000	
Virginia:			
Boyce, Shenandoah River.....		2,000,000	
Charlottesville, Maury's pond.....			125
Danville, Maple Grove Pond.....			200
Dinwiddie, Cat Tail Pond.....		300,000	
Dogue Creek, Potomac River.....		26,680,000	
Little Hunting Creek, Potomac River.....		4,550,000	
Pohick Creek, Potomac River.....		10,205,000	
Rockfish, Hardwick Lake.....		400,000	
Rockfish Lake.....			250
Scottsville, Spring Pond.....			100
Washington:			
Tacoma, American and Gravelly Lakes.....			500
West Virginia:			
Milton, Newman Springs.....			100
Rippon, Bull Shin Creek.....		1,000,000	
Wisconsin:			
Elkhart, Elkhart Lake.....			300
La Crosse, Mississippi River.....			4,000
Lake Mills, Rock Lake.....			600
Prairie du Chien, Mississippi River.....			37,750
Total ^a	5,200,000	326,885,000	108,439

^a Lost in transit. 856 fingerlings.

STRIPED BASS.

Disposition.	Eggs.	Fry.
Maryland:		
Havre de Grace, Chesapeake Bay.....		115,000
North Carolina:		
Weldon, Roanoke River.....	4,566,000	2,669,000
Total.....	4,566,000	2,784,000

WHITE BASS.

Disposition.	Fingerlings, yearlings, and adults.
Arkansas:	
Helena, Mississippi River.....	5,950
Wisconsin:	
Genoa, Mississippi River.....	34
LaCrosse, Mississippi River.....	33
Victory, Mississippi River.....	33
Total.....	6,050

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

WHITE PERCH.

Disposition.	Eggs.	Fry.
Connecticut:		
Seymour, Hemp Swamp Pond.....		400,000
Delaware:		
Nassau, Red Mill Pond.....		2,400,000
Wilmington, Brandywine Creek.....		800,000
Maryland:		
Bush River Station, Bush River.....		2,000,000
Chase, Dundee Creek.....		4,000,000
Havre de Grace, Chesapeake Bay.....		122,450,000
Elk River.....		18,250,000
Susquehanna River.....		66,800,000
Hendersons Point, Elk River.....		32,555,000
Locust Point, Chesapeake Bay.....		5,150,000
Swan Creek, Chesapeake Bay.....		17,100,000
Town Point, Elk River.....		37,750,000
Wild Duck Harbor, Susquehanna River.....		20,825,000
Massachusetts:		
Gardner, Stoddard Meadow Pond.....		400,000
Tilton Pond.....		400,000
Whitman Pond.....		400,000
Leominster, Spectacle Pond.....		800,000
South Sudbury, Bright's pond.....		400,000
New Hampshire:		
Baboosic, Baboosic Lake.....		800,000
Raymond, Pawtuckaway Lake.....		400,000
Winchester, Forest Lake.....		600,000
New Jersey:		
Boonton, Dixson Pond.....		600,000
New York:		
Albany, Forest, Fish and Game Commission.....	15,000,000	
Lake Waccabuc, Waccabuc Lake.....		800,000
Lewisboro, Trinity Lake.....		600,000
Middletown, Hennessey Lake.....		600,000
New York, New York Aquarium.....	1,500,000	
Pennsylvania:		
Annaville, Quittapahilla Creek.....		400,000
Vermont:		
Montpelier, Groton Lake.....		800,000
Total.....	16,500,000	338,480,000

YELLOW BASS.

Disposition.	Fingerlings, yearlings, and adults.
Arkansas:	
Helena, Mississippi River.....	250

SEA BASS.

Disposition.	Fry.
Massachusetts:	
Falmouth, Buzzards Bay.....	253,000
Quissett Harbor.....	555,000
Total.....	808,000

MACKEREL.

Massachusetts:	
Falmouth, Buzzards Bay.....	388,000
Great Harbor.....	338,000
Gosnold, Vineyard Sound.....	38,000
Total.....	764,000

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

FRESHWATER DRUM.

Disposition.	Fingerlings, yearlings, and adults.
Arkansas:	
Helena, Mississippi River.....	8,950
Iowa:	
North McGregor, Mississippi River.....	1,500
Wisconsin:	
Prairie du Chien, Mississippi River.....	1,500
Total.....	11,950

COD.

Disposition.	Eggs.	Fry.
Maine:		
Boothbay Harbor, Boothbay Harbor.....		6,310,000
Kinekins Bay.....		4,304,000
Cape Elizabeth, Casco Bay.....		4,274,000
Massachusetts:		
Beverly, Massachusetts Bay.....		38,658,000
Falmouth, Buzzards Bay.....		9,733,000
Gloucester, Atlantic Ocean.....	9,854,000	22,510,000
Ipswich Bay.....		29,060,000
Massachusetts Bay.....		9,305,000
Gosnold, Buzzards Bay.....		5,979,000
Vineyard Sound.....		44,423,000
Great Harbor, Vineyard Sound.....		163,000
Manchester, Massachusetts Bay.....		4,630,000
Marblehead, Massachusetts Bay.....		2,580,000
Provincetown, Provincetown Harbor.....		802,000
Rockport, Atlantic Ocean.....		18,250,000
Ipswich Bay.....		9,000,000
Woods Hole, Eel Pond.....		253,000
Total.....	9,854,000	210,354,000

HADDOCK.

Disposition.	Fry.
Maine:	
Boothbay Harbor, Boothbay Harbor.....	712,000

POLLOCK.

Disposition.	Fry.	Disposition.	Fry.
Massachusetts:		Massachusetts—Continued.	
Beverly, Massachusetts Bay.....	1,330,000	Manchester, Massachusetts Bay.....	14,510,000
Gloucester, Atlantic Ocean.....	12,400,000	Rockport, Atlantic Ocean.....	5,800,000
Ipswich Bay.....	1,180,000	Total.....	38,140,000
Massachusetts Bay.....	2,920,000		

DETAILS OF DISTRIBUTION OF FISH AND FISH EGGS—Continued.

FLATFISH.

Disposition.	Fry.*	Disposition.	Fry.
Maine:		Massachusetts—Continued.	
Boothbay Harbor, Boothbay Harbor	380,176,000	Manchester, Massachusetts Bay	61,020,000
Linekin Bay	4,591,000	Monument Beach, Monument Beach Harbor	5,751,000
Mill Cove	17,398,000	Provincetown, Provincetown Harbor	4,678,000
Massachusetts:		Quissett, Quissett Harbor	7,797,000
Beverly, Massachusetts Bay	18,210,000	Roekport, Roekport Harbor	5,080,000
Falmouth, Buzzards Bay	11,156,000	Waquoit, Waquoit Bay	23,655,000
Great Harbor	6,138,000	Wareham, Wareham River	4,142,000
Little Harbor	2,047,000	Woods Hole, Great Harbor	11,661,000
Quissett Harbor	6,579,000	Woods Hole Harbor	6,090,000
Gloucester, Annisquam River	111,170,000	Rhode Island:	
Gloucester Harbor	109,540,000	East Greenwich, East Greenwich Bay	12,134,000
Ipswich Bay	7,800,000	Newport, Narragansett Bay	13,254,000
Gosnold, Buzzards Bay	21,783,000	Wickford, Wickford Harbor	6,434,000
Hadley Harbor	17,264,000		
Lackey Bay	12,328,000		
Robinson Hole	7,063,000		
Tarpaulin Cove	17,006,000		
Vineyard Sound	18,810,000		
		Total	930,755,000


LOBSTERS.

Maine:		Maine—Continued.	
Biddeford Pool, Biddeford Pool Harbor	10,000,000	South Addison, Pleasant Bay	250,000
Wood Isle Harbor	2,000,000	South Hancock, Skillings River	2,000,000
Boothbay Harbor, Boothbay Harbor	6,000,000	Southport, Atlantic Ocean	4,500,000
Bristol, Johns Bay	3,000,000	Cape Harbor	1,500,000
Brooklin, Naskog Harbor	250,000	Deckers Cove	1,500,000
Camden, Camden Harbor	1,000,000	Ebencook Harbor	500,000
Cape Porpoise, Cape Porpoise Harbor	4,500,000	St. George, Martins Harbor	1,000,000
Damariscotta, Damariscotta River	500,000	Stonington, Stonington Harbor	500,000
Deer Isle, Eggemoggin Reach	500,000	Surry, Union Bay	250,000
Southwest Harbor	400,000	Swan Isle, Old Harbor	500,000
East Boothbay, Linekin Bay	1,000,000	Tennants Harbor, Owls Head Bay	1,000,000
Eastport, Broad Cove	5,250,000	Vinal Haven, Vinal Haven Harbor	3,000,000
Falmouth, Caseo Bay	4,000,000	Wells, Wells Bay	500,000
Frenchboro, Frenchboro Harbor	500,000	West Lubee, Grand Manan Channel	350,000
Long Isle Harbor	1,500,000	Winnegance, New Meadows River	1,500,000
Friendship, Friendship Harbor	3,500,000	Winter Harbor, Winter Harbor	500,000
Isleboro, Penobscot Bay	400,000	York, York Harbor	4,500,000
Isleford, Isleford Harbor	1,500,000	Massachusetts:	
Isle of Shoals, Gulf of Maine	1,600,000	Bakers Island, Massachusetts Bay	300,000
Isle of Shoals Harbor	1,000,000	Beverly, Massachusetts Bay	1,400,000
Piscataqua River	400,000	Boston, Boston Bay	3,700,000
Jonesport, Roque Isle Harbor	650,000	Cohasset, Massachusetts Bay	834,000
Kennebunk, Kennebunk Port Harbor	500,000	Falmouth, Buzzards Bay	493,000
Wells Bay	500,000	Quissett Harbor	874,000
Kittery Point, Peppers Cove	1,500,000	Vineyard Sound	341,000
Little Deer Isle, Billings Cove	200,000	Gloucester, Atlantic Ocean	2,800,000
Lowry, Delanos Cove	3,000,000	Gloucester Harbor	600,000
Milbridge, Pigeon Hill Bay	2,000,000	Ipswich Bay	500,000
Mount Desert, Bass Harbor	1,000,000	Gosnold, Buzzards Bay	2,721,000
Southwest Harbor	500,000	Cuttjunk Harbor	1,087,000
New Harbor, New Harbor	3,500,000	Hadley Harbor	827,000
North Haven, North Haven Harbor	1,500,000	Laekeys Bay	2,868,000
Pulpit Harbor	1,000,000	Vineyard Sound	6,165,000
Orrs Island, Lowells Cove	500,000	Lanesville, Ipswich Bay	1,100,000
Pemaquid, Pemaquid Harbor	3,500,000	Manchester, Massachusetts Bay	2,860,000
Port Clyde, Port Clyde Harbor	1,000,000	Marblehead, Boston Bay	300,000
Portland, Caseo Bay	5,000,000	Roekport, Atlantic Ocean	600,000
Peaks Isle Roads	3,500,000	Roekport Harbor	600,000
Portland Harbor	2,500,000	Swampscott Harbor, Massachusetts Bay	200,000
Prospect Harbor, Bunkers Harbor	3,000,000	Woods Hole, Coles Pond	192,000
Dyers Bay	12,000,000	Great Harbor	1,097,000
Roekland, Roekland Harbor	3,000,000	New Hampshire:	
Roekport, Roekport Harbor	1,000,000	Stratford, Little New Harbor	4,000,000
Small Point, Horse Isle Harbor	500,000	Oregon:	
Small Point Harbor	2,106,000	Yaquina, Yaquina Bay	a 1,532
		Total	162,505,000

* Adults, of which 520 were lost in transit.

**DREDGING AND HYDROGRAPHIC RECORDS OF THE
U. S. FISHERIES STEAMER ALBATROSS DURING
THE PHILIPPINE EXPEDITION, 1907-1910**

Bureau of Fisheries Document No. 741



DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES STEAMER ALBATROSS DURING THE PHILIPPINE EXPEDITION, 1907-1910.

The Philippine cruise of the *Albatross* covered a greater period of time than any single expedition previously undertaken by that vessel. The ship left San Francisco October 16, 1907, and, sailing by way of the Hawaiian Islands, Midway, and Guam, arrived at Manila November 28. The stop at Midway, occasioned by a requisition of the vessel to carry stores from Honolulu to the United States marines stationed on Midway, was made the opportunity to take a small collection of the reef fishes and shore fauna of that group of islands. The number of fishes was very small, owing to the inability to carry enough explosive to do effective work, only 10 pounds of dynamite being allowed for use here and at Guam. Small collections were similarly made at this latter place when the ship stopped there for coal.

A two months' delay in the arrival of the stores which had been shipped from New York direct to Manila limited the vessel's activity for that period to the immediate vicinity of Manila. Thereafter the work was done by a series of short cruises made to the different parts of the Archipelago with Manila as a base for supplies and the deposit of collections.

During the period between February 2 and June 9, 1908, cruises were made to the southward, the first along the southwest side of Mindanao, thence through the Sulu groups, extending as far as Sandakan, Borneo; the second through the central group, including Panay, Negros, Cebu, Leyte, Masbate, and Marinduque; the third about the east and southeast coasts of Mindanao.

After the return to Manila from the last of these cruises it had become apparent that the *Albatross* required extensive repairs, and in August the ship left for Hongkong to have these made. Upon conclusion of this work in October Pratas Reef was visited and a number of soundings and trawl hauls were later made in the China Sea between that reef and the Batan Islands. Some work was done in the Batan and Babuyan islands and on the northern end of Luzon. Contemplated stops along the northwesterly coast of Luzon were prevented by bad weather which culminated in a typhoon.

During December, 1908, and January, 1909, a cruise through the Calamianes and the western and southern regions of Palawan was completed, touching on the return trip at Sandakan, Cagayan Sulu, and Iloilo. Late in January and early in February a number of cod trawl sets were made in the vicinity of Mariveles, but with indifferent success. The succeeding month was spent along the southern coasts of Luzon and adjacent islands, continuing thence southerly along the small islands to Bohol, thence westerly by the Cagayanes to the east coast of Palawan and northward into the Cuyos, returning to Manila early in April.

After a short trip to Lingayen Gulf early in May, the ship cruised along the small islands north of Samar and on the southeast coast of Luzon as far as Maculabo Island above San Miguel Bay, returning to Manila late in June. The latter part of July and all of August and September were spent in cruising from the southern coast of Samar, along southeastern Leyte, thence along the northern coast of Mindanao as far as Dapitan, thence northerly to Cebu, where some time was lost in repairing the boilers. The latter part of the period was consumed in further work in the vicinity of Zamboanga and along the Sulu group as far as Borneo, touching at a few small islands adjacent to the Borneo coast. Early in November the ship undertook a supplementary trip through the Dutch East Indies, touching at Menado, Ternate, Amboina, and Macassar, as well as at many intermediate points. On this trip a number of trawl hauls were made, including some exploration of the waters of the gulfs of Tomini and Boni in Celebes.

The homeward trip from Manila was begun January 21, 1910. Bad weather and other difficulties prevented the execution of orders to continue the work in the vicinity of Formosa and the Loo Choo Islands; at only two stops in Formosa were any collections made. After further repairs to the vessel in Japan, sail was set for the United States and San Francisco was reached May 4, 1910, after an absence of over two and one-half years.

EXPLANATION OF TABLES.

The last previous dredging station of the *Albatross* was no. 5095, the last hydrographic station was no. 4896, occupied during the northwestern Pacific cruise of 1906. (See Bureau of Fisheries Document 621.) Five hundred and seventy-seven dredging and 41 hydrographic stations were occupied during the Philippine expedition, extending the series of dredging stations to no. 5622 and the hydrographic series to no. 4937. In the tables the series are distinguished by the prefixed letters D and H, respectively.

Only those stations where the ship's gear was used (i. e., with the ship as an instrument) to collect natural-history specimens have been designated in the records as dredging stations. At times specimens were taken with dip nets during the occupation of a hydrographic station, but on account of the irregularity of such collecting the station was not regarded as a collecting station. No numbers have been given to the numerous shore stations, nor to minor collections made with the ship at anchor. But numbers have been given in the dredging series to hauls of the large intermediate net when used in a tideway with the ship at anchor.

Since the shore work constitutes such an important part of the total, the data regarding shore stations is shown in chronological order with the dredging stations, the locality, apparatus, etc., appearing in the appropriate columns. To economize time most of the reef collections of fishes were made with dynamite. The method was to locate the desirable fishes in the coral growth by means of a view glass (a glass-bottomed box) used from a boat. A small charge of dynamite with electrical connections was carefully lowered and discharged. Such fishes as floated were at once collected with a dip net, and the place marked by a buoy. As soon as the bottom had cleared it was searched and the dead fish gathered by diving or more usually by means of long-handled spears.

The various kinds of apparatus used at each station are recorded in the tables in chronological order, each on a separate line, opposite the station number, or, in case of unnumbered stations, opposite the locality, in the column "Apparatus."

The "Position" of a station is that point occupied by the vessel, as determined by the navigator at the time of beginning the first operation at that station. The position of the subsequent operations under the same station number corresponds in a general way to the line as indicated under "Drift." The distance covered by all the operations of a station is usually, however, not greater than the negligible error of observation, except in stations near shore determined by bearings.

In relation to the hydrographic information obtained, the degree of accuracy with which positions are located is of greater importance, and a description of the methods is necessary to the proper use of this information. A great part of the region traversed is still unsurveyed; and even where surveyed, parts are incorrectly or incompletely charted. Owing to press of work and lack of time, no opportunity was afforded to correct such errors, and the best available charts were therefore used as the basis of all determinations of position when in sight of land; in the column "Chart" is noted the number and edition of the chart used at each station.

When in sight of land position was fixed by compass bearings, and from the position so obtained *on the chart in use* the latitude and longitude were pricked off and set down in the record as the position of the station. If these charts should hereafter be corrected in latitude and longitude, the positions assigned to the stations must be changed accordingly.

In conformity with previous practice, an additional position, by true bearing and distance, of some prominent shore feature is given for each station when practicable. As viewed from the ship; the nearest and most prominent objects on shore from which the ship's position was determined were often topographical features, inconspicuous and unnamed on the chart, and impossible of identification by a brief written description. Therefore the bearings given in the tables were laid off from the plotted position on the chart to some object prominent on the chart, whether the object could actually be seen from the ship or not; though whenever convenient one of the two points taken for bearings by the navigator in determining the position is used in the table as the point of reference. The letters (S.), (N.), (W.), or (E.) indicate, respectively, the south, north, west, or east tangent of the point of reference after which they are placed; e. g., Verde Id. (E.)=eastern tangent of Verde Island.

All bearings are true unless otherwise indicated.

The spelling of all geographic names in these tables is that found on the charts designated in the column "Chart." There is considerable variation in this respect in the different issues of charts.

"Time of day" in the case of soundings indicates the time the plummet struck bottom; in the case of dredgings, the time at which the apparatus began to tow on the bottom; in the case of intermediate nets, the time at which the nets started to tow at the depth indicated; in the case of surface hauls, the time at which they were lowered into the water and began to be towed or the current to pass through them.

"Depth" (in fathoms) is the depth obtained by the sounding when a sounding was made. In cases where no sounding was made the depth is estimated from the chart, unless the station immediately follows another, in which case the depth obtained at the preceding station is given. In seine hauls the depths given are approximate, and represent the greatest depth of water through which the seine was hauled.

"Temperatures." The air temperatures are taken from the ship's log for the hour nearest the hour entered in the time column; the same is true of the surface temperatures where the towing commenced near the hour mark, but in other cases the surface temperature was

taken at the time given. The bottom temperature was taken at the time of sounding. All readings by Fahrenheit thermometer.

"Density." The water density is in all cases reduced to 15° C. The density of bottom water was ascertained from a sample taken by the Sigsbee water bottle. Inability to secure an accurate working of this instrument led to the discontinuance of the trials.

In the double column "Trial" is indicated the depth at which apparatus was worked, as well as the duration of operation. In the case of bottom apparatus this latter is the time during which it is supposed to be dragging on the bottom, up to the beginning of reeling in; for intermediate nets the time occupied in towing at the depth shown in the depth column is indicated by the first quantity, the time occupied in hoisting by the second; for surface nets the time indicated is the time actually towed at the surface.

In the double column of "Drift" is shown approximately the general direction in which the gear was hauled as well as the distance. The state of the currents and of the wind, with the exigencies incident to the steering of the ship, make this more or less inaccurate.

The apparatus used consisted of the usual beam trawls for all work on the bottom. All intermediate and surface work was done with a large tow net and small plankton or Kofoid nets, except an unsuccessful trial of a triangular shear-board net.

ABBREVIATIONS AND SYMBOLS.

- 12' Ag.....12-foot Agassiz beam trawl. The Agassiz type of beam trawl was used more and with better results than any other used during the cruise. The runners now in use stand 4 feet in height and the usual type of net carries a taut headline, making the full opening available. For deep-sea work where the possibility of upsetting the frame is great, a reversible net is used, with a running bolt-rope passing through the clips forward of the middle of the shoes. The use of this net is indicated by the abbreviation "rev."
- 25' Ag.The same runners used in the 12-foot frame but spread by use of two light spars for beams to a 25-foot opening. Used successfully on smooth bottoms.
- 9' Alb.-Blk.....9-foot Albatross-Blake beam trawl.
- B. A.....British Admiralty.
- 3-bd. int.a net with triangular opening operated by 3 shear boards and handled by a 3-part bridle from dredging cable—in no case successfully.
- 2' Blk.....a 2-foot Blake trawl, generally used from a steam launch or rowboat; net made of $\frac{1}{2}$ -inch webbing.
- botm.....bottom.
- C. S.....Coast Survey.
- D.....dredging, or collecting, station.
- dip.....ordinary dip net on a 12-inch or 14-inch ring, with bamboo handle; used extensively in reef fishing with dynamite and from the gang plank with electric light.
- dyn.....dynamite.

- e. 1.....electric light.
- H.....hydrographic station.
- H. O.....U. S. Hydrographic Office.
- hbr.....harbor.
- int. 3.....intermediate 3. This is a large ship's net on a 5½-foot ring; net about 11 feet long made of no. 0000 grit gauze, with about 3 feet of the bottom of no. 3 silk, and a brass bucket at the bottom. The outside netting is ½-inch webbing for the protection of the silk.
- int. 4.....intermediate 4; same as intermediate 3, but with an extension of 6 feet of 1-inch webbing carried to a 10-foot ring, thus increasing the opening to 10 feet.
- int. 5.....intermediate 5; similar to intermediate 4, but with no. 14 grit gauze only in the bottom part from the 3-foot ring to the bucket; above this ½-inch webbing to the 5½-foot ring, and thence 6 feet of ¾-inch webbing to the 10-foot ring. Equipped with a funnel of ¾-inch webbing.
- 9' Jn. dr.....Johnston oyster dredge. This is an Albatross-Blake beam trawl with a rake bar bolted at the heel. Used also in 6-foot length.
- K. 1.....a small plankton or Kofoid net, made of no. 12 silk, on a 14-inch ring.
- K. 2.....same as above, but made of no. 20 silk.
- K. 4.....same as above, but made of no. 3 silk.
- K. 5.....same as K. 2, but made of no. 1 silk.
- K. 6.....a net of same length as other Kofoid nets, but provided with clamps on opposite sides of the ring to attach directly to the cable; also with a bail from the ring to the bucket. Designed to lower and hoist with the ship lying to and the cable running vertically, thus making no catch except while ship is underway and towing.
- Lt.....light.
- Luc. sdr.....Lucas sounding machine.
- m. b.....mud bag. When more than one mud bag is used the two supplementary bags are rigged one at either end of the trawl frame.
- 6' McC.....6-foot McCormick; an adaptation of the Blake trawl frame, with rear beam bolted to bottom shoe and serving as a spindle on which bent teeth of ½ by 2 inch iron work as a rake. Not successful.
- 2' o. p.....open plankton net on 2-foot ring; made of no. 1 silk.
- spec.....specimen.
- 12' Tnr.....12-foot Tanner beam trawl.
- Tnr.-Blish sdr...Tanner-Blish sounding machine.
- therm.....Negretti & Zambra thermometer, with Tanner case.
- wat. bot.....Sigsbee water bottle.
- * signifies depth as shown by chart when no sounding has been made.
- ** signifies depth and character of bottom as obtained by sounding at previous station.
- † signifies nets towed astern, from taffrail, side by side.
- § signifies apparatus towed (horizontally) at depth indicated, during number of minutes given in the first period; then hoisted (vertically) to surface, net open, in time next shown.

The letters (a), (b), (c), (d), (e), when used with the abbreviation for sounding apparatus, indicate the kind of sounding cup used; thus,

- (a)...Sigsbee sounding rod. (d)...bail-cutter.
- (b)...Lucas snapper. (e)...ordinary lead with tallow.
- (c)...Lucas 4-tube sounding rod.

“Character of bottom,” determined by the specimens from the sounding cup, is expressed by abbreviations, the key to which is appended. It will be noted that these abbreviations are arbitrarily capitalized for nouns. When used as adjectives, however, the noun abbreviations are not capitalized.

bk....black.	fne....fine.	M.....Mud.	sctrd...scattered.
bl.....blue.	For...Foraminifera.	mrgn...marginal.	Sh.....Shells.
br....brown.	G.....Gravel.	Mss...Masses.	sml.....small.
br-gn..brownish-green.	Glob..Globigerina.	Oz....Ooze.	Sp.....Specks.
brk....broken.	gn....green.	P.....Pebbles.	St.....Stones.
C.....Clay.	gn-br..greenish-brown.	Ptr....Pteropod.	vol...volcanic.
Clmps.Clumps.	gn-gy..greenish-gray.	R.....Rock.	W.....Seaweed.
Co.....Coral.	gy....gray.	Rf.....Reef.	wh.....white.
crs....coarse.	hrd...hard.	rky....rocky.	
dk....dark.	Lav....Lava.	S.....Sand.	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Between Honolulu and Manila.</i>					
	Midway Ids. Harbor.....		1907. Nov. 7	10.00 a. m.	<i>fms.</i>	co. Clmps.; S.....
	do.....		Nov. 8	9.00 a. m.		Co.....
	Guam; Apra Bay (rf).....		Nov. 19	1.00 p. m.		mrgn. co. Rf.....
	do.....		Nov. 20	1.00 p. m.		co. Mss.....
	do.....		Nov. 21	9.00 a. m.		mrgn. Rf; sml. stag-horn Clmps; S.....
	do.....		do.....	1.00 p. m.		co. Mss.; S.....
	<i>Manila Bay and vicinity.^a</i>					
	Manila Bay (Luneta Beach).	C. S. 4240; Feb., 1907.	Dec. 6	3.00 p. m.		M., S.....
	Manila Bay, inside break-water (anch.).	do.....	do.....	7.00 p. m.	3.5	M.....
	do.....	do.....	Dec. 7	7.00 p. m.	3.5	M.....
	do.....	do.....	Dec. 8	7.00 p. m.	3.5	M.....
	Manila Bay (Malate Beach)..	C. S. 4712... Sept., 1904.	Dec. 9	9.00 a. m.		fne. S.....
	Manila Bay, inside break-water (anch.).	do.....	do.....	7.00 p. m.	3.5	M.....
	Manila Bay, inside break-water.	do.....	Dec. 11	8.00 p. m.		M.....
	do.....	do.....	Dec. 12	10.00 a. m.	2	M.....
	Manila Bay, outside break-water.	do.....	do.....	1.30 p. m.		M., sml. R.....
	Manila Bay (Luneta Beach).	do.....	Dec. 30	9.00 a. m.		M., S.....
	Manila Bay (near anch.)....	do.....	do.....	4.00 p. m.	3.5	M.....
	<i>China Sea off southern Luzon.</i>					
D. 5096	Corregidor Lt., N. 2.70 miles (14° 20' 23" N., 120° 34' 15" E.).	C. S. 4240; Feb., 1907.	1908. Jan. 2	10.42 a. m.	28	gy. M., S., Sh.....
				11.01 a. m.	28	gy. M., S., Sh.....
D. 5097	Corregidor Lt., N. 6° E., 3.60 miles (14° 19' 15" N., 120° 33' 52" E.).	do.....	do.....	11.18 a. m.	*30	gy. M., S., Sh.....
D. 5098	Corregidor Lt., N. 21° E., 4.30 miles (14° 18' 40" N., 120° 32' 40" E.).	do.....	do.....	12.44 p. m.	*38	gy. M., S., Sh.....
D. 5099	Corregidor Lt., N. 36° E., 4.80 miles (14° 18' 55" N., 120° 31' 20" E.).	do.....	do.....	1.21 p. m.	*30	gy. M., S., Sh.....
D. 5100	Corregidor Lt., N. 16° E., 5.70 miles (14° 17' 15" N., 120° 32' 40" E.).	do.....	do.....	2.15 p. m.	35	gy. S.....
D. 5101	Corregidor Lt., S. 82° E., 10.50 miles (14° 24' 30" N., 120° 23' 20" E.).	do.....	Jan. 6	2.22 p. m.	35	gy. S.....
				1.16 p. m.	*43
D. 5102	Sueste Pt. Lt., S. 85° W., 1.20 miles (14° 45' N., 120° 12' 30" E.).	C. S. 4254; Sept., 1902.	do.....	4.20 p. m.	*33
	Subig Bay (Subig anch.)....	do.....	do.....	7.00 p. m.	11
	Subig Bay, Subig (beach)....	do.....	Jan. 7	9.00 a. m.		S.....
D. 5103	Calaclan Pt., S. 86° E., 2.50 miles (14° 49' 30" N., 120° 13' 30" E.).	do.....	do.....	1.46 p. m.	*20	gy. M.....

^a From December 16 to 21 a shore party made collections at the mouth of the Santa Cruz River and the adjacent shore of Laguna de Bay and visited the markets at Santa Cruz and Majajjay. A party visited Taal Lake December 24 to 29 and made collections by seining (45' seine) on the south side of Taal Id., and by purchase from natives on the Pansipit River, and at Taal December 31 and January 1 a shore party made collections on Mariguina River.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910.

Temperatures.			Density.		Apparatus.	Trawl.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
					dyn.....	8-12 ft..	1 00			Work interrupted by storm.
					dyn.....	20-30 ft.	2 00			
					dyn.....	6-20 ft..	4 00			
					dyn.....	6-20 ft..	3 00			
					dyn.....	3-10 ft..	3 00			Mostly on shore flat.
					dyn.....	6-20 ft..	2 00			
					100' seine.....	4 ft	2 30			5 hauls.
					dip; e. l.....	surf....	2 00			
					dip; e. l.....	surf....	2 00			
					dip; e. l.....	surf....	2 00			
					150' seine.....	6 ft	2 30			Do.
					dip; e. l.....	surf....	2 00			
					2' o. p.....	surf....	20			Towed from steam launch.
					2' Blk.....	botm....				Do.
					2' Blk.....	botm....				Several hauls from mouth of Pasig River to outer entrance through breakwater.
					150' seine.....	4 ft	2 30			5 hauls.
					2 wire traps...	botm....				Finally hauled on Jan. 4, 1908.
79.5	79				Tnr.-Blish.sdr. (b).					
79.7	79				9' Tnr.; m. b..	botm...	21	SW. <i>a</i>		Veered 5 fms. during haul, not on bottom; water-haul.
80	79				9' Tnr.; m. b..	botm...	19	NW.byW. <i>a</i>		Veered at 5 minute intervals from 75 to 94 and to 104 fms. Trawl capsize on bottom, but made a small catch.
82	79				9' Tnr.; m. b..	botm...	20	W.byN. <i>a</i>		Net capsize on bottom, but made a small catch.
81	80				9' Tnr.; m. b..	botm...	20	W. by N. <i>a</i>		
86	80				Tnr.-Blish.sdr. (b).					
					9' Tnr.; m. b..	botm...	20	NE. <i>a</i>		
82	78		1.02391		int. 4 §.....	37 fms..	20	NW. $\frac{1}{2}$ W.		70 fms. dredge cable out.
							4			
86.5	81		1.02447		int. 4 §.....	28 fms..	20	N. 11° E.		Cable veered from 45 to 57 fms. during haul.
							3			
					dip; e. l.....	surf....	2 00			
					250' seine.....	20 ft	2 30			5 hauls.
84	79				12' Tnr.; m. b.	botm...	20	S. 45° E.	0.6	

a Course steered by ship.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>China Sea off southern Luzon—Continued.</i>					
	Olongapo (beach).....	C. S. 4254;	1908, Jan. 7	2.00 p. m.	<i>fms.</i>	S.....
	Beach opposite Olongapo.....	do.....	do.....	3.30 p. m.		grassy.....
D. 5104	Olongapo (anch.).....	do.....	do.....	7.00 p. m.	13	
	Sueste Pt. Lt., S. 58° W., 1.30 miles (14° 45' 48" N., 120° 12' 20" E.).....	do.....	Jan. 8	10.20 a. m.	*33	(?).....
D. 5105	Sueste Pt. Lt., N. 57° W., 1.90 miles (14° 43' 55" N., 120° 12' 50" E.).....	do.....	do.....	11.06 a. m.	*25	(?).....
	Grande I. (rf.).....	do.....	do.....	1.00 p. m.		setrd. Clmps. Co.....
	Port Binanga (beach).....	do.....	do.....	2.00 p. m.		
	Port Binanga (anch.).....	do.....	do.....	7.00 p. m.	6	
	Port Binanga (rf.).....	do.....	Jan. 9	8.30 a. m.		setrd. Clmps. Co.....
D. 5106	Corregidor Lt., S. 57° E., 2.25 miles (14° 23' 55" N., 120° 32' 33" E.).....	C. S. 4240; Feb., 1907.	do.....	1.58 p. m.	*37	gy. M.....
D. 5107	Corregidor Lt., S. 17° E., 1.75 miles (14° 24' 30" N., 120° 33' 40" E.).....	do.....	do.....	2.38 p. m.	*28	gy. M.....
	Manila Bay (Luneta Beach). Limbones Cove (E. shore, beach).....	do.....	Jan. 13 Jan. 14	4.30 p. m.		fne. S S., P., Co.....
	Limbones Cove (SW. shore, rf.).....	do.....	do.....	4.30 p. m.		solid Co.....
	Limbones Cove (anch.).....	do.....	do.....	7.00 p. m.	10	
D. 5108	Corregidor Lt., N. 39° E., 22.50 miles (14° 05' 5" N., 120° 19' 45" E.).....	do.....	Jan. 15	8.01 a. m.	13	Co.....
				8.34 a. m.	13	Co.....
				8.47 a. m.	16	Co.....
				9.00 a. m.	16	Co.....
				9.20 a. m.	16	Co.....
D. 5109	Corregidor Lt., N. 42° E., 25.80 miles (14° 03' 45" N., 120° 16' 30" E.).....	do.....	do.....	10.26 a. m.	10	Co.....
				10.43 a. m.	12	Co.....
D. 5110	Corregidor Lt., N. 20° E., 25 miles (13° 59' 20" N., 120° 75' 45" E.).....	do.....	do.....	3.18 p. m.	135	dk. gy. M.....
				3.32 p. m.	135	dk. gy. M.....
	Nasugbu Bay (anch.).....	do.....	do.....	7.00 p. m.	10	
	Nasugbu Bay (beach near town).....	do.....	Jan. 16	9.00 a. m.		S.....
	Nasugbu Bay (Pillar Rock, rf.).....	do.....	do.....	9.00 a. m.		setrd. Clmps. Co.....
D. 5111	Sombrero Id., S. 41° E., 4.50 miles (13° 45' 15" N., 120° 46' 30" E.).....	do.....	do.....	2.38 p. m.	236	
				3.08 p. m.	236	gn. M.....
				7.00 p. m.	10	
D. 5112	Balayan Bay (Taal anch.).....	do.....	do.....	7.00 p. m.	10	
	Sombrero Id., S. 18° E., 6.75 miles (13° 48' 22" N., 120° 47' 25" E.).....	do.....	Jan. 17	2.06 p. m.	177	dk. gn. M.....
				2.33 p. m.	177	dk. gn. M.....
D. 5113	Sombrero Id., S. 7° W., 9.50 miles (13° 51' 30" N., 120° 50' 30" E.).....	do.....	do.....	3.43 p. m.	159	dk. gn. M.....
				4.02 p. m.	159	dk. gn. M.....
	<i>Balayan Bay and Verde Id. Passage.^a</i>					
	Balayan Bay (Ligpo Pt. rf.).....	C. S. 4240,	Jan. 18	10.00 a. m.		dense Co., S.....

^a Collecting trip to Taal Lake on Jan. 18. Dredging with hand dredge.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trawl.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>			<i>mi.</i>	
					250' seine.....	20 ft ...	1 15			1 haul.
					250' seine.....	8 ft ...	1 15			Do
81	78				dip.; e. l.	surf....	2 00	S. 22° W..	0.8	
					12' Tnr.; m. b.	botm...	20			
81	78				12' Tnr.; m. b.	botm...	20	N. 60° W..	(?)	
					dyn.....	6-20 ft..	4 30			4 hauls.
					150' seine.....	surf....	2 00			
					dip.; e. l.	surf....	2 00			
					dyn.....	6-15 ft..	2 00			
86.5	78		1.02393		12' Tnr.; m. b.	botm...	20	N. 48° E..	1.2	Tail lashing slipped; no catch except in mud bag.
84.5	78		1.02379		12' Tnr.; m. b.	botm...	20	N. 44° E..	1.7	
					150' seine.....	4 ft ...				
					250' seine.....	12 ft ...	1 30			
					dyn.....	6-12 ft..	1 30			
					dip.; e. l.	surf....	2 00			
80	80	80	1.02406		Tnr-Blish sdr. (b).					
80	80				9' Alb.-Blk.; m. b.	botm...	1	N. 36° E..		Dredging cable fouled gin block. Trawl not dragged on bottom.
81	80				Tnr-Blish sdr. (b).					
81	80				8 swabs.....	botm...	10	S.....	.3	
81	80				9 hand lines.	botm...	23			No catch.
82	80		1.02386		9' Alb.-Blk.	botm...	12		(?)	Trawl immediately torn on coral.
82	80				8 swabs.....	botm...	11		(?)	Soundings with hand lead.
89	80	59	1.02406		Tnr-Blish sdr. (b).					
85	80				12' Tnr.; m. b.	botm...	20	N. 20° E..	.6	20 fms. cable veered during haul.
					dip.; e. l.	surf....	2 00			
					130' seine.....	surf....	2 30			
					dyn.....	6-15 ft..	3 00			
84	80				Tnr-Blish sdr. (b).					Soundling cup lost; therm. did not trip.
84	80				12' Tnr.; m. b.	botm...	30	N. 22° E..	1.8	
					dip.; e. l.	surf....	1 30			
86	80	52.4	1.02416	1.02496	Tnr-Blish sdr. (b).					
84	80				12' Tnr.; m. b.	botm...	30	N. 13° E..	1.3	
82	80		1.02413		Tnr-Blish sdr. (e).					
80	80				12' Tnr.; m. b.	botm...	10	N. 9° E..	.6	Uneven bottom.
					dyn.....	6-20 ft..	5 00			

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Balayan Bay and Verde Id. Passage—Continued.</i>					
.....	Balayan Bay (near beach, Taal).	C. S. 4240; Feb., 1907.	1908. Jan. 19	3.00 p. m.	<i>fms.</i>	blk. S., M.
.....	Balayan Bay (Taal, anch.)do.....do.....	7.00 p. m.	10
.....	Maricaban Id. (rf. inside Sepoc Pt.).do.....	Jan. 20	9.00 a. m.	dense Co., S.
D. 5114	Sombrero Id. N. 36° E., 7.2 miles (13° 36' 11" N., 120° 45' 26" E.).do.....do.....	10.49 a. m. 11.17 a. m.	340 340	fne. S. fne. S.
D. 5115	Sombrero Id. N. 49° E., 7.30 miles (13° 37' 11" N., 120° 43' 40" E.).do.....do.....	1.08 p. m. 1.41 p. m.	340 340	(?) (?) (?) (?)
D. 5116	Sombrero Id. N. 69° E., 2.50 miles (13° 41' N., 120° 47' 05" E.).do.....do.....	2.53 p. m. 3.13 p. m.	200 200	(?) (?)
D. 5117	Sombrero Id. S. 17° E., 10.80 miles (13° 52' 22" N., 120° 46' 22" E.).do.....	Jan. 21	9.10 a. m.	118	(?) (?)
D. 5118	Sombrero Id. S. 47° E., 10 miles (13° 48' 45" N., 120° 41' 51" E.).do.....do.....	9.27 a. m. 10.41 a. m.	118 159	dk. gn. M. dk. gn. M.
D. 5119	Sombrero Id. S. 80° E., 18.90 miles (13° 45' 05" N., 120° 30' 30" E.).do.....do.....	11.00 a. m. 1.24 p. m. 1.56 p. m.	159 394 394	dk. gn. M. gn. M., S. gn. M., S.
D. 5120	Sombrero Id., S. 79° 30' E., 19.2 miles (13° 45' 30" N., 120° 30' 15" E.).do.....do.....	2.41 p. m. 3.10 p. m.	393 393	gn. M., S.
.....	Nasugbu Bay (anch.)do.....do.....	7.30 p. m.	10
.....	Manila Bay (inside break-water).do.....	Jan. 28	11.00 a. m.
.....	Manila Bay (inside break-water, anch.)do.....	Jan. 31	10.00 a. m.
	<i>East coast of Mindoro.</i>					
D. 5121	Malabrigo Lt., N. 14° W., 9 miles (13° 27' 20" N., 121° 17' 45" E.).	C. S. 4714; June, 1906.	Feb. 2	8.14 a. m.	108	dk. gn. M. dk. gn. M. gn. M.
D. 5122	Malabrigo Lt., N. 46° W., 20.60 miles (13° 21' 30" N., 120° 30' 33" E.).do.....do.....	8.30 a. m. 10.34 a. m.	108 220
D. 5123	Malabrigo Lt., N. 44° W., 32.50 miles (13° 12' 45" N., 121° 38' 45" E.).do.....do.....	10.59 a. m. 1.09 p. m.	220 283	gn. M. gn. M.
D. 5124	Pt. Origen (N.), S. 56° E., 20.75 miles (12° 52' N., 121° 48' 30" E.).do.....do.....	1.44 p. m. 5.04 p. m. 5.38 p. m.	283 281 281	gn. M. sft. gn. M. sft. gn. M.
	<i>Sulu Sea, vicinity southern Panay.</i>					
D. 5125	Nogas Id. (W.), S. 11° E., 24 miles (10° 48' N., 121° 48' 30" E.).	C. S. 4718, Dec., 1906.	Feb. 3	9.07 a. m. 9.41 a. m.	411 411	gn. M.
D. 5126	Nogas Id. (W.), S. 26° 30' E., 11.75 miles (10° 34' 45" N., 121° 47' 30" E.).do.....do.....	1.05 p. m. 2.00 p. m.	742 742	sft. gn. M. sft. gn. M.
.....	Naso Pt., Panay (anch.)do.....do.....	7.00 p. m.	10
.....	Naso Pt., Panay (near anch.)do.....do.....	7.00 p. m.
.....	Naso Pt., Panay (beach)do.....	Feb. 4	8.30 a. m.
.....	Naso Pt., Panay (shore, tide pools).do.....do.....	9.00 a. m.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Tempera- tures.			Density.		Apparatus.	Trawl.		Drift.		Remarks.
Air.	Surface.	Bottom.	Sur- face.	Bot- tom.		Depth.	Dura- tion.	Direction.	Distance.	
° F.	° F.	° F.				h. m.		mi.		
					250 fm. seine		2 00			Purse seine owned and hauled by native fishermen.
					dip; e. l.	surf.	1 30			
					dyn.	6-20 ft.	5 00			
81.5	79		1.02447		Luc. sdr. (e).					
84	80				12' Tnr.; m. b.	botm.	20	N. 54° E.	0.5	Cable veered from 400 to 520 fms. during haul.
82	80	(?)	1.02434	1.02454	Luc. sdr. (b).					Sounding cup did not close. Therm. not properly attached and fouled water bottle.
83	80				12' Tnr.; m. b.	botm.	20	N. 43° E.	1.0	Therm. not properly attached; fouled stray line. No specimen in sounding cup.
86	80	50.2	1.02426		Luc. sdr. (b).					
86	80				12' Tnr.; m. b.	botm.	20	N. 5° E.	0.5	
82	79		1.02475		Tnr.-Blish sdr. (b).					
82	79				12' Tnr.; m. b.	botm.	20	N. 31° W.	0.8	
81	79		1.02426		Tnr.-Blish sdr. (b).					
81	79				12' Tnr.; m. b.	botm.	30	N. 50° W.	0.8	
82	80	43.7	1.02386	1.02468	Luc. sdr. (b).					
82	80				12' Tnr.; m. b.	botm.	9	N. 23° E.	1.0	
82	80	43.7	1.02386	1.02480	Luc. sdr. (b).					
82	80				int. 4 §.	350 fms.	20 17	N. 5° W.	1.0	393 fms. dredge cable out.
					dip; e. l.	surf.	1 30			
					2' o. p.	surf.	15			Towed from steam launch.
					dyn. cap.; dip.	surf.				
76	79		1.02420		Tnr.-Blish sdr. (b).					
76	79				12' Tnr.; m. b.	botm.	20	S. 79° E.	1.0	Snapper failed to close.
78	79		1.02489		Tnr.-Blish sdr. (b).					Do.
79	79				12' Tnr.; m. b.	botm.	20	S. 59° E.	1.3	
80	79		1.02475		Tnr.-Blish sdr. (b).					
79	79				12' Tnr.; m. b.	botm.	20	S. 6° W.	1.3	Do.
82	79		1.02468		Tnr.-Blish sdr. (b).					
80.5	79				12' Tnr.; m. b.	botm.	17	S. 9° W.	1.5	
81	80	50	1.02444	1.02475	Luc. sdr. (b).					
83.5	80				int. 4 §.	365 fms.	20 26	N. 62° W.	1.5	550 fms. dredge cable out.
83	80	49.5	(?)	(?)	Luc. sdr. (a).					No specimen in water bottle.
84	80				12' Tnr.; m. b.	botm.	20	N. 81° W.	1.5	
					dip; e. l.	surf.	1 30			
					5 gill nets.	botm. and surf.				Set over night.
					130' seine.	10 ft	3 00			6 hauls.
					copper sulphate.		2 00			

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Sulu Sea, vicinity southern Panay—Continued.</i>					
D. 5127	Nogas Id. (W.), N. 11° 30' E., 22 miles (10° 02' 45" N., 121° 48' 15" E.).	C. S. 4718; Dec., 1906.	1908. Feb. 4.	2.57 p. m. 4.06 p. m.	fms. - 958 958	gy. M., Glob..... gy. M., Glob.....
D. 5128	Nogas Id. (W.), N. 6° E., 32.50 miles (9° 52' 10" N., 121° 49' 35" E.).do.....do.....	7.05 p. m.		
	<i>Sulu Sea off western Min- danao.</i>					
H. 4897	Dulunguin Pt., S. 70° E., 4.80 mile (7° 46' N., 122° E.).	C. S. 4723, Oct., 1905.	Feb. 5	11.43 a. m.	1,570	gy. M., Glob.....
H. 4898	Dulunguin Pt., N. 50° E., 1 mile (7° 43' 45" N., 122° 03' 45" E.).do.....do.....	1.13 p. m.	221	gy. M., Glob.....
D. 5129	Dulunguin Pt., N. 44° E., 3.80 miles (7° 41' 30" N., 122° 01' 45" E.).do.....do.....	2.04 p. m. 2.23 p. m.	0-100	
D. 5130	Dulunguin Pt., N. 1° W., 9.50 miles (7° 35' N., 122° 04' 45" E.).do.....do.....	3.29 p. m. 3.48 p. m.	102 102	
.....	Panabutan Bay (NW beach, near river).	C. S. 4644; July, 1905.do.....	5.00 p. m.		sft. M., S.....
.....	Panabutan Bay (anch.)do.....do.....	7.30 p. m.	11	
H. 4899	Id. off Panabutan Pt., S. 75° W., 3 miles.do.....	Feb. 6	8.48 a. m.	18	sft. gn. M.....
H. 4900	Id. off Panabutan Pt., W., 0.30 mile.do.....do.....	8.58 a. m.	19	sft. gn. M.....
H. 4901	Id. off Panabutan Pt., N. 52° W., 0.30 mile.do.....do.....	9.04 a. m.	21	gn. M., S.....
.....	Panabutan Bay (beach)do.....do.....	9.00 a. m.		S., M.....
.....	Panabutan Bay (Siriguay Pt., rf.).do.....do.....	9.00 a. m.		scftrd. Co.....
H. 4902	Id. off Panabutan Pt., N. 31° W., 0.50 mile.do.....do.....	9.10 a. m.	23	gn. M., fine S.....
H. 4903	Id. off Panabutan Pt., N. 15° W., 0.50 mile.do.....do.....		27	co. S.....
D. 5131	Id. off Panabutan Pt., N. 20° E., 0.40 mile.do.....do.....	9.14 a. m.	27	gn. M., co. S.....
D. 5132	Id. off Panabutan Pt., N. 15° W., 0.30 mile.do.....do.....	9.27 a. m. 9.54 a. m.	27 *26	gn. M., co. S..... gn. M., S.....
H. 4904	Id. off Panabutan Pt., N. 62° E., 0.30 mile.do.....do.....	10.23 a. m.	38	gn. M., S.....
D. 5133	Id. off Panabutan Pt., N. 52° E., 1.50 miles.do.....do.....	10.28 a. m.	38	gn. M., S.....
.....	Caldera Bay (anch.)do.....do.....	10.40 a. m. 7.30 p. m.	38	gn. M., S.....
	<i>Sulu Archipelago, near Ba- sitan Id.</i>					
D. 5134	Balukbaluk Id. (N.) S. 59° W., 6.25 miles (6° 44' 45" N., 121° 48' E.).	C. S. 4511; Dec., 1904.	Feb. 7	7.14 a. m. 7.22 a. m.	25 25	fine S..... fine S.....
D. 5134a	Balukbaluk Id. (N.), S. 59° W., 4.90 miles (6° 44' 12" N., 121° 46' 55" E.).do.....do.....	7.54 a. m. 8.05 a. m.	34 34	gy. S..... gy. S.....

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Tempera- tures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Sur- face.	Bot- tom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
84.5	80	50.1	1.02477	1.02516	Luc.sdr. (a).					
83	81				9' alb.-Blk.; 2 m. b.	botm...	20	N. 9° W...	(?)	4.25 mi. distance given by re- corder.
82	80				int. 4.	surf...	20	S. 6° E...	0.6	
82	80				Luc.sdr. (a).					
82	80				Tnr.-Blish sdr. (b).					First attempt re- sulted in loss of all the apparatus used.
81	80	57.6	1.02482		Tnr.-Blish sdr.					Density at 100 fms. 1.02495.
80	80				int. 4 §	100 fms.	20 8	S. 31° W...	1.3	193 fms. dredge cable out.
81.5	79.5	59.2	1.02447	1.02451	Luc.sdr. (a).					
80.5	80				9' alb.-Blk	botm...				Trawl fouled bot- tom and carried away.
					130' seine	12 ft	30			1 haul.
					dip; e. l.	surf...	2 00			
					Tnr.-Blish sdr. (e).					
					Tnr.-Blish sdr. (e).					
					Tnr.-Blish sdr. (e).					
					175' seine		2 00			
					dyn.	8-15 ft.	2 00			Water brackish. Coral unthrifty.
					Tnr.-Blish sdr. (e).					
					Tnr.-Blish sdr. (e).					
88	79		1.02447		Tnr.-Blish sdr. (e).					
88	79				9' Tnr.; m. b.	botm...	13	N. 43° E...	.3	
85	79		1.02447		9' Tnr.; m. b.	botm...	20	S. 69° W...	.7	
					Tnr.-Blish sdr. (e).					
85.5	79.5		1.02447		Tnr.-Blish sdr. (e).					
85	80				9' Tnr.; m. b.	botm...	16	S. 21° E...	.4	
					2' o. p.	surf...	20			Set in tide current at gangway.
82	78	?	1.02497		Tnr.-Blish sdr. (e).					Therm. not allow- ed time to set.
81	78				9' Tnr.; m. b.	botm...	20	S. 42° W...	.9	Ship drifted to po- sition of 5134a while getting ap- paratus ready.
83	78	76.2			Tnr.-Blish sdr. (e).					15 sec. allowed for therm. to set.
83	78				int. 4 §	25 fms.	20 2	N. 26° E...	.9	50 fms. dredge cable out.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>Vicinity of Jolo.</i>						
D. 5135	Jolo Lt., S. 46° W., 11.90 miles (6° 11' 50" N., 121° 08' 20" E.).	C. S. 4542; Apr., 1903.	1908. Feb. 7	2.29 p. m.	<i>fms.</i> 161	fne. co. S.
.....	Jolo (anch.)	do	do	2.50 p. m.	161	fne. co. S.
.....	do	do	do	7.30 p. m.	14
.....	do	do	Feb. 8	7.30 p. m.	14
.....	Marongas Id., S. side	do	Feb. 10	1.30 p. m.	sctrd. Co., S.
.....	Pangasinan Id., S. Pt. (rf.)	do	Feb. 13	3.00 p. m.	sctrd. Co.
.....	Jolo (anch.)	do	do	7.30 p. m.	14
D. 5136	Jolo Lt., S. 37° E., 0.70 mile (6° 04' 20" N., 120° 59' 20" E.).	do	Feb. 14	8.50 a. m.	22	S., Sh.
.....	9.07 a. m.	22	S., Sh.
D. 5137	Jolo Lt., S. 61° E., 1.30 miles (6° 04' 25" N., 120° 58' 30" E.).	do	do	9.44 a. m.	20	S., Sh.
.....	9.55 a. m.	20	S., Sh.
D. 5138	Jolo Lt., S. 19° E., 2.50 miles (6° 06' N., 120° 58' 50" E.).	do	do	10.50 a. m.	19	S., Co.
.....	10.55 a. m.	19	S., Co.
D. 5139	Jolo Lt., S. 51° W., 3.60 miles (6° 06' N., 121° 02' 30" E.).	do	do	1.02 p. m.	20	co. S.
.....	1.13 p. m.	20	co. S.
D. 5140	Jolo Lt., S. 33° W., 6.10 miles (6° 08' 45" N., 121° 03' E.).	do	do	1.58 p. m.	76	fne. co. S.
.....	2.09 p. m.	76	fne. co. S.
.....	Bubuan Id., S. Pt. (rf.)	do	do	4.00 p. m.	co. Mss
.....	Bubuan Id. (anch.)	do	do	7.30 p. m.	12
D. 5141	Jolo Lt., S. 17° E., 5.50 miles (6° 09' N., 120° 58' E.).	do	Feb. 15	8.39 a. m.	29	co. S.
.....	8.47 a. m.	29	co. S.
D. 5142	Jolo Lt., S. 50° W., 3.90 miles (6° 06' 10" N., 121° 02' 40" E.).	do	do	10.26 a. m.	21	co. S., Sh.
.....	10.33 a. m.	21	co. S., Sh.
D. 5143	Jolo Lt., S. 50° W., 3.40 miles (6° 05' 50" N., 121° 02' 15" E.).	do	do	11.05 a. m.	19	co. S.
.....	11.09 a. m.	19	co. S.
D. 5144	Jolo Lt., S. 50° W., 3.40 miles (6° 05' 50" N., 121° 02' 15" E.).	do	do	11.19 a. m.	19	co. S.
.....	11.26 a. m.	19	co. S.
D. 5145	Jolo Lt., S. 16° E., 0.85 mile (6° 04' 30" N., 120° 59' 30" E.).	do	do	1.37 p. m.	23	co. S., Sh.
.....	1.44 p. m.	23	co. S., Sh.
<i>Sulu Archipelago, vicinity of Siasi.</i>						
D. 5146	Sulade Id. (E.), N. 18° W., 3.40 miles (5° 46' 40" N., 120° 48' 50" E.).	C. S. 4542; Apr., 1903.	Feb. 16	10.04 a. m.	24	co. S., Sh.
.....	10.11 a. m.	24	co. S., Sh.
D. 5147	Sulade Id. (E.), N. 3° E., 8.40 miles (5° 41' 40" N., 120° 47' 10" E.).	do	do	11.20 a. m.	21	co. S., Sh.
.....	11.27 a. m.	21	co. S., Sh.
D. 5148	Sirun Id. (N.), S. 80° W., 3.80 miles (5° 35' 40" N., 120° 47' 30" E.).	C. S. 4544; Oct., 1906.	do	1.00 p. m.	17	co. S.
.....	1.07 p. m.	17	co. S.
H. 4905	Sirun Id. (W.), N. 33° E., 2.43 miles (5° 32' 50" N., 120° 42' 15" E.).	do	Feb. 18	10	S., Co., Sh.
.....
D. 5149	Sirun Id. (W.), N. 39° E., 2.40 miles (5° 33' N., 120° 42' 10" E.).	do	do	9.26 a. m.	10	Co., Sh.
.....	9.32 a. m.	10	Co., Sh.
D. 5150	Sirun Id. (W.), N. 34° E., 11.7 miles (5° 23' 20" N., 120° 35' 45" E.).	C. S. 4514; Jan., 1906.	do	11.37 a. m.	21	co. S., Sh.
.....	11.43 a. m.	21	co. S., Sh.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Tempera- tures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Sur- face.	Bot- tom.		Depth.	Dura- tion.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
80.5	80	57.4	1.02457		Tnr.-Blish sdr. (e)					
80.5	81				12' Tnr.; m. b.	botm...	20	S. 26° W..	1.0	
					dip. e. l.	surf.	2 00			
					dip. e. l.	surf.	2 00			
					diving	4-8ft.	3 00			Coral heads taken ashore.
					dyn.	5-12ft.	2 00			
					dip. c. l.	surf.	1 30			
84	80		1.02489		Tnr.-Blish sdr. (e)					
83	79				12' Agz.; 2 m. b.	botm...	20	N. 72° W..	0.6	Lead line carried away.
84	80				Tnr.-Blish sdr. (e)					
84	80				12' Agz.; 2 m. b.	botm...	20	N. 27° W..	0.6	
85	80				Tnr.-Blish sdr. (e)					
85	80				12' Agz.; 2 m. b.	botm...	20	N. 15° E..	0.6	1 mud bag carried away.
83	80		1.02457		Tnr.-Blish sdr. (e)					
83	80				12' Agz.; m. b.	botm...	04	S. 45° E..	0.2	
83	80		1.02477		Tnr.-Blish sdr. (e)					
83	82				12' Agz. rev.; m. b.	botm...	20	N. 70° W..	0.8	
					dyn.	8-20ft.	1 00			
					dip.; e. l.	surf.	1 30			
81	78		1.02461		Tnr.-Blish sdr. (e)					
81	78				12' Agz.; m. b.	botm...	18	N. 13° E..	0.5	
87	80		1.02503		Tnr.-Blish sdr. (e)					
88	80				12' Agz.; m. b.	botm...	11	W.....	0.5	1 bridle-stop car- ried away. Sounding lead carried away.
89	80		1.02442		Tnr.-Blish sdr. (e)					
89	80				12' Agz.; m. b.	botm...	4			Fouled bottom; mud bag torn; no distance made.
91	81		1.02514		Tnr.-Blish sdr. (e)					
91	81				12' Agz.; m. b.	botm...	20	N. 45° W..	0.6	
88	77		1.02482		Tnr.-Blish sdr. (e)					
88	77				12' Agz.; m. b.	botm...	15	S.....	.6	
82	80		1.02468		Tnr.-Blish sdr. (e)					
82	81				12' Agz.; m. b.	botm...	20	N. 77° W..	1.1	
85	80		1.02447		Tnr.-Blish sdr. (e)					
84	80				12' Agz.; m. b.	botm...	20	S. 72° E..	.4	
82.5	80		1.02523		Tnr.-Blish sdr. (e)					
82.5	80				12' Agz.; m. b.	botm...	18	S. 51° E..	.7	
					Tnr.-Blish sdr. (e)					
81	78		1.02509		Tnr.-Blish sdr. (e)					
84	78				12' Agz.; m. b.	botm...	20	N. 10° W..	.8	
82	78		1.02495		Tnr.-Blish sdr. (e)					
82	78				12' Agz.; m. b.	botm...				Net fouled bottom; 1 bridle stop car- ried away; no distance made.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>Sulu Archipelago, Tawi Tawi Group.</i>						
D. 5151	Sirun Id. (C.), N. 58° E., 19.3 miles (5° 24' 40" N., 120° 27' 15" E.).	C. S. 4514; Jan., 1906.	1908. Feb. 18	1.02 p. m.	<i>fms.</i> 24	co. S., Sh.....
D. 5152	Pajumajan Id. (W.), S. 2° W., 2 miles (5° 22' 55" N., 120° 15' 45" E.).do.....do.....	1.07 p. m. 3.21 p. m.	24 34	co. S., Sh..... wh. S.....
D. 5153	Dos Amigos Bay (anch.)..... Tocanhi Pt., S. 27° E., 2.10 miles (5° 18' 10" N., 120° 2' 55" E.).do.....do..... Feb. 19	3.28 p. m. 9.00 a. m.	34 49	wh. S..... co. S., Sh.....
D. 5154	Bakun Pt., S. 11° W., 0.70 mile (5° 14' 50" N., 119° 58' 45" E.).	H. O. 1852; Apr., 1900.do.....	9.08 a. m. 10.23 a. m.	49 12	co. S., Sh..... co. S.....
D. 5155	Bakun Pt., N. 70° E., 1.70 miles (5° 13' 40" N., 119° 57' 20" E.).do.....do.....	10.35 a. m. 11.00 a. m. 11.04 a. m.	12 12 12	co. S..... co. S..... co. S.....
.....	Tataan Pass, Simulac Id. (S. end Basun Channel).do.....do.....	2.00 p. m. 7.00 p. m.	mrgn. co. Rf..... co. S.....
.....	Simulac Id. (S. end Basun Channel).do.....	Feb. 20	8.30 a. m. 1.30 p. m.	mrgn. co. Rf..... mrgn. co. Rf.....
.....	Tataan Pass (anch.).....do.....do.....	7.30 p. m.	9
.....	Tataan Pass (Simulac Id., rf.)do.....	Feb. 21	8.30 a. m.	mrgn. co. Rf.....
D. 5156	Tinakta Id. (N.), S. 77° W., 3.40 miles (5° 12' 50" N., 119° 55' 55" E.).do.....do.....	8.35 a. m.	18	fne. S., Sh.....
D. 5157	Tinakta Id. (N.), S. 80° W., 3.30 miles (5° 12' 30" N., 119° 55' 50" E.).do.....do.....	8.43 a. m. 8.59 a. m.	18 18	fne. S., Sh..... fne. S.....
D. 5158	Tinakta Id. (N.), N. 89° W., 1.90 miles (5° 12' N., 119° 54' 30" E.).do.....do.....	9.04 a. m. 9.21 a. m.	18 12	fne. S..... crs. S., Sh.....
D. 5159	Tinakta Id. (N.), N. 82° W., 1.40 miles (5° 11' 50" N., 119° 54' E.).do.....do.....	9.28 a. m. 10.04 a. m.	12 10	crs. S., Sh..... co. S.....
.....	Simulac Id. (rf.).....do.....do.....	10.08 a. m.	10	co. S.....
.....	Tataan Pass (anch.).....do.....do.....	1.30 p. m. 7.30 p. m. 9	mrgn. co. Rf.....
D. 5160	Tinakta Id. (N.), S. 72° W., 2.75 miles (5° 12' 40" N., 119° 55' 10" E.).do.....	Feb. 22	8.26 a. m.	12	S.....
D. 5161	Tinakta Id. (E.), N. 12° W., 1.80 miles (5° 10' 15" N., 119° 53' E.).do.....do.....	8.29 a. m. 9.03 a. m.	12 16	S..... fne. S., blk. Sp.....
H. 4906	Tinagta Id. (S), N. 63° E., 4.10 miles (5° 09' 55" N., 119° 48' 55" E.).	C. S. 4514; Jan., 1906.do.....	9.07 a. m. 9.51 a. m.	16 55	fne. S..... S., brk. Sh.....
D. 5162	Tinagta Id. (S.), N. 71° W., 5.40 miles (5° 10' N., 119° 47' 30" E.).do.....do.....	10.10 a. m. 10.31 a. m.	230 230 crs. S., brk. Sh.....
.....	Bongao (anch.).....do.....do.....	7.30 p. m.	6
.....	Bongao (near anch.).....do.....do.....	7.30 p. m.
.....	Sanguisapo Id. (rf.).....do.....	Feb. 24	9.00 a. m.	sml. Clmps. Co., S.....
D. 5163	Observation Id., N. 79° W., 6.70 miles (4° 59' 10" N., 119° 51' E.).do.....do.....	9.36 a. m.	28	co. S.....
D. 5164	Observation Id., S. 82° W., 8 miles (5° 01' 40" N., 119° 52' 20" E.).do.....do.....	9.43 a. m. 10.16 a. m.	28 18	co. S..... gn. M.....
D. 5165	Observation Id., N. 70° W., 6.40 miles (4° 58' 20" N., 119° 50' 30" E.).do.....do.....	10.21 a. m. 1.19 p. m.	18 *9	gn. M..... Co.....
D. 5166	Observation Id., N. 20° W., 4.60 miles (4° 56' 10" N., 119° 46' E.).do.....do.....	2.54 p. m. 3.05 p. m.	97 97	co. S..... co. S.....
.....	Simonor Id., N. side (rf.).....do.....do.....	3.15 p. m.	solid Co.....
D. 5167	Observation Id., N. 11° W., 5.60 miles (4° 55' 10" N., 119° 45' 30" E.).do.....do.....	3.36 p. m. 3.53 p. m.	110 110	Co.*..... Co.*.....

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Alr.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
90	80		1.02489		Tnr.-Blish sdr. (e)					
90	80				12' Agz.; m. b.	botm...	20	N. 86° E	.4	
87	81		1.02457		Tnr.-Blish sdr. (e)					
86	81				12' Agz.; m. b. dip; e. l.	botm... surf...	15 30	S. 56° W	.5	
84	80		1.02437		Tnr.-Blish sdr. (e)					
85	80				12' Agz.; m. b.	botm...	14	N. 27° W	.4	
85	81		1.02437		Tnr.-Blish sdr. (e)					
88	81				12' Agz.; m. b.	botm...	15	S. 42° W	.3	
84	81		1.02437		Tnr.-Blish sdr. (e)					
84	81				int. 4 §.	8 fms	21	S. 58° W	.3	20 fms. dredge cable out.
					dyn.	5-30 ft.	3 00			Set over night.
					4 gill nets					
					dyn.	5-40 ft.	3 00			Channel between reefs.
					dyn.	5-40 ft.	3 00			
					dip; e. l.	surf.	30			
					dyn.	6-20 ft.	3 00			
79	79		1.02422		Tnr.-Blish sdr. (e)					
79	79				9' Jn. dr.	botm...	2	S. 28° E	.1	
79	79		1.02422		Tnr.-Blish sdr. (e)					
79	79				9' Jn. dr.	botm...	5	S. 29° W	.2	
80	79		1.02422		Tnr.-Blish sdr. (e)					
80	79				9' Jn. dr.	botm...	4	N. 80° W	.1	
83	80		1.02422		Tnr.-Blish sdr. (e)					
83	80				9' Jn. dr.	botm...	2	S. 14° E	.2	
					dyn.	6-20 ft.	3 00			
					dip. e. l.	surf.	1 00			
85	82				Tnr.-Blish sdr. (e)					
85	82				9' Jn. dr.	botm...	3	S. 67° W	.2	
90	82				Tnr.-Blish sdr. (e)					
90	82				9' Jn. dr.	botm...	1	S.	.1	Net fouled bottom.
94	82	63.5			Luc. sdr. (a)					
90	82	52.9	1.02447		Luc. sdr. (a)					
85	82				12' Agz.; m. b.	botm...	15	S. 9° E		
					dip; e. l.	surf.				
					4 gill nets					Final haul Feb. 24.
					dyn.	6-15 ft.	2 30			
91	77		1.02447		Tnr.-Blish sdr. (e)					
91	77				9' Jn. dr.	botm...	4	N. 63° W	.3	
89	80		1.02442		Tnr.-Blish sdr. (e)					
90	80				9' Jn. dr.	botm...	8	N. 30° E	.4	
84	80		1.02495		9' Jn. dr.	botm...	4	S.	.2	No sounding taken.
83	81	69.4	1.02644		Tnr.-Blish sdr. (e)					
83	81				12' Agz.; m. b.	botm...	2	S. 5° E	(?)	Distance recorded .7 mile; 1 bridle stop carried away.
					dyn.	6-15 ft.	2 00			
82	80		1.02406		Luc. sdr. (a)					
82	80				12' Agz.; m. b.	botm...	20	S. 14° W	1.4	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Sulu Archipelago, Tawi Tawi Group—Continued.</i>					
D. 5168	Observation Id. N. 17° W., 4.20 miles (4° 56' 30" N., 119° 45' 40" E.).	C. S. 4514; Jan., 1906.	1908. Feb. 25	7.09 a. m. 7.23 a. m.	<i>fms.</i> 80 80	co. S..... co. S.....
	<i>Sulu Archipelago, vicinity Sibutu Id.</i>					
.....	Sitanki (anch.).....	C. S. 4722; Apr., 1905.	Feb. 25	7.30 p. m.		
.....	Sitanki (near anch.).....	do.....	do.....	7.30 p. m.		
.....	Tumindao Reef S. end (rf.).....	do.....	Feb. 26	9.00 a. m. 1.30 p. m.		sctrd. Clmps. Co.. sctrd. Clmps. Co.. co. S.....
D. 5169	Sibutu Id. (S. E.), N. 38° E., 8 miles (4° 32' 15" N., 119° 22' 45" E.).	do.....	Feb. 27	8.36 a. m.	* 10	
.....	Sitanki wharf.....	do.....	do.....	10.00 a. m.		S., M., Co.....
D. 5170	Sibutu Id. (S. end), N. 38° E., 13.50 miles (4° 28' N., 119° 19' 30" E.).	do.....	do.....	11.06 a. m. 11.17 a. m.	128 128	crs. S..... crs. S.....
H. 4907	Sibutu Id. (S. end), N. 10° E., 13.50 miles (4° 26' N., 119° 25' 30" E.).	do.....	do.....	12.51 p. m.	850	gn. M.....
D. 5171	Omapui Id. (W.), S. 22° W., 12 miles (5° 05' N., 119° 28' E.).	do.....	Feb. 28	3.21 p. m. 3.47 p. m.	250 250	fne. co. S..... fne. co. S.....
	<i>Sandakan and vicinity.</i>					
.....	Sandakan (near anch.).....	B. A. 950.....	Feb. 29	8.15 p. m.	7	
.....	Sandakan (anch.).....	do.....	do.....	8.15 p. m.	7	
.....	Sandakan (beach above fish- ermen's village).	do.....	Mar. 1 Mar. 2	8.00 p. m. 2.00 p. m.	7	S., R.....
	<i>Vicinity of Jolo.</i>					
.....	Usada Id., S. end (rf.).....	C. S. 4722; Apr., 1905.	Mar. 5	9.00 a. m.		sctrd. Co.....
D. 5172	Jolo Lt., E., 24.75 miles (6° 03' 15" N., 120° 35' 30" E.).	do.....	do.....	10.06 a. m.	318	fne. S., Sh.....
H. 4908	Jolo Lt., N. 78° E., 7.50 miles (6° 02' 30" N., 120° 52' 20" E.).	C. S. 4542; Apr., 1903.	do.....	10.31 a. m. 2.27 p. m.	318 171	fne. S., Sh..... Sh., Co.....
D. 5173	Jolo Lt., N. 82° E., 6.75 miles (6° 02' 55" N., 120° 53' E.).	do.....	do.....	2.39 p. m. 2.57 p. m.	186 186	Sh., Co..... Sh., Co.....
D. 5174	Jolo Lt., E. 2.60 miles (6° 03' 45" N., 120° 57' E.).	do.....	do.....	3.46 p. m. 3.51 p. m.	20 20	crs. S..... crs. S.....
.....	Jolo (anch.).....	do.....	do.....	4.00 p. m.		sctrd. Co.....
.....	Jolo (rf. near anch., north).....	do.....	Mar. 6	9.00 a. m.		Co., S.....
.....	Jolo (beach, west of town).....	do.....	do.....	2.00 p. m.		S., Co., grassy.....
.....	Jolo (near anch.).....	do.....	do.....	4.00 p. m.	10	S.....
.....	Jolo (west of anch.).....	do.....	Mar. 7	9.00 a. m.		S., Co. (staghorn Mss.).

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trawl.		Drift.		Remarks.
Alr.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.								
79	79		1.02386		Luc. sdr. (e)		h. m.		mi.	
79.5	79				12' Agz.; m. b.	botm...	3	S.....	(?)	Net fouled bottom.
					dip; e. l.	surf....	1 00			
					4 gill nets					Set over night.
					dyn.....	9-15ft	3 00			
81	79		1.02509		9' Jn. dr.	botm...	5	S. 11° W.	.2	No sounding.
					dyn.....	12-15ft.	1 00			
0.5	78		1.02426		Luc. sdr. (e)					
81	78				12' Agz.; m. b.	botm...	2	S. 27° E.	(?)	Distance recorded, 0.5 mile; 1 bridle stop carried away.
82	79				Luc. sdr. (a)					
76	83	53.5	1.02373	1.02462	Luc. sdr. (a)					
76	83				12' Agz.; m. b.	botm...	20	S. 45° W.	(?)	Distance not obtainable on account of fog.
					2' o. p.	surf....	15			Towed from steam launch.
					dip; e. l.	surf....	1 30			
					dip; e. l.	surf....	1 30			
					130' seine.	12ft....	3 30			6 hauls.
					dyn.....	6-12ft.	2 00			
84	82		1.02447		Luc. sdr. (a)					Temperature at 277 fms. 53.3. Density at 277 fms. 1.02462.
85	82				12' Agz.; m. b.	botm...	20	N. 47° W.	1.0	Net slightly damaged.
96	84				Tnr.-Blish sdr (b).					
99	83		1.02518		Tnr.-Blish sdr (b).					
93	83				9' Jn. dr.	botm...	6	E.....	(?)	Distance recorded 0.7 mile.
100	82				Tnr.-Blish sdr (e).					
100	82				9' Jn. dr.	botm...	6	N. 58° E.	.4	Hauled and shifted about 7 p. m.; not found on following morning.
					4 gill nets					
					dyn.....	8ft....	3 00			
					130' seine.	4ft....	2 00			4 hauls; 1 at mouth of stream.
					4 traps.	botm...				Hauled following morning and at 1 p. m.
					dyn.....	4-10ft.	3 00			

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Sulu Sea, S. E. of Cagayanes Ids.</i>					
D. 5175	Manucan Id. (E.), N. 45° W., 23.25 miles (9° 21' N., 121° 37' 45" E.).	C. S. 4717; Feb., 1903.	1908. Mar. 8	7.22 p. m.	fms. *
	<i>Manila Bay.</i>					
.....	Manila Bay (Luneta beach).....	C. S. 4240; Feb., 1907.	Mar. 16	9.00 a. m.	S., M.
.....	Cavite (Sangley Pt. beach).....do.....	Mar. 23	9.00 a. m.	S.
	<i>Verde Id. Passage.</i>					
D. 5176	Escarceo Lt., S. 57° E., 7 miles (13° 35' 15" N., 120° 53' 20" E.).	C. S. 4240; Feb., 1907.	Mar. 24	7.01 p. m.	*260	*S.
D. 5177	Escarceo Lt., S. 53° E., 5.80 miles (13° 35' N., 120° 54' 36" E.).do.....do.....	7.33 p. m.	*260	*S.
	<i>Vicinity of Romblon.</i>					
D. 5178	Pt. Origon (N.), S. 5° E., 2.30 miles (12° 43' N., 122° 06' 15" E.).	C. S. 4714; June, 1906.	Mar. 25	8.35 a. m.	73	fne. S.
D. 5179	Romblon Lt., S. 56° E., 4.50 miles (12° 38' 15" N., 122° 12' 30" E.).do.....do.....	8.51 a. m. 10.41 a. m. 10.49 a. m.	78 37 37	fne. S. hrd. S. hrd. S.
.....	Romblon Harbor (rf. S. of Agbatan Pt.).....	C. S. 4442; Mar., 1907.do.....	2.00 p. m.	Mss. staghorn Co.
.....	Romblon (anch.).....do.....do.....	8.00 p. m.	20
.....	Romblon (beach at Binagon and Agpatan Pts.).....do.....	Mar. 26	9.00 a. m.	S., Co.
.....	Romblon (rf. E. of Sabang Pt.).....do.....do.....	9.00 a. m.	mrgn. Clmps. Co.
.....	Romblon (rf. E. side Rosas Pt.).....do.....do.....	1.00 p. m.	co. Clmps.
D. 5180	Romblon Lt., N. 6° 30' E., 7.10 miles (12° 28' 30" N., 122° 15' E.).	C. S. 4715; Apr., 1907.do.....	7.32 p. m.
	<i>Off eastern Panay.</i>					
D. 5181	Antonia Id. (S.), S. 63° W., 6.60 miles (11° 36' 40" N., 123° 26' 35" E.).	C. S. 4417; Feb., 1905.	Mar. 27	8.39 a. m.	26	M., fne. S.
D. 5182	Antonia Id. (S.), N., 43° W., 3.70 miles (11° 30' 40" N., 123° 23' 20" E.).do.....do.....	8.46 a. m. 9.43 a. m. 9.51 a. m.	26 24 24	M., fne. S. hrd. S., M. fne. S., M.
	<i>Between Panay and Negros.</i>					
D. 5183	Lusaran Lt., S. 29° E., 4 miles (10° 32' 48" N., 122° 26' E.).	C. S. 4718; Dec., 1906.	Mar. 30	10.27 a. m. 10.51 a. m.	96 96	sft. gn. M. sft. gn. M.
D. 5184	Lusaran Lt., N. 22° E., 11.25 miles (10° 18' 30" N., 122° 23' 30" E.).do.....do.....	1.09 p. m. 1.53 p. m.	565 565	gn. M. gn. M.
D. 5185	Lusaran Lt., N. 23° E., 25.50 miles (10° 05' 45" N., 122° 18' 30" E.).do.....do.....	4.39 p. m. 5.26 p. m.	638 638	gn. M. gn. M.
D. 5186	Lusaran Lt., N. 20° E., 37.80 miles (9° 53' 30" N., 122° 15' 30" E.).do.....do.....	8.01 p. m.
	<i>Tanon Strait, east coast of Negros.</i>					
D. 5187	Apo Id., S. 21° W., 12.50 miles (9° 16' 45" N., 123° 21' 15" E.).	C. S. 4718; Dec., 1906.	Mar. 31	1.06 p. m. 1.26 p. m.	225 225	sft. gn. M. sft. gn. M.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Tempera- tures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Sur- face.	Bot- tom.		Depth.	Dura- tion.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
82	82				int. 4.....	surf....	0 20	N. 7° E...	1.3	Chart indicates no bottom at 70 fms.
					130' seine.....	4ft....	1 30			4 hauls.
					130' seine.....	10ft....	2 30			5 hauls.
80	79				int. 4.....	surf....	0 21	S. 72° E...	1.0	
80	79				int. 4 §.....	25 fms..	0 20 1.5	E.....	0.9	40 fms. dredge cable out.
80	80		1.02515	1.02516	Luc. sdr. (a).....					Therm. failed to trip.
80	80				12' Agz.; 3 m.b.	botm...	0 20	N. 84° W..	2.0	
81	81	75.7			Luc. sdr. (a).....					
81	81				12' Agz.; 3 m.b.	botm...	0 15	N. 81° W..	1.3	
					dyn.....	8-15ft..	3 00			
					dip; e. l.....	surf....	2 00			
					150' seine.....					Several hauls.
					dyn.....	10-20 ft.	3 00			Few shots made.
					dyn.....	8-15 ft..	1 30			Interrupted by rain.
79	80		1.02530		int. 4.....	surf....	20	S. 5° E...	(?)	
80	80		1.02544		Tnr.-Blish sdr. (e).					
80	80				9' Jn. dr.....	botm...	4	S. 46° W..	.3	
81	80		1.02515		Tnr.-Blish sdr. (e).					
81	80				9' Jn. dr.....	botm...	8	S. 39° W..	.7	Veered from 43 to 55 fms.
83	81	63.4	1.02489	1.02551	Luc. sdr. (a).....					
84	81				12' Agz.; 3 m.b.	botm...	20	S. 78° W..	.7	Veered from 192 to 250 fms. during haul.
90	83	49.8	1.02489	1.02505	Luc. sdr. (a).....					
92	82				12' Agz.; 3 m.b.	botm...	20	S. 52° W..	2.0	
81	82	49.8	1.02481	1.02492	Luc. sdr. (b).....					
81	82				int. 4 §.....	550 fms.	20 48	S. 64° W..	2.5	1,000 fms. dredge cable out.
81	80		1.02530		int. 4.....	surf....	20	S. 4° W...	.8	
87	81	53.6	1.02475	1.02492	Luc. sdr. (a).....					
87	81				9' Jn. dr.....	botm...	11	S. 79° W..	.6	Lashing slipped; catch lost.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Tanon Strait, east coast of Negros—Continued.</i>					
	Port Bais (anch.)	C. S. 4718; Dec., 1906.	1908. Mar. 31	8.00 p. m.	fms. 9	
D. 5188	Pescador Id., N. 16° E., 14 miles (9° 44' N., 123° 14' 20" E.).do.....	Apr. 1	10.21 a. m. 10.44 a. m.	299 299	gn. M. gn. M.
D. 5189	Pescador Id., N. 72° E., 3.30 miles (9° 56' 30" N., 123° 15' E.).do.....do.....	1.08 p. m. 1.33 p. m.	300 300	gn. M. gn. M.
D. 5190	Pescador Id., S. 9° E., 10.70 miles (10° 08' 15" N., 123° 16' 45" E.).do.....do.....	4.16 p. m. 4.39 p. m.	295 295	gn. M. gn. M.
D. 5191	Guijulgán (beach). Refugio Id. (S.), S. 74° W., 5.50 miles (10° 29' 45" N., 123° 31' 15" E.).do.....	Apr. 2	8.00 a. m. 2.58 p. m. 3.26 p. m. 258 258	S., G., grassy. gn. M. gn. M.
	Balamban (anch.)do.....do.....	8.00 p. m.	12	S.
	<i>Off northern Cebu Id.</i>					
D. 5192	Jilantagan Id. (E.), N. 13° W., 3 miles (11° 09' 15" N., 123° 50' E.).	C. S. 4718; Dec., 1906.	Apr. 3	9.28 a. m.	32	gn. S.
D. 5193	Chocolate Id., N. 77° E., 8 miles (11° 16' 45" N., 123° 55' 45" E.).do.....do.....	9.40 a. m. 11.03 a. m.	32 71	gn. S. gn. M.
D. 5194	Chocolate Id., N. 66° W., 8 miles (11° 15' 30" N., 124° 11' E.).do.....do.....	11.12 a. m. 1.58 p. m. 2.15 p. m.	71 148 148	gn. M. gn. M. gn. M.
D. 5195	Capitancillo Id. Lt., N., 11.75 miles (10° 47' N., 124° 06' 30" E.).do.....do.....	7.03 p. m.		
D. 5196	Capitancillo Lt., N. 5° 30' W., 14.30 miles (10° 44' 30" N., 124° 07' 30" E.).do.....do.....	7.42 p. m.		
	Mactán Cove, S. E. shore (rf.).do.....	Apr. 6	10.00 a. m.		mrgn. Clmps. Co.
	Mactán Id. (shore, opposite Cebu).do.....	Apr. 7	8.00 a. m.		honey-combed Rf.
	<i>Vicinity western Bohol.</i>					
	Mantacao Id., S. side (rf.)	C. S. 4718; Dec., 1906.	Apr. 8	3.00 p. m.		mrgn. Mss. Co.
	Mantacao Id., S. side (beach).do.....do.....	3.00 p. m.		S.
	Mantacao Id. (anch.)do.....do.....	8.00 p. m.	10	
D. 5197	Baliscasag Id., S., 22 miles (9° 52' 30" N., 123° 40' 45" E.).do.....	Apr. 9	8.34 a. m. 8.55 a. m.	174 174	gn. M. gn. M.
D. 5198	Baliscasag Id., S. 6° E., 10.25 miles (9° 40' 50" N., 123° 39' 45" E.).do.....do.....	11.05 a. m. 11.25 a. m.	220 220	gn. M. gn. M.
	Tagbilaran Channel (beach on Bohol side near S. anch.)do.....do.....	3.00 p. m.		S., grassy.
D. 5199	Pamilacan Id. (E.), S. 61° W., 6.25 miles (9° 31' 50" N., 124° 40' E.).	C. S. 4719; Aug., 1904.do.....	7.36 p. m.		
D. 5200	Pamilacan Id. (E.), S. 66° W., 7.25 miles (9° 31' 50" N., 124° 02' 05" E.).do.....do.....	8.07 p. m.		
	<i>Sogod Bay, southern Leyte Id.</i>					
D. 5201	Limasaua Id. (E.), S. 1° E., 14.80 miles (10° 10' N., 125° 04' 15" E.).	C. S. 4719; Aug., 1904.	Apr. 10	8.24 a. m. 9.13 a. m.	554 554	gy. S., M. gy. S., M.
D. 5202	Limasaua Id. (E.), S. 2° E., 16.70 miles (10° 12' N., 125° 04' 10" E.).do.....do.....	10.31 a. m. 11.07 a. m.	502 502	gy. M. gy. M.
D. 5203	Limasaua Id. (S.), S. 38° W., 5.50 miles (9° 58' N., 125° 07' 40" E.).do.....do.....	2.21 p. m. 3.47 p. m.	775 775	gn. M. gn. M.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Sur- face.	Bot- tom.		Depth.	Dura- tion.	Direction.	Distance.	
° F.	° F.	° F.				h. m.		mi.		
					dip; e. l.	surf.	3 30			Brackish water.
82.5	81	62.6	1.02475	1.02502	Luc. sdr. (a) ..					
84.5	81				12' Agz.; 3 m.b.	botm.	20	N. 63° W.6	
85	82	62.8	1.02468	1.02495	Luc. sdr. (a) ..					
89	82				12' Agz.; 3 m.b.	botm.	20	N. 70° E.	1.0	
92.5	83	63	1.02468	1.02482	Luc. sdr. (a) ..					
90	83				int. 4 §	250 fms.	20	N. 43° W.8	400 fms. dredge cable out. 8 hauls.
					150' seine.	9ft.	3 00			
93	83	62.8	1.02497	1.02516	Luc. sdr. (a) ..					
91.5	83				12' Agz.; 3 m.b.	botm.	20	S. 88° W.9	
					dip; e. l.	surf.	1 30			
82	82		1.02518		Tnr.-Blisb sdr. (b) ..					
82	82				9' Jn. dr.	botm.	3	N. 45° W.2	
86	82		1.02503		Tnr.-Blisb sdr. (e) ..					
90	82				12' Agz.; 3 m.b.	botm.	20	N. 44° W.	1.3	
85	83	56.5	1.02447	1.02597	Luc. sdr. (a) ..					
84	83				12' Agz.; 3 m.b.	botm.	20	S. 25° W.8	
82.5	84		1.02514		int. 4.	surf.	20	S. 22° 30' E.	1.5	No sounding.
81.5	82		1.02518		int. 4.	surf.	20			Ship steered in circle.
					dyn.	10-20 ft.	2 00			High water.
					poison.		2 00			Tide pools.
					dyn.	10-30 ft.	2 30			
					130' seine.	5ft.	2 30			6 hauls.
					dip; e. l.	surf.	1 00			
89	81	54.3	1.02489	1.02513	Luc. sdr. (a) ..					
91	81				12' Agz.; 3 m.b.	botm.	20	N. 58° W.	1.0	
84	81	53.9	1.02434	1.02500	Luc. sdr. (a) ..					
					12' Agz.; 3 m.b.	botm.	20	S. 54° W.	1.1	
					130' seine.	6ft.	1 00			3 hauls.
83	79		1.02530		int. 4.	surf.	20	E.6	
82.5	79		1.02468		int. 4.	surf.	18			Ship steered in circle.
80	79	52.8	1.02440	1.02497	Luc. sdr. (a) ..					
85	80				12' Agz.; 3 m.b.	botm.	20	S. 24° W.	1.5	Veered 112 fms. cable during haul. Therm. failed to trip.
80	80	(?)	1.02440	1.02457	Luc. sdr. (a) ..					
79	80				12' Agz.; 3 m.b.	botm.	20	(?)	(?)	
82	80	52.9	1.02468	1.02606	Luc. sdr. (a) ..					
83	81				12' Agz.; 3 m.b.	botm.	31	N. 72° W.	2.7	Veered from 1,200 to 1,330 fms. dur- ing haul.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>Off east coast of Leyte Id.</i>						
D. 5204	Mariquitdaquit Id., N. 88° E., 3.50 miles (11° 04' 18" N., 125° 05' 30" E.).	C. S. 4719; Aug., 1904.	1908. Apr. 11	9.48 a. m.	<i>fms.</i> 15	gn. M.
.....	Tacloban (anch.)do.....	Apr. 12	8.00 p. m.	3	M., S.
.....	Tacloban (near anch.)do.....do.....	8.00 p. m.
D. 5205	Caguayan Pt., N. 2° E., 0.70 mile (11° 19' 30" N., 124° 58' 05" E.).do.....	Apr. 13	9.28 a. m.	8
.....	San Januico Strait, N. of Nababuy Id. (rf.). ^ado.....do.....	1.00 p. m.	staghorn Co., R.
<i>Off western Samar.</i>						
D. 5206	Badian Id. (N.), N. 27° E., 5.75 miles (11° 31' 40" N., 124° 42' 40" E.).	C. S. 4420; May, 1907.	Apr. 14	9.54 a. m.	32	gn. M.
D. 5207	Badian Id. (N.), S. 74° E., 4.70 miles (11° 38' 05" N., 124° 40' 45" E.).do.....do.....	10.02 a. m.	32	gn. M.
.....do.....do.....	11.22 a. m.	35	gn. M., S.
D. 5208	Taratara Id. (N.), S. 67° 30' E., 4.10 miles (11° 45' 53" N., 124° 42' 50" E.).	C. S. 4451; June, 1904.do.....	11.27 a. m.	35	gn. M., S.
.....do.....do.....	12.53 p. m.	26	sft. gn. M.
D. 5209	Taratara Id. (N.), S. 53° W., 1.80 miles (11° 45' 25" N., 124° 48' 05" E.).do.....do.....	12.59 p. m.	26	sft. gn. M.
.....do.....do.....	2.03 p. m.	20	gn. M.
.....	Catbalogan (Pamuntangan Rf.).do.....do.....	2.13 p. m.	20	gn. M.
.....	Catbalogan (near anch.)do.....do.....	2.13 p. m.	20	gn. M.
.....	Catbalogan (beach above Aguada Pt.).do.....	Apr. 15	4.00 p. m.	sft. Co., S.
.....	Catbalogan (Pamuntangan Rf.).do.....do.....	7.00 p. m.	5
.....	Catbalogan (Quinituay Rf.).do.....do.....	8.00 a. m.	S., M.
.....	Catbalogan (Lutao Rf. and Anas Pt.).do.....	Apr. 15	8.00 a. m.	S., M.
.....	Catbalogan (Quinituay Rf., beach).do.....do.....	8.00 a. m.	sft. Co., algæ
.....	Catbalogan (Quinituay Rf.).do.....do.....	8.00 a. m.	sft. Co., algæ
.....	Catbalogan (Quinituay Rf.).do.....do.....	1.30 p. m.	staghorn Clm ps., Co., R.
.....	Catbalogan (Lutao Rf. and Anas Pt.).do.....	Apr. 16	8.30 a. m.	staghorn Clm ps., Co., R.
.....	Catbalogan (Quinituay Rf., beach).do.....do.....	8.30 a. m.	S., Co.
.....	Catbalogan (Quinituay Rf.).do.....do.....	8.30 a. m.	S., Co.
.....	Catbalogan (Quinituay Rf.).do.....do.....	2.30 p. m.	staghorn Mss., Co., R.
D. 5210	Limbancuayan Id. (E.), N. 1° W., 3.60 miles (11° 49' 55" N., 124° 28' 05" E.).	C. S. 4420; May, 1907.	Apr. 17	10.17 a. m.	50	fine gy. S.
.....do.....do.....	10.30 a. m.	50	fine gy. S.
.....do.....do.....	10.30 a. m.	50	fine gy. S.
<i>East of Masbate Id.</i>						
D. 5211	Panalangan Pt., Talajit Id., N. 33° E., 5.25 miles (11° 51' 35" N., 124° 14' E.).	C. S. 4715; Apr., 1907.	Apr. 17	1.05 p. m.	155	gn. M., S.
.....do.....do.....	1.20 p. m.	155	gn. M., S.
.....	Cataingan Bay (upper rf., inside Dumurug Pt.).	C. S. 4455; Sept., 1904.do.....	4.00 p. m.	S., sctrd. Clmps. staghorn Co.
.....	Cataingan Bay, Dumurug Pt. (beach).do.....	Apr. 18	8.30 a. m.	S., sctrd. Clmps. staghorn Co.
.....	Cataingan Bay (upper rf., inside Dumurug Pt.).do.....do.....	8.30 a. m.	S., sctrd. Clmps. staghorn Co.
.....do.....	Apr. 19	3.00 p. m.	S., sctrd. Clmps. staghorn Co.
.....	Cataingan Bay (anch.)do.....do.....	8.00 p. m.	20
D. 5212	Panalangan Pt., S. 54° 30' E., 14.50 miles (12° 04' 15" N., 124° 04' 36" E.).	C. S. 4715; Apr., 1907.	Apr. 20	8.29 a. m.	108	gy. S., M.
.....do.....do.....	8.45 a. m.	108	gy. S., M.
D. 5213	Destacado Id. (S.), N. 87° E., 8.50 miles (12° 15' N., 123° 57' 30" E.).do.....do.....	10.38 a. m.	80	S., M., Sh.
.....	Masbate (rf. N. of town).do.....do.....	10.47 a. m.	80	S., M., Sh.
.....do.....do.....	3.00 p. m.	Co., R.

^a One boat made collections up the Silaga River for a few miles.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>			<i>mi.</i>	
84	82	1.02391	12' Agz.; 3 m. b.	botm...	21	N. 57° W..	1.0	Sounding with hand lead.
.....	dip; e. l.	surf.
.....	2 gill nets.
84	83	1.02448	12' Agz.; 3 m. b.	botm...	Hauled following morning.
.....	dyn.....	3-10 ft.	3 00	Fouled bottom; trawl lost; mud bag only recovered; sounding with hand lead. Brackish water.
83	83	1.02406	Tnr.-Blish sdr. (e).
83	83	12' Agz.; m. b.	botm...	20	N. 18° W..	.7
86	84	1.02395	Tnr.-Blish sdr. (e).
85	84	12' Agz.; m. b.	botm...	15	N. 16° E..	.5
84	84	1.02483	Tnr.-Blish sdr. (e).
84	84	12' Agz.; m. b.	botm...	20	N. 27° E..	.6
81	84	1.02493	Tnr.-Blish sdr. (e).
81	84	12' Agz.; m. b.	botm...	20	S. 28° E..	.6	Mud bag lost.
.....	K2.....	surf.	10	S. 28° E..	.3	Towed alongside.
.....	dyn.....	12-15 ft.	1 00
.....	2 gill nets.
.....	130' seine.....	6 ft.	3 00	Finally hauled on Apr. 17.
.....	dyn.....	12-15 ft.	3 00
.....	dyn.....	4-20 ft.	3 00
.....	dyn.....	8-30 ft.	3 00	Coral unthrifty.
.....	150' seine.....	6 ft.	2 hauls.
.....	dyn.....	4-30 ft.	2 30	2 boats used.
82	84	76.3	1.02406	1.02523	Luc. sdr. (a)
83	83	12' Agz.; m. b.	botm...	11	N. 1° W..	.2
83	83	K2.....	surf.	11	N. 1° W..	.2	Towed alongside.
83	84	56.6	1.02482	1.02509	Luc. sdr. (a)
84	84	int. 4 §.....	(?)	20	N. 31° W..	1.7	200 fms. dredge cable out.
84	84	K2.....	surf.	20	N. 31° W..	1.7	Towed alongside.
.....	dyn.....	6-10 ft.	1 30
.....	150' seine.....	2 30	5 hauls.
.....	dyn.....	6-10 ft.	3 00
.....	dyn.....	6-10 ft.	1 00
.....	dip; e. l.	surf.
82	80	59.9	1.02467	1.02476	Luc. sdr. (a)
83	80	12' Agz.; m. b.	botm...	20	N. 21° W..	.9	Veered 8 fms. during haul.
82	81	1.02489	Tnr.-Blish sdr. (e).
85	81	12' Agz.; m. b.	botm...	20	N. 22° W..	.8
.....	dyn.....	6-25 ft.	2 00

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>East of Masbate Id.—Cont'd.</i>					
	Masbate (near anch.)	C. S. 4715; Apr., 1907.	1908. Apr. 20	5.30 p. m.	<i>fms.</i>	
	Masbate (anch.)do.....do.....	5.30 p. m.		
D. 5214	Palanog Lt., Masbate, S. 17° W., 2.60 miles (12° 25' 18" N., 123° 37' 15" E.).do.....	Apr. 21	8.00 p. m. 8.59 a. m.	20 218	gn. M. gn. M.
D. 5215	Palanog Lt., S. 5° 30' E., 8.50 miles (12° 31' 30" N., 123° 35' 24" E.).do.....do.....	10.27 a. m. 11.32 a. m.	604 604	gn. M. gn. M.
	<i>Between Burias and Luzon.</i>					
	Port San Miguel (beach)	C. S. 4454; May, 1906.	Apr. 21	3.00 p. m.		S.
	Port San Miguel (rf. N. of Puro Id.).do.....do.....	3.00 p. m.		S., mrgn. Clmps. Co.
	Port San Miguel (anch.)do.....do.....	7.00 p. m.	19	
D. 5216	Anima Sola Id., N. 44° W., 29.50 miles (12° 52' N., 123° 23' 30" E.).	C. S. 4715; Apr., 1907.	Apr. 22	8.19 a. m. 8.36 a. m.	215 215	gn. M. gn. M.
D. 5217	Anima Sola Id., N. 42° W., 17.30 miles (13° 20' N., 123° 14' 15" E.).do.....do.....	10.31 a. m. 10.44 a. m.	105 105	crs. gy. S. crs. gy. S.
D. 5218	Anima Sola Id. (E.), N. 10° W., 2 miles (13° 11' 15" N., 123° 02' 45" E.).do.....do.....	12.58 p. m.	20	crs. S.
	Burias Id., Port Busin (pt. below fort rf.).	C. S. 4454; May, 1906.do.....	1.05 p. m.	20	crs. S.
	Burias Id., Port Busin (anch.)do.....do.....	3.00 p. m.		mrgn. co. Rf.
	Port Busin (pt. below fort, rf.).do.....	Apr. 23	8.00 p. m. 5.30 a. m.	12	mrgn. co. Rf.
	Port Busin (beach at fort pt.)do.....do.....	5.30 a. m.		S., R., Co.
	<i>Between Marinduque and Luzon.</i>					
D. 5219	Mompog Id. (NE.), N. 35° 30' W., 12.25 miles (13° 21' N., 122° 18' 45" E.).	C. S. 4715; Apr., 1907.	Apr. 23	1.57 p. m. 2.37 p. m.	530 530	gn. M. gn. M.
	Santa Cruz Harbor Marinduque (anch.)	C. S. 4453; July, 1908.do.....	8.00 p. m.	12	S.
	Santa Cruz Id. (SE.)do.....	Apr. 24	6.00 a. m.		mrgn. Co.
	Santa Cruz Id. (SE.)do.....do.....	8.30 a. m.		S., grassy
D. 5220	San Andreas Id. (W.), S. 57° W., 8.50 miles (13° 38' N., 121° 58' E.).	C. S. 4714; June, 1906.do.....	12.57 p. m.	50	sft. gn. M.
D. 5221	San Andreas Id. (W.), S. 27° E., 5.50 miles (13° 38' 15" N., 121° 48' 15" E.).do.....do.....	1.06 p. m. 3.05 p. m.	50 193	sft. gn. M. gn. M.
D. 5222	San Andreas Id. (W.), S. 57° E., 9.20 miles (13° 38' 30" N., 121° 42' 45" E.).do.....do.....	3.25 p. m.	193	gn. M.
D. 5223	Malabrigo Lt., W., 9.80 miles (13° 36' N., 121° 25' 30" E.).do.....do.....	4.33 p. m. 4.49 p. m.	195 195	gn. M. gn. M.
D. 5224	Malabrigo Lt., N. 79° W., 6.25 miles (13° 34' 50" N., 121° 21' 45" E.).do.....do.....	7.47 p. m.		
	<i>China Sea, south of Corregidor.</i>					
D. 5225	Corregidor Lt., N. 10° E., 9.50 miles (14° 13' 24" N., 120° 32' 36" E.).	C. S. 4240; Feb., 1907.	May 4	7.06 p. m.		
D. 5226	Corregidor Lt., N. 10° E., 10.70 miles (14° 12' 15" N., 120° 32' 24" E.).do.....do.....	7.45 p. m.		

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
					2 gill nets					Hauled following morning. Lost.
					2 wire traps	botm.				
					dip; e. l.	surface	1-30			
81	82	51.4	1.02475	1.02485	Luc. sdr. (a)					
81	81				12' Agz.; m. b.	botm.	20	N. 36° E.	1.0	
82	81	50.5	1.02440	1.02441	Luc. sdr. (a)					
82	82				12' Agz.; m. b.	botm.	20	S. 77° E.	1.2	
					150' seine	15 ft	2 30			5 hauls.
					dyn.	6-30 ft.	2 30			
					dip; e. l.	surface	3 00			
80	80	51.9	1.02481	1.02465	Luc. sdr. (a)					
80	80				12' Agz.; m. b.	botm.	20	N. 42° W.	1.5	
83	82	63.1	1.02489	1.02496	Luc. sdr. (a)					
85	81				12' Agz.; m. b.	botm.	20	N. 45° W.	1.2	
86	82		1.02538		Tnr.-Blish sdr.					
					(e)					
86	82				9' Jn. dr.	botm.	5	N. 16° W.	.2	
					dyn.	10-30 ft.	2 00			
					dip; e. l.	surface	2 00			
					dyn.	10-30 ft.	1 30			
					150' seine	6 ft	1 30			3 hauls.
84	86	50.8	1.02468	1.02467	Luc. sdr. (a)					
86	87				12' Agz.; m. b.	botm.	20	N. 27° E.	1.5	
					dip; e. l.	surface	2 00			
					dyn.	6-15 ft.	1 00			
					150' seine	4 ft				5 hauls; beach in- side reef.
87	85		1.02493		Tnr.-Blish sdr.					
					(e)					
87	85				12' Agz.; m. b.	botm.	14	N. 54° W.	.7	
85	84	52.4	1.02503	1.02467	Luc. sdr. (a)					
85	84				12' Agz.; m. b.	botm.	20	N. 21° W.	1.0	
85	85	52.8	1.02470	1.02447	Luc. sdr. (a)					
86	85				12' Agz.; m. b.	botm.	20	N. 20° W.	1.7	
83	84				int. 4	surface	20	S. 69° W.	1.8	
83	84				int. 4	surface	10	N. 80° W.	.4	
85	84		1.02448		int. 4 §	40 fms.	20 (?)	S.9	Record incom- plete.
85	83		1.02514		int. 4	surface	20	S.8	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>East of Mindoro.</i>					
D. 5227	Pt. Orion, S. 44° E., 18.30 miles (12° 53' 45" N., 121° 52' 30" E.).	C. S. 4714; June, 1906.	1908. May 5	1.04 p. m. 1.30 p. m.	fms. 322 322	gn. M.
	<i>South of Romblon.</i>					
D. 5228	Romblon Lt., N. 3° E., 6.25 miles (12° 29' 30" N., 122° 15' 45" E.).	C. S. 4715; Apr., 1907.	May 5	7.02 p. m. 7.02 p. m.	
	<i>Between Cebu and Leyte.</i>					
D. 5229	Talong Id. (E.), S. 17° W., 5.75 miles (10° 48' 45" N., 124° 21' 15" E.).	C. S. 4719; Aug., 1904.	May 7	9.34 a. m. 9.55 a. m.	*290 *290
	<i>Between Bohol and Leyte.</i>					
D. 5230	Limasaua Id. (S.), S. 68° E., 22.50 miles (10° 01' 50" N., 124° 42' 30" E.).	C. S. 4719; Aug., 1904.	May 7	7.03 p. m. 7.13 p. m. 7.13 p. m.	118 118 118	gy. S.
D. 5231	Limasaua Id. (S.), S. 68° E., 21.70 miles (10° 01' 15" N., 124° 43' 15" E.).do.....do.....	7.48 p. m.	
D. 5232	Limasaua Id. (S.), S. 69° E., 20.60 miles (10° 00' 45" N., 124° 44' 06" E.).do.....do.....	8.25 p. m.	
D. 5233	Limasaua Id. (S.), S. 70° E., 19.50 miles (10° 00' 22" N., 124° 45' 06" E.).do.....do.....	9.00 p. m.	
D. 5234	Limasaua Id. (S.), S. 70° 30' E., 18.50 miles (10° N., 124° 46' 06" E.).do.....do.....	9.42 p. m.	
	<i>Pacific Ocean, east coast Mindanao.</i>					
.....	Surigao (beach near Bilan Bilan).	C. S. 4644; July, 1905.	May 8	8.30 a. m.		M., S., Co., grassy.
.....	Surigao (rf. above Bilan Bilan).do.....do.....	1.30 p. m.		R., co. Clmps.
D. 5235	Nagubat Id. (S.), S. 58° W., 7 miles (9° 43' N., 125° 48' 15" E.).	C. S. 4719; Aug., 1904.	May 9	9.24 a. m. 9.30 a. m.	44 44	sft. M. sft. M.
.....	Generale Id. (S. W. shore, beach).do.....do.....	3.00 p. m.		S., Co., grassy.
.....	Generale Id. (rf.).do.....do.....	3.00 p. m.		mrgn. Co.
.....	Generale Id. (Capunuyyugan Pt., rf.).do.....	May 10	8.30 a. m.		mrgn. Co.
.....	Generale Id. (rf.).do.....do.....	3.00 p. m.		
D. 5236	Magabao Id. (S.), N. 85° W., 9.10 miles (8° 50' 45" N., 126° 26' 52" E.).do.....	May 11	10.27 a. m. 11.02 a. m.	494 494	fne. gy. S. fne. gy. S.
.....	Liang Bay (rf. S. of town).do.....do.....	4.00 p. m.		co. Mss., algæ
.....	Liang Bay (anch.).do.....do.....	8.00 p. m.	15 (?)
D. 5237	Sanco Pt. Id. (N.), N. 69° W., 5.75 miles (8° 09' 06" N., 126° 31' 45" E.).	C. S. 4724; Oct., 1909.	May 12	10.11 a. m. 10.42 a. m.	249 249	gn. M.
D. 5238	Pt. Lambajon, S. 65° W., 4.30 miles (7° 34' 45" N., 126° 38' 15" E.).do.....do.....	3.00 p. m. 3.28 p. m.	380 380	gn. M. gn. M.
.....	Baganga Bay (rf. inside Pt. Lacud).do.....	May 13	8.30 a. m.		mrgn. Co.
.....	Baganga Bay (S. W. shore, beach).do.....do.....	8.30 a. m.		S.
.....	Baganga Bay (W. shore, beach).do.....do.....	1.00 p. m.		S., G.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				h. m.		mi.		
86	86		1.02498		Luc. sdr. (a)					
85	87				Int. 4 §	290 fms.	20 18	S. 30° E	0.6	400 fms. dredge cable out.
84	85		1.02519		Int. 4	surface.	20	S. 30° E	.6	
84	85				K2, K5 †	surface.	20	S. 30° E	.6	
86	85		1.02525		Tnr.-Blsh sdr. (c)					
86	85				Int. 4; K2, K5 §		20 11	S. 17° W	.5	225 fms. dredge cable out.
84	84	57.6	1.02477	1.02496	Luc. sdr. (a)					
84	84				Int. 4	surface.	20	S. 63° E	.6	
84	84				K2, K5 †	surface.	20	S. 63° E	.6	
85	84		1.02531		Int. 4; K2, K5 §	80 fms.	20 7	S. 63° E	.4	125 fms. dredge cable out.
83.5	84		1.02531		Int. 4	surface.	20	S. 63° E	.6	
83	84		1.02514		Int. 4; K2, K5 §	100 fms.	20 9	S. 63° E	.8	150 fms. dredge cable out.
83	84		1.02531		Int. 4; K2, K5 §	15 fms.	20 2	S. 63° E	.4	25 fms. dredge cable out.
					150' seine	6-30 ft.	3 00			5 hauls.
					dyn.	6-15 ft.	4 00			
84	86		1.02475		Tnr.-Blsh sdr. (c)					
84	86				12' Agz.; 3 m. b.	botm.	20	S. 56° E	.6	1 bridle stop carried away.
					150' seine	6-8 ft.	2 00			5 hauls.
					dyn.	12-20 ft.	2 00			
					dyn.	12-20 ft.	3 00			
					dyn.	4-15 ft.	2 00			
87	85	41.2	1.02453	1.02522	Luc. sdr. (a)					
86	86				12' Agz.; 3 m. b.	botm.	20	S. 4° E	2.5	Bridle stops carried away; net capsized; catch saved.
					dyn.	12 ft.	30			Seining party failed to find suitable beach.
					dip; e. l.	surface.	30			
85	85	46.4	1.02477	1.02482	Luc. sdr. (a)					
85	85				12' Agz.; 3 m. b.	botm.	17	S. 3° E	2.1	Veered at intervals from 450 to 550 fms.
91	86	43.0	1.02453	1.02459	Luc. sdr. (a)					
85	86				12' Agz.; 3 m. b.	botm.	20	S. 15° W	2.5	
					dyn.	4-20 ft.	2 00			Roily, brackish water.
					130' seine	10-20 ft.				7 hauls.
					250' seine	30 ft.				3 hauls. River explored.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Pujada Bay and vicinity.</i>					
D. 5239	Univan Id. (N.), N. 78° E., 2.25 miles (6° 49' 08" N., 126° 15' 12" E.).	C. S. 4646; Jan., 1905.	1908. May 14	12.44 p. m. 1.02 p. m.	<i>fms.</i> 171 171	sft. gy. M. sft. gy. M.
D. 5240	Univan Id. (N.), E., 2.40 miles (6° 49' 36" N., 126° 15' E.).do.....do.....	1.33 p. m. 1.49 p. m.	145 145	sft. gy. M. sft. gy. M.
D. 5241	Univan Id. (N.), S. 68° E., 3 miles (6° 50' 45" N., 126° 14' 38" E.).do.....do.....	2.24 p. m. 3.05 p. m.	215 215	sft. gy. M. sft. gy. M.
D. 5242	Univan Id. (N.), S. 56° E., 4 miles (6° 51' 53" N., 126° 14' 10" E.).do.....do.....	3.46 p. m. 4.03 p. m.	191 191	sft. gy. M. sft. gy. M.
.....	Pujada Bay (rf. S. of Taitadaga Pt.).do.....	May 15	9.00 a. m.	S., co. Clmps.
.....	Pujada Bay (beach both sides Mati).do.....do.....	9.00 a. m.	Co., R., S.
D. 5243	Univan Id. (N.), S. 66° E., 3.10 miles (6° 50' 55" N., 126° 14' 35" E.).do.....do.....	12.54 p. m. 1.12 p. m.	218 218	gy. M. gy. M.
D. 5244	Univan Id. (N.), S. 52° 30' E., 4 miles (6° 52' 05" N., 126° 14' 15" E.).do.....do.....	1.48 p. m.	171	gy. M.
D. 5245	Univan Id. (N.), S. 41° E., 4 miles (6° 52' 36" N., 126° 14' 52" E.).do.....do.....	2.05 p. m. 2.47 p. m. 3.02 p. m.	171 135 135	gy. M. gy. M. gy. M.
	<i>Pacific Ocean, east of Mindanao.</i>					
D. 5246	Luban Id. (N.), S. 58° W., 4.6 miles (6° 29' 15" N., 126° 18' 45" E.).	C. S. 4724; Oct., 1909.	May 15	7.10 p. m.
	<i>Gulf of Davao.</i>					
.....	Beach east of Davao town...	C. S. 4724; Oct., 1909.	May 16	9.00 a. m.	M., S.
D. 5247	Dumalag Id. (S.), S. 78° W., 3.8 miles (7° 02' N., 125° 38' 45" E.).do.....	May 18	8.47 a. m.	135	M.
D. 5248	Lanang Pt., S. 33° W., 0.40 mile (7° 07' 25" N., 125° 40' 24" E.).	C. S. 4648; Sept., 1907.do.....	9.08 a. m. 10.30 a. m. 10.38 a. m.	135 18 18	M. Co. Co.
D. 5249	Lanang Pt., N. 1 mile (7° 06' 06" N., 125° 40' 08" E.).do.....do.....	10.57 a. m. 11.02 a. m.	23 23	Co., S. Co., S.
D. 5250	Linao Pt., N. 22° E., 1.1 miles (7° 05' 07" N., 125° 39' 45" E.).do.....do.....	11.20 a. m.	23	Co., S.
D. 5251	Linao Pt., N. 32° E., 1.1 miles (7° 05' 12" N., 125° 39' 35" E.).do.....do.....	11.24 a. m. 1.07 p. m.	23 20	Co., S. Co.
D. 5252	Linao Pt., N. 22° E., 1.5 miles (7° 04' 48" N., 125° 39' 38" E.).do.....do.....	1.10 p. m. 1.22 p. m.	20 28	Co. Co.
D. 5253	Linao Pt., N. 22° E., 1.5 miles (7° 04' 48" N., 125° 39' 38" E.).do.....do.....	1.25 p. m. 1.34 p. m.	28 28	Co. Co.
D. 5254	Linao Pt., N. 44° E., 0.7 mile (7° 05' 42" N., 125° 39' 42" E.).do.....do.....	1.47 p. m. 2.22 p. m.	28 21	Co. S., Co.
D. 5255	Dumalag Id. (S.), S. 65° W., 4.5 miles (7° 03' N., 125° 39' E.).do.....do.....	2.26 p. m. 6.03 p. m. 6.13 p. m.	21 100 100	S., Co. sft. M. sft. M.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—(Continued.)

Tempera- tures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Sur- face.	Bot- tom.		Depth.	Dura- tion.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
84	86		1.02417		Tnr.-Blis sdr. (e).					
84	86				12' Agz.; 3 m. b.	botm...	7	N. 13° W.	0.5	Bridle and trip- ping stops car- ried away; net torn; frame twisted; 1 mud bag lost.
84	86		1.02448		Tnr.-Blis sdr. (e).					
84	86				int. 4 §	115 fms.	20 7	N. 16° W.	1.1	175 fms. dredge ca- ble out.
85	85		1.02453		Tnr.-Blis sdr. (e).					
84	85				9' alb. Blk.; m. b.	botm...	20	N. 15° W.	1.1	Veered from 506 to 540 fms.
84	85	64.1	1.02457	1.02489	Luc. sdr. (a).					
83.5	85				9' alb. Blk.; m. b.	botm...	20	N. 13° W.	1.0	
					dyn.	6-20 ft.	2 30			
					150' seine.	10 ft.	2 00			5 hauls.
84	84	63.6	1.02453	1.02468	Luc. sdr. (a).					
85	85				12' Agz.; m. b.	botm...	20	N. 15° W.	1.1	
84	85		1.02497		Tnr.-Blis sdr. (e).					
84	85				12' Agz.; m. b.	botm...	20	N. 46° E.	.7	
84	84		1.02468		Tnr.-Blis sdr. (e).					
84	84				12' Agz.; m. b.	botm...	20	N. 2° W.	.8	Net damaged.
83	82		1.02477		int. 4 §	100 fms.	20 8	S. 6° E.	1.8	150 fms. dredge ca- ble out.
					150' seine.	6 ft.	2 00			3 hauls.
80	83		1.02417		Tnr.-Blis sdr. (e).					
81	83				12' Agz.; m. b.	botm...	20	N. 76° W.	.5	
84.5	83		1.02453		Tnr.-Blis sdr. (e).					
84.5	83				6' Jn. dr.	botm...	4	(?)	(?)	Veered from 27 to 30 fms.
85	84		1.02453		Tnr.-Blis sdr. (e).					
85	84				6' Jn. dr.	botm...	7	(?)	(?)	Veered from 30 to 36 fms.
84	84		1.02457		Tnr.-Blis sdr. (e).					
84	84				6' Jn. dr.	botm...	3	(?)	(?)	
86	83		1.02433		Tnr.-Blis sdr. (e).					
86	83				6' Jn. dr.	botm...	5	(?)	(?)	
85	83		1.02417		Tnr.-Blis sdr. (e).					
85	83				6' Jn. dr.	botm...	4	S. 29° E.	.2	
83	84		1.02433		Tnr.-Blis sdr. (e).					
83	84				6' Jn. dr.	botm...	11	N. 11° E.	1.0	
83	83		1.02417		Tnr.-Blis sdr. (e).					
83	83				6' Jn. dr.	botm...	5	N.	.3	
83	84		1.02227		Tnr.-Blis sdr. (e).					
83	84				12' Agz.; m. b.	botm...	20	(?)	(?)	Made after dark.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date	Time of day.	Depth.	Character of bottom.
	<i>Southern Mindanao, eastern Iilana Bay.</i>			1908.	<i>fms.</i>	
.....	Cotabato (beach outside Panalisan Pt.). ^a	C. S. 4723; Oct., 1905.	May 20	2.30 p. m.	S., M.....
.....	Cotabato (near anch. outside Panalisan Pt.).do.....do.....	7.00 p. m.	30	S.....
.....	Malabang (beach below river). ^bdo.....	May 21	8.30 a. m.	S.....
.....	Malabang (river).....do.....do.....	3.00 p. m.	S.....
.....	Malabang (anch.).....do.....do.....	8.00 p. m.	13
D. 5256	Utara Pt., Bongo Id., N. 76° W., 2.80 miles (7° 21' 45" N., 124° 07' 15" E.)	C. S. 4619; Apr., 1907.	May 22	9.39 a. m.	158	M.....
.....do.....do.....	9.54 a. m.	158	M.....
D. 5257	Utara Pt., Bongo Id., N. 88° W., 7.70 miles (7° 22' 12" N., 124° 12' 15" E.).do.....do.....	10.07 a. m.	28	M.....
.....	Polloc (Marigabato Pt., rf.)do.....do.....	1.00 p. m.	S., setrd. Co.....
.....	Parang (Lalayanga Pt., rf.)do.....	May 23	8.30 a. m.	setrd. Co., co. R.....
.....	Parang (beach in front of village).	C. S. 4723; Oct., 1905.do.....	8.30 a. m.	S.....
	<i>Vicinity of Zamboanga.</i>					
.....	Zamboanga (W. end Little Sta. Cruz Id., rf.)	C. S. 4723; Oct., 1905.	May 26	10.10 a. m.	sft. Co., co. heads.
.....	Zamboanga (Little Sta. Cruz Id., rf.)do.....	May 28	7.00 a. m.	sft. Co., mrgn. Rfs..
	<i>Iloilo.</i>					
.....	E. of mouth of Iloilo River (beach).do.....	June 2	9.00 a. m.	S.....
	<i>Off southern Panay.</i>					
D. 5258	Juraojuro Id. (S.), S. 75° W., 16.25 miles (10° 27' 45" N., 122° 12' 30" E.).	C. S. 4717; Feb., 1903.	June 2	7.08 p. m.
	<i>Off northwestern Panay.</i>					
D. 5259	Caluya Id. (S.), S. 73° W., 12 miles (11° 57' 30" N., 121° 42' 15" E.).	C. S. 4714; June, 1906.	June 3	10.06 a. m. 10.31 a. m.	312 312	gy. M., Glob..... gy. M., Glob.....
	<i>Off southeastern Mindoro.</i>					
D. 5260	Balanja Pt., N. 28° W., 7.20 miles (12° 25' 35" N., 121° 31' 35" E.).	C. S. 4311; July, 1904.	June 3	3.14 p. m. 3.32 p. m.	234 234	gn. M., S..... gn. M., S.....
.....	Mansalay (anch.).....do.....do.....	9.00 p. m.	9
.....	Balanja Pt. (rf.).....do.....	June 4	7.30 a. m.	mrgn. Co.....
.....	Mansalay Bay (W. shore, beach).do.....do.....	7.30 a. m.	S., Co.....
.....	Mansalay Bay (NE. shore, rf.)do.....do.....	1.00 p. m.	setrd. Co.....
H. 4912	Balanja Pt., N. 73° W., 3.70 miles (12° 30' 55" N., 121° 31' 50" E.).do.....do.....	5.34 p. m.	56	bl. M., S.....
D. 5261	Balanja Pt., N. 80° W., 6 miles (12° 30' 55" N., 121° 34' 24" E.).do.....do.....	6.00 p. m. 6.11 p. m.	145 145	S., M..... S., M.....
	<i>Off eastern Mindoro.</i>					
D. 5262	Pt. Orilon, N. 83° E., 28.50 miles (12° 37' 30" N., 121° 37' 30" E.).	C. S. 4714; June, 1906.	June 4	7.39 p. m. 7.45 p. m.

^a On May 20 collecting party went up Mindanao River to Cotabato; visited market.

^b May 22 to 24 shore party made collections at Lake Lanao; visited market at Vicar.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.								
					130' seine.....	6 ft	h. m. 3 00			7 hauls.
					2 gill nets.....					Set over night.
					150' seine.....	20 ft	3 00			5 hauls.
					130' seine.....	5 ft				3 hauls.
					dip; e. l.....	surf.	1 30			
83	86		1.02262		Tnr.-Blish sdr. (e).					
83	86				12' Agz.; m. b.	botm...	20	N. 49° E..	0.6	
83	86		1.02277		Tnr.-Blish sdr. (e).					
83	86				12' Agz.; m. b.	botm...	20	S. 66° E...	.6	
					dyn.....	4-25ft.	3 00			
					dyn.....	6-25 ft.	3 00			
					150' seine.....	20 ft	3 00			8 hauls.
					dyn.....	12 ft	1 30			
					dyn.....	5-30 ft..	4 00			
					150' seine.....	5 ft	2 00			5 hauls.
84	84		1.02587		int. 5.....	surface.	20	S. 67° 30' W.	.3	
84.5	85	49.3	1.02489	1.02484	Luc. sdr. (a)..					
84	85				12' Agz.; m. b.	botm...	20	N. 6° W...	1.0	
85	85	51.4	1.02484	1.02484	Luc. sdr. (a)..					
85	83				12' Agz.; m. b.	botm...	20	N. 14° W..	2.2	
					dip; e. l.....	surf.	1 00			
					dyn.....	8-15ft.	4 00			
					150' seine.....	5-10ft.	3 00			5 hauls; many medu- sae.
					dyn.....	5-15 ft..	2 00			
			1.02463		Tnr.-Blish sdr. (e).					
85	84		1.02448		Tnr.-Blish sdr. (e).					
85	83				12' Agz.; m. b.	botm...	20	N. 29° E..	.4	
85	83		1.02448		int. 5.....	surface.	20	N.....	.5	
85	83				K2, K5†.....	surface.	15	N.....	.4	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Off eastern Mindoro—Cont'd.</i>					
D. 5263	Pt. Origan, N. 85° E., 28.3 miles (12° 38' 30" N., 121° 37' 30" E.).	C. S. 4714; June, 1906.	1908. June 4	8.17 p. m.	<i>fms.</i>	
	Naujan River (anch.) ^a	do	June 5	8.00 p. m.	17	S.
	<i>Verde Id. Passage and Batangas Bay.^b</i>					
D. 5264	Malabrigo Lt., N. 86° 30' E., 7.30 miles (13° 35' 30" N., 121° 08' E.).	C. S. 4240; Feb., 1907.	June 6	8.19 a. m.	181	S., P.
				8.38 a. m.	181	S., P.
D. 5265	Matocot Pt., Luzon, S. 17° E., 3.30 miles (13° 41' 15" N., 120° 00' 50" E.).	do	do	10.49 a. m.	135	S., M.
D. 5266	Matocot Pt., S. 22° E., 7 miles (13° 44' 36" N., 120° 59' 15" E.).	C. S., 4240; Feb., 1907.	June 8	11.09 a. m. 9.08 a. m.	135 100	S., M. M.
D. 5267	Matocot Pt., S., 39° E., 5.50 miles (13° 42' 20" N., 120° 58' 25" E.).	do	do	9.18 a. m. 10.08 a. m.	100 170	M. P., S., Sh.
D. 5268	Matocot Pt., S., 50° E., 5.80 miles (13° 42' N., 120° 57' 15" E.).	do	do	10.25 a. m. 10.59 a. m.	170 170	P., S., Sh. S., P.
D. 5269	Matocot Pt., S., 54° E., 3 miles (13° 39' 50" N., 120° 59' 30" E.).	do	do	11.14 a. m. 1.08 p. m.	170 220	S., P. fne. S., P.
D. 5270	Escarceo Lt., S. 9° E., 4.25 miles (13° 35' 45" N., 120° 58' 30" E.)	do	do	1.34 p. m. 3.07 p. m.	220 235	fne. S., P. gy. S., blk. Sp.
				3.27 p. m.	235	
	Port Galera (anch.)	do	do	8.30 p. m.	13	
	Port Galera (Paniquian Id., beach).	do	June 9	8.30 a. m.		S., Co.
	Port Galera (Medio Id., rf.)	do	do	8.30 a. m.		mrng. Clmps. Co.
	<i>Manila Bay.</i>					
	Cavite (anch.)	C. S. 4240; Feb., 1907.	June 9	8.00 p. m.	4	
	Bacoor (beach)	do	June 15	10.00 a. m.		S.
	<i>China Sea. vicinity southern Luzon.</i>					
	Jamelo Cove (rf.)	C. S. 4240; Feb., 1907.	July 13	8.00 a. m.		Co. unthrifty and sparse.
	Jamelo Cove (beach)	do	do	8.00 a. m.		S.
	Jamelo Cove (E. side), (rf.)	do	do	2.00 p. m.		Co. unthrifty and sparse.
	Jamelo Cove (beach)	do	do	2.00 p. m.		S.
D. 5271	Corregidor Lt., N. 17° E., 20.70 miles (14° 03' N., 120° 27' 45" E.).	do	July 14	8.08 a. m.	56	S.
				8.30 a. m.	56	S.
D. 5272	Corregidor Lt., N. 26° E., 25.50 miles (14° N., 120° 22' 30" E.).	do	do	9.32 a. m.	118	M., Sh., co. S.
				10.05 a. m.	118	M., Sh., co. S.
D. 5273	Corregidor Lt., N. 27° E., 27.25 miles (13° 58' 45" N., 120° 21' 35" E.).	do	do	10.34 a. m.	114	M., Sh., co. S.
				10.47 a. m.	114	M., Sh., co. S.
	Tilig Bay (beach inside vil- lage).	do	do	2.30 p. m.		S., M.

^a On June 5 a shore party went about 4 miles up the Naujan River in boats.

^b On June 7 a collecting trip was made up the Batangas River for about 2 miles; several hauls with a 15-foot seine.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—(Continued.)

Tempera- tures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Sur- face.	Bot- tom.		Depth.	Dura- tion.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
4	83	int. 5; K2, K5 §.	65 fms..	20 5	N.....	0.5	
.....	dip.; e. l.....	surface.	1-30	
84	84	1.02453	Tnr.-Blish sdr. (e).	
84	84	12' Agz.; m. b..	botm...	4	S. 37° E...	.5	Cable parted while heaving in; trawl lost with 20 fms. cable.
87	85	1.02489	Tnr.-Blish sdr. (e).	
89	85	12' Agz.; m. b.	botm...	20	N. 46° W...	1.0	
83	84	1.02448	Tnr.-Blish sdr. (e).	
84	85	12' Agz.; m. b.	botm...	20	N. 86° W...	1.1	
85	85	1.02448	Tnr.-Blish sdr. (e).	
85	85	12' Agz.; m. b.	botm...	20	S. 65° W...	1.3	
83	85	1.02433	Tnr.-Blish sdr. (e).	
85	85	12' Agz.; m. b.	botm...	20	N. 3° W...	1.0	
84	85	1.02417	1.02509	Tnr.-Blish sdr. (e).	Therm. failed to register.
85	85	12' Agz.; m. b.	botm...	20	N. 18° E...	1.5	
85	84	1.02448	Tnr.-Blish sdr. (e).	Water bottle failed to work.
80.5	83	int. 5; K2, K5 §.	140 fms.	20 8	N. 1° W...	1.1	200 fms. dredge cable out.
.....	dip; e. l.....	surface.	45	
.....	150' seine.....	7 ft	2 00	5 hauls.
.....	dyn.....	8-20ft..	4 00	
.....	dip; e. l.....	surface.	1 00	
.....	45' seine.....	4 ft	8 hauls.
.....	dyn.....	8-15 ft..	3 00	4 shots.
.....	150' seine.....	10 ft	3 00	7 hauls.
.....	dyn.....	8-15ft..	3 00	
.....	150' seine.....	6 ft	4 00	3 hauls.
83	85	1.02552	Tnr.-Blish sdr. (e).	First attempt at sounding re- sulted in loss of all apparatus used.
83	85	12' Agz.....	botm...	20	S.....	.7	
83	84	57.4	1.02453	Tnr.-Blish sdr. (e).	
83	84	12' Agz.....	botm...	26	S. 37° E...	.3	
83	84	Tnr.-Blish sdr. (e).	
83	84	12' Agz.....	botm...	30	N. 8° E....	1.7	
.....	130' seine.....	8 ft	2 30	4 hauls.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>China Sea, vicinity southern Luzon—Continued.</i>						
	Tilig Bay (rf. outside village).	C. S. 4240; Feb., 1907.	1908. July 14	3.00 p. m.	<i>fms.</i>	mrgn. rf.
	Tilig Bay (anch.).	do	do	8.30 p. m.		
	Tilig Bay (rf.).	do	July 15	9.00 p. m.		dense co. growth.
	do.	do	do	1.15 p. m.		mrgn. Co.
D. 5274	Malavatuan Id. (N.), S. 73° 30' E., 17.50 miles (13° 57' 30" N., 120° 03' 25" E.).	do	July 16	9.18 a. m.	525	gy. M., S.
				9.59 a. m.	525	gy. M., S.
D. 5275	Malavatuan Id. (N.), S. 71° E., 10.75 miles (13° 55' 55" N., 120° 10' 15" E.).	do	do	12.51 p. m.	117	fne. S.
H. 4913	Malavatuan Id. (N.), S. 67° E., 9.30 miles (13° 56' N., 120° 11' 40" E.).	do	do	1.05 p. m.	117	fne. S.
				1.28 p. m.	117	S., Sh., P.
	Balikias Bay (rf.).	do	July 17	5.30 a. m.		mrgn. Rf.
D. 5276	Malavatuan Id. (NW.), N. 61° 30' E., 6.50 miles (13° 49' 15" N., 120° 14' 45" E.).	do	do	8.44 a. m.	18	Sh., P., S.
D. 5277	Malavatuan Id. (N.), S. 56° E., 8 miles (13° 56' 55" N., 120° 13' 45" E.).	do	do	8.51 a. m.	18	Sh., P., S.
				10.02 a. m.	80	fne. S.
D. 5278	Malavatuan Id. (N.), S. 23° E., 8.50 miles (14° 00' 10" N., 120° 17' 15" E.).	do	do	10.19 a. m.	80	fne. S.
				11.34 a. m.	102	fne. S., M., Sh.
				11.53 a. m.	102	fne. S., M., Sh.
D. 5279	Malavatuan Id. (W.), S. 18° W., 5.40 miles (13° 57' 30" N., 120° 22' 15" E.).	do	do	1.13 p. m.	117	gn. M.
				1.26 p. m.	117	gn. M.
D. 5280	Malavatuan Id. (N.), S. 60° W., 6.10 miles (13° 55' 20" N., 120° 25' 55" E.).	do	do	2.42 p. m.	193	gy. S.
				3.05 p. m.	193	gy. S.
D. 5281	Malavatuan Id. (N.), S. 84° W., 4.30 miles (13° 52' 45" N., 120° 25' E.).	do	July 18	10.17 a. m.	201	dk. gy. S.
				10.40 a. m.	201	dk. gy. S.
D. 5282	Malavatuan Id. (N.), S. 84° W., 6.20 miles (13° 53' N., 120° 26' 45" E.).	do	do	11.21 a. m.	248	dk. gy. S.
				11.44 a. m.	248	dk. gy. S.
D. 5283	Malavatuan Id. (N.), N. 64° W., 8.75 miles (13° 48' 30" N., 120° 28' 40" E.).	do	do	1.06 p. m.	280	dk. gy. S.
				1.36 p. m.	280	dk. gy. S.
	Looc Bay (anch.).	do	do	8.45 p. m.		
D. 5284	Malavatuan Id. (S.), N. 46° W., 14.25 miles (13° 42' 05" N., 120° 30' 45" E.).	do	July 20	8.07 a. m.	422	gy. M., Glob.
				8.45 a. m.	422	gy. M., Glob.
D. 5285	Malavatuan Id. (S.), N. 45° W., 17.50 miles (13° 39' 36" N., 120° 32' 55" E.).	do	do	10.05 a. m.	272	sft. M.
				10.33 a. m.	272	sft. M.
H. 4914	Malavatuan Id. (S.), N. 42° W., 18.70 miles (13° 38' 05" N., 120° 33' E.).	do	do	11.35 a. m.	464	gy. M., S.
D. 5286	Malavatuan Id. (S.), N. 45° W., 19.50 miles (13° 38' 15" N., 120° 34' 20" E.).	do	do	12.31 p. m.	450	gy. S., M.
				1.09 p. m.	450	gy. S., M.
D. 5287	Sombro Id., N. 68° E., 11.25 miles (13° 37' 40" N., 120° 39' E.).	do	do	2.30 p. m.	379	gy. S.
				2.58 p. m.	379	gy. S.
	Port Maricaban (anch.).	do	do	8.15 p. m.		
	do.	do	do	9.00 p. m.		
	Port Maricaban (rf.).	do	July 21	6.00 a. m.		staghorn Clmps. S.
D. 5288	Matocot Pt., Luzon, S. 20° E., 5.70 miles (13° 43' 30" N., 121° E.).	do	July 22	8.14 a. m.	*140	S., M.*
D. 5289	Matocot Pt., S. 42° E., 5 miles (13° 41' 50" N., 120° 58' 30" E.).	do	do	9.03 a. m.	172	brk. Sh., S.
				9.25 a. m.	172	brk. Sh., S.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.								
					dyn.....	15 ft	<i>h. m.</i> 3 00		<i>mi.</i>	6 shots.
					dip; e. l.....	surface	1 00			
					dyn.....	12-20ft.	2 00			2 shots.
					dyn.....	15 ft	4 15			10 shots.
82	83	41.3	1.02497	1.02577	Luc. sdr. (a)...					
82	83				12' Agz.....	botm...	30	N. 63° E...	1.7	
82	83		1.02453		Tnr.-Blish sdr. (e).....					
82	84				12' Agz.....	botm...	20	N. 84° E...	1.5	
					Tnr.-Blish sdr. (e).....					Terminal sound- ing of D. 5275.
					dyn.....	6-12ft..	2 00			7 shots.
80	82				Tnr.-Blish sdr. (e).....					
80	82				12' Agz.; m. b.	botm...	15	N. 22° W...	.7	Net badly torn.
82.5	83	58.6	1.02442		Tnr.-Blish sdr. (e).....					
81	83				12' Agz.; m. b.	botm...	20	S. 70° E...	1.2	
82	82	59.6	1.02457		Tnr.-Blish sdr. (e).....					
83	82				12' Agz.; m. b.	botm...	4	S. 80° E...	.6	Belly of net carried away by weight of mud when hoisted from water.
83	84		1.02452		Tnr.-Blish sdr. (e).....					
83	83				12' Agz.; m. b.	botm...	9	N. 60° E...	.8	Net torn; 1 bridle stop carried away
81	83	49.6	1.02422	1.02517	Luc. sdr. (a)...					
81	83				12' Agz.; m. b.	botm...	18	N. 38° E...	.6	
81.5	84	50.4	1.02402	1.02538	Luc. sdr. (a)...					
82	83				12' Agz.; m. b.	botm...	20	N. 86° E...	1.3	
82	83	47.4	1.02437	1.02517	Luc. sdr. (a)...					
82	83				12' Agz.; m. b.	botm...	20	N. 85° E...	.7	
79	83	46.8	1.02417	1.02517	Luc. sdr. (a)...					
80	83				12' Agz.; m. b.	botm...	24	S. 83° E...	1.7	
					dip; e. l.....	surface	2 15			
83	84	42.3	1.02437	1.02566	Luc. sdr. (a)...					
84	84				12' Agz.; m. b.	botm...	25	S. 24° E...	1.1	
85	84	46.5	1.02497	1.02421	Luc. sdr. (a)...					
84	84				12' Agz.; m. b.	botm...	30	S. 21° E...	1.7	Sounding cup lost.
84	84	46.5	1.02473		Luc. sdr. (a)...					
84.5	84	42.5	1.02503	1.02556	Luc. sdr. (a)...					
85	85				12' Agz.; m. b.	botm...	20	N. 78° E...	1.8	Net wrecked.
84	85	43.4	1.02433	1.02521	Luc. sdr. (a)...					
84	84				int. 5 §.....	310 fms.	20	S. 73° E...	2.2	550 fms. dredge cable out.
					dip; e. l.....	surface	2 45			
					K2; K5.....	surface	15			Towed from row boat.
					dyn.....	12-20 ft.	4 00			9 shots.
82	83		1.02477		int. 5 §.....	115 fms.	19 8	N 76° W...	.7	200 fms. dredge cable out.
82	83		1.02497	1.02359	Tnr.-Blishsdr. (e).....					
82	84				12' Agz.; m. b.	botm...	20	S. 52° E...	1.0	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>China Sea, vicinity southern Luzon—Continued.</i>					
D. 5290	Matocot Pt., S. 50° E., 3.10 miles (13° 40' 09" N., 120° 59' 30" E.).	C. S. 4240; Feb., 1907.	1908. July 22	10.54 a. m.	fms. *214	Lav., G.
.....	Verde Id., San Augustine Vill. (rf.).do.....do.....	1.00 p. m.	setrd. Clmps. Co. on sloping bottom.
.....	Verde Id. (E. side) (rf.).do.....do.....	4.00 p. m.	dead Co.; S.
.....	Varadero Bay (anch.).do.....do.....	8.00 p. m.
.....	Varadero Bay (N. side) (rf.).do.....	July 23	6.00 a. m.	setrd. Clmps., sft. Co.
.....	Varadero Bay (beach).do.....do.....	8.15 a. m.	S., grassy
D. 5291	Escarceo Lt., N. 39° W., 2.20 miles (13° 29' 40" N., 121° 00' 45" E.).do.....do.....	1.27 p. m.	173	fne. bk. S.
.....do.....do.....	1.45 p. m.	173	fne. bk. S.
D. 5292	Escarceo Lt., N. 30° W., 3.25 miles (13° 28' 45" N., 121° 01' 12" E.).do.....do.....	2.23 p. m.	162	fne. bk. S.
.....do.....do.....	2.37 p. m.	162	fne. bk. S.
D. 5293	Escarceo Lt., N. 59° W., 6 miles (13° 28' 15" N., 121° 04' 30" E.).do.....do.....	3.42 p. m.	180	fne. bk. S.
.....do.....do.....	3.59 p. m.	180	fne. bk. S.
.....	Varadero Bay (fresh-water stream).do.....	July 24	6.00 a. m.	M.
D. 5294	Escarceo Lt., S. 71° W., 2.75 miles (13° 32' 15" N., 121° 02' E.).do.....do.....	8.54 a. m.	244	S., P.
.....do.....do.....	9.13 a. m.	244	S., P.
D. 5295	Escarceo Lt., S. 20° W., 2 miles (13° 33' 15" N., 121° E.).do.....do.....	10.06 a. m.	231	gy. S.
.....do.....do.....	10.26 a. m.	231	gy. S.
D. 5296	Matocot Pt., S. 63° E., 4.50 miles (13° 40' 09" N., 120° 57' 45" E.).do.....do.....	12.47 p. m.	*210	M., S.*
D. 5297	Matocot Pt., S. 50° E., 5.10 miles (13° 41' 20" N., 120° 58' E.).do.....do.....	1.55 p. m.	*198	M., S.*
D. 5298	Matocot Pt., S. 38° E., 6.70 miles (13° 43' 25" N., 120° 57' 40" E.).do.....do.....	3.09 p. m.	*140	S.*
D. 5299	(20° 05' N., 116° 05' E.).	H. O. 798; June, 1885.	Aug. 8	8.10 a. m.	524
.....do.....do.....	8.53 a. m.	524	gy. M., S.
D. 5300	(20° 31' N., 115° 49' E.).do.....do.....	2.07 p. m.	265	gy. M., S.
.....do.....do.....	2.29 p. m.	265	gy. M., S.
	<i>China Sea, vicinity Hongkong.</i>					
D. 5301	(20° 37' N., 115° 43' E.).	H. O. 798; June, 1885.	Aug. 8	5.06 p. m.	208	gy. M., S.
.....do.....do.....	5.29 p. m.	208	gy. M., S.
D. 5302	(21° 42' N., 114° 50' E.).do.....	Aug. 9	6.43 a. m.	38	sft. gy. M.
.....do.....do.....	6.51 a. m.	38	sft. gy. M.
D. 5303	(21° 44' N., 114° 48' E.).do.....do.....	8.21 a. m.	34	bl. M.
.....do.....do.....	8.27 a. m.	34	bl. M.
D. 5304	(21° 46' N., 114° 47' E.).do.....do.....	9.06 a. m.	*34	bl. M.
D. 5305	(21° 54' N., 114° 46' E.).do.....	Oct. 24	8.07 p. m.	*37	sft. gy. M.
.....	Pratas Id. (SW. side, beach).do.....	Oct. 25	3.00 p. m.	S., Co., grass
.....	Pratas Id. (SW. side, rf.).do.....do.....	3.00 p. m.	setrd. Clmps. Co., S.
D. 5306	(20° 55' N., 116° 40' E.).do.....	Oct. 26	8.09 a. m.	170	Co., S.
.....do.....do.....	8.35 a. m.	170	Co., S.
D. 5307	(21° 08' N., 116° 45' E.).do.....do.....	10.39 a. m.	186	Glob.
.....do.....do.....	11.04 a. m.	186	Glob.
D. 5308	(21° 54' N., 115° 42' E.).do.....	Nov. 4	6.35 a. m.	62	S., M.
.....do.....do.....	6.43 a. m.	62	S., M.
D. 5309	(21° 50' N., 115° 51' E.).do.....do.....	8.20 a. m.	62	gn. M.
.....do.....do.....	8.32 a. m.	62	gn. M.
.....do.....do.....	3.32 a. m.	62	gn. M.
D. 5310	(21° 33' N., 116° 13' E.).do.....do.....	12.36 p. m.	100	S., Sh.
.....do.....do.....	12.51 p. m.	100	S., Sh.
D. 5311	(21° 33' N., 116° 15' E.).do.....do.....	1.52 p. m.	88	crs. S., Sh.
.....do.....do.....	1.39 p. m.	88	crs. S., Sh.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Tempera- tures.			Density.		Apparatus.	Tide.		Drift.		Remarks.
Alr.	Surface.	Bottom.	Sur- face.	Bot- tom.		Depth.	Dura- tion.	Direction.	Dis- tance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>m.</i>		
84	84		1.02482	1.02354	12' Agz.; m. b.	botm...	20	S. 36° E...	1.3	Sounding failed on account of too light lead. Net slightly torn.
					dyn.....	12-25 ft.	1 30			4 shots.
					dyn.....	12-25 ft.	1 00			Do.
					dip; e. l.....	surface	6 00			
					dyn.....	6-15 ft.	4 00			8 shots.
					150' seine.....	8 ft	3 00			7 hauls.
86	84	51.5	1.02462	1.02468	Luc. sdr. (a)					
85	84				12' Agz.; m. b.	botm...	20	S. 28° E...	1.0	
83	84	52.4	1.02473	1.02421	Luc. sdr. (a)					
83	84				12' Agz.; m. b.	botm...	20	S. 13° E...	.9	
84	84	57.4	1.02457	1.02510	Luc. sdr. (a)					
84.5	84				12' Agz.; m. b.	botm...	30	W.....	.8	
					20' seine.....	3 ft				6 hauls.
82	83	48.4	1.02580	1.02482	Luc. sdr. (a)					
83	83				12' Agz.; m. b.	botm...	17	N. 86° W...	.6	Mud bag torn.
83	84	51.3	1.02457	1.02513	Luc. sdr. (a)					
83	84				12' Agz.; m. b.	botm...	20	N. 59° W...	1.2	
84	84		1.02473		12' Agz.; m. b.	botm...	20	S. 63° E...	1.2	
85	85		1.02477		12' Agz.; m. b.	botm...	20	S. 69° E...	1.0	
83	84				12' Agz.; m. b.	botm...	10	S. 31° E...	.5	Do.
85.5	83	42.5	1.02396	1.02538	Luc. sdr. (a)					
83.5	84				12' Agz.; m. b.	botm...	22			Ship steered circular course.
86	85		1.02350	1.02430	Luc. sdr. (a)					Therm. failed to trip.
87	85				12' Agz.; m. b.	botm...	20			
85	84	50.5	1.02433	1.02456	Luc. sdr. (a)					
85	84				12' Agz.; m. b.	botm...	20			
84	83	72.1	1.02288		Tnr.-Blshsdr.					
					(e).					
84	84				12' Agz.; m. b.	botm...	15			
85	83	71.6	1.01960	1.02386	Tnr.-Blshsdr.					
					(e).					
84	84				12' Agz.; m. b.	botm...	20			
85.5	84				12' Agz.	botm...	20			
79	78				12' Tnr.	botm...	20			
					130' seine.....	15 ft	2 00			3 hauls.
					dyn.....	10-25 ft.	2 00			3 shots.
80	80	51.4		1.02489	Luc. sdr. (a)					
79.5	80				12' Tnr.	botm...	20			
80	80	51.6	1.02434	1.02510	Luc. sdr. (a)					
80.5	80				12' Tnr.	botm...	20			
77	77		1.02461		Tnr.-Blshsdr.					
					(e).					
77	78				12' Tnr.	botm...	15			
78	79	73.3			Tnr.-Blshsdr.					
					(e).					
79	79				12' Tnr.	botm...	20			
79	79				K2	surface	20			Towed from horse block.
80	80	65.5			Tnr.-Blshsdr.					
					(e).					
80	80				12' Tnr.	botm...	20			
81	80				Tnr.-Blshsdr.					
					(e).					
81	80				12' Tnr.; m. b.	botm...	20			

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>China Sea, vicinity Hong-kong—Continued.</i>						
D. 5312	(21° 30' N., 116° 32' E.).....	H. O. 798; June, 1885.	1908. Nov. 4	4.05 p. m. 4.27 p. m. 6.20 p. m.	<i>fms.</i> 140 140 150	S., sml. Sh..... S., sml. Sh..... S.....
D. 5313	(21° 30' N., 116° 43' E.).....do.....do.....	6.45 p. m. 6.05 a. m. 6.25 a. m.	150 122 122	S..... S., brk. Sh..... S., brk. Sh.....
D. 5314	(21° 41' N., 116° 46' E.).....do.....	Nov. 5	6.25 a. m. 6.25 a. m.	122 122	S., brk. Sh..... S., brk. Sh.....
<i>China Sea, vicinity Formosa.</i>						
D. 5315	(21° 40' N., 116° 58' E.).....	H. O. 798; June, 1885.	Nov. 5	8.21 a. m. 8.42 a. m.	148 148	S., Sh..... S., Sh.....
D. 5316	(21° 39' N., 117° 07' E.).....do.....do.....	10.37 a. m. 10.57 a. m.	159 159	S., Sh..... S., Sh.....
D. 5517	(21° 36' N., 117° 27' E.).....do.....do.....	2.05 p. m. 2.31 p. m.	230 230	S., sml. Sh..... S., sml. Sh.....
D. 5318	(21° 32' N., 117° 46' E.).....do.....do.....	5.03 p. m.	340	S., br. C.....
				5.32 p. m.	340	S., br. C.....
D. 5319	(21° 31' N., 117° 53' E.).....do.....do.....	7.23 p. m.		
H. 4915	(21° 23' N., 118° 30' E.).....do.....	Nov. 6	12.11 a. m.	(?)689	
H. 4916	(21° 14' N., 119° 02' E.).....do.....do.....	4.32 a. m.	1,498	
H. 4917	(21° 06' N., 119° 38' E.).....do.....do.....	10.15 a. m.	1,758	sft. br. M.....
D. 5320	(20° 58' N., 120° 03' E.).....do.....do.....	2.25 p. m.	1,804	gy. M.....
				3.18 p. m.	1,804	
H. 4918	(20° 46' N., 120° 52' E.).....do.....do.....	9.32 p. m.	1,220	sft. M.....
	Santo Domingo, Batan Id. (rf.)	C. S. 4710; July, 1905.	Nov. 7	8.00 a. m.		Co., Lav.....
	Sabtan Id. (SW. side) (rf.)do.....	Nov. 8	1.00 p. m.		Co., R.....
H. 4919	Ibugos Id. (S. end) N. 77° W., 1 mile (20° 19' 15" N., 121° 51' E.)do.....	Nov. 9	6.00 a. m.		Co., R.....
				(?)*	64	
H. 4920	Ibugos Id. (S. end) N. 81° W., 1.25 miles (20° 19' 15" N., 121° 51' 20" E.)do.....do.....	11.18 a. m.	46	
D. 5321	Ibugos Id. (S. end) S. 89° W., 1.25 miles (20° 19' 30" N., 121° 51' 15" E.)do.....do.....	11.23 a. m.	26	wh. S., Co., brk. Sh.....
				11.25 a. m.	26	wh. S., Co., brk. Sh.....
D. 5322	Ibugos Id. (S. end) S. 84° W., 1.25 miles (20° 19' 36" N., 121° 51' 15" E.)do.....do.....	11.42 a. m.	21	wh. S., Co., brk. Sh.....
<i>China Sea, vicinity of Batanes.</i>						
D. 5323	Ibugos Id. (S. end), N. 0° 30' W., 12 miles (20° 07' 15" N., 121° 50' E.)	C. S. 4710; July, 1905.	Nov. 9	1.39 p. m. 2.12 p. m.	303 303	
D. 5324	Ibugos Id. (S. end), N. 15° E., 10.50 miles (20° 09' N., 121° 47' E.)do.....do.....	3.19 p. m. 4.10 p. m.	564 564	rky..... rky.....
	Port San Pio Quinto, Cami- guin Id. (rf.)	C. S. 4711; May, 1907.	Nov. 10	9.30 a. m.		Co., R.....
	Port San Pio Quinto (beach)do.....do.....	1.30 p. m. 1.30 p. m.		Co., R..... S., P.....

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trawl.		Drift.		Remarks.
Alr.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.								
80	80	57.5	1.02461	1.02482	Luc. sdr. (a)		<i>h. m.</i>		<i>mi.</i>	
81	80				12' Tnr.; m. b.	botm...	17			
78	80	53.6	1.02461	1.02513	Luc. sdr. (a)					
77	80				12' Tnr.; m. b.	botm...	15			
78	78	59.5	1.02461	1.02526	Luc. sdr. (a)					
78	79				12' Tnr.; m. b.	botm...	20			
78	79				K2	surface.	20			Towed from horse block.
79	79	54.4	1.02500	1.02506	Luc. sdr. (a)					
80	79				12' Tnr.; m. b.	botm...	20			
80	80	53.4	1.02481	1.02517	Luc. sdr. (a)					
82	80				12' Tnr.; m. b.	botm...	25			Mud bag torn.
82	80	50.6	1.02474		Luc. sdr. (a)					
81	80				12' Tnr.; m. b.	botm...	20			
81	79				Luc. sdr. (a)					Sounding outfit lost with 340 fms. wire.
80	79				12' Tnr.; m. b.	botm...	6			Bridle stop carried away; net came up, upside down.
79	79				int. 4 §	20 fms.	27			40 fms. dredge cable out.
79	78				Luc. sdr. (a)		6			Sounding outfit lost with 689 fms. wire. May not have reached bottom.
79	78				Luc. sdr. (a)					Outfit and stray line lost while heaving in.
80	80				Luc. sdr. (a)					Strong current.
80	80	36.2		1.02574	Luc. sdr. (a)					Therm. possibly tripped at 930 fms.
80	80				int. 4, 2; K2 §	500 fms.	20			
80	80	36.4			Luc. sdr. (a)		33			
					dyn.	10-20ft.	3	30		8 shots.
					dyn.	10-25ft.	4	00		9 shots.
					dyn.	10-25ft.	2	00		2 shots.
					Tnr.-Blishsdr. (e)					
					Tnr.-Blishsdr. (e)					
82	81				Tnr.-Blishsdr. (e)					
82	81				9' Jn. dr	botm...	4	N.	0.2	
82	81				9' Jn. dr	botm...	9	N.	.2	Sounding with hand lead.
81	82	58.4	1.02558		Luc. sdr. (a)					
81	82				12' Tnr.; m. b.	botm...	20	N. 62° W.	3.2	
82	82	40.9	1.02523	1.02533	Luc. sdr. (a)					
78	81				12' Tnr.; m. b.	botm...	2			Trawl lost; bridle and mud bag recovered.
					dyn.	12-20 ft.	2	30		2 shots.
					dyn.	12-25ft.	3	00		3 shots.
					130' seine	10 ft.	3	00		5 hauls.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>China Sea, vicinity of Batanes—Continued.</i>					
.....	Port San Pio Quinto (beach at head of bay).	C. S. 4711; May, 1907.	1908. Nov. 11	9.00 a. m.	fms.	fine S.
.....	Port San Pio Quinto (rf.)	do	do	8.30 a. m.	strd. Clmps. Co.
.....			Nov. 12	1.30 p. m.	strd. Clmps. Co.
.....				6.00 a. m.	strd. Clmps. Co.
	<i>Off northern Luzon.</i>					
D. 5325	Hermanos Id. (N.), N. 86° E., 16.75 miles (18° 34' 15" N., 121° 51' 15" E.).	C. S. 4711; May, 1907.	Nov. 12	10.45 a. m. 11.13 a. m.	224 224	gn. M. gn. M.
D. 5326	Hermanos Id. (N.), N. 69° E., 8 miles (18° 32' 30" N., 122° 01' E.).	do	do	1.00 p. m. 1.28 p. m.	230 230	M. M.
D. 5327	Hermanos Id. (N.), N. 55° E., 6.80 miles (18° 31' 30" N., 122° 03' E.).	do	do	2.16 p. m.	198	sft. M., fine S.
.....	Port San Vicente, Luzon side (beach). ^a	do	Nov. 13 Nov. 18	2.29 p. m. 2.00 p. m. 8.00 a. m.	198	sft. M., fine S. M., S., grass, etc. M., S., sticks, leaves.
.....	Channel bet. Palaui and San Vicente Islands, Palaui side (beach).	do	do	3.00 p. m.	S., M., grass.
.....	Palaui Id. (W. side) (rf.)	do	do	10.00 a. m.	strd. Co., S.
.....	Palaui Id. (W. side), small stream.	do	do	2.00 p. m.
D. 5328	Hermanos Id., N. 79° E., 28.40 miles (18° 29' 45" N., 121° 39' E.).	do	Nov. 19	9.23 a. m. 9.44 a. m.	150 150	bl. M. bl. M.
D. 5329	Font Id. (W.), N. 28° E., 24.25 miles (18° 33' N., 121° 37' 30" E.).	do	do	10.58 a. m. 11.25 a. m.	212 212	bl. M. bl. M.
D. 5330	Font Id. (W.), N. 24° E., 23.30 miles (18° 33' 30" N., 121° 39' 15" E.).	do	do	1.12 p. m. 1.33 p. m.	178 178	br. M. br. M.
	<i>Off western Luzon.</i>					
D. 5331	Hermana Menor Id. (E.), N. 13° E., 7.30 miles (15° 36' 45" N., 119° 47' 45" E.).	C. S. 4712; Sept., 1904.	Nov. 22	8.12 a. m. 8.41 a. m.	178 178	S., Sh., M. S., Sh., M.
.....	Port Matalvi (rf.)	do	do	10.30 a. m.	strd. Co., S.
.....	Port Matalvi (anch.)	do	do	1.30 p. m. 7.45 p. m.	strd. Co., S.
.....	Port Matalvi (rf.)	do	Nov. 23	6.00 a. m.	strd. Co., S.
.....	Port Matalvi (E. side San Salvador Id.) (beach).	do	do	8.30 a. m.	S., Co., grass.
.....	Port Matalvi (E. side Macalaba Id.) (beach).	do	do	1.30 p. m.	S., M., grass.
.....	Port Matalvi (rf.)	do	do	1.30 p. m.	strd. Co.
	<i>Mindoro Strait.</i>					
.....	Paluan Bay, Pantocomi Pt.	C. S. 4345; Feb., 1905.	Dec. 11	7.15 a. m.	R., Co.
.....	Paluan Bay, Lipa Beach	do	do	8.30 a. m.	S., P.
.....	Paluan Bay, Paluan River	do	do	9.00 a. m.	M.
.....	Paluan Bay, Malugao River	do	do	2.00 p. m.	M., sticks, leaves.
.....	Paluan Bay, beach N. of Malugao River.	do	do	3.00 p. m.	M.
.....	Paluan Bay, Caluagan River.	do	do	3.00 p. m.
.....	Paluan Bay, anch.	do	do	7.00 p. m.

^a On November 14 a party went up Palaui River about 3 or 4 miles, in prahm, seining with 25-foot and 45-foot seines at intervals along entire distance.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Alr.	Surface.	Bottom.	Surface.	Bot- tom.		Depth.	Dura- tion.	Direction.	Distance.	
° F.	° F.	° F.								
					130' seine.	5 ft.	<i>h. m.</i> 3 00			7 hauls.
					25' seine.	3 ft.	1 30			12 hauls in small stream.
					45' seine.	3 ft.	1 30			10 hauls in small stream.
					dyn.	12-20 ft.	3 30			4 shots.
					dyn.	12-20 ft.	4 00			2 shots.
					dyn.	12-25 ft.	1 00			
81	82	53.2	1.02491	1.02525	Luc. sdr.(a)					
81	82				12' Tnr.; m. b.	botm.	19	S. 50° E.	1.0	
82	81	55.4	1.02437	1.02496	Luc. sdr.(a)					
81	81				12' Tnr.; m. b.	botm.	20	S. 60° E.	2.0	
82	82	(?)	1.02434	1.02468	Luc. sdr.(a)					Therm. failed to trip.
81	81				12' Tnr.; m. b.	botm.	20			7 hauls.
					130' seine.	5 ft.	3 00			Do.
					130' seine.	5 ft.	4 00			
					130' seine.	2-4 ft.	1 30			4 hauls.
					dyn.	10-20 ft.	5 00			7 shots in a. m.; several in p. m.
					45' seine.		2 00			3 hauls.
78	79	53.9	1.02464	1.02513	Luc. sdr.(a)					
78	78				12' Tnr.; m. b.	botm.	20	N. 52° W.	1.2	
79	78	51.4	1.02492	1.02593	Luc. sdr.(a)					
79	78				12' Tnr.; m. b.	botm.	10	N. 50° W.	2.2	
78	78	53.4	1.02516	1.02523	Luc. sdr.(a)					
78	78				12' Tnr.; m. b.	botm.	20	(?)	(?)	
80.5	80	54.7	1.02422	1.02496	Luc. sdr.(a)					
80.5	80				12' Tnr.; m. b.	botm.	20	S. 49° E.	2.0	
					dyn.	10-20 ft.	1 30			3 shots.
					dyn.	10-20 ft.	3 30			4 shots.
					dip; e. l.	surf.	1 30			2 dynamite caps exploded at gang-way.
					dyn.	10-30 ft.	5 00			8 shots.
					130' seine.	4-10 ft.	3 00			7 hauls.
					130' seine.	2-4 ft.	2 00			4 hauls.
					dyn.	8-20 ft.	2 00			4 shots.
					dyn.	10-20 ft.	4 15			4 shots.
					130' seine.	8 ft.	3 00			10 hauls.
					25' seine.	2 ft.	2 00			Do.
					130' seine.	5 ft.	2 30			5 hauls.
					25' seine.	3 ft.	3 30			4 hauls.
					16' seine					
					dip; e. l.	surface.	1 30			

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Mindoro Strait—Continued.</i>					
	Sablayan Bay, near Sablayan.	C. S. 4345; Feb., 1905.	1908. Dec. 12	10.00 a. m.	<i>fms.</i>	Co.
D. 5332	Apo Lt., S. 66° W., 18.2 miles (12° 47' 15" N., 120° 41' E.).	C. S. 4714; June, 1906.	do	10.39 a. m.	745	gn. M.
H. 4921	Apo Lt., S. 65° W., 19.4 miles.	do	do	11.50 a. m.		gy. M., crs. S.
	Sablayan Bay, Sablayan Pt.	C. S. 4345; Feb., 1905.	do	1.50 p. m.	584	
	Sablayan Bay, anch.	do	do	3.30 p. m.		
	Sablayan Bay, Sablayan Pt.	do	Dec. 13	7.00 p. m.		co. R.
	Sablayan Bay, Pandan Id.	do	do	10.00 a. m.		Co.
	Sablayan Bay, Bagaong River.	do	do	10.00 a. m.		
	Sablayan Bay, anch.	do	do	9.00 p. m.		
D. 5333	Apo Lt., N. 45° W., 19 miles (12° 26' 30" N., 120° 37' 45" E.).	C. S. 4714; June, 1906.	Dec. 14	7.40 a. m.	310	S.
				8.26 a. m.		
D. 5334	Apo Lt., N. 44° W., 19.7 miles (12° 25' 40" N., 120° 38' E.).	do	do	9.18 a. m.	612	gy. M.
				9.58 a. m.		
				10.17 a. m.		
	Tara Id., west.	do	do	3.00 p. m.		dense Co.
	Tara Id., anch.	do	do	7.00 p. m.		
	Tara Id., west.	do	Dec. 15	7.30 a. m.		
	Tara Id., bayou near village.	do	do	7.30 a. m.		sft. M.
	Tara Id., beach near village.	do	do	9.00 a. m.		S., Co., grass
	<i>Busuanga Id.</i>					
	Port Caltom.	C. S. 4714; June, 1906.	Dec. 15	2.00 p. m.		setrd. Co.
	Port Caltom, beach near village.	do	do	2.00 p. m.		S., Co., W.
	Port Caltom, anch.	do	do	7.00 p. m.		
	Port Caltom, Pangauran River.	do	Dec. 16	7.00 a. m.		
	Port Uson, Malbato River.	C. S. 4345; Feb., 1905.	Dec. 17	1.00 p. m.		
	Port Uson, Mayanpayan Id.	do	do	2.00 p. m.		setrd. Co.
	Port Uson, anch.	do	do	8.00 p. m.		
	<i>Linapacan Strait.</i>					
D. 5335	Observatory Id. (N.), S. 55° W., 10.7 miles (11° 37' 15" N., 119° 48' 45" E.).	C. S. 4716; Jan., 1903.	Dec. 18	12.22 p. m.	46	S., M.
D. 5336	Observatory Id. (N.), S. 42° W., 9 miles (11° 37' 45" N., 119° 46' E.).	do	do	12.43 p. m.	46	S., M.
				1.16 p. m.		
				1.26 p. m.		
	Linapacan Id., Malcochin Harbor.	do	do	3.30 p. m.		S., W., Co.
	Linapacan Id., Malcochin Harbor, anch.	do	do	8.00 p. m.		
	Linapacan Id., Malcochin Harbor, beach.	do	Dec. 19	8.00 a. m.		S. Co.
	Linapacan Id., Malcochin Harbor reef.	do	do	8.00 a. m.		setrd. Co.
	Observatory Id., west beach.	do	do	2.30 p. m.		S., Co., W.
	Observatory Id., west.	do	do	2.30 p. m.		setrd. Co.
	<i>Palawan Passage.</i>					
D. 5337	Observatory Id. (N.), S. 80° E., 13.8 miles (11° 34' N., 119° 29' E.).	C. S. 4716; Jan., 1903.	Dec. 20	7.31 a. m.	43	fne. Co., S., M.
				7.40 a. m.		
D. 5338	Observatory Id. (N.), S. 82° E., 15 miles (11° 33' 45" N., 119° 24' 45" E.).	do	do	8.04 a. m.	43	Co., S., M.
				8.12 a. m.		
H. 4922	Cauayan Id. (N.), S. 37° E., 11.5 miles (11° 25' 45" N., 119° 14' E.).	do	do	8.15 a. m.	21	Co., S., Sh.
				10.01 a. m.		
D. 5339	Cauayan Id. (N.), S. 59° E., 10 miles (11° 22' N., 119° 12' E.).	do	do	10.32 a. m.	52	M.
				10.43 a. m.		
	North Guntao Id.	do	do	1.00 p. m.		Co., S.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Tide.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.								
					dyn.....	6-12 ft..	<i>h. m.</i> 4 30			9 shots.
84	81	38.2	1.02385	1.02548	Luc. sdr. (a)...					
82	81				12' Tnr.; m. b.	botm...	20			
83	82	40.2	1.02401	1.02535	Luc. sdr. (a)...					
					dyn.....		1 30			4 shots.
					dip; e. l.....	surface.	1 30			
					copper sulphate.		2 00			Work done in tide pools.
					dyn.....	6-9 ft..	2 00			5 shots.
					16' seine.....		6 00			
					dip; e. l.....	surface.	1 00			
79	80	73.8	1.02406	1.02543	Luc. sdr. (a)...					
81	80				12' Agz.; m. b.	botm...	22			
					dip; e. l.....	surface.	1 02			
81	80	43.2	1.02385	1.02516	Luc. sdr. (a)...					
82	80				K. 2.....	surface.	1 02			
					12' Agz.; m. b.	botm...	7	S. 60° W..	2.0	All gear but mud bag lost.
					dyn.....	10-20 ft.	2 00			3 shots.
					dip; e. l.....	surface.	1 00'			
					dyn.....	10-20 ft.	4 00			5 shots.
					130' seine.....	3 ft.....	1 30			2 hauls.
					25' seine.....	2 ft.....	1 30			12 hauls.
					dyn.....	10-20 ft.	3 00			8 shots.
					130' seine.....	4 ft.....	3 00			6 hauls.
					dip; e. l.....	surface.	1 30			
					dyn.; 25' seine.		9 00			10 shots.
					dyn.....		4 00			
					dyn.....	10-20 ft.	2 30			
					dip; e. l.....	surface.	1 00			
					Tnr. sdr. (e)...					Therm. failed to trip.
82	80				9' Tnr.; m. b.	botm...	17	N. 77° W..	1.2	No therm. used.
83	81				Tnr. sdr. (e)...					Lost bottom of net.
83	81				9' Tnr.; m. b.	botm...	6	N. 80° W..	1.2	
					130' seine.....	3 ft.....	1 30			3 hauls.
					dip; e. l.....	surface.	1 00			
					130' seine.....	3 ft.....	4 00			7 hauls.
					dyn.....	10-20 ft.	4 00			11 shots.
					130' seine.....	4 ft.....	2 30			6 hauls.
					dyn.....	15 ft.....				1 shot.
					Tnr. sdr. (e)...					No therm. used.
81	80		1.02427		9' Tnr.; m. b.	botm...	9	S. 82° W..	1.0	
					Tnr. sdr. (e)...					Do.
81	80				9' Tnr.; m. b.	botm...	20	N. 70° W..	1.3	
81	80				K. 2.....	surface.	20			Do.
					Tnr. sdr. (e)...					
					Tnr. sdr. (e)...					
83	81		1.02406		9' Tnr.; m. b.	botm...	20	S. 58° W..	2.2	
84	81				ynd.....	24-30 ft.	2 30			7 shots.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Malampaya Sound, Palawan Id.</i>					
	Bolalo Bay, anch.	C. S. 4349;	1908. Dec. 20	8.30 p. m.	<i>fms.</i>	
	Bolalo Bay, flats near shore.	Aug., 1908.	Dec. 21	8.00 a. m.		S., Co., W.
	Bolalo Bay, mouth of bay.	do.	do.	8.00 a. m.		Co., W.
	Bolalo Bay, head of bay.	do.	do.	1.00 p. m.		
	Bolalo Bay, anch.	do.	do.	7.30 p. m.		
	Bolalo Bay, near anch.	do.	do.	9.00 p. m.		
D. 5340	Cone Id., N. 2° E., 1.5 miles (10° 55' 51" N., 119° 14' 12" E.).	do.	Dec. 22	8.22 a. m.	19-24	
	Endeavor Strait, near Relinquish Head.	do.	do.	9.00 a. m.		
	Endeavor Strait, Chase Head.	do.	do.	2.00 p. m.		Co., S.
	Endeavor Strait, Limunancong.	do.	do.	2.00 p. m.		S.
	Endeavor Strait, Relinquish Head to Nalinbungan Pt.	do.	Dec. 23	8.00 a. m.		Co., S.
D. 5341	Endeavor Pt. (W.), S. 18° E., 1.2 miles (10° 57' 51" N., 119° 17' 20" E.).	do.	do.	2.03 p. m.	19-22	gy. M.
D. 5342	Endeavor Pt. (S.), S. 58° E., 0.5 miles (10° 56' 55" N., 119° 17' 24" E.).	do.	do.	2.35 p. m.	14-25	gy. M.
	Endeavor Strait, anch. bet. Bando and Endeavor points.	do.	do.	8.00 p. m.		
	Endeavor Strait, anch. bet. Bando and Endeavor points.	do.	do.	8.30 p. m.		
	Malapna Id., N. W.	do.	Dec. 24	8.00 a. m.		Co., S., W.
	Inner Sound, near Pancel.	do.	Dec. 25	1.00 p. m.		S., R.
D. 5343	Cliff Id., S. 22° E., 5.2 miles (10° 51' 35" N., 119° 23' 24" E.).	do.	Dec. 26	7.46 a. m.	*5	M.
D. 5344	Cliff Id., S. 34° E., 4.7 miles (10° 50' 40" N., 119° 22' 32" E.).	do.	do.	8.22 a. m.	6	M.
	Inner Sound, Malampaya River.	do.	do.	9.00 a. m.		sft. M.
D. 5345	Cliff Id., S. 43° E., 4.4 miles (10° 50' N., 119° 22' 03" E.).	do.	do.	9.16 a. m.	7	M.
D. 5346	Cliff Id., S. 37° E., 4.6 miles (10° 50' 30" N., 119° 22' 20" E.).	do.	do.	10.18 a. m.	*7	M.
D. 5347	Cliff Id., S. 26° E., 4.5 miles (10° 50' 44" N., 119° 23' 09" E.).	do.	do.	10.58 a. m.	5	M.
	<i>Palawan Passage.</i>					
H. 4923	Pt. Tabonan, S. 87° E., 11.4 miles (10° 57' 15" N., 119° 1' E.).	C. S. 4716;	Dec. 27	6.32 a. m.	51	Co., S.
		Jan., 1903.				
H. 4924	Pt. Tabonan, East, 16.3 miles (10° 57' N., 118° 55' 45" E.).	do.	do.	7.10 a. m.	62	S.
H. 4925	Pt. Tabonan, S. 87° E., 24.3 miles (10° 58' 15" N., 118° 47' 15" E.).	do.	do.	8.05 a. m.	184	fine. Co., S.
D. 5348	Pt. Tabonan, S. 89° E., 33.5 miles (10° 57' 45" N., 118° 38' 15" E.).	do.	do.	9.28 a. m. 10.09 a. m.	375	Co., S.
D. 5349	Pt. Tabonan, N. 85° E., 45.2 miles (10° 54' N., 118° 26' 20" E.).	do.	do.	12.41 p. m. 1.40 p. m.	730	Co., S.
D. 5350	Pt. Tabonan, N. 76° E., 43.7 miles (10° 46' 40" N., 118° 29' E.).	do.	do.	4.10 p. m. 5.14 p. m.	515	gy. M.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
					dip; e. l.	surface.	1 00			
					130' seine.....	2-4 ft...	3 30			11 hauls.
					dyn.....	6-9 ft...	3 30			5 shots.
					dyn.....		4 00			3 shots.
					dip; e. l.	surface.	1 00			
					K2, K5.....	surface.	20			Tow'd from wherry.
					hand line.....					
81	80				int. 3 §.....	17-22 fms.	20 1	N. 3° W...	0.4	
					dyn.....		2 00			5 shots.
					dyn.....	9-12 ft...	2 00			3 shots.
					25' and 130' seines.	5 ft...	2 00			13 hauls.
					dyn.....	18-20 ft.	6 00			13 shots.
					hand line.....					
83	82				9' Tnr.; m. b. .	botm...	15	S. 2° E...	.7	
83	82				hand line.....					
83	82				9' Tnr.....	botm...	19	S. 25° W...	.7	Net slightly torn.
					K2; 2' o. p.	surface.	20			Towed from steam launch.
					dip; e. l.	surface.	1 30			
					dyn.....		3 30			11 shots.
					dyn.....		4 00			3 shots.
80	81				6' McC.....	botm...	15	S. 78° W...	.4	
					hand line.....					
81	81				6' McC.....	botm...	26	S. 18° W...	.7	
					dyn.; 130' seine	3-6 ft...	6 00			6 shots, 4 hauls.
					hand line.....					
80	81				9' Tnr.....	botm...	20	N. 47° W...	.6	
81	80				9' Tnr.....	botm...	10	S. 72° E...	1.0	
					hand line.....					
81	81				9' Tnr.....	botm...	10	N. 36° E...	.5	
					Tnr. sdr. (e)...					
					Tnr. sdr. (e)...					
					Lue. sdr. (a)...					
82	81	56.4	1.02422	1.02576	Lue. sdr. (a)...					
82	81				12' Tnr.; m. b. .	botm...	20	N. 80° W...	1.5	No land in sight; latitude and longitude ap- proximate.
		40.6	1.02406	1.02564	Lue. sdr. (a)...					
83	81				12' Tnr.; m. b. .	botm...	20	S. 80° W...	1.5	Do.
			1.02381	1.02523	Lue. sdr. (a)...					
82	80				12' Tnr.; m. b. .	botm...	20	S. 85° W...	3.0	Do.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Palawan Passage—Cont'd.</i>					
D. 5351	Pt. Tabonan, N. 62° E., 47 miles (10° 35' N., 118° 30' E.).	C. S. 4716; Jan., 1903.	1908. Dec. 27	8.43 p. m. 8.53 p. m.	<i>fms.</i> 50	Co., S.
	<i>Ulugan Bay, Palawan Id.</i>					
	Oyster Inlet.....	C. S. 4346; Aug., 1905.	Dec. 28	9.00 a. m.		S., Co.
	Baheli River to Wood Pt.....	do.....	do.....	9.30 a. m.		M., S., W.
	Magsiapo Reef.....	do.....	do.....	1.00 p. m.		Co.
	Sagumay Pt.....	do.....	do.....	1.00 p. m.		S., Co.
	Anchorage (near Tidepole Pt.).....	do.....	do.....	8.30 p. m.		
	Rita Id. (W. and S.).....	do.....	Dec. 29	8.00 a. m.		S., Co.
	Cañholo River.....	do.....	do.....	11.00 a. m.		G., bowlders.
D. 5352	Tidepole Pt., S. 34° W., 0.4 mile (10° 04' 30" N., 119° 05' E.).	do.....	Dec. 30	6.18 a. m.	25	M.
	<i>Nakoda Bay, Palawan Id.</i>					
	Siriniao Id. (S.W.).....	C. S. 4346; Aug., 1905.	Dec. 30	3.00 p. m.		S., W.
	River (unnamed), SE. of Maricaban Id.	do.....	Dec. 31	6.00 a. m.		M., S., G.
	<i>Balabac Strait.</i>					
D. 5353	Cape Melville Lt., S. 85° E., 16.8 miles (7° 50' 45" N., 116° 43' 15" E.).	C. S. 4309; Nov., 1906.	1909. Jan. 1	6.33 a. m. 7.10 a. m.	148	
D. 5354	Cape Melville Lt., N. 85° E., 16.8 miles (7° 47' 50" N., 116° 43' 15" E.).	do.....	do.....	8.33 a. m. 9.55 a. m.	117	M.
	<i>North Balabac Strait.</i>					
	Caxisigan Id. (W.).....	C. S. 4347; Dec., 1905.	Jan. 2	1.00 p. m.		Co., S.
	Port Ciego, Martinez Pt.....	do.....	Jan. 3	9.00 a. m.		W., Co.
	Port Ciego, Paz Id.....	do.....	do.....	9.00 a. m.		W., Co.
	Candaraman Id. (E.).....	do.....	Jan. 4	8.30 a. m.		S., Co.
	Bugsuk Id. (S.).....	C. S. 4309; Nov., 1906.	Jan. 5	8.00 a. m.		S., Co.
D. 5355	Balabac Lt., S. 61° W., 16.6 miles (8° 08' 10" N., 117° 19' 15" E.).	do.....	do.....	9.40 a. m. 9.52 a. m.	44	Co., S.
D. 5356	Balabac Lt., S. 64° W., 15.5 miles (8° 06' 40" N., 117° 18' 45" E.).	do.....	do.....	10.21 a. m. 10.36 a. m.	58	S., Sh.
D. 5357	Balabac Lt., S. 65° W., 14.3 miles (8° 06' N., 117° 17' 10" E.).	do.....	do.....	11.13 a. m. 11.27 a. m.	68	Co., S.
	<i>Jolo Sea.</i>					
	Taganak Id. (SE.).....	C. S. 4720; Jan., 1904.	Jan. 7	1.00 p. m.		Co.
D. 5358	Sandakan Lt., S. 34° W., 19.7 miles (6° 06' 40" N., 118° 18' 15" E.).	do.....	do.....	7.20 p. m. 7.29 p. m.	39	M.
	Cagayan de Jolo (S.).....	C. S. 4348; June, 1905.	Jan. 8	8.30 a. m.		Co., S.
	Cagayan de Jolo, Singuan Lake.	do.....	do.....	9.00 a. m. 3.00 p. m.		S., Co., W. M.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Alr.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				h. m.		mi.		
81	80				Tnr. sdr. (e) 12' Tnr.; m. b.	botm...	2			Net wrecked; latitude and longitude approximate.
					dyn.....		3 00			12 shots.
					130' seine.....	2-5ft...	5 00			9 hauls.
					dyn.....		2 30			2 shots.
					dyn.....		2 00			Do.
					dip; e. l.....	surface.	1 00			
					250' seine; dyn.	20-40ft.	2 00			2 hauls, 6 shots.
					25' seine.....		3 00			
					hand line					
80	81				int. 4. §.....	24 fms.	20 2	N. 4° E...	0.9	
					130' seine.....	4-10ft..	1 30			5 hauls.
					dyn.; 16'-45' seine.		10 00			
					Luc. sdr. (a)...					148 fms. sounding wire lost.
75	80				9' Tnr.; m. b.	botm...	34	SE.		Foggy; latitude and longitude approximate.
75	80				Tnr. sdr. (e) 9' Tnr.; m. b.	botm...	25	SE.		Do.
					dyn.....	15ft.				
					dyn.....	12ft....	4 00			5 shots.
					dyn.....		4 30			6 shots.
					dyn.....	9-15ft..	2 30			15 shots.
					dyn.....	9-18ft..	4 00			Do.
82	82		1.02518		Tnr. sdr. (e) 6' McC.....	botm...	19	S. 14° W..	1.6	
85	82				Tnr. sdr. (e) 6' McC.....	botm...	16	S. 50° W..	1.3	
85	82				Tnr. sdr. (e) 9' Tnr.; m. b.	botm...	01	N. 45° E..	.6	Net torn.
					dyn.....	15ft....	4 00			10 shots.
80	82				Tnr. sdr. (e) 12' Agz.; m. b.	botm...	14	N. 56° E..	.7	
					dyn.....		3 00			5 shots.
					130' seine.....	2-4ft...	2 30			4 hauls.
					dyn.....	10-40ft.	1 00			5 shots.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Jolo Sea—Continued.</i>					
H. 4926	7° 39' N., 120° 04' 45" E.	C. S. 4721; Jan., 1903.	1909. Jan. 9	6.11 a. m.	fms. 460	M.
D. 5359	8° 12' 45" N., 120° 37' 15" E.	do.	do.	12.52 p. m. 3.31 p. m.	2, 275	
	<i>Iloilo Strait.</i>					
	Anilao River, Passi, Panay.		Jan. 13			G.
	Guimaras Id., vicinity of Buena Vista.	C. S. 4416; Dec., 1907.	Jan. 14			
	<i>Manila Bay.</i>					
	Mariveles Bay.	C. S. 4249; Apr., 1904.	Jan. 28	1.00 p. m.		S.
	Boca Chica (mouth of North Channel).	C. S. 4240; Feb., 1907.	Jan. 29	a. m.—p. m.		
	Pucot River (near Mariveles)	C. S. 4249; Apr., 1904.	do.	8.00 a. m.		
	Mariveles River.	do.	Jan. 30	1.00 p. m.		
	Mariveles Bay and Pucot River.	do.	do.	3.00 p. m.		
	Luzon Point.	C. S. 4240; Feb., 1907.	Jan. 31	7.30 a. m.		
	Mariveles wharf.	C. S. 4249; Apr., 1904.	Feb. 1	8.00 a. m.		
	Mariveles Bay (west).	do.	do.	2.00 p. m.		
	La Monja (Id.).	C. S. 4240; Feb., 1907.	Feb. 7	— a. m.		
	Luzon Pt.	do.	do.	— p. m.		
D. 5360	Corregidor Lt., N. 74° W., 6.9 miles (14° 21' N., 120° 41' E.).	do.	do.	7.25 p. m.	12	hrd.
	Limbones Cove.	do.	Feb. 8	— p. m.		
	do.	do.	do.	— p. m.		setrd. Co.
D. 5361	Corregidor Lt., S. 89° W., 7.2 miles (14° 24' 15" N., 120° 41' 30" E.).	do.	do.	8.48 p. m.	*12	
	<i>China Sea, off western Luzon.</i>					
D. 5362	Cape Santiago Lt., S. 35° E., 14.6 miles (13° 58' 20" N., 120° 30' 30" E.).	C. S. 4240; Feb., 1907.	Feb. 19	3.57 p. m.	*125	
	Pagapas Bay, Luzon.	do.	Feb. 20	8.00 a. m.		Co.
	Pagapas Bay, Santiago River	do.	do.	8.00 a. m.		M., G.
	<i>Balayan Bay, Luzon.</i>					
D. 5363	C. Santiago Lt., S. 79° W., 4.5 miles (13° 47' 20" N., 120° 43' 30" E.).	C. S. 4240; Feb., 1907.	Feb. 20	9.27 a. m.	*180	
D. 5364	C. Santiago Lt., S. 68° W., 5.4 miles (13° 48' 30" N., 120° 43' 45" E.).	do.	do.	2.40 p. m.	*160	
	Taal anchorage.	do.	do.	7.30 p. m.		
D. 5365	C. Santiago Lt., N. 73° W., 6.7 miles (13° 44' 24" N., 120° 45' 30" E.).	do.	Feb. 22	9.04 a. m.	*214	
	<i>Batangas Bay, Luzon.</i>					
D. 5366	Escareco Lt., S. 5° E., 7.7 miles (13° 39' N., 120° 58' 30" E.).	C. S. 4240; Feb., 1907.	Feb. 22	1.40 p. m.	*240	

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
					Luc. sdr. (a).....					Sounding wire carried away. Latitude and longitude approximate.
83	82				Luc. sdr. (a)..... 12' Agz. rev.....					Sounding wire lost. Longitude and latitude approximate.
					dyn.....	12-18 ft.				15 shots; 1 day's work.
					130' seine.....	20-30 ft.				11 hauls; all-day expedition.
					130' seine.....	4-10 ft.	4 00			10 hauls.
					4 trawl lines.....					Half of one trawl went adrift.
					25' and 130' seines; dyn.....			3.0		All-day expedition.
					25' seine; dyn.....					Half-day expedition.
					dyn.....		2 00			13 shots.
					3 trawl lines.....		10 00			
					dyn.....		2 00			3 shots.
					1 trawl line.....					
					cod trawls.....					
					cod trawls.....					
					hand line.....					
					25' Agz.....	botm...	1 00	N. 48° E...	1.3	
					cod trawls.....					
76	78				dyn.....	15-20 ft.	2 00			5 shots.
					25' Agz.....	botm...	9 08	N. 29° E...	12.0	
					3-bd. int. tr.....	60 fms.	12	N. 58° W...	1.0	
					dyn.....	15 ft.	6 00			8 shots.
					130' seine.....	4 ft.	3 00			5 hauls.
					25' Agz.....	botm...	1 15	N. 25° E...	3.0	
					25' Agz.....	botm...	43	N. 45° E...	2.8	
					dip; e. l.....	surface.	1 30			
					25' Agz.....	botm...	36	N. 10° W...	3.0	
80	79				3-bd. int. tr.....	150 fms.	20	N. 6° E...	2.5	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>Verde Island Passage.</i>						
D. 5367	Malabrigo Lt., N. 81° E., 8 miles (13° 34' 37" N., 121° 07' 30" E.).	C. S. 4240; Feb., 1907.	1909. Feb. 22	5.10 p. m.	fms. *180	S.*
<i>Marinduque Id. and vicinity.</i>						
.....	Port Banalacan, Marinduque	C. S. 4453; July, 1904.	Feb. 23	7.30 a. m.	Co., S.
D. 5368	Tayabas Lt. (outer), N. 32° W., 21.8 miles (13° 35' 30" N., 121° 48' E.).	C. S. 4714; June, 1906.	do....	2.08 p. m. 2.45 p. m.	181	gy. M.
.....	Capulaan Bay, Pagbilao, Chica Id.	do....	Feb. 24	7.00 a. m.	Co.
.....	Tayabas River (3 branches)	do....	do....	7.00 a. m.
D. 5369	Tayabas Lt. (outer), N. 50° W., 8.8 miles (13° 48' N., 121° 43' E.).	C. S. 4267; Aug., 1907.	do....	8.04 a. m. 8.30 a. m.	106	bk. S.
D. 5370	Tayabas Lt. (outer), N. 32° W., 11.6 miles (13° 44' 15" N., 121° 42' 30" E.).	C. S. 4714; June, 1906.	do....	9.35 a. m. 9.58 a. m.	159	sft. M.
D. 5371	Tayabas Lt. (outer), N. 43° W., 6 miles (13° 49' 40" N., 121° 40' 15" E.).	C. S. 4267; Aug., 1907.	do....	2.32 p. m.	*83	gn. M. (m. b.)
D. 5372	Tabayas Lt. (outer), N. 3° W., 4.5 miles (13° 49' 12" N., 121° 36' 09" E.).	do....	do....	3.42 p. m.	*150	gn. M. (m. b.)
.....	Tayabas Bay, Lucena anchorage.	do....	do....	8.00 p. m.
D. 5373	Tayabas Lt. (outer), N. 20° E., 15 miles (13° 40' N., 121° 31' 10" E.).	C. S. 4714; June, 1906.	Mar. 2	9.38 a. m. 10.15 a. m.	338	sft. M.
D. 5374	Tayabas Lt. (outer), N. 9° E., 7.4 miles (13° 46' 45" N., 121° 35' 08" E.).	do....	do....	11.57 a. m.	* 190	gy. M. (m. b.)
D. 5375	Tayabas Lt. (outer), N. 49° W., 18.2 miles (13° 42' 15" N., 121° 50' 15" E.).	do....	do....	3.05 p. m. 3.25 p. m.	107	gn. M.
D. 5376	Tayabas Lt. (outer), N. 53° W., 18.7 miles (13° 42' 50" N., 121° 51' 30" E.).	do....	do....	4.19 p. m.	*90	gy. M., S. (m. b.)
.....	Pitogo Anchorage, Luzon.	do....	Mar. 3	6.00 a. m.	Co.
.....	Mompog Id. (S.).	do....	do....	10.00 a. m.	Co., S.
D. 5377	Mompog Id. (E.), N. 55° W., 9 miles (13° 26' N., 122° 19' E.).	C. S. 4715; Apr., 1907.	Mar. 4	7.09 a. m. 8.03 a. m.	400	sft. gn. M.
D. 5378	Mompog Id. (E.), N. 38° W., 17 miles (13° 17' 45" N., 122° 22' E.).	do....	do....	10.02 a. m. 10.40 a. m.	395	sft. gn. M.
H. 4927	Mompog Id. (E.), N. 37° W., 25.6 miles (13° 10' 35" N., 122° 27' 30" E.).	do....	do....	1.06 p. m.	730
D. 5379	Mompog Id. (E.), N. 30° W., 37 miles (12° 59' 15" N., 122° 30' 40" E.).	do....	do....	2.46 p. m. 4.02 p. m.	920
D. 5380	Mompog Id. (E.), N. 31° W., 33 miles (13° 02' 45" N., 122° 29' E.).	do....	do....	7.26 p. m.
<i>Burias Id.</i>						
.....	Alimango Bay	C. S. 4715; Apr., 1907.	Mar. 5	8.00 a. m.	Co.
.....	Alimango River	do....	do....	9.00 a. m.	S., M.
<i>Ragay Gulf, Luzon.</i>						
.....	Alibijaban Id.	C. S. 4715; Apr., 1907.	Mar. 6	9.00 a. m.	Co.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Alr.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
83	80				25' Agz.....	botm...	<i>h. m.</i> 26	N. 63° E...	<i>mi.</i> 0.9	Rear beam broken and iron frame twisted.
					dyn.....	12-24 ft.				8 shots.
87	82				Luc. sdr. (a)... 12' Agz.; m. b.	botm...	37	N. 22° W...	6.0	
					dyn.....		4 00			10 shots.
					sml. seines; dyn. Tnr. sdr. (e)... 12' Agz.; m. b.	botm...	20	S. 9° W...	1.7	All-day expedition by 3 parties.
80	79	54.3			Luc. sdr. (a)... 12' Agz.; m. b.	botm...	20	S. 31° W...	3.3	
80	80				12' Agz.; m. b.	botm...	20	S. 31° W...	3.3	
83	80				12' Agz.; m. b.	botm...	22	S. 87° W...	.9	
82	81				12' Agz.; m. b.	botm...	21	S. 74° E...	1.5	
					dip; e. l.....	surface.	1 00			
82	80	51.8	1.02550		Luc. sdr. (a)... 12' Tnr.; m. b.	botm...	20	N. 32° E...	4.5	
81	80				12' Tnr.; m. b.	botm...	33	N. 29° E...	2.0	
82	80				Tnr. sdr. (e)... 12' Agz.; m. b.	botm...	20	N. 39° W...	1.5	
82	80				12' Agz.; m. b.	botm...	22	N. 11° W...	1.5	Net torn in two places near mouth.
					dyn.....	10-20ft.	1 00			1 shot.
					dyn.....	12-18ft.	6 00			15 shots.
79	80	49.6			Luc. sdr. (a)... 12' Agz.; m. b.	botm...	13	S. 31° E...	2.5	Net completely wrecked.
80	80	50.4			Luc. sdr. (a)... 12' Agz.; m. b.	botm...	20	S. 40° E...	3.5	Net wrecked; pieces recovered.
85	81	50.4			Luc. sdr. (a)...					
83	81	50.5	1.02443		Luc. sdr. (a)... 12' Agz.; m. b.	botm...	30	N. 43° W...	5.3	
82	81				int. 4.....					Net lost while veering out.
					dyn.....	12-24 ft.	9 00			20 shots.
					130' seine; dyn.	4ft.....	3 00			2 hauls, 5 shots.
					dyn.....	12-30ft.	5 00			20 hauls.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Ragay Gulf, Luzon—Cont'd.</i>					
D. 5381	Arena Pt. (Luzon), S. 68° W., 2.8 miles (13° 14' 15" N., 122° 44' 45" E.).	C. S. 4715; Apr., 1907.	1909. Mar. 6	9.15 a. m. 9.35 a. m.	<i>fms.</i> 88	co. S
D. 5382	Arena Pt. (Luzon), S. 55° W., 3.8 miles (13° 15' 20" N., 123° 45' 30" E.).do.....do.....	10.02 a. m. 10.23 a. m.	128	M.
	<i>Burias Id.</i>					
.....	Port Busin	C. S. 4454; May, 1906.	Mar. 6	8.00 p. m.
.....do.....do.....	Mar. 7	6.00 a. m.	Co.
.....do.....do.....	Mar. 8	6.00 a. m.	Co.
.....do.....do.....do.....	8.00 a. m.
D. 5383	Arena Pt. (Luzon), S. 66° W., 22 miles (13° 22' N., 123° 02' 30" E.).	C. S. 4715; Apr., 1907.do.....	3.08 p. m. 3.35 p. m.	127	gn. M.
D. 5384	Arena Pt. (Luzon), S. 64° W., 20.7 miles (13° 22' 15" N., 123° 01' 15" E.).do.....do.....	4.03 p. m. 4.32 p. m.	220
.....	Port Busin	C. S. 4454; May, 1906.do.....	7.00 p. m.
	<i>Ragay Gulf, Luzon.</i>					
.....	Refugio Id., Pasacao Anchorage.	C. S. 4454; May, 1906.	Mar. 9	8.00 a. m.	R.
D. 5385	Arena Pt. (Luzon), S. 61° W., 23.7 miles (13° 24' 50" N., 123° 03' 70" E.).	C. S. 4715; Apr., 1907.do.....	9.22 a. m. 9.54 a. m.	327	gy. M.
.....	Galvaney Id. (near Caima Bay).do.....do.....	3.00 p. m.	Co.
D. 5386	Arena Pt. (Luzon), S. 5° W., 25.3 miles (13° 38' 30" N., 122° 44' 30" E.).do.....do.....	3.25 p. m. 3.55 p. m.	287
.....	Ragay Bay (anchorage).do.....do.....	7.00 p. m.
.....	Ragay Riverdo.....	Mar. 10	7.30 a. m.	S.
.....	Ragay Baydo.....do.....	7.30 a. m.	Co., S.
	<i>Between Burias and Luzon.</i>					
.....	Canmahala Bay, Luzon	C. S. 4715; Apr., 1907.	Mar. 11	8.00 a. m.	Co., S.
D. 5387	Bagatao Id. Lt. (outer), S. 80° E., 27 miles (12° 54' 40" N., 123° 20' 30" E.).do.....do.....	1.06 p. m. 1.42 p. m.	209	soft gn. M.
D. 5388	Bagatao Id. Lt. (outer), S. 86° E., 21 miles (12° 51' 30" N., 123° 26' 15" E.).do.....do.....	2.51 p. m. 3.27 p. m.	226	soft gn. M.
.....	Bagatas Id. (anchorage)do.....do.....	7.15 p. m.
	<i>Between Ticao Id. and Luzon.</i>					
D. 5389	Bagatao Id. Lt. (outer), N. 3° W., 14 miles (12° 35' 45" N., 123° 48' 18" E.).	C. S. 4219; Dec., 1904.	Mar. 12	1.46 p. m.	*109-80	S.*
D. 5390	Bagatao Id. Lt. (outer), N. 12° W., 19 miles (12° 30' 54" N., 123° 51' 30" E.).do.....do.....	2.56 p. m.	*54	fne. S.*
	<i>Between Samar and Masbate.</i>					
.....	Escarpada Id., Bagacay Bay.	C. S. 4220; May, 1907.	Mar. 13	6.00 a. m.	Co., S.
.....	Destacado Id., Lode Baydo.....do.....	8.00 a. m.	R., Co.
D. 5391	Tubig Pt. (Destacado Id.), N. 31° E., 3 miles (12° 13' 15" N., 124° 05' 03" E.).do.....do.....	9.07 a. m.	*118
D. 5392	Tubig Pt., N. 49° E., 5 miles (12° 12' 35" N., 124° 02' 48" E.).do.....do.....	9.54 a. m. 10.10 a. m.	135	gn. M., S.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
82	80				Tnr. sdr. (e)... 12' Agz.; m. b.	botm...	15	N. 13° E..	1.3	
83	79				Tnr. sdr. (e)... 12' Agz.; m. b.	botm...	15	N. 18° E..	1.5	
					2 gill nets....	surface..				Hauled 6 a. m. on 8th.
					dyn.....	10-18ft.	2 30			4 shots.
					dyn.....	10-20ft.	5 00			9 shots.
					copper sul- phate.		2 00			Beach and tide pools.
84	80	62.5	1.02293		Luc. sdr. (a)... 12' Agz.; m. b.	botm...	20	N. 70° W..	1.3	
84	80	62.4			Luc. sdr. (a)... 12' Agz.; m. b.	botm...	25	N. 74° W..	2.7	
					dip; e. l.....	surface..	1 00			
					dyn.....	12-30ft.	4 00			12 shots.
82	78	62.4			Luc. sdr. (a)... 12' Agz.; m. b.	botm...	13	N. 47° W..	1.6	
					dyn.....	10-25ft.	2 00			7 shots.
83	82	62.4	1.02487		Luc. sdr. (a)... 12' Agz.; m. b.	botm...	8	N. 30° E..	1.3	Net badly torn.
					dip; e. l.....	surface..	1 00			
					16, 130 seines; dyn.	3-5 ft..	2 30			Half-day trip.
					dyn.....	4-20ft..	4 00			10 shots.
					dyn.....	4-30ft..	3 30			8 shots.
85	79	52.4	1.02503		Luc. sdr. (a)... 12' Agz.; m. b.	botm...	20	N. 44° E..	.8	
					K2.....	surface..	20	N. 44° E..	.8	
84	78	51.4			Luc. sdr. (a)... 12' Agz.; m. b.	botm...	26	N. 67° E..	1.5	
					K2.....	surface..	26	N. 67° E..	1.5	
					dip; e. l.....	surface..	45			
78	78				3-bd. int. tr...	40 - 55 fms.	17	N. 79° E..	1.6	
79	78				3-bd. int. tr...	50 fms..	26	N. 58° E..	1.5	
					dyn.....	5-30ft..	1 00			2 shots.
					dyn.....	18ft....	4 00			7 shots.
77	77				12' Agz.; m. b.	botm...	20	S. 88° W..	1.3	
					K2.....	10ft....	20	S. 88° W..	1.3	
78	77				Tnr. sdr. (e)... 12' Agz.; m. b.	botm...	5	S. 36° W..	.5	Net slightly torn.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>Between Samar and Masbate—Continued.</i>						
D. 5393	Panganalan Pt., Talajit Id., S. 59° E., 14.8 miles (12° 03' 30" N., 124° 03' 36" E.).	C. S. 4418; Apr., 1906.	1909. Mar. 13	1.44 p. m. 2.04 p. m.	<i>fms.</i> 136	hrd. S.
D. 5394	Panalangan Pt., Talajit Id., S. 68° E., 8.1 miles (12° 00' 30" N., 124° 05' 36" E.).	do.	do.	2.56 p. m. 3.13 p. m.	153	gn. M.
<i>Masbate Island.</i>						
.....	Port Cataingan.	C. S. 4418; Apr., 1906.	Mar. 14	9.00 a. m.		Co.
<i>Between Samar and Masbate.</i>						
.....	Buang B., Talajit Id.	C. S. 4418; Apr., 1906.	Mar. 15	8.00 a. m.		rky.
D. 5395	Panalangan Pt., Talajit Id., S. 81° E., 2.9 miles (11° 56' 40" N., 124° 14' E.).	do.	do.	8.38 a. m. 8.55 a. m.	140	hrd. gn. M. (m. b.).
D. 5396	Panalangan Pt., Talajit Id., S. 78° E., 4.5 miles (11° 57' N., 124° 12' 24" E.).	do.	do.	9.30 a. m. 9.45 a. m.	137	hrd. gn. M. (m. b.).
D. 5397	Panalangan Pt., Talajit Id., S. 78° E., 6 miles (11° 57' 27" N., 124° 10' 42" E.).	do.	do.	10.21 a. m. 10.36 a. m.	134	gn. M.
<i>Between Masbate and Leyte.</i>						
.....	Gigantangan Id. (west).	C. S. 4418; Apr., 1906.	Mar. 15	3.00 p. m.		limestone.
D. 5398	Gigantangan Id. (S.), S. 45° E., 2.7 miles (11° 35' 12" N., 124° 13' 48" E.).	do.	do.	3.03 p. m. 3.21 p. m.	114	gn. M.
<i>North of Cebu.</i>						
.....	Malapaseua Id. (west).	C. S. 4718; Dec., 1906.	Mar. 16	6.00 a. m.		R., Co.
D. 5399	Tanguingui Id. Lt., N. 70° W., 22.8 miles (11° 21' 45" N., 124° 05' E.).	do.	do.	8.54 a. m. 9.01 a. m.	32	S., Sh.
D. 5400	Tanguingui Id. Lt., N. 77° W., 22.5 miles (11° 24' 24" N., 124° 05' 30" E.).	do.	do.	9.34 a. m. 9.50 a. m.	25	S., Sh.
D. 5401	Tanguingui Id. Lt., N. 79° W., 23 miles (11° 24' 45" N., 124° 06' E.).	do.	do.	9.58 a. m. 10.05 a. m.	30	fne. S.
<i>Between Leyte and Cebu.</i>						
D. 5402	Capitancillo Id. Lt., S. 37° W., 16.1 miles (11° 11' 45" N., 124° 15' 45" E.).	C. S. 4718; Dec., 1906.	Mar. 16	1.54 p. m. 2.16 p. m.	188	gn. M.
.....	Calangaman Id. (north).	do.	do.	2.30 p. m.		setd. Co., R.
D. 5403	Capitancillo Id. Lt., S. 46° W., 15.7 miles (11° 10' N., 124° 17' 15" E.).	do.	do.	2.56 p. m. 3.14 p. m.	182	gn. M.
<i>Dupon Bay (Leyte) and vicinity.</i>						
.....	Sacaysacay Pt.	C. S. 4426; May, 1904.	Mar. 17	8.30 a. m.		Co.
.....	Guint River.	do.	do.	8.30 a. m.		
D. 5404	Ponson Id. (N.), S. 79° E., 6.8 miles (10° 50' N., 124° 26' 18" E.).	do.	do.	8.37 a. m. 8.58 a. m.	190	M.
D. 5405	Ponson Id. (N.), S. 86° E., 8.5 miles (10° 49' 26" N., 124° 24' 23" E.).	do.	do.	9.46 a. m. 10.09 a. m.	262	hrd.
D. 5406	Ponson Id. (N.), S. 88° E., 10.2 miles (10° 49' 03" N., 124° 22' 30" E.).	do.	do.	11.13 a. m. 11.41 a. m.	298	M.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Tempera- tures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Sur- face.	Bot- tom.		Depth.	Dura- tion.	Direction.	Dis- tance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
82	78				Tnr. sdr. (e) . . . 12' Agz.; m. b.	botm...	8	S. 11° W.	1.0	
80	78				Tnr. sdr. (e) . . . 12' Agz.; m. b.	botm...	9	S. 41° W.	1.1	
					dyn.....	12 ft ...	2 00			4 shots.
					dyn.....	18-30 ft.	4 00			15 shots.
79	78		1.02466		Luc. sdr. (e) . . . 12' Agz.; m. b.	botm...	19	N. 75° W.	1.2	
					K. 2.....	surface.	19	N. 75° W.	1.2	
79	79				Luc. sdr. (e) . . . 12' Agz.; m. b.	botm...	20	N. 66° W.	1.5	
					K. 2.....	surface.	20	N. 66° W.	1.5	
79	79				Luc. sdr. (e) . . . 12' Agz.; m. b.	botm...	16	N. 69° W.	1.2	
					K. 2.....	surface.	16	N. 69° W.	1.2	
					dyn.....	12-15 ft.	1 00			3 shots.
81	80				Tnr. sdr. (e) . . . 12' Agz.; m. b.	botm...	7	N. 49° W.	.5	
					dyn.....	10-20 ft.	3 30			14 shots.
79	79				Tnr. sdr. (e) . . . 6' McC.....	botm...	9	N. 22° E.	.5	
80	80		1.02458		Tnr. sdr. (e) . . . 6' McC.....	botm...	12	N. 10° E.	.4	
80	80				Tnr. sdr. (e) . . . 6' McC.....	botm...	27	N. 61° E.	.9	
81	81	55.8			Luc. sdr. (a) . . . 12' Agz.; m. b.	botm...	22	S. 45° E.	1.9	
					K. 2.....	surface.	22	S. 45° E.	1.9	
					dyn.....	8-25 ft.	2 00			7 shots.
81	81	55.7			Luc. sdr. (a) . . . 12' Agz.....	botm...	29	S. 55° E.	1.8	
					dyn.....	12-30 ft.	7 30			16 shots.
					dyn.....		7 00			
81	78	55.4			Luc. sdr. (a) . . . 12' Agz.....	botm...	26	S. 74° W.	1.8	
82	80				Luc. sdr. (e) . . . 12' Agz.....	botm...	20	S. 82° W.	1.9	
83	81				Luc. sdr. (e) . . . 12' Agz.....	botm...	27	N. 81° W.	2.0	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>Dupon Bay (Leyte) and vicinity—Continued.</i>						
D. 5407	Ponson Id. (N.), S. 76° E., 12.2 miles (10° 51' 38" N., 124° 20' 54" E.).	C. S. 4426; May, 1904.	1909. Mar. 17	12.57 p. m. 1.28 p. m.	fms. 350	gn. M.
.....	Anchorage, Dupon Bay.....do.....do.....	7.00 p. m.
<i>Between Cebu and Leyte.</i>						
D. 5408	Capitancillo Lt., N. 25° W., 20.8 miles (10° 40' 15" N., 124° 15' E.).	C. S. 4718; Dec., 1906.	Mar. 18	8.05 a. m. 8.23 a. m.	159	gn. M.
D. 5409	Capitancillo Lt., N. 19° W., 22 miles (10° 38' N., 124° 13' 08" E.).do.....do.....	9.16 a. m. 9.51 a. m.	189	gn. M.
D. 5410	Bagacay Pt. Lt., S. 37° W., 7.2 miles (10° 28' 45" N., 124° 05' 30" E.).do.....do.....	11.21 a. m. 11.56 a. m.	385	gn. M.
<i>Between Cebu and Bohol.</i>						
D. 5411	Lauis Pt. Lt., N. 35° E., 4.7 miles (10° 30' 31" N., 123° 51' 15" E.).	C. S. 4718; Dec., 1906.	Mar. 23	8.18 a. m. 8.48 a. m.	145	gn. M.
D. 5412	Lauis Pt. Lt., N. 21° E., 5.5 miles (10° 09' 15" N., 123° 52' E.).do.....do.....	9.36 a. m. 9.58 a. m.	162	gn. M.
.....	Pandanon Id. (south).....do.....do.....	2.30 p. m.	Co., S.
.....do.....do.....do.....	2.30 p. m.	S., Co.
.....	Reef opposite Pandanon Id.do.....	Mar. 24	7.30 a. m.	Co., S.
D. 5413	Lauis Pt. Lt., N. 68° W., 10 miles (10° 10' 35" N., 124° 03' 15" E.).do.....do.....	11.34 a. m.	* 42
D. 5414	Lauis Pt. Lt., N. 67° W., 9.5 miles (10° 10' 40" N., 124° 02' 45" E.).do.....do.....	12.04 p. m.
D. 5415	Lauis Pt. Lt., N. 24° W., 7.2 miles (10° 07' 50" N., 123° 57' E.).do.....do.....	1.21 p. m. 1.41 p. m.	88	fne. S.
D. 5416	Lauis Pt. Lt., N. 12° E., 2.9 miles (10° 11' 30" N., 123° 53' 30" E.).do.....	Mar. 25	7.20 a. m. 7.43 a. m.	150	gn. M.
D. 5417	Lauis Pt. Lt., N. 10° E., 3.5 miles (10° 10' N., 123° 53' 15" E.).do.....do.....	8.18 a. m. 8.40 a. m.	165	gy. M., S.
D. 5418	Lauis Pt. Lt., N. 16° E., 5.6 miles (10° 08' 50" N., 123° 52' 30" E.).do.....do.....	9.28 a. m. 9.48 a. m.	159	gy. M., S.
D. 5419	Lauis Pt. Lt., N. 27° E., 17.8 miles (9° 58' 30" N., 123° 46' E.).do.....do.....	1.35 p. m. 1.55 p. m.	175	gn. M.
D. 5420	Cruz Pt. (Bohol), S. 20° E., 6 miles (9° 49' 35" N., 123° 45' E.).do.....do.....	3.33 p. m. 3.48 p. m.	127
<i>Bohol Island.</i>						
.....	Maribojoc Bay (anchorage).....	C. S. 4718; Dec., 1906.	Mar. 24	7.30 p. m.
.....	Maribojoc Bay (E. of Cruz Pt.)do.....	Mar. 26	6.00 a. m.	Co., R.
<i>Between Panay and Guimaras.</i>						
D. 5421	Lusaran Pt. Lt., S. 27° E., 5 miles (10° 33' 30" N., 122° 26' E.).	C. S. 4718; Dec., 1906.	Mar. 30	5.38 p. m. 6.10 p. m.	137	gn. M.
D. 5422	Lusaran Pt. Lt., S. 80° E., 9.7 miles (10° 31' N., 122° 18' 45" E.).do.....do.....	7.17 p. m.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
82	81				Luc. sdr. (e). 12' Agz.	botm...	20	S. 49° E. . .	1.6	
					dip; e. l.; dyn. caps.	surface.	3 00			2 shots.
83	80	55.4	1.02462		Luc. sdr. (a). 12' Agz.; m. b. K. 2.	botm... surface.	20 20	S. 46° W. . . S. 46° W. . .	1.3 1.3	
81	80				Luc. sdr. (e). 12' Agz.; m. b. K.	botm... surface.	29 29	S. 51° W. . . S. 51° W. . .	2.0 2.0	Record incomplete.
82	80				Luc. sdr. (e). 12' Agz.; m. b.	botm...	14	S. 3° W. . .	1.2	
80	81	55.2			Luc. sdr. (e). 12' Agz.; m. b. K. 2.	botm... surface.	24 24	S. 33° W. . . S. 33° E. . .	1.7 1.7	
81	81	54.8			Luc. sdr. (e). 12' Agz.	botm...	22	S. 67° E. . .	1.7	
					dyn.	6-12 ft.	2 30			4 shots.
					130' seine.	5 ft	5 30			11 hauls.
					dyn.	10-12 ft.	1 00			3 shots.
82	82				6' McC.	botm...	6	N. 30° W. . .	.6	
82	82				6' McC.	botm...	9	N. 23° W. . .	1.2	
83	81	62.4			Luc. sdr. (a). 12' Agz.; m. b.	botm...	19	N. 81° W. . .	1.5	
81	80	54.4			Luc. sdr. (a). 12' Agz.	botm...	20	South.	1.5	
81	80	54.4			Luc. sdr. (a). 12' Agz.	botm...	20	S. 18° W. . .	1.2	
81	81	54.4			Luc. sdr. (a). 12' Agz.	botm...	20	S. 82° W. . .	.8	
83	81	54.5			Luc. sdr. (a). 12' Agz.	botm...	20	S. 74° W. . .	1.3	
83	81	59			Luc. sdr. (a). 12' Agz.; m. b.	botm...	17	S. 54° W. . .	1.2	
					dip; e. l.	surface.	1 30			
					dyn.	10-20 ft.	2 00			6 shots.
84	82	58.4			Luc. sdr. (a). 12' Agz.; m. b.	botm...	19	S. 70° W. . .	1.5	
84	82				int. 3.	surface.	20	W. by S. . .	1.5	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Jolo Sea.</i>					
.....	Cagayan Id., Cagayanes Ids. (NW.).	C. S. 4717; Feb., 1903.	1909. Mar. 31	9.00 a. m.	<i>fms.</i>	mg. n. Rf.
D. 5423	Cagayan Id. (S.), S. 11° E., 4.8 miles (9° 38' 30" N., 121° 11' E.).	do.....	9.16 a. m. 9.55 a. m.	508	gy. M., co. S.
D. 5424	Cagayan Id. (S.), S. 11° W., 3.4 miles (9° 37' 05" N., 121° 12' 37" E.).	do.....	12.52 p. m. 1.24 p. m.	340	co. S.
D. 5425	Cagayan Id. (S.), S. 14° E., 4 miles (9° 37' 45" N., 121° 11' E.).	do.....	2.20 p. m. 2.57 p. m.	495	gy. M., co. S.
	<i>Eastern Palawan and vicinity.</i>					
.....	Mantaquin Bay (Palawan)..	C. S. 4716; Feb., 1903.	Apr. 1	3.00 p. m.	S.
.....	Rasa Id. (southwest).....	do.....	3.00 p. m.	sft. Co., R
.....	Malinao River (Palawan)....	Apr. 2	8.00 a. m.	Co.
.....	Rasa Id. (southwest).....	do.....	9.00 a. m.	S., G.
D. 5426	Mantaquin Bay.....	do.....	2.30 p. m.	fne. gy. S.
.....	30th of June Id., N. 29° E., 12.2 miles (9° 12' N., 118° 28' E.).	Apr. 3	6.42 a. m. 6.44 a. m.	27
D. 5427	30th of June Id., N. 16° W., 11.5 miles (9° 11' 30" N., 118° 37' 08" E.).	do.....	8.04 a. m. 8.09 a. m.	37	S, Sh.
D. 5428	30th of June Id., N. 62° W., 19.5 miles (9° 13' N., 118° 51' 15" E.).	do.....	10.14 a. m. 11.23 a. m.	1,105	gy. M.
H. 4928	Fondeado Id. (SE.), N. 29° E., 23 miles (9° 34' 48" N., 118° 45' E.).	do.....	3.28 p. m.	902	gy. M., fne. co. S.
H. 4929	Fondeado Id. (SE.), N. 19° E., 19 miles (9° 37' 30" N., 118° 48' 30" E.).	do.....	4.39 p. m.	554	gy. M.
.....	Iwahig River and tributaries (Pta. Princesa).	C. S. 4343; July, 1903.	Apr. 4	7.00 a. m.
.....	Puerta Princesa (west of Bancaobancaon Pt.).	Apr. 5	6.30 a. m.	S., R., Co.
D. 5429	Fondeado Id. (SE.), N. 18° E., 15 miles (9° 41' 30" N., 118° 50' 22" E.).	C. S. 4716; Feb., 1903.	do.....	7.32 a. m. 8.14 a. m.	766	gn. M.
.....	Machesi Id. (southwest).....	do.....	1.00 p. m.	S., M., Co.
D. 5430	Fondeado Ids. (W.), N. 57° W., 10.5 miles (9° 49' 40" N., 119° 03' 20" E.).	Apr. 6	10.07 a. m. 10.54 a. m.	464	glob. Oz.
.....	Verde del Sur Id. (south)....	do.....	2.00 p. m.	Co., G., S.
.....	do.....	do.....	2.00 p. m.	S.
.....	do.....	do.....	8.00 p. m.
.....	Port Langean, Dumarán Id. (east).	Apr. 7	4.00 p. m.	R., Co.
.....	do.....	do.....	5.30 p. m.
.....	Port Langean, Dumarán Id. (anch.).	do.....	7.30 p. m.
.....	Port Langean, Dumarán Id. (Green Pt.).	Apr. 8	7.00 a. m.	S., Co., G.
.....	Wreck Bay, Dalaganem Id..	C. S. 4717; Feb., 1903.	do.....	2.30 p. m.	R., S., Co.
D. 5431	Corandagos Id. (NW.), N. 28° E., 4.8 miles (10° 38' 45" N., 120° 12' 45" E.).	do.....	2.49 p. m. 2.54 p. m.	51	S.
D. 5432	Corandagos Id. (NW.), N. 30° E., 5.7 miles (10° 37' 50" N., 120° 12' E.).	do.....	3.26 p. m. 3.34 p. m.	51	S.
D. 5433	Corandagos Id. (NW.), N. 35° E., 6.5 miles (10° 37' 30" N., 120° 11' 05" E.).	do.....	4.04 p. m. 4.16 p. m.	54	gn. M., co. S.
D. 5434	Corandagos Id. (N.), S. 63° W., 7.6 miles (10° 46' 45" N., 120° 22' 45" E.).	do.....	7.50 p. m.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Tempera- tures.			Density.		Apparatus.	Tria].		Drift.		Remarks.
Air.	Surface.	Bottom.	Sur- face.	Bot- tom.		Depth.	Dura- tion.	Direction.	Dis- tance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
		49.8			dyn.....	2-50ft..	3 00			6 shots.
82	82				Luc. sdr. (a)...					
		50.4			12' Agz.; m. b. .	botm...	27	N. W.	1.5	
81	82				Luc. sdr. (a)...					
		49.4			12' Agz.; m. b. .	botm...	20	N. 67° W.	1.5	
82	83				Luc. sdr. (a)...					
					12' Agz.; m. b. .	botm...	20	N. 62° W.	1.2	
					130' seine.....	4 ft	2 00			6 hauls.
					dyn.....	6-12ft..	2 00			8 shots.
					dyn.....	4 00			5.0	
					dyn.....	8-10 ft..	3 00			4 shots.
					500' seine.....	10 ft..	2 30			3 hauls.
81	82				Tnr. sdr. (e)...					
					6' McC.	botm...	9	N. 20° E.	.3	
81	82				Tnr. sdr. (e)...					
					6' McC.	botm...	5			Net lost.
		49.7			Luc. sdr. (a)...					
85	83				12' Agz.; m. b. .	botm...	21	N. by W.	1.0	
86	83	49.4			Luc. sdr. (a)...					
83	82	49.4			Luc. sdr. (a)...					
					dyn.....		12 00			
					dyn.....	4-20ft..	2 30			6 shots.
82	83				Luc. sdr. (a)...					
					12' Agz.; m. b. .	botm...	18	N. 73° W.	1.9	
		50			dyn.....	6-12ft..	4 00			10 shots.
84	83				Luc. sdr. (a)...					
					12' Agz.	botm...	25	N	1.5	
					K. 2.	surface	25	N	1.5	
					dyn.....	8-10ft..	3 00			6 shots.
					130' seine.....	2-4 ft..	3 00			20 hauls.
					gill nets.....		12 00			2 lines.
					dyn.....	6-15ft..	1 30			5 shots.
					gill nets.....		12 00			2 lines.
					dip; e. l.	surface.	20			
					dyn.....	8 ft	4 30			17 shots.
					dyn.....	12-18ft.	3 00			6 shots.
					Tnr. sdr. (e)...					
84	83				6' McC.	botm...	20	S. 46° W.	.8	
84	83				Tnr. sdr. (e)...					
					6' McC.	botm...	20	S. 68° W.	1.3	
					Tnr. sdr. (e)...					
83	83				6' McC.	botm...	20	S. 44° W.	1.2	
83	83				int. 3.	surface.	20	N. 70° E.	.2	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Cuyos Islands.</i>					
	Cuyo Id. (west).....	C. S. 4345; Feb., 1905.	1909, Apr. 9	8.30 a. m.	fms.	R., Co.....
	do.....	do.....	do.....	8.30 a. m.		S.....
D. 5435	Bisucay Id. (northeast).....	do.....	do.....	2.00 p. m.		Co., R.....
	Bisucay Id. (NE.), S. 55° E., 1 mile (10° 50' N., 120° 58' 10" E.).	C. S. 4717; Feb., 1903.	do.....	7.50 p. m.		
	<i>West coast of Luzon, Manila Bay to Lingayen Gulf.</i>					
D. 5436	Corregidor Lt., N. 83° E., 5.2 miles (14° 22' 37" N., 120° 29' E.).	C. S. 4240; Feb., 1907.	May 7	7.03 p. m.	*32	
	Hermana Mayor Id. (west).....	C. S. 4712; Sept., 1904.	May 8	9.00 a. m.		S., Co.....
	Caiman Cove.....	C. S. 4210; Sept., 1907.	do.....	3.30 p. m.		S., Co., R.....
	do.....	do.....	do.....	7.00 p. m.		M.....
D. 5437	Hermana Mayor Lt., N. 69° E., 4.9 miles (15° 45' 54" N., 119° 42' 45" E.).	do.....	do.....	10.27 a. m.		
				12.07 p. m.		
D. 5438	Hermana Mayor Lt., S. 21° E., 7.5 miles (15° 54' 42" N., 119° 44' 42" E.).	do.....	do.....	3.50 p. m.	297	gn. M.....
				4.20 p. m.		
D. 5439	Caiman Cove.....	do.....	May 9	6.00 a. m.		S., Co.....
	Hermana Mayor Lt., S. 33° E., 12.6 miles (15° 58' 15" N., 119° 40' 20" E.).	do.....	do.....	9.44 a. m.	940	gn. M.....
				10.49 a. m.		
	Bolinao Bay (north of Boli- nao).	C. S. 4238; Feb., 1905.	do.....	8.00 p. m.		
	Bolinao Bay (east of village).	do.....	May 10	6.00 a. m.		S., Co., R.....
	do.....	do.....	do.....	8.00 a. m.		S.....
D. 5440	S. Fernando Pt. Lt., N. 82° E., 23.1 miles (16° 33' 52" N., 119° 52' 54" E.).	C. S. 4209; Oct., 1905.	do.....	1.35 p. m.	172	fine gy. S., Glob.....
				2.01 p. m.		
D. 5441	S. Fernando Pt. Lt., S. 87° E., 18.7 miles (16° 38' N., 119° 57' 18" E.).	do.....	do.....	3.20 p. m.	186	
				3.47 p. m.		
D. 5442	S. Fernando Pt. Lt., N. 39° E., 8.4 miles (16° 30' 36" N., 120° 11' 06" E.).	do.....	do.....	6.48 p. m.	45	co. S.....
				6.58 p. m.		
	Lingayen G. (east of Pt. Guecet).	do.....	May 11	10.00 a. m.		S.....
	<i>East coast of Luzon, San Bernardino Strait to San Miguel Bay.</i>					
	Matnog Bay.....	C. S. 4258; Jan., 1903.	May 31	2.00 p. m.		Co., R.....
	do.....	do.....	do.....	2.00 p. m.		S., Co.....
	do.....	do.....	do.....	6.00 p. m.		
	Balicuatro Ids., Biri Chan- nel (southern Biri Id.).	C. S. 4220; May, 1907.	June 1	8.00 a. m.		mgn. Rf.....
				7.00 p. m.		
	Batag Id. (west, near Leung Pt.).	C. S. 4449; Jan., 1907.	June 2	6.00 a. m.		mgn. Rf.....
				4.00 p. m.		Co., co. R.....
				5.00 p. m.		
D. 5443	Atalaya Pt., Batag Id., S. 64° E., 3.6 miles (12° 43' 05" N., 125° 01' E.).	do.....	June 3	8.00 a. m.	241	Co.....
				8.50 a. m.		co. S., Sh.....
				9.19 a. m.		
D. 5444	Atalaya Pt., Batag Id., S. 65° E., 5.1 miles (12° 43' 51" N., 124° 58' 50" E.).	do.....	do.....	9.57 a. m.	308	gn. M.....
				10.32 a. m.		
D. 5445	Atalaya Pt., Batag Id., S. 56° E., 5.3 miles (12° 44' 42" N., 124° 59' 50" E.).	do.....	do.....	11.25 a. m.	383	gn. M., S.....
				12.01 p. m.		

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
					dyn.....	4-16 ft..	3 0			7 shots.
					130' seine.....	3-4 ft..	2 30			10 hauls.
83	83				dyn.....	6-18 ft..	3 00			9 shots.
					int. 3.....	surface.	21	W. x N.	0.7	
85	86				int. 4.....	surface.	15	W.....	.5	
					dyn.....	8-10 ft..	5 00			5 shots.
					dyn.....	5-12 ft..	2 00			4 shots.
					2 gill nets.....	9 fms..	11 00			
88	86				6 K. 6.....	100-600 fms.	36	N. 61° W.	.9	
87	86				Int. 4 §.....	450 fms.	27 22			
87	87	46.2			Luc. sdr. (a)..					
					12' Agz.; m. b.	botm...	21	S. 5° E...	1.2	
					dyn.....	10-12 ft..	2 00			8 shots.
89	87	36.7			Luc. sdr. (a)..					
					12' Agz.; m. b.	botm...	14	N. 16° W.	2.5	Net slightly torn.
					dip; e. l.....	surface.	1 00			
					dyn.....	10-12 ft..	4 00			7 shots.
					130' seine.....	4 ft.....	3 00			5 hauls.
86	87	53.2			Luc. sdr. (a)..					
					12' Agz.; m. b.	botm...	20	N. 22° E..	1.8	
86	87	52.2			Luc. sdr. (a)..					
					25' Agz.....	botm...	20	N. 64° E..	1.8	
82	85				Tnr. sdr. (e)..					
					25' Agz.....	botm...	10 34	S. 12° E..	15.5	
					500' seine.....	4-12 ft..	4 30			5 hauls.
					dyn.....	10-12 ft..	3 00			5 shots.
					430' seine.....	5 ft.....	2 30			3 hauls.
					3 gill nets.....	12 00	12 00			
					dyn.....	12-24 ft..	7 00			13 shots.
					3 gill nets.....	12 00	12 00			
					dyn.....	10-12 ft..	3 00			7 shots.
					dyn.....	6-15 ft..	1 30			6 shots.
					3 gill nets.....	13 00	13 00			
					dyn.....	5-15 ft..	7 30			17 shots.
82	83	51.3			Luc. sdr. (a)..					
					12' Agz.....	botm...	20	N. 70° W.	1.9	
85	83	45.3			Luc. sdr. (a)..					
					12' Agz.....	botm...	17	N. 65° E..	1.1	
85	83	44.3			Luc. sdr. (a)..					
					12' Agz.....	botm...	37	S. 73° E..	1.5	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>East coast of Luzon, San Bernardino Strait to San Miguel Bay—Continued.</i>					
D. 5446	Atalaya Pt., Batag Id., S. 64° E., 5.3 miles (12° 43' 51" N., 124° 59' 18" E.).	C. S. 4449; Jan., 1907.	1909. June 3	1.25 p. m. 1.58 p. m.	<i>fms.</i> 300	gn. M.
D. 5447	S. Miguel Pt., S. 7° W., 3.5 miles (13° 28' N., 123° 46' 18" E.).	C. S. 4221; June, 1905.	June 4	5.37 a. m. 6.14 a. m.	310	gn. M.
	Tabaco Bay (west of S. Miguel Pt.).	C. S. 4237; Mar., 1905.	do	8.00 a. m.		co. S.
D. 5448	S. Miguel Pt., N. 23° E., 1.5 miles (13° 23' 10" N., 123° 45' 19" E.).	do	do	8.55 a. m.	*47	
	Batan Id. (north, west of Camisog Pt.).	C. S. 4259; Aug., 1906.	do	1.00 p. m.		S., Co.
D. 5449	East Pt. (Batan Id.), S. 43° E., 7.9 miles (13° 21' 36" N., 124° 00' 30" E.).	C. S. 4221; June, 1905.	do	2.38 p. m.	*300	
D. 5450	East Pt. (Batan Id.), S. 36° E., 9.2 miles (13° 23' 15" N., 124° 00' 30" E.).	do	do	3.19 p. m. 3.52 p. m.	408	gn. M., Co.
D. 5451	East Pt. (Batan Id.), S. 38° E., 8.2 miles (13° 22' 22" N., 124° 00' 48" E.).	do	June 5	7.34 a. m.	*380	
	Batan Id. (southwest, of Batan).	C. S. 4259; Aug., 1906.	do	8.00 a. m. 1.00 p. m.		S., Co. tide pools.
	Rapurapu Id. (Bahayon Pt.).	do	do	1.00 p. m.		Co.
	Albay G., Yaua River.	C. S. 4237; Mar., 1905.	June 7	6.00 a. m.		
D. 5452	Legaspi Lt., S. 38° W., 3 miles (13° 11' 54" N., 123° 47' 10" E.).	C. S. 4221; June, 1905.	do	8.51 a. m.	*110	
D. 5453	Legaspi Lt., S. 58° W., 4.5 miles (13° 12' N., 123° 49' 18" E.).	do	do	9.44 a. m.	*146	
D. 5454	Legaspi Lt., S. 64° W., 5.7 miles (13° 12' N., 123° 50' 30" E.).	do	do	10.46 a. m.	*153	
D. 5455	Legaspi Lt., S. 70° W., 6.7 miles (13° 11' 51" N., 123° 51' 42" E.).	do	do	11.57 a. m.	*165	
D. 5456	Legaspi Lt., S. 76° W., 6.7 miles (13° 11' 10" N., 123° 51' 52" E.).	do	do	12.55 p. m.	*142	
D. 5457	Legaspi Lt., S. 60° W., 5 miles (13° 12' N., 123° 49' 40" E.).	do	June 8	9.40 a. m.	*146	
	Batan Id., Caracaran Bay.	C. S. 4259; Aug., 1906.	do	1.00 p. m.		S., Co.
D. 5458	Legaspi Lt., S. 84° W., 14 miles (13° 10' 54" N., 123° 59' 38" E.).	C. S. 4221; June, 1905.	do	2.04 p. m.	*200	
D. 5459	Legaspi Lt., S. 88° W., 14.3 miles (13° 10' 21" N., 123° 59' 54" E.).	do	do	3.41 p. m.	*201	
	Catanduanes Id., Cabugao Bay (east).	C. S. 4269; Feb., 1909.	June 9	9.00 a. m.		R., Co., grass.
	Catanduanes Id., Cabugao River.	do	do	9.00 a. m.		
	Catanduanes Id., Cabugao Bay.	do	do	7.00 p. m.		
	Catanduanes Id., Agojo Pt.	C. S. 4222; Jan., 1909.	June 10	8.30 a. m.		co. S.
D. 5460	Sialat Pt. Lt., N. 24° E., 8.2 miles (13° 32' 30" N., 123° 58' 06" E.).	do	do	8.37 a. m.	565	gy. M.
	Palumbanes Ids., Porongpong Id. (southwest).	do	do	9.22 a. m. 3.00 p. m.		S., Co.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.					<i>h. m.</i>		<i>mi.</i>	
84	83				Luc. sdr. (a)...	botm...	28	S. 83° E...	1.6	Therm. failed to register.
		45.3			12' Agz.	botm...	21	N. 64° E ..	1.5	
					dyn.	10-15 ft.	3 00			7 shots.
86	86				12' Agz.; m. b.	botm...	21	S. 64° E...	.8	
					dyn.	8-10 ft.	4 30			6 shots.
85	86				12' Agz.; m. b.	botm...	21	N.	1.4	
		42.3			Luc. sdr. (a)...	botm...	28	N.	1.9	
85	86				12' Agz.; m. b.	botm...	28	N.	1.9	
79	84				int. 5 §.....	280 fms.	21 12	S. 61° E...	1.0	
					dyn.	10 ft.	8 00			10 shots.
					copper sulph'te.		2 00			
					dyn.	8-12 ft.	4 30			4 shots.
					25' seine; dyn.		9 30			
85	85				12' Agz.	botm...	14	N. 48° E...	1.0	
85	86				12' Agz.	botm...	20	E.	1.1	
86	86				12' Agz.	botm...	21	S. 79° E...	1.2	
86	86				12' Agz.	botm...	14	S. 63° E...	1.1	
87	86				int. 4 §.....	120 fms.	19 7	N. 88° W..	1.3	
85	85				12' Agz.	botm...	20	S. 72° E...	1.4	
					dyn.	6-10 ft.	3 30			13 shots.
87	85				12' Agz.	botm...	23	S. 56° E...	.6	
85	85				12' Agz.; m. b.	botm...	20	N. 86° W..	.8	
					dyn.	10-18 ft.	2 30			6 shots.
					dyn.; 25' seine.		8 00			
					dip; e. l.	surface.	1 30			
					dyn.	12 ft.	2 30			13 shots.
					Luc. sdr. (a)...					Therm. failed to register.
86	85				12' Agz.; m. b.	botm...	14	N. 43° W.	2.0	
					dyn.	8-20 ft.	2 30			5 shots.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>East coast of Luzon, San Bernardino Strait to San Miguel Bay—Continued.</i>					
			1909.		<i>fms.</i>	
	Palumbanes Ids., "West Id." (west).	C. S. 4222; Jan., 1909.	June 11	7.00 a. m.		co. R
	Lahuy Id., Pocket Bay (west).	do.	do.	1.00 p. m.		co. S
	Quinalasag Id., Masamat Bay.	do.	do.	7.00 p. m.		
	Quinalasag Id., Masamat Bay (east).	do.	June 12	6.00 a. m.		S., Co.
	Butaanan Id. (west and south).	C. S. 4223; June, 1908.	do.	2.30 p. m.		S., Co.
	Butaanan Id. (south).	do.	June 13	6.30 a. m.		Co., S
	Maculabo Id. (west).	C. S. 4715; Apr., 1907.	do.	3.30 p. m.		Co.
	do.	do.	June 14	7.30 p. m.		Co.
				6.30 a. m.		tide pools
				9.00 a. m.		
	S. Miguel Bay, Colasi Pt.	C. S. 4223; June, 1908.	do.	7.00 p. m.		
D. 5461	Carango Id. (W.), N. 12° W., 4.9 miles (13° 57' 42" N., 123° 06' 42" E.).	do.	do.	7.10 p. m.	11	
	Canimo Pass, Daet Pt.	do.	June 15	9.00 a. m.		Co., S.
	Canimo Pass, Basut River.	do.	do.	9.00 a. m.		
D. 5462	Sialat Lt., S. 80° E., 5 miles (13° 40' 42" N., 123° 56' 30" E.).	C. S. 4222; Jan., 1909.	June 16	5.50 a. m.	469	gy. M.
				6.44 a. m.		
	Lagonoy G., Palag Bay (east)	do.	do.	9.00 a. m.		Co., R.
D. 5463	Sialat Pt. Lt., S. 74° E., 3.9 miles (13° 40' 57" N., 123° 57' 45" E.).	do.	do.	10.28 a. m.	*300	S.*
D. 5464	Sialat Pt. Lt., N. 82° E., 4.1 miles (13° 39' 15" N., 123° 57' 15" E.).	do.	do.	2.14 p. m.	*400	
	Lagonoy G., Alto Pt. anch.	do.	do.	7.30 p. m.		
	Lagonoy G., Rosa Id.	do.	do.	7.30 a. m.		S., Co.
	Lagonoy G., Bato River.	do.	June 17	7.30 a. m.		
D. 5465	Atulayan Id. (E.), S. 50° W., 7.3 miles (13° 39' 42" N., 123° 40' 39" E.).	do.	do.	8.39 a. m.	*500	gy. M. (m. b.).
D. 5466	Atulayan Id. (E.), S. 62° W., 7.7 miles (13° 38' 36" N., 123° 41' 45" E.).	do.	do.	10.40 a. m.	*540	gy. M. (m. b.).
	Lagonoy G., Atulayan Bay (south).	do.	do.	3.00 p. m.		S., R.
	Lagonoy G., Atulayan Bay (west).	do.	do.	6.30 p. m.		
	Lagonoy G., Atulayan Bay (anch.).	do.	do.	8.00 p. m.		
	Lagonoy G., Nato River.	do.	June 18	6.30 a. m.		
	Lagonoy G., Atulayan Id. (east).	do.	do.	7.00 a. m.		Co., S.
D. 5467	Atulayan Id. (S.), S. 79° W., 2.5 miles (13° 35' 27" N., 123° 37' 18" E.).	do.	do.	7.52 a. m.	*480	gy. M. (m. b.).
D. 5468	Atulayan Id. (S.), S. 83° W., 5.7 miles (13° 35' 39" N., 123° 40' 28" E.).	do.	do.	9.58 a. m.	*569	gn. M. (m. b.).
D. 5469	Atulayan Id. (E.), S. 63° W., 4 miles (13° 36' 48" N., 123° 38' 24" E.).	do.	do.	1.29 p. m.	*500	gn. M. (net).
D. 5470	Atulayan Id. (E.), S. 68° W., 6.7 miles (13° 37' 30" N., 123° 41' 09" E.).	do.	do.	3.26 p. m.	*560	M.*
	Lagonoy G., Nato anch.	do.	do.	7.30 p. m.		
D. 5471	Sialat Pt. Lt., N. 71° E., 15 miles (13° 34' 57" N., 123° 47' 06" E.).	do.	June 19	9.17 a. m.	*568	
D. 5472	Sialat Pt. Lt., N. 63° E., 13.6 miles (13° 33' 36" N., 123° 49' E.).	do.	do.	11.12 a. m.	*550	

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>			<i>mi.</i>	
					dyn.....	8-10 ft..	2 00			3 shots.
					dyn.....	12-15 ft.	3 00			2 shots.
					dip; e. l.....	surface.	1 00			
					dyn.....	10 ft....	3 30			10 shots.
					dyn.....	8 ft....	3 00			9 shots.
					dyn.....	10 ft....	4 30			11 shots.
					dyn.....	15-25 ft.	1 30			7 shots.
					dip; e. l.....	surface.	1 30			
					dyn.....	8-18 ft..	4 30			11 shots.
					copper sulphate.		2 00			
					4 gill nets.....		12 00			
84	86				25' Agz.....	botm...	17	E.....	2.5	
					dyn.....	5-10 ft..	2 45			5 shots.
		41.3			small seines.....		10 00		6.0	
					Luc. sdr. (a).....					
83	85				25' Agz.....	botm...	17	S. 35° E..	1.5	Bridle stops and one preventer carried away.
					dyn.....	8-25 ft..	5 30			24 shots.
83	84				12' Agz.; m. b.	botm...	16	S. 82° W..	.8	
84	85				12' Agz.; m. b.	botm...	10	S. 40° W..	.2	Bridle stops carried away; net badly torn.
					dip; e. l.....	surface.	1 00			
					dyn.....	8-10 ft..	4 30			6 shots.
					dyn.....	4 30			1.5	
83	84				12' Agz.; m. b.	botm...	20	S. 39° E..	1.6	
84	86				12' Agz.; m. b.	botm...	22	S. 63° E..	1.6	
					130' seine, 2 wings.	15 ft....	2 30			3 hauls.
					4 gill nets.....		11 00			
					dip; e. l.....	surface.	1 00			
					25' seine.....		11 00		4.5	
					dyn.....	8-10 ft..	5 00			10 shots.
83	85				12' Agz.; m. b.	botm...	42	N. 89° E..	2.7	
85	86				12' Agz.; m. b.	botm...	33	E.....	2.1	
84	86				12' Agz.....	botm...	42	N. 86° E..	2.8	
84	86				12' Agz.....	botm...	34	S. 50° E..	1.6	
					dip; e. l.....	surface.	1 00			
80	84				12' Agz.....	botm...	29	S. 60° E..	2.1	
83	85				12' Agz.....	botm...	25	S. 62° E..	1.7	Bridle stops and lashing carried away; load lost.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>East coast of Luzon, San Bernardino Strait to San Miguel Bay—Continued.</i>					
D. 5473	East Pt. (Batan), S. 20° E., 8.9 miles (13° 24' 15" N., 124° 02' 48" E.).	C. S. 4221; June, 1905.	1909. June 19	2.05 p. m. 2.49 p. m.	<i>fms.</i> 545	gy. M., S.
.....	Albay G., between Paron and Jesus Pt.do.....	June 21	1.00 p. m.	Co.
.....	Batan Id., East Pt.	C. S. 4250; Aug., 1906.	June 22	8.00 a. m.	Co., S.
.....	Rapurapu Id.do.....do.....	1.00 p. m.	S., Co.
.....	Batan Id., Batan anch.do.....do.....	8.00 p. m.
.....	Port Gubat (Luzon).	C. S. 4258; Jan., 1903.	June 23	1.00 p. m.	Co., S.
D. 5474	S. Bernardino Lt., S. 6° W., 8.4 miles (12° 53' 48" N., 124° 18' E.).	C. S. 4220; May, 1907.	June 24	7.18 a. m. 7.37 a. m.	124	Co.
D. 5475	S. Bernardino Lt., S. 27° W., 11 miles (12° 55' 26" N., 124° 22' 12" E.).do.....do.....	8.51 a. m. 9.15 a. m.	195	Sh.
D. 5476	S. Bernardino Lt., S. 37° W., 13.5 miles (12° 56' 24" N., 124° 25' 24" E.).do.....do.....	10.29 a. m. 11.02 a. m.	270	fne. S.
.....	Langao Pt. (extreme south- ern Luzon).do.....do.....	3.30 p. m.	Co.
	<i>Between Samar and Leyte, vicinity of Surigao Strait.</i>					
.....	Bito Lake and River (Leyte).	C. S. 4423; June, 1905.	July 26	5.30 a. m.
H. 4930	Abuyog (Leyte).do.....do.....	8.00 a. m.	S.
.....	Taebuc Pt. (Leyte), S. 81° W., 16 miles (10° 46' 24" N., 125° 17' 33" E.).do.....	July 27	7.02 a. m.	93	S.
H. 4931	Pagbabaenan Pt. (Malhon Id.), S. 79° E., 16.5 miles (10° 45' 10" N., 125° 27' 48" E.).do.....do.....	8.12 a. m.	63	crs. S., Sh.
.....	Casogoran (Malhon Id.).do.....do.....	10.30 a. m.	S., Co.
.....	Gigoso Pt., Quinapundan Bay (Samar).do.....	July 28	11.00 a. m.	S., Co.
.....	San Roque (Leyte).	C. S. 4719; Aug., 1907.	July 29	9.30 a. m.	Co., S.
H. 4932	Taebuc Pt. (Leyte), N. 79° W., 9.5 miles (10° 42' 10" N., 125° 10' 36" E.).do.....do.....	10.02 a. m.	44	gy. M.
D. 5477	Taebuc Pt. (Leyte), S. 87° W., 11 miles (10° 44' 45" N., 125° 12' 30" E.).do.....do.....	10.23 a. m. 10.33 a. m.	48	gy. M.
D. 5478	Taebuc Pt. (Leyte), S. 80° W., 15.2 miles (10° 46' 24" N., 125° 16' 30" E.).	C. S. 4423; June, 1905.do.....	11.33 a. m. 11.44 a. m.	57	Sh.
D. 5479	Taebuc Pt. (Leyte), S. 78° W., 16.5 miles (10° 47' 15" N., 125° 17' 50" E.).do.....do.....	1.02 p. m. 1.16 p. m.	62	gy. M.
D. 5480	Taebuc Pt. (Leyte), S. 87° W., 17.3 miles (10° 44' 36" N., 125° 19' E.).do.....do.....	2.03 p. m. 2.12 p. m.	62	fne. S.
.....	Hinunangan Bay (Leyte)....	C. S. 4719; Aug., 1907.	July 30	7.30 a. m.	Co., S.
D. 5481	Cabugan Grande Id. (N.), N. 86° W., 3.8 miles (10° 27' 30" N., 125° 17' 10" E.).do.....do.....	8.18 a. m. 8.28 a. m.	61	S., Sh., G.
D. 5482	Cabugan Grande Id. (N.), N. 87° W., 4.5 miles (10° 27' 30" N., 125° 18' E.).do.....do.....	8.56 a. m. 9.11 a. m.	67	brk. Sh., S., gn. M.
D. 5483	Cabugan Grande Id. (N.), N. 88° W., 5.7 miles (10° 27' 30" N., 125° 19' 15" E.).do.....do.....	9.48 a. m. 10.00 a. m.	74	S., brk. Sh.
D. 5484	Cabugan Grande Id. (N.), S. 88° W., 6.4 miles (10° 28' N., 125° 20' E.).do.....do.....	10.33 a. m. 10.44 a. m.	76	S., brk. Sh.
H. 4933	Cabugan Grande Id. (N.), N. 70° W., 9.1 miles (10° 24' 37" N., 125° 22' 15" E.).do.....do.....	12.02 p. m.	90	gn. M., S., brk. Sh.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.								
85	86	40.3			Luc. sdr. (a)		<i>h. m.</i>		<i>m.</i>	
					12' Agz.	botm.	15	S. 41° E.	1.2	Bridle stops carried away.
					dyn.	10 ft.	4 00			12 shots.
					dyn.	7-12ft.	3 30			5 shots.
					dyn.	10-15ft.	5 00			14 shots.
					dip; e. l.		1 00			
					dyn.	8-15ft.	4 00			11 shots.
					Tnr. sdr. (e)					
82	82				12' Agz.	botm.	16	S. 58° W.	.8	
		59.3			Luc. sdr. (a)					
85	82				12' Agz.	botm.	16	N. 82° W.	1.2	
		48.3			Luc. sdr. (a)					
84	83				12' Agz.	botm.	26	N. 84° W.	1.0	
					dyn.	6-15ft.	2 00			8 shots.
					d y n., s m l		12 00		4.0	
					seines.					
					130' seine.	12 ft.	4 00			3 hauls.
					Tnr. sdr. (e)					
					Tnr. sdr. (e)					
					dyn.	9-18ft.	6 00			18 shots.
					dyn.	6-10ft.	5 15			17 shots.
					dyn.	8-15ft.	5 45			25 shots.
					Tnr. sdr. (e)					
					Tnr. sdr. (e)					
86	83				12' Agz.	botm.	20	S. 64° E.	1.0	
					Tnr. sdr. (e)					
87	83				12' Agz.	botm.	14	S. 74° E.	.7	
					Tnr. sdr. (e)					
87	84				12' Agz.	botm.	20	S. 51° E.	.8	
					Tnr. sdr. (e)					
88	84				12' Agz.	botm.	20	E.	.7	
					dyn.	10-15ft.	8 00			20 shots.
					Tnr. sdr. (e)					
84	83				12' Agz.	botm.	20	E. by S.	1.0	
					Tnr. sdr. (e)					
84	83				12' Agz.	botm.	24	E. ¼ S.	1.2	
					Tnr. sdr. (e)					
84	83				12' Agz.	botm.	21	N. 58° E.	1.2	
					Tnr. sdr. (e)					
85	83				12' Agz.	botm.	30	N. 70° E.	1.2	
					Tnr. sdr. (e)					

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Between Samar and Leyte, vicinity of Surigao Strait—Continued.</i>					
D. 5485	Cabugan Grande Id. (N.), N. 59° W., 10.5 miles (10° 22' 15" N., 125° 22' 30" E.).	C. S. 4719; Aug., 1907.	1909. July 30	12.42 p. m. 12.57 p. m.	fms. 103	gn. M.
	<i>Between Leyte and Mindanao.</i>					
D. 5486	Botobolo Pt. (Panaon Id.), S. 19° W., 6 miles (10° 02' N., 125° 19' 20" E.).	C. S. 4719; Aug. 1907.	July 31	8.37 a. m. 9.20 a. m.	585
D. 5487	San Ricardo Pt. (Panaon Id.), S. 50° E., 11.2 miles (10° 02' 45" N., 125° 05' 33" E.).do.....	do.....	1.11 p. m. 2.03 p. m.	732	gn. M.
D. 5488	San Ricardo Pt. (Panaon Id.), S. 59° E., 9 miles (10° N., 125° 6' 45" E.).do.....	do.....	3.59 p. m. 4.52 p. m.	772	gn. M.
D. 5489	San Ricardo Pt. (Panaon Id.), N. 42° E., 6.6 miles (9° 50' 30" N., 125° 10' E.).do.....	do.....	7.21 p. m.		
D. 5490	San Ricardo Pt., N. 9° E., 23.9 miles (9° 32' N., 125° 11' E.).do.....	Aug. 1..	5.10 a. m. 6.20 a. m.	830	gn. M.
D. 5491	Diuata Pt. (W.), S. 9° W., 19.3 miles (9° 24' N., 125° 12' E.).do.....	do.....	8.25 a. m. 10.12 a. m.	736	gn. M., Co.
D. 5492	Diuata Pt. (W.), S. 45° W., 15.2 miles (9° 12' 45" N., 125° 20' E.).do.....	do.....	12.42 p. m. 1.31 p. m.	735	gy. M.
D. 5493	Diuata Pt. (N.), N. 84° W., 5.5 miles (9° 04' N., 125° 20' E.).do.....	Aug. 2..	6.13 a. m. 7.03 a. m.	478	gn. M.
D. 5494	Diuata Pt. (N.), N. 74° W., 4.2 miles (9° 06' 30" N., 125° 18' 40" E.).do.....	do.....	8.30 a. m. 9.17 a. m.	678	gn. M., S.
D. 5495	Diuata Pt. (N.), S. 76° E., 9.4 miles (9° 06' 30" N., 125° 00' 20" E.).do.....	do.....	12.44 p. m. 1.54 p. m.	976	gy. M.
	Mahing River, Camiguin Id. (mouth).do.....	Aug. 3..	2.30 a. m.		
	Mahing, Camiguin Id.do.....	do.....	6.30 a. m.		S., Co.
D. 5496	Bantigui Id., N. 64° W., 7 miles (9° 08' 26" N., 124° 57' E.).do.....	do.....	7.40 a. m. 8.46 a. m.	788
D. 5497	Bantigui Id., N. 64° W., 10 miles (9° 07' 15" N., 124° 59' 30" E.).do.....	do.....	9.55 a. m. 10.59 a. m.	960	gn. M., fne. S.
D. 5498	Bantigui Id., N. 64° W., 10 miles (9° 07' 15" N., 124° 59' 30" E.).do.....	do.....	2.50 p. m.	960	gn. M., fne. S.
	<i>Northern Mindanao and vicinity.</i>					
D. 5499	Macabalan Pt. Lt. (Mindanao), S. 20° E., 11.6 miles (8° 41' 30" N., 124° 35' 40" E.).	C. S. 4719; Aug., 1907.	Aug. 4..	9.10 a. m. 9.50 a. m.	554	gn. M., fne. S.
D. 5500	Macabalan Pt. Lt. (Mindanao), S. 20° E., 7.9 miles (8° 37' 45" N., 124° 36' 45" E.).do.....	do.....	11.05 a. m. 11.25 a. m.	267	gn. M.
	Opol, Macajalar Bay (Mindanao).	C. S. 4644; July, 1905.	do.....	1.00 p. m.		S., Co.
D. 5501	Macabalan Pt. Lt. (Mindanao), S. 35° E., 8.2 miles (8° 37' 37" N., 124° 35' E.).	C. S. 4719; Aug., 1907.	do.....	1.50 p. m. 2.28 p. m.	214	fne. S., gy. M.
D. 5502	Macabalan Pt. Lt. (Mindanao), S. 35° E., 8.2 miles (8° 37' 37" N., 124° 35' E.).do.....	do.....	3.28 p. m.	**214

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
85	83				Tnr. sdr. (c) 12' Agz.	botm...	20	N. 40° E...	1.7	
84	82	52.1			Luc. sdr. (a) 12' Agz.	botm...	21	S. 37° E...	3.0	
84	84	52.3			Luc. sdr. (a) 12' Agz.	botm...	23	S. 65° E...	2.7	
85	83	52.3			Luc. sdr. (a) 12' Agz.	botm...	46	S. 43° E...	3.5	
84	83				K. 2.....	10 ft....	46	S. 43° E...	3.5	
					int. 4 §.....	50 fms..	20	S.....	.5	
					K. 5.....	surface.	4 20	S.....	.5	
83	84	52.5			Luc. sdr. (a) 12' Agz.	botm...	12	S. 28° E...	.9	Whole apparatus carried away.
84	83	52.3			Luc. sdr. (a) 12' Agz.	botm...	31	S. 45° E...	2.7	
84	85	52.3			Luc. sdr. (a) 12' Agz.	botm...	28	S. 14° E...	2.3	
80	83	52.1			Luc. sdr. (a) 12' Agz.	botm...	45	N. 32° W..	3.0	
82	83	53.3			Luc. sdr. (a) 12' Agz.	botm...	35	N. 5° E...	3.2	
					K. 5.....	surface.	35	N. 5° E...	3.2	
84	83	52.3			Luc. sdr. (a) 12' Agz.	botm...	35	S. 17° E...	2.7	
					K. 5.....	600 fms.	33	S. 17° E...	2.0	
					25' seine; dyn		3 00			Mouth of river. 21 shots.
80	83	52.3			Luc. sdr. (a) 12' Agz.	botm...	16	S. 52° E...	2.5	Lost apparatus and 1,000 fms. wire.
80	83	52.3			Luc. sdr. (a) int. 4 §.....		800 fms. 20 35	S. 60° E...	2.6	
82	84				12' Agz.	botm...	27	S. 48° E...	3.4	
83	84	52.3			Luc. sdr. (a) 12' Agz.	botm...	5	N. 76° E...	1.9	Bridle stops lost frame twisted.
87	84	53.5			Luc. sdr. (a) int. 4 §.....	200 fms.	18 13	S. 67° E...	1.0	
					dyn.....	5-12 ft.	4 30			20 shots.
85	86	54.3			Luc. sdr. (a) 12' Tnr.	botm...	20	S. 38° E...	1.5	
84	86				12' Tnr.	botm...	20	S. 38° E...	1.7	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Northern Mindanao and vicinity—Continued.</i>					
D. 5503	Macabalan Pt. Lt. (Mindanao), S. 31° E., 6.6 miles (8° 36' 26" N., 124° 36' 08" E.).	C. S. 4719; Aug., 1907.	1909. Aug. 4...	4.10 p. m. 4.38 p. m.	fms. 226	gn. M.
D. 5504	Macabalan Pt. Lt. (Mindanao), S. 39° E., 6 miles (8° 35' 30" N., 124° 36' E.).do.....	Aug. 5...	5.50 a. m. 6.15 a. m.	200	gn. M.
D. 5505	Macabalan Pt. Lt. (Mindanao), S. 31° E., 7.7 miles (8° 37' 15" N., 124° 36' E.).do.....do.....	7.25 a. m.	*220	
D. 5506	Macabalan Pt. Lt. (Mindanao), S. 41° E., 12.2 miles (8° 40' N., 124° 31' 45" E.).do.....do.....	8.40 a. m. 9.12 a. m.	262	gn. M.
D. 5507	Camp Overton Lt., Iligan Bay (Mindanao), S. 11° E., 8.6 miles (8° 21' 12" N., 124° 12' 06" E.).	C. S. 4613; June, 1906.do.....	1.09 p. m. 1.44 p. m.	425	gn. M., fne. S.
D. 5508	Camp Overton Lt., Iligan Bay, S. 6° E., 4.9 miles (8° 17' 24" N., 124° 11' 42" E.).do.....do.....	2.53 p. m. 3.17 p. m.	270	gn. M., fne. S.
.....	Camp Overton, Iligan Bay (Mindanao).do.....	Aug. 6...	8.00 a. m.		Co., S.
.....	Nonucan R., Iligan Bay (near Camp Overton).do.....do.....	8.00 a. m.		
D. 5509	Camp Overton Lt., S. 61° E., 5.7 miles (8° 15' 24" N., 124° 07' 18" E.).do.....	Aug. 7...	8.06 a. m. 8.36 a. m.	377	gy. M.
D. 5510	Camp Overton Lt., S. 68° E., 9.1 miles (8° 16' N., 124° 03' 50" E.).do.....do.....	9.53 a. m. 10.31 a. m.	423	gy. M., fne. S.
D. 5511	Camp Overton Lt., S. 80° E., 15.3 miles (8° 15' 20" N., 123° 57' E.).do.....do.....	11.46 a. m. 12.18 p. m.	410	gy. M., S.
D. 5512	Camp Overton Lt., S. 76° E., 14 miles (8° 16' 02" N., 123° 58' 26" E.).do.....do.....	1.09 p. m. 1.46 p. m.	445	gy. M., fne. S.
D. 5513	Camp Overton Lt., S. 67° E., 10.3 miles (8° 16' 45" N., 124° 02' 48" E.).do.....do.....	3.07 p. m. 3.53 p. m.	505	gy. M., fne. S.
D. 5514	Camp Overton Lt., S. 34° E., 24.3 miles (8° 32' 42" N., 123° 58' 36" E.).do.....	Aug. 8...	7.58 a. m. 8.50 a. m.	697	gn. M., S.
D. 5515	Camp Overton Lt., S. 26° E., 24.6 miles (8° 34' 48" N., 124° 01' 24" E.).do.....do.....	10.42 a. m.		
.....	Inamucan Bay (Mindanao).do.....do.....	2.30 p. m.		R., Co.
.....do.....do.....	Aug. 9...	5.30 a. m.		S.
.....	Murcielagos Bay (Mindanao).	C. S. 4641; Apr., 1902.do.....	9.30 a. m.		Co., S.
D. 5516	Pt. Tagolo Lt. (Mindanao), S. 80° W., 9.7 miles (8° 40' N., 123° 32' 30" E.).	C. S. 4723; Oct., 1905.do.....	9.57 a. m. 10.21 a. m.	175	Glob.
D. 5517	Pt. Tagolo Lt., S. 83° W., 10.5 miles (8° 45' 30" N., 123° 33' 45" E.).do.....do.....	11.00 a. m. 11.21 a. m.	169	Glob.
D. 5518	Pt. Tagolo Lt., S. 64° W., 8.7 miles (8° 48' N., 123° 31' E.).do.....do.....	12.36 p. m. 12.55 p. m.	200	gy. M., Glob.
D. 5519	Pt. Tagolo Lt., S. 71° W., 8.7 miles (8° 47' N., 123° 31' 15" E.).do.....do.....	1.38 p. m. 1.56 p. m.	182	Glob., S.
D. 5520	Pt. Tagolo Lt., N. 48° E., 4.5 miles (8° 41' 15" N., 123° 18' 30" E.).do.....	Aug. 10...	6.02 a. m. 6.20 a. m.	102	
D. 5521	Pt. Tagolo Lt., S. 11° E., 3 miles (8° 47' N., 123° 22' 30" E.).do.....do.....	7.24 a. m. 7.51 a. m.	221	fne. S.
.....	Silino Id. (west).do.....do.....	8.40 a. m.		S., Co.
D. 5522	Pt. Tagolo Lt., S. 39° W., 6 miles (8° 49' N., 123° 20' 30" E.).do.....do.....	9.11 a. m. 9.57 a. m.	230	Glob.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Tempera- tures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Sur- face.	Bot- tom.		Depth.	Dura- tion.	Direction.	Distance.	
° F.	° F.	° F.					<i>h. m.</i>		<i>mi.</i>	
83	86	53.3			Luc. sdr. (a) 12' Tnr.	botm.	20	S. 2° E.	1.2	
77	83	54.3			Luc. sdr. (a) 12' Tnr.	botm.	20	N. 7° W.	1.7	
79	83				12' Tnr.	botm.	24	N. 18° W.	1.4	
84	82	53.3			Luc. sdr. (a) 12' Tnr.	botm.	14	N. 24° W.	1.7	
85	84	52.8			Luc. sdr. (a) 12' Tnr.	botm.	20	S. 8° W.	1.0	
84	85	53.3			Luc. sdr. (a) 12' Tnr.	botm.	24	S. 2° E.	1.8	
					dyn.	6-12 ft.	8 00			10 shots.
					dyn.		8 30			3.5
79	82	53.0			Luc. sdr. (a) 12' Tnr.	botm.	23	N. 34° W.	1.4	
83	84	53.0			Luc. sdr. (a) 12' Tnr.	botm.	7	S. 44° W.	1.6	Net badly torn.
84	85	53.0			Luc. sdr. (a) 12' Tnr.	botm.	20	N. 64° E.	1.9	
91	86	52.8			Luc. sdr. (a) 12' Tnr.	botm.	20	N. 74° E.	2.2	
84	85	52.8			Luc. sdr. (a) 12' Tnr.	botm.	15	S. 83° E.	1.7	Beam frame sprung; net torn.
81	83	52.3			Luc. sdr. (a) 12' Tnr.	botm.	27	N. 47° E.	3.0	Net fouled over beam.
85	83				12' Tnr.	botm.	28	S. 20° W.	1.6	No sounding, depth about 700 fms.
					dyn.	8-15 ft.	3 00			11 shots.
					430' seine.	3 ft.	2 00			3 hauls.
					dyn.	6-25 ft.	4 15			15 shots.
85	84	54.3			Luc. sdr. (a) 12' Tnr.	botm.	20	S. 63° E.	1.2	
83	85	54.3			Luc. sdr. (a) 12' Tnr.	botm.	18	S. 50° E.	1.1	
84	85	54.0			Luc. sdr. (a) 12' Tnr.	botm.	21	S. 9° E.	1.2	
83	85	54.3			Luc. sdr. (a) 12' Tnr.	botm.	43	S. 14° E.	1.6	
79	84	61.3			Luc. sdr. (a) 12' Tnr.	botm.	24	N. 13° E.	1.3	No bottom sam- ple in sounding cup.
81	84	53.3			Luc. sdr. (a) 12' Tnr.	botm.	4	N. 52° E.	.9	Whole apparatus carried away. 13 shots.
81	84	52.3			dyn. Luc. sdr. (a) 12' Tnr.	10-20 ft. botm.	3 00			Net fouled over beam.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>Northern Mindanao and vicinity—Continued.</i>						
D. 5523	Pt. Tagolo Lt., S. 48° W., 6.7 miles (8° 48' 44" N., 123° 27' 35" E.).	C. S. 4723; Oct., 1905.	1909, Aug. 10	10.49 a. m.	<i>fms.</i>	
D. 5524	Pt. Tagolo Lt., S. 34° W., 17 miles (8° 58' 07" N., 123° 32' 45" E.).	do	do	1.06 p. m. 1.51 p. m.	360	S
<i>Between Siquijor and Bohol Ids.</i>						
D. 5525	Balicasag Id. (C.), N. 11° W., 18.2 miles (9° 12' 30" N., 123° 44' 07" E.).	C. S. 4718; Dec., 1906.	Aug. 11	8.28 a. m.	405	gy. M.
D. 5526	Balicasag Id. (C.), N. 15° W., 18.4 miles (9° 12' 45" N., 123° 45' 30" E.).	do	do	9.29 a. m. 10.56 a. m.	805	gn. M., Glob.
D. 5527	Balicasag Id. (C.), N. 14° W., 8.2 miles (9° 22' 30" N., 123° 42' 40" E.).	do	do	1.07 p. m. 1.38 p. m.	392	glob. Oz.
D. 5528	Balicasag Id. (C.), N. 15° E., 5.8 miles (9° 24' 45" N., 123° 39' 15" E.).	do	do	3.08 p. m. 3.42 p. m.	439	glob. Oz.
D. 5529	Balicasag Id. (C.), N. 11° E., 6.9 miles (9° 23' 45" N., 123° 39' 30" E.).	do	do	4.44 p. m. 5.19 p. m.	441	gy. M., Glob.
D. 5530	Balicasag Id. (C.), N. 32° E., 4.3 miles (9° 26' 45" N., 123° 38' 30" E.).	do	do	7.14 p. m.		
D. 5531	Balicasag Id. (C.), N. 43° E., 4.2 miles (9° 27' 30" N., 123° 38' 00" E.).	do	do	7.49 p. m.		
<i>Between Masbate and Leyte.</i>						
D. 5532	Gigantangan Id. (S.), S. 33° E., 3.8 miles (11° 36' 39" N., 124° 13' 30" E.).	C. S. 4718; Dec., 1906.	Aug. 13	7.14 p. m.		
<i>Between Cebu and Siquijor.</i>						
D. 5533	Balicasag Id. (C.), N. 71° E., 9.4 miles (9° 27' 15" N., 123° 31' 48" E.).	C. S. 4718; Dec., 1906.	Aug. 19	5.30 a. m. 6.08 a. m.	432	gn. M., S
D. 5534	Balicasag Id. (C.), N. 72° E., 14.7 miles (9° 26' 00" N., 123° 26' 37" E.).	do	do	8.23 a. m. 8.53 a. m.	333	gy. glob. Oz.
D. 5535	Apo Id. (C.), S. 24° W., 17 miles (9° 20' 30" N., 123° 23' 45" E.).	do	do	10.38 a. m. 11.07 a. m.	310	gy. glob. Oz.
<i>Between Negros and Siquijor.</i>						
D. 5536	Apo Id. (C.), S. 26° W., 11.8 miles (9° 15' 45" N., 123° 22' 00" E.).	C. S. 4718; Dec., 1906.	Aug. 19	12.50 p. m. 1.36 p. m.	279	gn. M.
D. 5537	Apo Id. (C.), S. 46° W., 8.7 miles (9° 11' 00" N., 123° 23' 00" E.).	do	do	3.15 p. m. 3.39 p. m.	254	gn. M.
D. 5538	Apo Id. (C.), S. 64° W., 7.3 miles (9° 08' 15" N., 123° 23' 20" E.).	do	do	4.55 p. m. 5.20 p. m.	256	gn. M., S
D. 5539	Apo Id. (C.), N. 78° W., 8.2 miles (9° 03' 20" N., 123° 24' 45" E.).	do	do	7.11 p. m.		
D. 5540	Apo Id. (C.), N. 76° W., 8.1 miles (9° 03' 00" N., 123° 24' 30" E.).	do	do	7.42 p. m.		

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Tempera- tures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Sur- face.	Bot- tom.		Depth.	Dura- tion.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
82	84				12' Tnr.; m. b.	botm.	20	S. 22° E.	1.2	No sounding.
83	84	52.8			Luc. sdr. (a) 12' Tnr.	botm.	25	S. 16° W.	1.2	
82	82	53.3			12' Tnr.	botm.	22	N. 85° E.	1.7	
82	82	52.3			Luc. sdr. (a) 12' Tnr.	botm.	17	E.	1.8	
87	84	53.3			Luc. sdr. (a) 12' Tnr.	botm.	20	S. 14° E.	1.2	
87	85	53.3			Luc. sdr. (a) 12' Tnr.	botm.	29	S. 17° E.	1.3	
85	85	53			Luc. sdr. (a) 12' Thr.; m. b.	botm.	35	S. 17° E.	1.6	
84	84				int. 4.	surface.	20			
83	84				int. 4.	surface.	28			
86	84				int. 4.	surface.	14			
80	81	53.3			Luc. sdr. (a) 12' Tnr.	botm.	23	S. 30° E.	1.3	
82	82	53.3			Luc. sdr. (a) 12' Tnr.	botm.	20	S. 64° W.	1.8	
83	84	53.3			Luc. sdr. (a) 12' Tnr.	botm.	09	S. 69° W.	1.5	Bridle carried away at surface, causing loss of most of catch.
84	85	53.5			Luc. sdr. (a) 12' Tnr.	botm.	20	S. 60° W.	2.7	
87	84	53.5			K. 5. Luc. sdr. (a)	botm.	20	S. 75° W.	2.0	
85	84				12' Tnr.	botm.	20	S. 75° W.	2.0	
83	83	53.3			Luc. sdr. (a) 12' Tnr.	botm.	22	S. 80° W.	1.3	
83	83				int. 4.	surface.	19			
83	83				int. 4.	surface.	16			

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>Northern Mindanao and vicinity.</i>						
D. 5541	Tagolo Lt., S. 65° W., 12.7 miles (8° 49' 38" N., 123° 34' 30" E.).	C. S. 4723; Oct., 1905.	1909. Aug. 20	5.25 a. m. 5.51 a. m.	fms. 219	fne. S., brk. Sh....
D. 5542	Tagolo Lt., S. 70° W., 13.2 miles (8° 48' 30" N., 123° 35' 30" E.).	do.....	do.....	6.34 a. m. 6.56 a. m.	200	fne. S., brk. Sh....
D. 5543	Tagolo Lt., S. 75° W., 12.5 miles (8° 47' 15" N., 123° 35' 00" E.).	do.....	do.....	8.46 a. m. 9.04 a. m.	162	S.....
.....	Murcielagos Bay (Mindanao).	C. S. 4641; Apr., 1902.	do.....	1.00 p. m.	S., Co.....
.....	Cascade River, Murcielagos Bay.	do.....	do.....	1.00 p. m.
D. 5544	Coronado Pt., S. 37° W., 21.5 miles (8° 16' 30" N., 122° 26' 30" E.).	C. S. 4723... Oct., 1905.	Sept. 6	10.34 a. m. 11.17 a. m.	759	gn. M., fne. S.....
<i>East of Zamboanga.</i>						
.....	Tictauan Id., east.....	C. S. 4511; Dec., 1904.	Sept. 8	7.45 a. m.	S., Co., R.....
.....	Malanipa Id., northeast.....	do.....	do.....	1.00 p. m.	S., R., Co.....
.....	Sacol Id., northeast.....	do.....	do.....	7.00 p. m.
.....	Tulnalutan Id., north.....	do.....	Sept. 9	6.00 a. m. 1.00 p. m.	Co..... Co., S., R.....
<i>South of Zamboanga.</i>						
.....	Isabel Channel, Basilan Id...	C. S. 4543; May, 1907.	Sept. 11	8.30 a. m.	S., Co.....
.....	Lampinigan Id., north and east.	do.....	do.....	1.30 p. m. 7.30 p. m.	Co., S.....
.....	Balukbaluk Id., west.....	C. S. 4511; Dec., 1904.	Sept. 12	8.30 a. m.	Co., S.....
.....	Pilas Id., northeast.....	do.....	do.....	2.00 p. m.	Co.....
.....	Tapiantana Id., north.....	C. S. 4512; Sept., 1906.	Sept. 13	9.30 a. m.	S., Co.....
.....	Bulan Id., north.....	do.....	do.....	3.00 p. m. 7.30 p. m.	Co.....
.....	Tonquil Id., Gumila Reef...	do.....	Sept. 14	8.30 a. m.	Co., S.....
.....	Tonquil Id., northwest.....	do.....	do.....	2.00 p. m.	Co., S.....
<i>Jolo I. and vicinity.</i>						
.....	Tulayan Id.....	C. S. 4512; Sept., 1906	Sept. 15	9.00 a. m.	Co., S.....
D. 5545	Noble Pt., Tulayan Id. (E.), S. 19° W., 3 miles (6° 04' 45" N., 121° 20' 20" E.).	do.....	do.....	9.26 a. m. 9.43 a. m.	114	fne. co. S.....
D. 5546	Noble Pt., Tulayan Id. (E.) S. 13° W., 5 miles (6° 06' 48" N., 121° 20' 32" E.).	do.....	do.....	10.34 a. m. 10.52 a. m.	138	fne. co. S.....
D. 5547	Noble Pt., Tulayan Id. (E.), S. 38° E., 9.5 miles (6° 09' 20" N., 121° 13' 40" E.).	C. S. 4542; Apr., 1903.	do.....	1.31 p. m. 1.51 p. m.	155	fne. S.....
D. 5548	Jolo Lt. (Jolo), N. 77° E., 14.9 miles (6° 00' 20" N., 120° 45' 35" E.).	do.....	Sept. 17	7.55 a. m. 8.20 a. m.	232	S., brk. Sh.....
D. 5549	Jolo Lt. (Jolo), N. 80° E., 15.8 miles (6° 01' 15" N., 120° 44' 20" E.).	do.....	do.....	9.09 a. m. 9.36 a. m.	263	S., Glob., For....
D. 5550	Jolo Lt. (Jolo), N. 83° E., 15.5 miles (6° 02' 00" N., 120° 44' 40" E.).	do.....	do.....	10.20 a. m. 10.46 a. m.	258	fne. S., Sh.....
.....	Sulade Id., north.....	do.....	do.....	1.00 p. m.	Co., S.....
D. 5551	Jolo Lt. (E.), N. 60° E., 18 miles (5° 54' 48" N., 120° 44' 24" E.).	do.....	do.....	1.46 p. m. 2.07 p. m.	193	fne. S.....

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
°F.	°F.	°F.					<i>h. m.</i>		<i>mi.</i>	
81	83	53.3			Luc. sdr. (a) 12' Tnr.	botm...	21	S. 17° E...	1.0	
		54.3			Luc. sdr. (a) 12' Tnr.	botm...	20	S. 25° W...	1.4	Net came up torn and tangled.
		54.5			Luc. sdr. (a) 12' Tnr.	botm...	17	S. 20° W...	.7	Bridle stops carried away, frame bent, net badly torn.
					dyn.....	4-12 ft.	8 00			22 shots.
					dyn.....		4 30		1.5	
82	83	49.8			Luc. sdr. (a) int. 4 §	600 fms.	20 33	N. 49° W...	1.5	
					dyn.....	10-15 ft.	3 30			12 shots.
					dyn.....	10-18 ft.	2 45			10 shots.
					dip; e. l.....		1 00			Do.
					dyn.....	12-15 ft.	4 00			16 shots.
					dyn.....	9-20 ft.	3 30			
					dyn.....	10-30 ft.	2 30			6 shots.
					dyn.....	6-18 ft.	4 00			18 shots.
					dip; e. l.....		1 15			
					dyn.....	8-18 ft.	3 00			9 shots.
					dyn.....	10-20 ft.	3 30			10 shots.
					dyn.....	8-10 ft.	2 00			8 shots.
					dyn.....	10-15 ft.	2 30			12 shots.
					dip; e. l.....		1 00			
					dyn.....	4-6 ft.	3 00			14 shots.
					dyn.....	8-10 ft.	3 15			12 shots.
					dyn.....	8-20 ft.	3 00			7 shots.
82	82				Luc. sdr. (e) 9' Tnr.	botm...	16	S. 34° E...	1.1	
83	82	58.3			Luc. sdr. (a) 9' Tnr.	botm...	19	S. 49° E...	1.4	
84	82	56.3			Luc. sdr. (a) 9' Tnr.	botm...	20	S. 32° E...	1.5	
82	82	53.5			Luc. sdr. (a) 9' Tnr.; m. b.	botm...	29	N. 55° W...	1.5	
83	83	52.3			Luc. sdr. (a) 9' Tnr.; m. b.	botm...	21	N. 23° E...	1.1	
85	83	52.3			Luc. sdr. (a) 9' Tnr.	botm...	28	S. 60° E...	1.2	
					dyn.....	10-15 ft.	4 00			14 shots.
84	83	53.3			Luc. sdr. (a) 9' Tnr.	botm...	20	S. 15° E...	1.1	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>Jolo I. and vicinity—Cont'd.</i>						
D. 5552	Jolo Lt. (E.), N. 60° E., 18.3 miles (5° 54' 30" N., 120° 44' 15" E.).	C. S. 4542; Apr., 1903.	1909. Sept. 17	3.18 p. m.	<i>fms.</i>	
D. 5553	Sulade Id. (NW.), S. 4° E., 0.5 mile (5° 51' 00" N., 120° 46' 30" E.).do.....do.....	7.28 p. m.		
D. 5554	Cabalian Pt. (Jolo), N. 76° E., 3.8 miles (5° 52' 27" N., 120° 52' 18" E.).do.....	Sept. 18.	9.19 a. m. 9.29 a. m.	25	Co., S
D. 5555	Cabalian Pt. (Jolo), N. 50° W., 3.3 miles (5° 51' 15" N., 120° 58' 35" E.).do.....do.....	10.59 a. m. 11.09 a. m.	34	crs. S
D. 5556	Cabalian Pt., N. 59° W., 4.5 miles (5° 50' 55" N., 121° 00' 00" E.).do.....do.....	11.36 a. m.	15	
D. 5557	Teomabal Id. (N.)..... Cabalian Pt., N. 70° W., 5.2 miles (5° 51' 30" N., 121° 01' 00" E.).do.....do.....do.....do.....	1.30 p. m. 2.58 p. m.	13	scrtd. Co., S. S., Co.*
D. 5558	Cabalian Pt., S., 1.1 miles (5° 51' 33" N., 121° 00' 58" E.).do.....do.....	3.17 p. m.	15	Co.*
D. 5559	Cabalian Pt., N. 66° W., 5.1 miles (5° 51' 36" N., 121° 00' 45" E.).do.....do.....	3.35 p. m.	13	Co.*
D. 5560	Cabalian Pt., N. 76° W., 5 miles (5° 52' 00" N., 121° 01' 06").do.....do.....	4.04 p. m.	14	
D. 5561	Teomabal Id. (NW.), S. 36° W., 0.2 mile (5° 50' 45" N., 121° 01' 15" E.).do.....do.....	6.13 p. m.	*10	
D. 5562	Tutu Bay (Jolo), N. 87° E., 17.2 miles (5° 54' 20" N., 121° 13' 12" E.).do.....	Sept. 19.	8.15 a. m. 1.45 p. m. 6.07 p. m.		Co., S. Co., S.
<i>Between Jolo and Tawi Tawi.</i>						
.....	Siasi Id., north.....	C. S. 4544; Oct., 1906.	Sept. 20.	10.30 a. m.		Co., S
.....	Tara Id., Panpan Pt.....do.....do.....	1.00 p. m.		Co., S
.....	Bolipongpong Id., south.....	C. S. 4722; Jan., 1909.do.....	3.30 p. m.		S., Co., R
D. 5563	Singaan Id., north..... Dammi Id. (N.), N. 79° W., 6.1 miles (5° 48' 12" N., 120° 30' 48" E.).do.....do.....	Sept. 21.do.....	10.00 a. m. 10.25 a. m. 10.47 a. m.	224	Co. fine. co. S
D. 5564	Dammi Id. (N.), S. 85° W., 6.1 miles (5° 50' 00" N., 120° 31' 00" E.).do.....do.....	11.24 a. m. 11.45 a. m.	236	fine. Co., S
D. 5565	Dammi Id. (N.), S. 69° W., 6 miles (5° 51' 42" N., 120° 30' 30" E.).do.....do.....	12.32 p. m. 1.00 p. m.	243	S., ptr. Sh
D. 5566	Dammi Id. (N.), S. 67° W., 6.8 miles (5° 52' 12" N., 120° 31' 00" E.).do.....do.....	1.42 p. m. 2.07 p. m.	244	fine. S., Sh
<i>North of Tawi Tawi.</i>						
D. 5567	Dammi Id. (N.), N. 81° W., 9 miles (5° 48' 00" N., 120° 33' 45" E.).	C. S. 4722; Jan., 1909.	Sept. 21.	3.36 p. m. 4.05 p. m.	268	fine. S
D. 5568	Singaan Id. (N.), West, 0.9 mile (5° 45' 50" N., 120° 26' 00" E.).do.....do.....	6.35 p. m.	13	S., Co.
D. 5569	Simaluc Id. (SE.), S. 8° W., 6.4 miles (5° 33' 15" N., 120° 15' 30" E.).do.....	Sept. 22.	8.19 a. m. 8.49 a. m.	303	co. S
D. 5570	Simaluc Id. (SE.), S. 17° E., 5.7 miles (5° 32' 15" N., 120° 12' 57" E.).do.....do.....	9.55 a. m. 10.27 a. m.	330	fine. S., Glob

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
83	83				9' Tnr.; m. b.	botm.	21	S. 23° E.	1.5	Depth about as previous station.
82	83				int. 4.	surface.	10 41			Ship at anchor.
83	84				Tnr. sdr. (e). 6' McC.	botm.	6	N. 74° W.	2	Net torn.
82	83				Tnr. sdr. (e). 6' McC.	botm.	4	N. 75° E.	5	
82	83				hand lead. 6' McC.	botm.	3	N. 68° E.	3	Trawl and 15 fms cable lost. 7 shots.
83	82				dyn. 10-25 ft. hand lead. 6' McC.	botm.	3 30 5	S. W.	1	
83	82				hand lead. 6' McC.	botm.	3	S. 44° W.	4	
83	82				hand lead. 6' McC.	botm.	7	S. W.	6	
83	82				hand lead. 6' McC.	botm.	9	S. 20° E.	5	Everything carried away except bridle.
81	82				int. 4.	surface.	11 47			Ship at anchor.
					dyn. 10-20 ft. dyn. 2-20 ft.		2 15 2 30			7 shots. 10 shots.
84	82				int. 4.	surface.	11 41			Ship at anchor.
					dyn. 15 ft.		1 30			5 shots.
					dyn. 8-15 ft. dyn. 8-20 ft.		1 30 1 45			Do. Do.
					dyn. 9-25 ft.		8 00			17 shots.
83	83	52.3			Luc. sdr. (a). 9' Tnr.; m. b.	botm.	18	N. 6° W.	1.3	
84	83	52.3			Luc. sdr. (a). 9' Tnr.	botm.	28	N. 9° E.	1.5	
86	84	52.3			Luc. sdr. (a). 9' Tnr.; m. b.	botm.	21	N. 45° E.	7	
84	84	52.5			Luc. sdr. (a). 9' Tnr.	botm.	27	N. 56° E.	1.6	
85	83	52.0			Luc. sdr. (a). 9' Tnr.; m. b.	botm.	21	N. 71° E.	1.2	
82	83				int. 4.	surface.	11 20			Ship at anchor.
84	83	52.3			Luc. sdr. (a). 9' Tnr.; m. b.	botm.	10	S. 73° E.	1.0	Net torn.
87	83	52.3			Luc. sdr. (a). 9' Tnr.; m. b.	botm.	17	N. 45° W.	1.0	Net came up fouled on bolt head.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>North of Tawi Tawi—Cont'd.</i>				<i>fms.</i>	
	Simaluc Id., north.....	C. S. 4722; Jan., 1909.	1909. Sept. 22.	12.30 p. m.		S., Co.....
D. 5571	Simaluc Id. (N.), S. 66° E., 5.8 miles (5° 30' 45" N., 120° 07' 57" E.).	do.....	do.....	1.31 p. m. 2.00 p. m.	340	S., Sh.....
D. 5572	Simaluc Id. (N.), S. 51° E., 4.7 miles (5° 31' 26" N., 120° 09' 45" E.).	do.....	do.....	3.02 p. m. 3.34 p. m.	334	S.....
D. 5573	Simaluc Id. (N.), S. 86° E., 0.4 mile (5° 28' 30" N., 120° 13' 00" E.).	do.....	do.....	6.03 p. m.	12	
D. 5574	Simaluc Id. (N.), S. 66° E., 5.8 miles (5° 30' 45" N., 120° 07' 57" E.).	do.....	Sept. 23.	7.20 a. m.	340	
D. 5575	Mt. Dromedario (Tawi Tawi), S. 16° W., 19.2 miles (5° 28' 30" N., 120° 02' 27" E.).	C. S. 4514; Jan., 1906	do.....	9.07 a. m. 9.43 a. m.	315	Co., S.....
D. 5576	Mt. Dromedario, S. 22° W., 17.2 miles (5° 25' 56" N., 120° 03' 39" E.).	C. S. 4722; Jan., 1909.	do.....	10.50 a. m. 11.22 a. m.	277	S.....
	Bacun River (Tawi Tawi)....	C. S. 4514; Jan., 1906.	do.....	1.30 p. m.		
	Simaluc Sibi Sibi Id.....	do.....	do.....	1.30 p. m.		Co., wh. S.....
D. 5577	Mt. Dromedario, S. 9° W., 10.9 miles (5° 20' 36" N., 119° 58' 51" E.).	do.....	do.....	2.38 p. m. 3.01 p. m.	240	crs. S.....
D. 5578	Mt. Dromedario, S. 9° W., 4.8 miles (5° 14' 38" N., 119° 57' 57" E.).	do.....	do.....	8.00 p. m.	10	
	<i>Vicinity of Darvel Bay, Borneo.</i>					
	Reef NW. of Tumindao Id.....	C. S. 4722; Jan., 1909.	Sept. 24.	1.00 p. m.		Co., S.....
D. 5579	Sibutu Id. peak, S. 77° E., 20.3 miles (4° 54' 15" N., 119° 09' 52" E.).	do.....	Sept. 25.	8.03 a. m. 8.25 a. m.	175	fne. S., Co.....
D. 5580	Sibutu Id. peak, S. 82° E., 23.2 miles (4° 52' 45" N., 119° 06' 45" E.).	do.....	do.....	9.20 a. m. 9.40 a. m.	162	br. S., Co.....
D. 5581	Bumbum Id., north Bumbum Id. (NW.), S. 83° W., 3.5 miles (4° 30' 25" N., 118° 41' 30" E.).	do..... H. O. 2117; June, 1903.	do..... do.....	2.30 p. m. 5.55 p. m.		Co., S..... S., Co.....
D. 5582	Si Amil Id. (N.), S. 82° W., 6.2 miles (4° 19' 54" N., 118° 58' 38" E.).	do.....	Sept. 26.	10.11 a. m. 11.15 a. m.	890	gy. M., fne. S.....
	Danawan Id.....	do.....	do..... Sept. 27.	2.00 p. m. 8.15 a. m.		S., Co..... S., Co.....
	<i>Sibuko Bay, Borneo, and vicinity.</i>					
D. 5583	Si Amil Id. (N.) N. 88 W., 3.2 mile (4° 19' 00" N., 118° 56' 20" E.).	H. O. 2117; June, 1903.	Sept. 27	1.48 p. m. 2.33 p. m.	447	fne. S.....
D. 5584	Si Amil Id. (N.) N. 74° W., 5.4 miles (4° 17' 40" N., 118° 57' 42" E.).	do.....	do.....	3.28 p. m. 4.02 p. m.	292	fn. S., gn. M.....
D. 5585	Sipadan Id. (M.) S. 89° W., 12 miles (4° 07' 00" N., 118° 49' 54" E.).	do.....	Sept. 28	8.49 a. m. 9.31 a. m.	476	gy. M.....
D. 5586	Sipadan Id. (M.) West, 9.4 miles (4° 06' 50" N., 118° 47' 20" E.).	do.....	do.....	11.09 a. m. 11.44 a. m.	347	gy. M.....
D. 5587	Sipadan Id. (N.)..... Sipadan Id. (W.) S. 12° E., 3.8 miles (4° 10' 35" N., 118° 37' 12" E.).	do..... do.....	do..... do.....	2.00 p. m. 2.35 p. m. 3.11 p. m.	415	Co., S..... gn. M., S., Co.....
D. 5588	Mabul Id. (S.) N. 81° E., 1.7 miles (4° 14' 20" N., 118° 30' 48" E.).	do.....	do.....	6.10 p. m.	11	

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
°F.	°F.	°F.				<i>h. m.</i>		<i>mi.</i>		
					dyn.....	5-18 ft..	4 00			11 shots.
81	84	52.3			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm...	21	N. 67° E...	1.4	
82	84	52.3			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm...	26	N. 82° E...	1.9	
83	83				int. 4.....	surface.	11 42			Ship at anchor.
81	82				9' Tnr.; m. b..	botm...	24	N. 58° E...	1.2	
83	83	52.3			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm...	20	S. 86° E...	2.2	
84	84	53.3			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm...	08	S. 2° E...	1.7	
					d y n.; s m l.		3 30			4 shots.
					seines.					
					d y n.....	5-20 ft..	3 30			8 shots.
79	82	54.3			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm...	18	S. 61° E...	1.8	Mud bag lost.
77	82				int. 4.....	surface.	9 49			Ship at anchor.
					d y n.....	5-25 ft..	4 00			17 shots.
80	82	55.3			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm...	20	S. 37° W...	1.5	
82	83	55.8			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm...	17	S. 16° W...	1.0	
82	83				d y n.....	4-15 ft..	3 00			13 shots.
82	83				int. 4.....	surface.	11 55			Ship at anchor.
81	82	38.3			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm...	17	S. 17° E...	3.3	
					d y n.....	3-20 ft..	3 30			13 shots.
					d y n.....	5-20 ft..	8 30			27 shots.
84	85	40.3			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm ..	28	S. 46° E...	2.0	
80	84	44.3			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm ..	21	S. 56° W...	1.3	Net badly torn and T a n e r beam lost.
84	82	41.1			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm ..	20	S. 53° W...	1.9	
83	84	44.0			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm ..	33	N. 42° W...	.8	
					d y n.....	8-20 ft..	2 15			13 shots.
85	85	42.3			Luc. sdr. (a)...					
					9' Tnr.; m. b..	botm ..	21	S. 15° E...	1.5	
83	82				int. 4.....	surface.	11 35			Ship at anchor.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Sibuko Bay, Borneo, and vicinity—Continued.</i>					
.....	Mabul Id. (S).....	H. O. 2117; June, 1903.	1909, Sept. 29	7.00 a. m.	<i>fms.</i>	Co.....
D. 5589	Mabul Id. (NW.) N. 3° W., 2.8 miles (4° 12' 10" N., 118° 38' 08" E.).do.....do.....	7.16 a. m. 7.44 a. m.	260	fine gy. S., gy. M.
D. 5590	Mabul Id. (NW.) N. 22° W., 4.3 miles (4° 10' 50" N., 118° 39' 35" E.).do.....do.....	8.33 a. m. 9.02 a. m.	310	gn. M., S.....
D. 5591	Mabul Id. (NW.) N. 6° W., 3.1 miles (4° 11' 48" N., 118° 38' 20" E.).do.....do.....	10.54 a. m.	260
D. 5592	Silungan Id. (M.) N. 1° W., 6.4 miles (4° 12' 44" N., 118° 27' 44" E.).do.....do.....	3.33 p. m. 4.00 p. m.	305	gn. M.....
D. 5593	Mt. Putri (sea tangent) Borneo, N. 52° W., 17.2 miles (4° 02' 40" N., 118° 11' 20" E.).	B. A. 2099; Apr., 1895.do.....	7.25 p. m. 7.34 p. m.	38	fine S.....
.....	Tawao River.....	B. A. 2576; Oct., 1882, cor. to Aug., 1905.	Sept. 30	9.30 a. m.	M., S.....
D. 5594	Mt. Putri (sea tangent) S. 82° E., 5.9 miles (4° 14' 20" N., 117° 53' 12" E.).	B. A. 2099; Apr., 1895.do.....	7.24 p. m.	11
.....	Silimpopon River.....	Oct. 2	8.00 a. m.
	<i>Off Zamboanga, Mindanao, P. I.</i>					
D. 5595	Zamboanga Lt. N. 31° W., 0.1 mile (6° 54' 00" N., 122° 04' 30" east).	C. S. 4645; July, 1907.	Oct. 6	7.13 p. m.	9
D. 5596do.....do.....	Oct. 10	6.00 p. m.	9
D. 5597do.....do.....	Oct. 12	11.45 a. m.	9
D. 5598do.....do.....do.....	3.10 p. m.	9
D. 5599do.....do.....do.....	6.20 p. m.	9
	<i>North of Celebes.</i>					
D. 5600	Menado (town) S. 58° E., 68 miles (2° 05' 00" N., 123° 52' 30" E.).	H. O. 1727; Apr., 1909.	Nov. 7	7.06 p. m.
.....	Talisse Id., east.....	B. A. 930; May, 1866, cor. to May, 1907.	Nov. 9	6.00 a. m.	Co.....
.....	Limbe Strait, vicinity of Strait Id.	Nov. 10	4.30 p. m. 6.00 a. m. 1.00 p. m.	Co. S., Co. S., Co.....
	<i>Gulf of Tomini, Celebes.</i>					
.....	Kema (town).....	B. A. 1727.	Nov. 13	8.45 a. m.	S.....
D. 5601	Limbe Id. (NE.), N., 20.7 miles (1° 13' 10" N., 125° 17' 05" E.).do.....do.....	1.15 p. m. 2.18 p. m.	765	S., Glob., Ptr.....
D. 5602	Gorontalo pier, N., 7.1 miles (0° 22' 00" N., 132° 03' 30" E.).	B. A. 942a; Oct., 1868, cor. to Mar., 1906.	Nov. 14	9.01 a. m. 10.15 a. m.	962	gy. M.....
D. 5603	Gorontalo pier N. 6° W., 5.7 m. (00° 24' 00" N., 123° 03' 45" E.).do.....	Nov. 15	1.12 p. m. 2.37 p. m.	803	S.....
D. 5604	Bilatu (town), N. 26° W., 8.7 miles (0° 22' 30" N., 122° 42' 30" E.).do.....do.....	7.25 p. m.
.....	Dodepo and Pasejogo Ids.	B. A. 900; Mar., 1901; cor. to Mar., 1907.	Nov. 16	8.00 a. m.	Co.....

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.								
					dyn.....	7-25 ft..	<i>h. m.</i> 5 00		<i>mi.</i>	15 shots.
81	82	45.7			Luc. sdr. (a). 9' Tnr.; m. b..	botm..	20	S. 49° E..	2.0	
82	83	44.3			Luc. sdr. (a). 9' Tnr.; m. b..	botm..	21	S. 55° E..	2.1	
84	84				9' Tnr.....	botm..	21	S. 58° E..	1.8	Depth estimated from dredging wire angle.
83	85	43.3			Luc. sdr. (a). 9' Tnr.....	botm..	10	N. 65° E..	.7	
84	83				Tnr. sdr. (a). 9' Tnr.....	botm..	15	West.....	1.4	Frame badly bent.
					dyn.....		8 30			
76	83				int. 4.....	surface.	1 37			Ship at anchor. Net badly torn.
					dyn.....		9 00			
80	80				int. 4.....	surface.	10 50			Ship at anchor.
80	81				int. 4.....	do..	12 30			Do.
83	82				int. 4.....	do..	2 20			Do.
85	82				int. 4.....	do..	3 00			Do.
84	82				int. 5.....	do..	11 15			Do.
80	82				int. 4.....	surface.	26			No bearings ob- tainable.
					dyn.....	10-18 ft.	5 30			16 shots.
					dyn.....	8-10 ft..	1 30			2 shots.
					dyn.....	8-10 ft..	4 30			11 shots.
					dyn.....	8-15 ft..	4 00			12 shots.
					380' seine.....	7 ft....	2 00			2 hauls.
81	83				Luc. sdr. (a). 12' Agz.; m. b.	botm..	21	S. 29° E..	1.8	
81	84				Luc. sdr. (a). 12' Agz.....	botm..	20	S.....	2.0	Net torn; bridle ropes torn loose.
84	84				Luc. sdr. (a). 12' Agz.....	botm..	13	E.....	1.0	One bridle stop carried away.
83	83				int. 4.....	surface.	25			No bearings ob- tainable.
					dyn.....	8-20 ft..	4 00			18 shots.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>Gulf of Tomini, Celebes—Con.</i>						
D. 5605	Dodepo Id. (W.) N. 14° W., 5.9 miles (0° 21' 33" N. 121° 34' 10" E.).	B. A. 900; Mar., 1901; c o r. t o Mar., 1907.	1909. Nov. 16	9.27 a. m. 10.25 a. m.	fms. 647
.....	Papajatu (Celebes).....do.....do.....	2.00 p. m.	M., Co.....
.....	Sadaa Id., north.....do.....	Nov. 17	6.00 a. m.	Co., R., S.....
D. 5606	Dodepo Id. (W.) N. 3° W., 10.8 miles (0° 16' 28" N., 121° 33' 30" E.).do.....do.....	9.09 a. m. 10.07 a. m.	834	gn. M.....
.....	Binang Unang Id., east.....	B. A. 942a; Oct., 1868, c o r. t o Mar., 1906.do.....	4.00 p. m.	Co., S.....
D. 5607	Binang Unang Id. (E.) S. 36° E., 5 miles (0° 04' 00" S., 121° 36' 00" E.).do.....	Nov. 18	8.25 a. m. 9.20 a. m.	761	fne. S.....
D. 5608	Binang Unang Id. peak, S. 87° E., 19 miles (0° 08' 00" S., 121° 19' 00" E.).do.....do.....	12.48 p. m. 2.02 p. m.	1,089	gy. M.....
D. 5609	Binang Unang Id. (N) N. 80° E., 21 miles (00° 11' 00" S., 121° 16' 00" E.).do.....do.....	3.37 p. m. 4.51 p. m.	1,092	gn. M.....
.....	Togian Bay, Togian Id.....do.....	Nov. 19	7.45 a. m.	Co.....
D. 5610	Batu Daka Id. (S.) N. 87° W., 20.9 miles (0° 36' 00" S., 122° 01' 00" E.).do.....do.....	3.59 p. m. 4.50 p. m.	678	gy. M.....
D. 5611	Buka Buka Id. (E.) S. 43° W., 6.4 miles (0° 40' 30" S., 121° 50' 00" E.).do.....do.....	7.14 p. m.
D. 5612	Buka Buka Id. (E.) S. 3° E., 7 miles (0° 38' 00" S., 121° 45' 40" E.).do.....	Nov. 20	6.04 a. m. 7.22 a. m.	750
.....	Buka Buka Id., north.....do.....do.....	9.15 a. m.	Co.....
D. 5613	Buka Buka Id. (E.) S. 28° 4 miles (0° 42' 00" S., 121° 44' 00" E.).do.....do.....	10.16 a. m. 11.14 a. m.	752	gy. M.....
.....	Malibagu Pt. (Celebes).....do.....	Nov. 21	10 00 a. m.	Co.....
<i>Molucca Passage.</i>						
D. 5614	Tifori Id. (C.) N. 19° E., 30.5 miles (0° 31' 00" N., 125° 58' 45" E.).	B. A. 942a; Oct., 1868, c o r. t o Mar., 1906.	Nov. 22	6.44 a. m. 7.58 a. m.	1,100	gy. M., S., Glob.....
D. 5615	Tifore Id. (C.) N. 40° W., 35 miles (0° 32' 30" N., 126° 31' 30" E.).do.....do.....	1.16 p. m. 2.37 p. m.	1,021	G.....
D. 5616	Tifore Id. (C.) N. 62° W., 50 miles (0° 36' 00" N., 126° 52' 20" E.).do.....do.....	6.44 p. m.
<i>Dodinga Bay, Gillolo Id.</i>						
.....	Tidore Id., north.....	B. A. 942a; Oct., 1868, c o r. t o Mar., 1906.	Nov. 24	8.00 a. m.	Co.....
.....	Maitara Id., north.....do.....	Nov. 26	8.15 a. m.	Co.....
D. 5617	Ternate Id. (SE.) S. 45° W., 7 miles (00° 49' 30" N., 127° 25' 30" E.).do.....	Nov. 27	10.42 a. m. 11.01 a. m.	131
H. 4934	Ternate Id. (SE.) S. 33° W., 7.8 miles (0° 51' 00" N., 127° 25' 10" E.).do.....do.....	11.37 a. m.	139	S., Lav.....

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Tide.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
82	82				Luc. sdr. (a) 12' Agz.	botm ..	21	S. 63° W ..	1.7	Net slightly torn.
					dyn.	15-20 ft.	1			2 shots.
					dyn.	10-20 ft.	1			10 shots.
83	83				Luc. sdr. (a) 12' Agz.	botm ..	20	S. 28° E ..	2.5	
					dyn.	10-12 ft.	2			11 shots.
81	83				Luc. sdr. (a) 12' Agz.	botm ..	20	S. 50° W ..	1.5	
80	82	36.3			Luc. sdr. (a) 12' Agz.	botm ..	20	S. 40° W ..	3.5	
83	83	36.3			Luc. sdr. (a) 12' Agz.	botm ..	33	S. 39° E ..	2.0	
					dyn.	5-18 ft.	3 30			Do.
84	87				Luc. sdr. (a) 12' Agz.	botm ..	27	N. 63° W ..	2.0	
83	84				Int. 4	surface	20			
80	83				Luc. sdr. (a) 12' Agz.	botm ..	22	S. 5° E ..	1.5	Therm., sounding cup, stray line and lead, and 70 fms. wire lost.
					dyb.	5-15 ft.	3 00			21 shots.
85	84				Luc. sdr. (c) 12' Agz.	botm ..	19	N. 20° E ..	1.8	
					dyn.	10-20 ft.	1 30			7 shots.
82	84				Luc. sdr. (c) 12' Agz.	botm ..	12	N. W ..	1.5	Shot failed to detach. Bridle stop carried away; net torn.
84	84				Luc. sdr. (c) 12' Agz.	botm ..	20	S. W ..	1.5	
80	84				int. 4 §.	20-30 fms.	18 2			
					dyn.	6-18 ft.	4 00			8 shots.
					dyn.	8-18 ft.	3 45			13 shots.
84	84				Luc. sdr. (c) 12' Agz.	botm ..	10	N. 71° W ..	1.0	
					Tnr. sdr. (e)					

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>Molucca Passage.</i>						
D. 5618	Mareh Id., S. 69° E., 7.8 miles (0° 37' 00" N., 127° 15' 00" E.).	B. A. 942a; Oct., 1868; cor. to Mar., 1906.	1909. Nov. 27	2.07 p. m. 2.44 p. m.	<i>fms.</i> 417	gy. M.
D. 5619	Mareh Id. (S.) S. 78° E., 7 miles (0° 35' 00" N., 127° 14' 40" E.).do.....do.....	3.36 p. m. 4.12 p. m.	435	fne. gy. S., M.
D. 5620	Makyan Id. (S.), S. 44° E., 7 miles (0° 21' 30" N., 127° 16' 45" E.).do.....	Nov. 28	5.48 a. m. 6.24 a. m.	358	gy. M.
<i>Between Gillolo and Makyan islands.</i>						
.....	Makyan Id. (SE.).....	B. A. 942a; Oct., 1868; cor. to Mar., 1906.	Nov. 28	8.30 a. m.	S., Co
D. 5621	Makyan Id. (S.), N. 54° W., 3 miles (0° 15' 00" N., 127° 24' 35" E.).do.....do.....	9.21 a. m. 9.50 a. m.	298	gy. and bk. S. (m. b.)
.....	Powati Anchorage (Makyan).	B. A. 912; Mar., 1885; cor. to Oct., 1906.	Nov. 29	6.00 a. m.	S., Co
D. 5622	Makyan Id. (NE.), N. 66° W., 4.1 miles (0° 19' 20" N., 127° 28' 30" E.).	B. A. 942a; Oct., 1868; cor. to Mar., 1906.do.....	7.36 a. m.	275	gy. M.
D. 5623	Makyan Id. (S.), S. 88° W., 7.5 miles (0° 16' 30" N., 127° 30' 00" E.).do.....do.....	8.03 a. m. 8.56 a. m. 9.22 a. m.
D. 5624	Makyan Id. (S.), N. 67° W., 8.9 miles (0° 12' 15" N., 127° 29' 30" E.).do.....do.....	10.30 a. m. 10.58 a. m.	288	fne. S., M.
<i>Between Gillolo and Kayoa islands.</i>						
.....	Kayoa Id. (northeast).....	B. A. 942a; Oct., 1868; cor. to Mar., 1906.	Nov. 29	1.30 p. m.	Co
D. 5625	Kayoa Id. (SE.), S. 3° W., 6 miles (0° 07' 00" N., 127° 28' 00" E.).do.....do.....	1.49 p. m. 2.16 p. m.	230	gy. M., fne. S.
D. 5626	Kayoa Id. (SE.), S. 5° W., 6.7 miles (0° 07' 30" N., 127° 29' 00" E.).do.....do.....	3.09 p. m. 3.34 p. m.	265	gy. M., fne. S.
D. 5627	Kayoa Id. (SE.), S. 15° E., 4.5 miles (0° 06' 00" N., 127° 26' 00" E.).do.....do.....	6.02 p. m.	22	M.
<i>Patiente Strait and southward.</i>						
D. 5628	St. Lamo Id. (SE.), N. 9° W., 7 miles (0° 28' 30" S., 127° 45' 00" E.).	B. A. 942a; Oct., 1868; cor. to Nar., 1906.	Nov. 30	11.22 a. m. 12.45 p. m.	1,291	gy. M.
.....	Gane (Gillolo).....	B. A. 912; Mar., 1885; cor. to Oct., 1906.	Dec. 1	8.00 a. m.	mrgn. Co., S.
D. 5629	Doworra Id. (S.), S. 62° W., 6 miles (0° 50' 00" S., 128° 12' 00" E.).	B. A. 942a; Oct., 1868; cor. to Mar., 1906.	Dec. 2	6.14 a. m. 6.43 a. m.	205	co. S.
.....	Doworra Id. (south).....do.....do.....	8.00 a. m.	Co.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
82	84				Luc. sdr. (c) 12' Agz.	botm...	20	S. 13° W..	2.0	
83	84				Luc. sdr. (c) 12' Agz.	botm...	29	S. 22° E..	1.8	
80	82				Luc. sdr. (c) 12' Agz.	botm...	21	South....	1.0	
					dyn.....	8-18ft..	3 00			17 shots.
81	84				Luc. sdr. (c) 12' Agz.; m. b.	botm...	20	S. 28° E..	1.2	
					dyn.....	10-20ft.	1 00			4 shots.
					Luc. sdr. (c).....					
80	83				12' Agz.; m. b.	botm...	21	S. 10° E..	1.0	
81	83				Luc. sdr. (c) 12' Agz.	botm...	20	South....	1.0	
83	83				Luc. sdr. (c) 12' Agz.	botm...	20	S. 15° E..	1.5	
					dyn.....	8-30ft..	3 00			20 shots.
83	84				Luc. sdr. (c) 12' Agz.	botm...	21	S. 5° W..	1.8	
84	84				Luc. sdr. (c) 12' Agz.	botm...	18	West.....	1.0	
83	83				hand lead int. 4.	5 fms.	11 40			Ship at anchor.
					Luc. sdr. (c).....					Stray line carried away.
86	84				12' Agz.	botm...	20	S. 20° E..	2.5	One bridle stop carried away.
					dyn.....	10-25ft.	7 00			24 shots.
80	83				Luc. sdr. (c) 12' Agz.	botm...	02			Dredge frame runner badly bent; lead rope broken; bridle stops lost.
					dyn.....	10-20ft.	3 30			13 shots.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>South of Patiente Strait.</i>						
D. 5630	Doworra Id. (N.), N. 3° W., 4.5 miles (0° 56' 30" S., 128° 05' 00" E.).	B. A. 942a, Oct., 1868; cor. to Mar., 1906.	1909. Dec. 2	8.51 a. m. 9.36 a. m.	fms. 569	co. S., M
D. 5631	Doworra Id. (N.), N. 38° E., 10.5 miles (0° 57' 00" S., 127° 56' 00" E.).	do.	do.	1.11 p. m. 2.16 p. m.	809	gn. M. (in net)
D. 5632	Selang Pt. (Bachian Id.), N. 56° W., 12.5 miles (1° 00' 00" S., 127° 50' 00" E.).	do.	do.	4.12 p. m. 5.08 p. m.	845	
D. 5633	Selang Pt., N. 24° W., 11.8 miles (1° 03' 00" S., 127° 44' 00" E.).	do.	do.	7.14 p. m.		
<i>Pitt Passage.</i>						
D. 5634	Gomomo Id. (E.), N. 41° E., 3 miles (1° 54' 00" S., 127° 36' 00" E.).	B. A. 942a, Oct., 1868; cor. to Mar., 1906.	Dec. 3	6.27 a. m. 7.02 a. m.	329	
D. 5635	Gomomo Id. (S.), Gomomo Id. (E.), N. 14° W., 2.5 miles (1° 53' 30" S., 127° 39' 00" E.).	do.	do.	8.15 a. m. 9.24 a. m. 9.56 a. m.	400	co. S hrd Co., R., soapstone.
D. 5636	Gomomo Id. (E.), N. 46° W., 6 miles (1° 55' 00" S., 127° 42' 30" E.).	do.	do.	11.51 a. m. 1.18 p. m.	1,262	gy. M., fine. S.
<i>Bouro Id. (south) and vicinity.</i>						
.....	Uki Id.	B. A. 942a; Oct., 1868, cor. to Mar., 1906.	Dec. 9	8.00 a. m.		mrgrn. Co.
.....	Uki River	do.	do.	8.00 a. m.		
.....	Uki Id.	do.	do.	1.00 p. m.		S., R.
D. 5637	Ambblau Id. (N.), N. 80° E., 21 miles (3° 53' 20" S., 126° 48' 00" E.).	do.	Dec. 10	7.06 a. m. 7.57 a. m.	700	gy. M.
.....	Tifu Bay (Bouro Id.)	do.	do.	1.00 p. m.		S., M., R., Co.
H. 4935	Tifu Bay entrance (W.), N. 4° E., 2.2 miles (3° 46' 15" S., 126° 24' 40" E.).	do.	do.	1.30 p. m.	198	
D. 5638	Tifu Bay entrance (W.), N. 17° E., 3.2 miles (3° 47' 15" S., 126° 23' 40" E.).	do.	do.	2.00 p. m. 2.36 p. m.	517	fine. gy. S
.....	Tomahu Id.	do.	Dec. 11	1.00 p. m. 7.00 p. m.		Co., S.
<i>Molucca Sea.</i>						
D. 5639	Cape Pamali (Wowoni Id.), (N.), S. 77° W., 27 miles (3° 54' 50" S., 123° 27' 20" E.).	B. A. 3616; May, 1907.	Dec. 13	5.23 a. m. 7.11 a. m.	1,560	gy. M.
<i>Buton Strait.</i>						
D. 5640	Labuan Blanda Id., N. 88° E., 1 mile (4° 27' 00" S., 122° 55' 40" E.).	B. A. 3470; Apr., 1906.	Dec. 13	5.02 p. m. 5.10 p. m.	24	S., brk. Sh.
D. 5641	Labuan Blanda Id. (S.) Kalono Pt. (W.), N. 61° W., 3.4 miles (4° 29' 24" S., 122° 52' 30" E.).	do.	Dec. 14	6.00 a. m. 9.30 a. m. 9.41 a. m.	39	mrgrn. Co. S., Sh.
D. 5642	Tikola Peninsula (N.), N. 38° W., 6.5 miles (4° 31' 40" S., 122° 49' 42" E.).	do.	do.	10.50 a. m. 11.00 a. m.	37	gy. M.
.....	Great Tobea Id.	do.	do.	1.00 p. m.		S., Co.
.....	Pendek Id., north	do.	Dec. 15	3.15 p. m.		Co., S.
D. 5643	Pendek Id. (N.), S. 77° E., 1.7 miles (5° 11' 45" S., 122° 42' 36" E.).	do.	do.	3.42 p. m. 4.06 p. m.	215	gn. M.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.				<i>h. m.</i>		<i>mi.</i>		
82	84				Luc. sdr. (c). 12' Agz.	botm...	24	S. S. W.	1.8	
84	86				Luc. sdr. (c). 12' Agz.	botm...	20	N. by W.	1.5	Sounding cup lost.
83	85				Luc. sdr. (c). 12' Agz.	botm...	22	S. E. by E.	2.0	
82	84				int. 4.	surface.	19			No bearings obtainable.
81	84				Luc. sdr. (c). 12' Agz.	botm...	13	S.W. by S.	1.0	
					dyn.	6-20ft.	7 30			23 shots.
82	83				Luc. sdr. (c). 12' Agz.	botm...	05	S. S. E.	.5	Bridle stops lost; frame bent.
83	83				Luc. sdr. (c). 12' Agz.	botm...	20	S. by E.	2.5	
					dyn.	10-30ft.	6 00			19 shots.
					dyn.		9 00		7.0	
					75' seine	12 ft	3 30			9 hauls.
79	83				Luc. sdr. (c). 12' Agz.	botm...	12	S. 21° W.	1.3	Net fouled on bottom. 18 shots.
					dyn.	2-20ft.	3 15			
					Luc. sdr. (c).					
84	86				Luc. sdr. (c). 12' Agz.	botm...	20	S. 78° E.	1.0	
					dyn.	3-15ft.	4 30			13 shots.
					dip; e. l.		1 30			
82	84				Luc. sdr. (c). 9' Agz. rev.	botm...	31	N. 36° W.	.8	
84	84				Tnr. sdr. (e). 12' Agz.	botm...	12	N. 52° W.	.3	
					dyn.	5ft.	1 45			5 shots.
83	84				Tnr. sdr. (e). 12' Agz.	botm...	17	S. 81° W.	.6	
84	85				Tnr. sdr. (e). 12' Agz.	botm...	17	N. 75° W.	1.4	
					dyn.	5-18 ft.	3 30			11 shots.
					dyn.	15-25 ft.	1 45			12 shots.
82	84				Luc. sdr. (c). 12' Agz.	botm...	17	S. 45° W.	.7	

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
<i>Buton Strait—Continued.</i>						
D. 5644	Makasser Id. (E.), N. 4° E., 1.3 miles (5° 27' 24" S., 122° 38' 00" E.).	B. A. 3470; Apr., 1906.	1909. Dec. 16	8.02 a. m.	fms. 22	
D. 5645	North Id. (N.E.), S. 10° W., 1.6 miles (5° 29' 06" S., 122° 36' 06" E.).do.....do.....	9.37 a. m. 9.54 a. m.	206	
D. 5646	North Id. (S.), S. 68° E., 7.5 miles (5° 31' 30" S., 122° 22' 40" E.).	B. A. 3616; May, 1907.do.....	11.36 a. m. 12 10 p. m.	456	gn. M.
D. 5647	North Id. (S.), S. 87° E., 11.6 miles (5° 34' 00" S., 122° 18' 15" E.).do.....do.....	2.07 p. m. 2.44 p. m.	519	gn. M.
D. 5648	North Id. (S.), N. 87° E., 10.2 miles (5° 35' 00" S., 122° 20' 00" E.).do.....do.....	3.47 p. m. 4.29 p. m.	559	gn. M.
D. 5649	North Id. (S.), N. 87° E., 22 miles (5° 36' 00" S., 122° 07' 36" E.).do.....do.....	7.23 p. m.		
<i>Gulf of Boni.</i>						
.....	Basa Id.	B. A. 3616; May, 1907.	Dec. 17	8.00 a. m.		Co. tide pools.
D. 5650	Lamulu Pt., N. 5° W., 12.5 miles (4° 53' 45" S., 121° 29' 00" E.).do.....do.....	8.34 a. m. 9.22 a. m.	540	gn. M.
D. 5651	Buginkali Pt., S. 67° E., 21 miles (4° 43' 50" S., 121° 23' 24" E.).do.....do.....	1.39 p. m. 2.32 p. m.	700	gn. M.
D. 5652	Lamulu, S. 36° E., 7.5 miles (4° 35' 00" S., 121° 23' 06" E.).do.....do.....	4.39 p. m. 5.24 p. m.	525	gn. M.
D. 5653	Lamulu, S. 40° E., 18 miles (4° 27' 36" S., 121° 16' 36" E.).do.....do.....	7.23 p. m.		
D. 5654	C. Tabako, N. 17° E., 21.5 miles (3° 42' 00" S., 120° 45' 50" E.).do.....	Dec. 18	5.41 a. m. 6.47 a. m.	805	
D. 5655	Labuandata Bay, C. Tabako, N. 7° E., 13 miles (3° 34' 10" S., 120° 50' 30" E.).do.....do.....	9.00 a. m. 10.20 a. m. 11.00 a. m.	608	Co., S. gy. M., fine S.
H. 4936	C. Tabako, N. 47° E., 9 miles (3° 28' 00" S., 120° 45' 40" E.).do.....do.....	1.40 p. m.	667	gy. M.
D. 5656	Olang Pt., N. 67° W., 14.5 miles (3° 17' 40" S., 120° 36' 45" E.).do.....	Dec. 19	7.36 a. m. 8.37 a. m.	484	gy. M.
D. 5657	Olang Pt., N. 61° W., 15.5 miles (3° 19' 40" S., 120° 36' 30" E.).do.....do.....	10.29 a. m. 11.08 a. m.	492	gy. M.
D. 5658	C. Loko Loko, S. 31° W., 12 miles (3° 32' 40" S., 120° 31' 30" E.).do.....do.....	1.38 p. m. 2.23 p. m.	510	gy. M.
D. 5659	C. Lassa, S. 78° W., 19 miles (5° 33' 20" S., 120° 47' 10" E.).do.....	Dec. 20	6.10 a. m. 6.57 a. m.	702	S. M.
<i>Flores Sea.</i>						
H. 4937	C. Lassa, S. 78° W., 20.5 miles (5° 32' 50" S., 120° 49' 10" E.).	B. A. 3616; May, 1907.	Dec. 20	8.12 a. m.	885	gy. M.
D. 5660	C. Lassa, S. 88° W., 20.5 miles (5° 36' 30" S., 120° 49' 00" E.).do.....do.....	9.14 a. m. 10.05 a. m.	692	gy. M., S.
D. 5661	C. Lassa, N. 21° E., 12.5 miles (5° 49' 40" S., 120° 24' 30" E.).do.....do.....	4.05 p. m. 4.24 p. m.	180	hrd.
D. 5662	Tana Keke Id. (W.), N. 17° W., 12.5 miles (5° 43' 00" S., 119° 18' 00" E.).	B. A. 2637, June, 1885; cor. to Oct., 1904.	Dec. 21	5.40 a. m. 6.12 a. m.	211	
.....	Tana Keke Id. (S.)do.....do.....	8.30 a. m.		Co.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.


Temperatures.		Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F.	° F.	° F.	° F.						
80	83			hand lead		<i>h. m.</i>		<i>mi.</i>	
				12' Agz.	botm.	11	S. 81° W.	.4	
79	83			Luc. sdr. (c)					
				12' Agz.	botm.	01	N. 34° W.	.7	
79	83			Luc. sdr. (c)					
				12' Agz.	botm.	20	East	1.1	
83	83			Luc. sdr. (c)					
				12' Agz.	botm.	20	S. 40° E.	1.0	
83	83	39.2		Luc. sdr. (c)					
				12' Agz.	botm.	23	S. 55° E.	.8	
83	83			int. 4	surface	21			No bearings obtainable.
				dyn.	12-20 ft.	3 00			10 shots.
				copper sulphate		3 00			
84	84	40.1		Luc. sdr. (c)					
				12' Agz.	botm.	10	S. 45° W.	.7	Bridle stops carried away.
85	84	38.7		Luc. sdr. (c)					Sounding cup carried away.
				12' Agz.	botm.	20	N. 11° W.	2.9	
84	84	41.2		Luc. sdr. (c)					
				12' Agz.	botm.	20	N. 61° W.	2.1	
82	82			int. 4	surface	20			No bearings obtainable.
79	83	38.3		Luc. sdr. (c)					
				12' Agz.	botm.	28	N. 1° W.	2.0	
84	84	39.2		dyn.	5-18 ft.	3 00			12 shots.
				Luc. sdr. (c)					
				12' Agz.	botm.	20	S. 45° E.	1.5	
				Luc. sdr. (c)					Therm. failed to register.
80	83	41.2		Luc. sdr. (c)					
				12' Agz.	botm.	05	S. 41° W.	1.8	
82	84	41.3		Luc. sdr. (c)					
				12' Agz.	botm.	20	S. 19° W.	2.0	
83	85	41.2		Luc. sdr. (c)					
				12' Agz.	botm.	20	S. 35° E.	1.2	
83	82	39.0		Luc. sdr. (c)					
				12' Agz.	botm.	21	S. 62° E.	1.0	
		38.2		Luc. sdr. (c)					
83	83	39.2		Luc. sdr. (c)					
				12' Agz.	botm.	20	S. 58° E.	1.8	
86	83	50.5		Luc. sdr. (c)					
				12' Agz.	botm.	03	N. 50° E.	1.1	Net torn below lead line.
82	83	48.8		Luc. sdr. (c)					
				12' Agz.	botm.	20	West	1.8	No bottom specimen.
				dyn.	9-18 ft.	2 45			16 shots.

DREDGING AND HYDROGRAPHIC RECORDS OF THE U. S. FISHERIES

Station No.	Position.	Chart.	Date.	Time of day.	Depth.	Character of bottom.
	<i>Macassar Strait.</i>					
D 5663	Kapoposang Id. (E.), N. 11° E., 1.7 miles (4° 43' 22" S., 118° 57' 35" E.).	Dutch 123; Sept., 1901.	1909. Dec. 27	7.20 p. m.	fms. 10
D. 5664	Kapoposang Lt., N. 66° E., 3.8 miles (4° 43' 22" S., 118° 53' 18" E.).do.....	Dec. 28	9.09 a. m. 9.43 a. m.	400	hrd.
D. 5665	Kapoposang Lt., S. 40° E., 18.8 miles (4° 27' 00" S., 118° 44' 00" E.).	B. A. 2637; June, 1885, cor.to Oct., 1904.do.....	1.51 p. m. 2.59 p. m.	1,008	M.
.....	Libani Bay, Celebes (W.).....do.....	Dec. 29	8.00 a. m.	Co.
D. 5666	Onkona Pt., S. 1° W., 11 miles (2° 54' 30" S., 118° 47' 00" E.).do.....do.....	8.39 a. m. 9.18 a. m.	272	gn. M.
D. 5667	Onkona Pt., S. 5° W., 11 miles (2° 56' 00" S., 118° 47' 30" E.).do.....do.....	9.55 a. m. 10.25 a. m.	367	gy. S., M.
D. 5668	Mamuju Id. (E.), S. 31° E., 10.6 miles (2° 28' 15" S., 118° 49' 00" E.).do.....do.....	3.41 p. m. 4.45 p. m.	901	gy. M.
D. 5669	Mamuju Id. (E.), S. 14° E., 18.5 miles (2° 19' 30" S., 118° 50' 00" E.).do.....do.....	7.25 p. m.
D. 5670	Chenoki Pt., S. 60° E., 40 miles (1° 19' 00" S., 118° 43' 00" E.).	B. A. 941b, Nov., 1867; cor.to Aug., 1907.	Dec. 30	7.03 a. m. 8.18 a. m.	1,181	gy. M.
D. 5671	Chenoki Pt., S. 31° E., 42.5 miles (1° 05' 00" S., 118° 56' 00" E.).do.....do.....	12.41 p. m. 1.45 p. m.	960	gy. M.
D. 5672	Dongala Lt., S. 80° E., 54 miles (0° 29' 00" S., 118° 51' 00" E.).	B. A. 2636; Apr., 1878, cor.to Apr., 1907.do.....	7.26 p. m.
.....	Birabirahan (west).....	B. A. 941b; Nov., 1867, cor.to Aug., 1907.	Dec. 31	8.45 a. m.	Co.
	<i>Trusan Tando Bulong, B. N. Borneo.</i>					
.....	Daisy Islet, 4° 27' 53" N., 118° 38' 25" E.	H. O. 2117; June, 1903.	1910. Jan. 6	1.45 p. m.	Co.
	<i>Sulu Sea.</i>					
.....	Doc Can Id., southwest.....	C. S. 4722.....	Jan. 7	10.15 a. m.	S., Co.
	<i>China Sea.</i>					
.....	Kwa Siang Bay, Formosa.....	Jan. 25	8.30 a. m.
.....	So Wan Bay, Formosa.....	Jan. 29	7.30 a. m.

STEAMER ALBATROSS IN THE PHILIPPINE ISLANDS, 1907-1910—Continued.

Temperatures.			Density.		Apparatus.	Trial.		Drift.		Remarks.
Air.	Surface.	Bottom.	Surface.	Bottom.		Depth.	Duration.	Direction.	Distance.	
° F. 83	° F. 84	° F.			hand line int. 4.	surface.	<i>h. m.</i> 10 40		<i>mi.</i>	Ship at anchor.
81	84	43.3			Luc. sdr. (c) 12' Agz.	botm.	21	S. 67° W.	2.5	No bottom sample in net. No bearings ob- tainable.
80	82				Luc. sdr. (c) 12' Agz.	botm.	05	SW	2.0	Entire net carried away on bottom 20 shots.
80	82	47.5			dyn Luc. sdr. (c) 12' Agz.	6-18 ft. botm.	3 30 12			
82	83	41.7			Luc. sdr. (c) 12' Agz.	botm.	20	N. 34° W.	1.5	
81	83	38.2			Luc. sdr. (c) 12' Agz.	botm.	19	S. 47° E.	2.8	Shot did not de- tach.
83	84				int. 4.	surface.	24	North.	1.0	
82	82	38.2			Luc. sdr. (c) 12' Agz.	botm.	20	South.	2.0	Shot did not de- tach. One bridle stop parted.
83	84	38.2			Luc. sdr. (c) 12' Agz.	botm.	23	S. 63° E.	2.0	
82	83				int. 4.	surface.	20	N. 10° W.		No bearings ob- tainable.
					dyn.	10-20 ft.	2 15			12 shots.
					dyn.	10-15 ft.	45			6 shots.
					dyn.	10-30 ft.	1 00			10 shots.
					dyn.	10-25 ft.	3 00			13 shots.
					dyn.	10-30 ft.	3 30			27 shots.



CONDITION AND EXTENT OF THE NATURAL OYSTER BEDS OF DELAWARE

BY H. F. MOORE

Assistant, U. S. Bureau of Fisheries

Bureau of Fisheries Document No. 745

CONTENTS.

	Page.
Introduction.....	3
Methods of the survey.....	4
Description of oyster grounds.....	7
Bombay bed.....	7
Thrum-cap bed.....	8
Over-the-Bar bed.....	9
Patches between Over-the-Bar and Sand beds.....	10
Sand bed.....	11
Leipsic Rock.....	11
Bed north of Silver bed.....	12
Between Silver bed and Simons Creek.....	12
Silver bed.....	13
Lumps between Silver and Ridge beds.....	14
Drum bed.....	15
Ridge bed.....	16
Small beds northeast of Ridge bed.....	18
Old bed.....	19
Outside of Old bed.....	19
Scattered patches between Ridge and Southwest beds.....	20
Southwest bed.....	20
Stone bed.....	22
East Line bed.....	22
Flogger bed.....	23
The beds in summary.....	23
Physical and biological conditions.....	27
Tides and currents.....	27
Salinity of the water.....	27
Enemies of the oyster.....	28

CONDITION AND EXTENT OF THE NATURAL OYSTER BEDS OF DELAWARE.

By H. F. MOORE,
Assistant, United States Bureau of Fisheries.

INTRODUCTION.

At the solicitation of the Delaware Oyster Survey Commission the Bureau of Fisheries during the summer of 1910 undertook a survey of the natural oyster beds of Delaware Bay within the jurisdiction of the State of Delaware. The State, which was making a survey of the planted beds under the supervision of Mr. C. C. Yates, of the United States Coast and Geodetic Survey, furnished the triangulation and made a small appropriation for the payment of two temporary employees during part of the work, but the Bureau of Fisheries furnished all other personnel, in addition to launches, boats, and equipment.

The steamer *Fish Hawk* was detailed for the work from June 1 to July 10, though, owing to unexpected delays in securing a launch able enough for the execution of hydrography in the open waters of the bay, she did not actually reach the field of operations until June 18. Part of the civilian personnel was ordered to the ship on May 26, in order to have the equipment in readiness for the anticipated commencement of work on June 1, on which date the entire party was assembled.

The purpose of the survey was the accurate location and charting of the natural oyster beds and the investigation of their present condition and productiveness. No previous survey or investigation of the beds of this region has been made, and although their approximate location is known to the local oystermen with reference to certain more or less indefinable natural landmarks, it is difficult for them to indicate, even roughly, their general position on the charts. Concerning some of the beds, and especially the southern extension of Flogger bed, the information obtained from the various sources was extremely contradictory.

METHODS OF THE SURVEY.

The methods employed were those pursued in former surveys of like character, and are explained in detail in a description of the beds of the James River,¹ from which some of the following is repeated:

A "boat sheet" was prepared, on which were accurately platted the positions, as determined by triangulation, of lighthouses and the towers erected as shore signals. These data were furnished by the State and were based on a development of the triangulation employed in the survey of the planted or leased beds.

The oyster beds were discovered by soundings with a lead line, but principally by means of a length of chain dragged over the bottom at the end of a copper wire running from the sounding boat. The wire was wound on a reel and its unwound length was adjusted to the depth of water and the speed of the launch, so that the chain was always on the bottom. Whenever the chain touched a shell or an oyster the shock or vibration was transmitted up the wire to the hand of a man whose sole duty it was to give heed to such signals and report them to the recorder.

The launches from which the soundings were made were run at a speed of between 3 and 4 miles per hour, usually on ranges ashore to insure the rectitude of the lines. At intervals of three minutes—in some cases two minutes—the position of the boat was determined by two simultaneous sextant observations of the angles between a set of three signals, the middle one of which was common to the two angles, the position being immediately platted on the boat sheet. At regular intervals of twenty seconds, as measured by a clock under the observation of the recorder, the leadsman made a sounding and reported to the recorder the depth of water and the character of the bottom, immediately after which the man at the wire reported the character of the chain indications since the last sounding—that is, whether they showed barren bottom or dense, scattering, or very scattering growths of oysters.

With the boat running at 3 miles per hour the soundings were between 80 and 90 feet apart, and, as the speed of the boat was uniform, the location of each was determinable within a yard or two by dividing the platted distance between the positions determined by the sextant by the number of soundings. The chain, of course, gave a continuous indication of the character of the bottom, but the record was made at the regular twenty-second intervals observed in sounding.

The chain, while indicating the absence or the relative abundance of objects on the bottom, gives no information as to whether they are shells or oysters, nor, if the latter, their size and condition. To obtain these data it was necessary to supplement the observations

¹ Moore, H. F.: Condition and extent of the oyster beds of James River, Virginia. Bureau of Fisheries Document No. 729.

already described by others more definite in respect to the desired particulars. Whenever, in the opinion of the officer in charge of the sounding boat, such information was required, a numbered buoy was dropped, the time and number being entered in the sounding book. Another launch, following the sounding boat, anchored alongside the buoy, and a quantity of the oysters and shells were tonged up, separated by sizes, and counted.

This boat at each station made a known number of "grabs" with the oyster tongs, exercising care to clean the bottom of oysters as thoroughly as possible at each grab. In a given depth of water and using the same boat and tongs, an oysterman will cover practically the same area of the bottom at each grab, but, other factors remaining the same, the area of the grab will decrease with an increase in the depth.

Careful measurements were made and tabulated showing the area per grab covered by the tonger employed on the work at each foot of depth of water and for each pair of tongs and boat used. With these data, and knowing the number of "grabs," the number of oysters of each size per square yard of bottom was readily obtainable by simple calculation. The following example will illustrate the data obtained and the form of the record:

DEPARTMENT OF COMMERCE AND LABOR.	
BUREAU OF FISHERIES.	
FIELD RECORD OF EXAMINATIONS OF OYSTER BEDS.	
General locality, <i>Delaware Bay, Delaware.</i>	
Local name of oyster ground, <i>Over-the-Bar.</i>	
Date, <i>July 9, 1910.</i> Time, <i>8.50 a. m.</i>	
Angle, <i>B 146-B 147.</i> Buoy No. <i>6.</i>	
Depth, <i>18 feet.</i> Bottom, <i>soft.</i>	
Condition of water, <i>clear.</i>	
Density, <i>1.008.</i> Temperature, <i>25° C.</i>	
Current, Stage of tide, <i>one hour flood.</i>	
Tongman, <i>M. A. Duffield.</i>	
No grabs made, <i>8.</i> Tongs, <i>20 feet.</i>	
Total area covered, <i>2.5 sq. yds.</i>	
No oysters taken $\left\{ \begin{array}{l} 1 \text{ in.}, 13. \quad 1 \text{ in.}-3 \text{ in.}, 129. \\ 3 \text{ in.}-4 \text{ in.}, 59. \quad 4 \text{ in.}, 11. \end{array} \right.$	
Quantity shells, <i>14.</i>	
Result $\left\{ \begin{array}{l} \text{Spat per square yard, } 5.2. \\ \text{Culls per square yard, } 51.6. \\ \text{Counts per square yard, } 28.0. \end{array} \right.$	

This furnishes an exact statement of the condition of the bed at a spot which can be platted on the chart with error in position of not more than a few yards. From the data obtained a close estimate may be formed of the number of bushels of oysters and shells per acre in the vicinity of the examination and, by multiplying the observations, for the bed as a whole. In the course of the survey 590 observations were made at various places, principally on the natural rocks, but some on the barren bottoms also.

In estimating the relative productiveness of the bottoms it appeared advisable to depart from the methods employed in the James River survey on account of the difference in the conditions under which the industry is prosecuted. Where tongs are used exclusively, a bed with a given quantity of oysters lying in shoal water is more valuable, commercially, than one with the same quantity of oysters in deep water, owing to the fact that the labor of the tonger is more efficient on the former. As has been pointed out, the area covered by a "grab" decreases with the depth, other factors being the same, and moreover the deeper the water the greater is the labor involved in making the grab and the smaller is the number of grabs which can be made in a given time.

In Delaware Bay, while there is a certain amount of tonging during the fall and at such times as the weather will permit in winter and early spring, the most important and productive fishing is by means of dredges, the use of which is permitted from April 15 to June 30, inclusive. In dredging, the effects of varying depths of water, within reasonable limits, are practically negligible so far as the catch is concerned. The time required for winding in from deep water is greater than from shallow water, but as the dredge is approximately equally efficient whatever the depth, and as the difference in the time required in winding is small as compared with the period during which the dredge is on the bottom, the factor of depth, so important in tonging, is practically inconsiderable.

The classification adopted in this report is as follows:

Depleted bottom.....	Less than 25 bushels per acre.
Very scattering growth.....	Between 25 and 75 bushels per acre.
Scattering growth.....	Between 75 and 150 bushels per acre.
Dense growth.....	Over 150 bushels per acre.

As the region is important for the production of seed rather than market oysters, all sizes are included in the estimates of the density of oyster growth, but all loose shells and other debris commonly dredged are excluded. "Depleted bottom" is not necessarily that which was formerly productive but now practically barren, but is merely an expression of the present impoverishment of the bed without respect to its past. In some cases it may be a formerly barren area slowly coming into productiveness.

The bottom rated as bearing a "very scattering growth" is the least productive bottom capable of furnishing a livelihood to the dredgers.

In the course of the survey 16,435 acres, or over 25 square miles, were explored with sounding lines and chains. Of this area 2,144 acres were found to be included in oyster beds of varying degrees of productiveness. In the survey the chain was dragged over 124 miles of the bottom, soundings were made at 5,772 places, and the position of the boat was instrumentally determined at 819 points.

DESCRIPTION OF OYSTER GROUNDS.

BOMBAY BED.

This is the northernmost public oyster bed within the confines of Delaware. Its northern limit is opposite the upper pier at Woodland Beach, and its southern end is a little below the small creek known locally as Tombstone. Its inner or southwestern edge is from 200 to 400 yards from shore, the average width of the bed is about one-fourth mile, and the total length slightly in excess of 1 mile.

The estimated area, density of growth, and contents of the bed are as follows:

OYSTER GROWTH ON BOMBAY BED.

Character of oyster growth.	Area.	Oysters per acre.			Estimated content of oysters.
		Under 3 inches.	Over 3 inches.	Total.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense	111	250	115	365	40,515
Scattering	12	103	23	126	2,512
Very scattering.....	6	22	5	27	162
Depleted.....	26	0	0	0	0
Total.....	155				43,189

The dense area comprises a broad strip running along the entire inshore edge of the bed. The scattering areas are two, the larger lying near the middle of the outer edge of the bed and the smaller, a very narrow strip, on the offshore edge of the lower end. Both merge more or less gradually into the dense area with which they are continuous. The area of very scattering growth is a small patch situated near the offshore part of the upper end of the bed, in the midst of the depleted bottom. The latter appears to be a formerly moderately productive area which has become covered by a deposit of mud and now produces no oysters, although there are numerous buried shells lying on a hard bottom about 6 inches beneath the present surface. This bed differs from all others of the region treated in this report in being founded on a stony bottom, a considerable proportion of the oysters taken being attached to rock fragments. The oysters are in small clusters, with thin, sharp shells. Small oysters predominate, not only numerically but by measure. No drills were found and, reasoning from the low salinity of the water, probably do not occur. The specific gravity of the water at the time of examination, July 10, 1910, was about 1.005, and it is likely that the bed suffers periodically during freshets. The average depth of water is about 8 to 10 feet.

It was reported that there were oysters between the piers, but none were found, although there were a few attached to the piling and lying on the bottom in its vicinity.

The details of the examination of this bed are shown in the following table:

DETAILS OF EXAMINATIONS OF BOMBAY BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
	1910.	<i>Feet.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
183.....	July 10..	10	Dense.....	1.6	35.2	2.6	129	26	155
184.....	do.....	10	do.....	10.5	42.0	12.6	184	126	310
189.....	do.....	10	do.....	11.0	65.8	19.5	269	195	464
191.....	do.....	11	do.....	15.8	34.2	3.2	175	32	207
192.....	do.....	10	do.....	28.4	54.2	20.5	299	205	504
195.....	do.....	12	do.....	9.5	17.9	9.5	96	95	191
197.....	do.....	12	do.....	52.0	58.4	13.7	387	137	524
199.....	do.....	11	do.....	74.2	57.4	10.0	461	100	561
194.....	do.....	12	Scattering.....	11.0	12.6	3.7	83	37	120
198.....	do.....	12	do.....	35.2	0.0	1.0	123	10	133
186.....	do.....	10	Very scattering.....	0.0	6.3	0.5	22	5	27
185.....	do.....	11	Depleted.....	0.0	0.0	0.0	0	0	0
187.....	do.....	10	do.....	0.0	0.0	0.0	0	0	0
188.....	do.....	10	do.....	0.0	0.0	0.0	0	0	0
190.....	do.....	11	do.....	0.0	0.0	0.0	0	0	0

THRUM-CAP BED.

For a distance of about 5 miles below Bombay bed the bottom is reported to be barren, with the possible exception of a few patches of insignificant size, and it was not deemed warrantable to incur the expense of an examination.

Thrum-cap bed is a somewhat triangular area lying about 1 mile offshore opposite the small stream known to the oystermen as Hay Ditch. It covers an area of about 78 acres, of which it is estimated 6 are covered by a dense growth, 14 by scattering, and 55 by very scattering, and 3 acres are characterized by a total absence of oysters, but with scattered shells buried in the mud.

The areas of dense and scattering growth form a narrow strip on the inshore edge of the bed, with the denser area at the upper end. The bottom covered with very scattering growth stretches in gradually decreasing productiveness from the outer edge of this strip toward the deeper water. The depleted area is a small patch where the dense growth shades off into the surrounding barren bottom. The depth of water on the bed varies from about 18 feet at the inshore edge to 22 feet on the outer border.

It is estimated that the bed contained at the time of examination 4,195 bushels of oysters of all sizes, of which the dense area bore 1,164 bushels, the scattering 1,106 bushels, and the very scattering 1,925 bushels.

There were comparatively few dead oysters, and no indications of the presence of drills were observed. In July the specific gravity of the water varied from about 1.003 at low water to 1.011 at high tide.

The results of the examinations of this bed are shown in the following table:

DETAILS OF EXAMINATIONS OF THRUM-CAP BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
	1910.	<i>Fect.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
179.....	July 9	19	Dense.....	4.4	28.4	8.0	114	80	194
181.....	do.....	20	Scattering.....	5.5	12.2	1.7	62	17	79
178.....	do.....	22	Very scattering.....	1.7	3.3	1.7	18	17	35
180.....	do.....	19	Depleted.....	0.0	0.0	0.0	0	0	0

OVER-THE-BAR BED.

This bed, like the preceding, from which it is separated by a distance of a little over one-eighth of a mile, lies just beyond the edge of the shifting sands, which extend to about the 12-foot curve. It is about 1½ miles from shore, and takes its name from its position some distance outside of a long sand bar, which, according to the navigational charts, is covered by about 4 feet of water at low tide, but on which the present survey found water a little deeper. The depth on the bed itself varies from 15 to 20 feet.

The extent and general condition of the bed in July, 1910, is shown in the following table:

OYSTER GROWTH ON OVER-THE-BAR BED.

Character of oyster growth.	Area.	Oysters per acre.			Estimated content of oysters.
		Under 3 inches.	Over 3 inches.	Total.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	109	103	162	275	29,975
Very scattering.....	15	41	0	41	615
Depleted.....	39	0	0	0	0
Total.....	163				30,590

The dense growth is found on two areas, 41 and 68 acres in extent, respectively, separated by a depleted area containing nothing but buried shells. The upper area is long and narrow and contains a large preponderance of oysters over 3 inches long. The northern end of the lower area is similar, with four or five times as many large oysters as small ones, but in the southern the two are in approximately equal quantity, and the average of both sizes is about 335 bushels per acre. The area of very scattering growth is found at the inshore edge of the southern part of the bed, and was apparently formed by a recent strike on a previously depleted area. The three depleted areas lie at the ends and the middle of the bed, the latter in reality separating the rock into two distinct parts. The depleted

bottom bears no oysters and but few exposed shells and, apparently, has been formed either by the silting of sparsely productive bottom or by shells dragged by dredging from the rock on to the adjacent muddy bottom.

The oysters throughout the entire bed are long, narrow, sharp-edged, and inferior in quality, and are almost invariably in clusters, whose bases are buried in soft mud. The bottom throughout is soft, and there is apparent nowhere any depth of shell deposits such as are found on Silver bed and the Ridge.

The details of the examinations made on this bed are shown in the following table:

DETAILS OF EXAMINATIONS OF OVER-THE-BAR BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
	1910.	<i>Fect.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
169.....	July 9	18	Dense.....	15.2	28.8	14.8	154	148	302
170.....	do	18	do.....	12.0	28.0	8.4	140	84	224
171.....	do	18	do.....	5.2	51.6	28.0	198	250	478
172.....	do	20	do.....	9.2	5.6	10.4	52	104	156
174.....	do	19	do.....	0.8	9.2	15.6	35	156	191
177.....	do	21	do.....	5.6	6.1	26.1	41	261	302
167.....	do	17	Very scattering.....	0.8	10.8	0.0	41	0	41
168.....	do	18	Depleted.....	0.0	0.0	0.0	0	0	0
173.....	do	20	do.....	0.0	0.0	0.0	0	0	0
176.....	do	20	do.....	0.0	0.0	0.0	0	0	0
182.....	do	21	do.....	0.0	0.0	0.0	0	0	0

PATCHES BETWEEN OVER-THE-BAR AND SAND BEDS.

In the area between these beds are several small scattered patches of oysters, but two of which were examined to determine their character. One of these has an area of about 16 acres and is estimated to contain about 1,000 or 1,200 bushels of oysters. The other is about 5 acres in extent and contains probably about 200 bushels of oysters. On both beds and probably on other small patches in the vicinity the oysters are long, thin, and narrow, and are found in scattered clusters.

The following table exhibits the data obtained from the examinations:

DETAILS OF EXAMINATIONS OF PATCHES BETWEEN OVER-THE-BAR BED AND SAND BEDS:

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
	1910.	<i>Fect.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
160.....	July 8	19	Very scattering.....	0	2.8	3.2	10	32	42
162.....	do	15	do.....	0	3.0	6.3	11	63	74

SAND BED.

Sand bed lies nearly north of the Ridge and northeast of Silver bed, being separated from the latter by a distance of about one-third of a mile. It covers an area of about 54 acres, of which 16 acres are covered by a dense growth of oysters and 11 acres by a scattering growth, the remaining 27 acres being depleted.

The productive bottom forms a zone along the inner edge of the bed, the southern and middle portions bearing the denser growth. The depleted bottom occupies the outer half of the bed. It is estimated that the bed contained about 4,600 bushels of oysters of all sizes at the time of examination, and that of these 3,700 bushels were on the area of dense growth, 700 bushels on the very scattered growth, and 200 bushels on the depleted bottom. Oysters over 3 inches long preponderated on the productive portions of the bed, but were inferior in quantity on the depleted area.

The oysters are superior in shape to those found on the bars north of this, being in smaller clusters and rounder. Dead oysters were comparatively few, and no indications of the drill were noted.

Several boats were observed working on Sand bed during the latter part of June, and it is reported that the bed was dredged to some extent earlier in the season.

The following examinations were made:

DETAILS OF EXAMINATIONS OF SAND BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
	1910.	<i>Fect.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
154.....	July 8	20	Dense.....	1.2	24.4	14.4	90	144	234
159.....	do.....	19	Very scattering.....	4.0	1.2	4.8	18	48	66
155.....	do.....	18	Depleted.....	0.0	0.0	0.0	0	0	0
156.....	do.....	19	do.....	0.4	2.4	0.4	10	4	14
157.....	do.....	19	do.....	1.6	0.4	6	4	10

LEIPSIK ROCK.

This is a small but exceedingly prolific bed lying in the mouth of Leipsic Creek within one-eighth of a mile of the shore. It is approximately circular in outline and consists of about 4 acres of very dense growth. It is estimated that the bed bears nearly 3,000 bushels of oysters, practically none of which is over 3 inches in length, and it is probable that it represents a recent rejuvenescence of an old bed. There is a deep deposit of shells forming the core of the bed, but around the edges this is covered by a deposit of mud which appears to be encroaching on and causing a gradual contraction of the productive area. It is probable that the oysters are subject to periodical destruction from fresh water and mud carried by freshets.

So far as could be learned the rock has not been worked for several years.

The following examinations were made:

DETAILS OF EXAMINATIONS OF LEIPSIC ROCK.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
	1910.	<i>Feet.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
140.....	July 7	11	Dense.....	41.0	114.5	1.4	544	14	558
144.....	do.....	12	do.....	0.0	14.8	0.4	52	4	56
145.....	do.....	10	do.....	118.0	300.0	1.6	1,460	16	1,476

BED NORTH OF SILVER BED.

North of the western end of Silver bed and separated from it by about one-eighth of a mile of soft bottom in which scattering shells are buried is a nameless bed covering about 25 acres. There are about 8 acres covered by scattering growth estimated to contain about 900 bushels of oysters and about 17 acres of very scattering oysters containing about 750 bushels. The northern part of the bed, which bears the heaviest growth, has a substratum of shells, but the southern edge lies on sandy bottom. The proportion of large oysters is greater than on Silver bed.

The following observations were made:

DETAILS OF EXAMINATIONS OF BED NORTH OF SILVER BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
	1910.	<i>Feet.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
158.....	July 8	14	Scattering.....	4.5	6.7	7.8	39	78	117
152.....	do.....	13	Very scattering.....	4.0	2.2	2.2	22	22	44

BETWEEN SILVER BED AND SIMONS CREEK.

Almost continuous with Silver bed and stretching for a distance of nearly one-half of a mile toward the mouth of Simons Creek is a bed of about 17 acres lying on the mud and sand. Its most productive area is nearest Silver bed, and the opposite end is bare except of scattered shells. The best part, about 5 acres in extent, bears a scattering growth of oysters estimated to contain about 375 bushels, and the area of very scattering growth which adjoins it bears about the same quantity on its 7 acres. The depleted bottom is practically bare at present, but is in a condition to catch a small set under favorable conditions.

The following table shows the results of examinations:

DETAILS OF EXAMINATIONS OF BED BETWEEN SILVER BED AND SIMONS CREEK.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
166.....	1910. July 8	<i>Fect.</i> 9	Scattering.....	<i>No.</i> 2.9	<i>No.</i> 3.4	<i>No.</i> 5.4	<i>Bu.</i> 22	<i>Bu.</i> 54	<i>Bu.</i> 76
165.....do.....	9	Very scattering.....	0.0	4.3	3.7	15	37	52
111.....	June 29	14	Depleted.....	0.0	0.0	0.0	0	0	0

SILVER BED.

This bed, which is said to derive its name from the silvery color of the shells found on the hard rock, is, excepting the Ridge, the largest and most important natural bed in Delaware. It lies about 1 mile east of the mouth of Dona River, locally known as Simons Creek. The bed has a maximum extent of about a mile east and west and slightly over a half mile north and south, and it lies in a depth of water varying from 8 to 12 feet.

The following table shows its general extent and condition in July, 1910:

OYSTER GROWTH ON SILVER BED.

Character of oyster growth.	Area.	Oysters per acre.			Estimated content of oysters.
		Under 3 inches.	Over 3 inches.	Total.	
Dense.....	<i>Acres.</i> 65	<i>Bushels.</i> 171	<i>Bushels.</i> 74	<i>Bushels.</i> 245	<i>Bushels.</i> 15,925
Scattering.....	20	82	27	109	2,180
Very scattering.....	45	25	21	46	2,070
Depleted.....	140	8	2	10	1,400
Total.....	270				21,575

The most productive parts of the bed lie in its northeast half and include a belt of dense and scattering growth about one-half mile long and varying from one-eighth to one-third mile in width.

A considerable part of the bottom covered by the bed is macadamized with a dense accumulation of shells, or probably two such areas separated by a belt of muddy bottom. In places the bottom was so hard with compacted shells and so smooth that a boat anchor would not take hold. Although this bed is not now raised above the surrounding barren bottom, it is probable that it originally formed a knoll, the crest of which has been cut away by dredging and tonging.

The area of dense growth lies in a compact body occupying the middle of the eastern half of the bed, gradually merging with two

small areas of scattering growth at the northwest and southeast ends, respectively. There is a third area of scattering growth near the western end of the bed. The very scattering growth forms a zone around the western and part of the southern side of the more prolific bottom, lying on a substratum of compacted shells. Most of the western half of the bed is composed of depleted bottom, which also extends as a narrow strip around practically the entire circumference of the rest of the bed, the bottom being generally hard and shelly with occasional patches of mud.

In general the present condition of the bed indicates a former greater extent of productive bottom. There is every indication that it has been closely dredged during the past season, and the present content of oysters is probably but a small proportion of the quantity on the bottom at the beginning of the season. The shells are in excellent condition to receive a set of spat, and under favorable circumstances the bed should speedily recuperate. There were comparatively few dead oysters, and drills or borers do not appear to be troublesome.

The following observations were made:

DETAILS OF EXAMINATIONS OF SILVER BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
	1910.	<i>Feet.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
58.....	June 25	14	Dense.....	23.7	65.0	12.2	310	122	432
110.....	June 29	14	do.....	5.9	21.1	6.7	94	67	161
149.....	July 8	13	do.....	1.5	29.6	7.8	109	78	187
163.....	do.....	10	do.....	8.7	40.0	2.9	170	29	199
55.....	June 25	14	Scattering.....	12.2	7.4	1.9	69	19	88
147.....	July 8	14	do.....	1.4	22.2	3.3	83	33	116
164.....	do.....	9	do.....	4.3	22.9	2.9	95	29	124
59.....	June 25	13	Very scattering.....	0.4	3.3	1.2	13	12	25
100.....	June 27	11	do.....	2.4	7.9	1.7	36	17	53
150.....	July 8	13	do.....	2.6	4.5	3.3	25	33	58
52.....	June 25	Depleted.....
53.....	do.....	13	do.....	0.3	4.5	0.0	17	0	17
60.....	do.....	13	do.....	0.0	0.0	0.0	0	0	0
98.....	June 27	11	do.....	0.0	2.8	0.3	10	3	13
99.....	do.....	11	do.....	1.4	1.0	0.7	8	7	15
109.....	June 29	14	do.....	0.0	0.0	0.0	0	0	0
148.....	July 8	13	do.....	1.1	2.2	0.0	11	0	11

LUMPS BETWEEN SILVER AND RIDGE BEDS.

Lying between Sand and Silver beds on the north and Ridge and Drum beds on the south are a number of small lumps and patches surrounded by a considerable area of barren bottom. Eight of these areas were located by the survey, most of them covering areas of 3 or 4 acres, and there are probably a number of others, as on account of their small size and irregular distribution but little time was spent in

looking for them. But three of these places were examined in detail, and their location may be determined by an inspection of the chart. One of them was about 3 acres in extent and was estimated to contain about 2,500 bushels of long, sharp-edged oysters in large clusters, growing on a soft, muddy bottom. The other two spots examined bore a very scattering growth. The largest of these, about one-fourth mile inshore of the upper end of Drum bed, was estimated to be about 8 acres in extent and to contain about 300 bushels of oysters. The other, just south of the middle of Silver bed, has an area of about 4 acres and contained at the time of examination about 120 bushels of oysters.

The five areas located but not examined varied in extent from about 1 to 14 acres, and are situated variously. They are shown on the chart as unshaded places surrounded by red lines. Judging from the chain readings none of them is particularly productive.

The following observations were made in this region:

DETAILS OF EXAMINATIONS OF LUMPS BETWEEN SILVER AND RIDGE BEDS.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
	1910.	<i>Feet.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
96.....	June 27	18	Dense.....	1.4	28.0	75.2	103	752	855
86.....	do.....	15	Very scattering.....	0.7	4.1	1.9	17	19	36
97.....	do.....	13	do.....	0.0	2.2	2.2	8	22	30

DRUM BED.

Drum bed lies west of and very close to the depleted edge of the ridge and about 1 mile from shore. It has a length of over one-half mile, a width of about one-fourth mile, and a total area of approximately 68 acres. Its condition and the relative extent of oyster growths of the several degrees of productiveness are shown in the following table:

OYSTER GROWTH ON DRUM BED.

Character of oyster growth.	Area.	Oysters per acre.			Estimated content of oysters.
		Under 3 inches.	Over 3 inches.	Total.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	16	139	83	222	3,552
Scattering.....	21	30	65	95	1,995
Very scattering.....	19	32	18	50	950
Depleted.....	12	1	6	7	84
Total.....	68				6,581

The most prolific part of the bed is an area about one-fourth mile square extending across its middle, consisting of an area of dense growth flanked on each side by one bearing a scattering growth. The northern end of the bed is composed of a gradually narrowing area of very scattering growth, and there is a small patch of similar character at the inside corner of the southern end.

The depleted bottom is in two patches, one adjoining the scattering and very scattering growths at the lower end and the other interposed between the dense scattering and very scattering oyster deposits just above the middle. The bottom is soft on the areas of very scattering growth and on part of the northernmost depleted area, but is elsewhere hard and shelly.

Small oysters exceed in quantity those over 3 inches long, excepting on the area of scattering growth, where there are about twice as many large as small ones. Loose shells are in fair abundance and of a character to catch a good set under favorable conditions.

The following observations were made:

DETAILS OF EXAMINATIONS OF DRUM BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
	1910.	<i>Fect.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
47.....	June 25	16	Dense.....	27.2	18.8	3.6	161	36	197
48.....	do.....	16½	do.....	16.0	30.0	7.6	161	76	237
107.....	June 29	17	do.....	4.0	23.2	13.6	95	136	231
51.....	June 25	17	Scattering.....	7.2	13.2	3.2	71	32	103
84.....	June 27	17	do.....	1.4	10.0	8.2	4	82	86
106.....	June 29	18	do.....	1.6	2.8	8.2	15	82	97
105.....	do.....	18	Very scattering.....	1.6	9.2	1.6	37	16	53
108.....	do.....	17	do.....	0.0	7.6	2.0	27	20	47
73.....	June 27	15	Depleted.....	0.0	0.4	0.4	1	4	5
85.....	do.....	17	do.....	0.0	0.0	0.8	0	8	8

RIDGE BED.

The Ridge bed, known to the oystermen as "The Ridge," is at present the most important natural bed in Delaware, and during the period of the present survey it sustained by far the heaviest dredging. During the latter half of June numerous vessels were at work daily and until the end of the month, when the dredging season closed, there appeared to be a fair catch.

The Ridge lies about 1½ miles from the nearest shore, midway between Dona River and Mahon River. It is triangular in shape, with a deep indentation or slough of muddy bottom projecting deeply into its base at the southern end. It has an extent of slightly over 1 mile north and south and its southern end is almost of equal extent east and west. It has a total area of 371 acres and the most productive bottom, that which is rated in this report as bearing dense and

scattering growths, stretches from the northern apex to about the middle of the bed, where it divides into two limbs astride the slough before alluded to.

It is evident that this bed, like Silver bed, is an old one, and without doubt its central portions, those which now bear the heaviest growth of oysters, were formerly elevated above the surrounding bottom to form a shoal or ridge which has been pulled down and in large part carried away by the oystermen, particularly the dredgers, until at present the water over it shoals but little as compared with the surrounding barren areas. The great deposit of shells which originally existed has been taken up and the bottom so denuded that in places the originally underlying mud has been brought to the surface. Many little patches of bare mud were found where there was every reason to expect a deposit of shells and oysters and it was apparent that the bed was being overworked.

The general condition and extent of the bed at the end of June, 1910, is shown in the following table:

OYSTER GROWTH ON RIDGE BED.

Character of oyster growth.	Area.	Oysters per acre.			Estimated content of oysters.
		Under 3 inches.	Over 3 inches.	Total.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	49	160	23	183	8,967
Scattering.....	86	96	25	121	10,406
Very scattering.....	65	36	21	57	3,705
Depleted.....	171	4	1	5	855
Total.....	371				23,933

The dense areas are two in number, separated by an area of scattered growth. The smaller of these areas lies at the northern apex of the bed and the larger one is a long belt along most of its eastern side. More or less soft mud is to be found in the former, especially near its upper edge, but the latter rests on a solid substratum of shells.

The lower end of the larger dense area gradually verges into a small spot of scattering growth, but most of the bottom bearing a growth of this character is embraced in a long, somewhat S-shaped strip running from near the northern end of the bed almost to its southwest corner. The northern end, especially between and adjacent to the dense growths, is most productive.

The very scattering growth is all confined to the southern edge of the bed, most of it being between the mud slough and the dense and scattering growth. Excepting close to the more productive areas there is much muddy bottom in this area. Most of the depleted

bottom lies on the west side of the bed, but there is a narrow strip along the eastern edge and embracing the southern end of the dense and scattering growth. Much of the depleted area is in reality denuded or barren, and although most of it lies on hard bottom there are numerous muddy spots, especially near the southern edge.

On this bed as a whole and especially on the more productive areas small oysters are in great preponderance. In many cases there were quantities of oysters so small that they fell between the teeth of the tongs.

The following observations were made on this bed:

DETAILS OF EXAMINATIONS OF RIDGE BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
	1910.	<i>Feet.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
67.....	June 26	17	Dense.....	20.0	20.0	3.9	140	39	179
91.....	June 27	18	do.....	31.2	21.2	1.6	183	16	199
93.....	do.....	16	do.....	29.1	16.0	1.6	158	16	174
62.....	June 26	16	Scattering.....	19.2	7.6	2.0	94	20	114
65.....	do.....	16	do.....	16.0	12.0	4.0	98	40	138
69.....	do.....	18	do.....	17.2	13.6	2.8	108	28	136
92.....	June 27	16	do.....	28.8	4.4	2.0	116	20	136
101.....	do.....	15	do.....	7.4	10.7	1.9	63	19	82
61.....	June 26	14	Very scattering.....	4.8	1.1	0.7	21	7	28
63.....	do.....	15	do.....	2.2	3.7	5.2	21	52	73
79.....	June 27	16	do.....	0.8	10.4	2.4	39	24	63
90.....	do.....	17	do.....	12.4	5.2	0.0	62	0	62
64.....	June 26	16	Depleted.....	0.0	0.0	0.0	0	0	0
66.....	do.....	16	do.....	1.2	2.0	0.0	11	0	11
70.....	do.....	17	do.....	2.0	2.4	0.0	15	0	15
80.....	June 27	15	do.....	0.0	0.0	0.0	0	0	0
81.....	do.....	14	do.....	2.6	0.7	0.4	12	4	16
82.....	do.....	16	do.....	0.0	0.0	0.0	0	0	0
83.....	do.....	17	do.....	0.4	0.4	0.0	3	0	3
87.....	do.....	16	do.....	0.0	0.0	0.0	0	0	0
88.....	do.....	16	do.....	0.0	0.0	0.0	0	0	0
89.....	do.....	16	do.....	0.0	0.0	0.0	0	0	0
102.....	do.....	13	do.....	0.0	0.0	0.0	0	0	0
103.....	do.....	13	do.....	0.0	0.4	1.1	1	11	12

SMALL BEDS NORTHEAST OF RIDGE BED.

Northeast of the Ridge is a small patch of about 7 acres of very scattering growth which is estimated to contain about 200 bushels of oysters, most of them over 3 inches in length.

The following results were obtained from an examination of this area:

DETAILS OF EXAMINATIONS OF SMALL BEDS NORTHEAST OF RIDGE BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
	1910.	<i>Feet.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
94.....	June 27	18	Very scattering.....	0	1.4	2.4	5	24	29

OLD BED.

Old bed lies close to the southeastern edge of the Ridge, from which it is separated by a narrow strip of mud with many buried shells. It is stated that the dredgers sometimes haul across the barren bottom from one bed to the other.

The condition and extent of the bed as determined by the survey were as follows:

OYSTER-GROWTH ON OLD BED.

Character of oyster growth.	Area.	Oysters per acre.			Estimated content of oysters.
		Under 3 inches.	Over 3 inches.	Total.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Very scattering.....	20	40	2	42	840
Depleted.....	17	10	3	13	221
Total.....	37				1,061

Although the bed is at present not very productive it has the appearance of former greater value. It lies on a dense bed of shells and is undoubtedly the remnant of an old accumulation. There are very few large oysters to be found, but the young growth is fair in places and the conditions for a new set are good. The bed evidently has been subjected to severe dredging.

The following observations were made:

DETAILS OF EXAMINATIONS OF OLD BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Feet.</i>		<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>Bu.</i>	<i>Bu.</i>	<i>Bu.</i>
76.....	1910. June 27	16	Very scattering.....	10.4	2.8	0.0	46	0	46
130.....	June 30	17	do.....	6.0	1.4	0.0	26	0	26
131.....	do.....	19	do.....	0.8	5.6	0.8	22	8	30
132.....	do.....	19	do.....	10.4	11.6	0.0	77	0	77
133.....	do.....	19	do.....	4.8	4.4	0.0	32	0	32
74.....	June 27	17	Depleted.....	0.0	2.0	0.0	7	0	7
75.....	do.....	18	do.....	1.2	2.0	0.8	11	8	19
78.....	do.....	17	do.....	0.0	1.2	0.4	4	4	8
134.....	June 30	20	do.....	0.6	4.4	0.0	18	0	18

OUTSIDE OF OLD BED.

Immediately outside of Old bed is an area of about 16 acres, surrounded by sand, for which the oystermen appear to have no name, if, even, they are aware of its existence. But one observation was made at this place, where a dense growth of young oysters was found. If the other parts of the bed are equally productive this patch contains about 6,800 bushels of oysters, practically all of them under 3 inches in length. The present growth is apparently of recent origin.

The following results were obtained from the examination:

DETAILS OF EXAMINATIONS OF BEDS OUTSIDE OF OLD BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
77.....	1910. June 27	<i>Feet.</i> 15	Dense.....	<i>No.</i> 35.0	<i>No.</i> 83.4	<i>No.</i> 15.0	<i>Bu.</i> 414	<i>Bu.</i> 15	<i>Bu.</i> 429

SCATTERED PATCHES BETWEEN RIDGE AND SOUTHWEST BEDS.

On the soft bottom lying between these two beds are a number of little patches of oyster growth, of which five were located with the chain and three were examined by tonging. The latter were all highly productive, and they probably represent the possibilities of oyster production in this vicinity on beds not frequented by the dredgers.

The three beds examined covered a total of 11 acres, and it is estimated that they contained about 5,300 bushels of oysters, of which nearly three-fourths were over 3 inches long. Based on the results of the examination, and assuming that the other beds found are equally productive, the five beds probably contain about 11,000 bushels, and it is probable that at least 20,000 bushels are scattered in little 2 to 5 acre patches in the vicinity.

The following table shows the data obtained from examinations:

DETAILS OF EXAMINATIONS OF SMALL SCATTERED PATCHES BETWEEN RIDGE AND SOUTHWEST BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
40.....	1910. June 22	<i>Feet.</i> 12	Dense.....	<i>No.</i> 20.3	<i>No.</i> 21.1	<i>No.</i> 21.1	<i>Bu.</i> 145	<i>Bu.</i> 211	<i>Bu.</i> 356
42.....do.....	14do.....	15.2	30.4	20.7	159	207	366
71.....	June 27	14do.....	15.2	14.4	71.5	104	715	819

SOUTHWEST BED.

Southwest bed lies in the southeastern part of the present productive natural oyster grounds of the State and its southern edge is about one-fourth mile north of the "east line" which separates the private beds from the public ones. It has a north and south extent of upward of one-half mile and a maximum width of about one-third mile, containing all told about 106 acres.

The extent and relative productiveness of the bottoms, as classified in this report, are shown in the table following.

OYSTER GROWTH ON SOUTHWEST BED.

Character of oyster growth.	Area.	Oysters per acre.			Estimated content of oysters.
		Under 3 inches.	Over 3 inches.	Total.	
	Acres.	Bushels.	Bushels.	Bushels.	Bushels.
Dense.....	11	40	744	784	8,624
Scattering.....	8	99	48	147	1,376
Very scattering.....	31	18	13	31	961
Depleted.....	56	4	1	5	280
Total.....	106				11,241

The area of dense growth is near the southern end of the bed and is flanked on the east and west sides by a very scattering growth, and on the north and south by depleted bottom. Most of the oysters are over 3 inches long and they appear to be in numerous small patches on the soft mud. The place has the appearance of bottom which has been overlooked by the oystermen and may as a whole be somewhat smaller in area than is indicated in the preceding table.

The bottom bearing scattering growth lies at the northeast edge of the bed and at its southwestern limits merges into a strip of very scattering growth running along the western edge of the bed as far as the densely covered bottom first described. There is another small patch of very scattering growth near the southeast corner of the bed.

The depleted bottom lies in three patches, one at each end of the bed and the other at the middle of the eastern edge.

Although it is not known whether Southwest bed was dredged during the past season, it bears every evidence that it has been over-worked. Excepting on the small area of dense growth there are few marketable oysters, and bare or almost bare muddy spots are of frequent occurrence. Many oysters had been killed by drills and many of these animals and their egg cases were found.

The following table shows the results of examinations:

DETAILS OF EXAMINATIONS OF SOUTHWEST BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Fect.</i>		No.	No.	No.	Bu.	Bu.	Bu.
126.....	1910. June 30	14	Dense.....	1.5	10.0	74.4	40	744	784
31.....	June 22	13	Scattering.....	5.6	22.7	4.8	99	48	147
121.....	June 30	13	Very scattering.....	0.4	5.6	0.4	21	4	25
122.....	do.....	12	do.....	1.5	7.5	0.4	31	4	35
128.....	do.....	15	do.....	0.0	1.4	3.0	5	30	35
129.....	do.....	14	do.....	0.0	4.4	1.5	15	15	30
32.....	June 22	13	Depleted.....	0.0	0.0	0.0	0	0	0
120.....	June 30	14	do.....	0.4	4.4	0.0	17	0	17
123.....	do.....	12	do.....	0.0	1.4	0.4	5	4	9
124.....	do.....	13	do.....	0.0	0.0	0.4	0	4	4
125.....	do.....	15	do.....	0.0	0.0	0.0	0	0	0
127.....	do.....	15	do.....	0.0	0.0	0.0	0	0	0

STONE BED.

This bed possibly takes its name from the quantity of hard, sandy worm tubes, known to the oystermen as "stone coral," which are found attached to and overgrowing the oysters. It is probable that a good many of the latter are stifled and killed by this growth, which is even more abundant on a small depleted patch lying between the Stone bed and the mouth of Mahon River.

The bed covers an area of about 33 acres of very scattering growth, on which there is an average of about 53 bushels of oysters per acre. It is estimated that about July 1, 1910, there were on the entire bed about 1,750 bushels of oysters, the large and small being in about equal quantities.

The following observations were made:

DETAILS OF EXAMINATION OF STONE BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
36.....	1910. June 22	<i>Feet.</i> 13	Very scattering.....	<i>No.</i> 0.7	<i>No.</i> 4.8	<i>No.</i> 2.6	<i>Bu.</i> 19	<i>Bu.</i> 26	<i>Bu.</i> 45
104.....	June 29	18do.....		10.0	2.4	35	24	59

EAST LINE BED.

This bed lies just at the line which marks the southern limits of the public grounds, and it appears that for that reason it has a sentimental interest to the oystermen. It has a diameter not much greater than the length of a boat and is too small to plot on the chart, on which its position is indicated by a circle.

Numerous examinations were made in its vicinity over an area of 6 or 8 acres, but at only one place were oysters found, and there they were very dense and mostly of marketable size.

The data obtained at this station are shown in the following table:

DETAILS OF EXAMINATIONS OF EAST LINE BED.

Station number.	Date of examination.	Depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
118.....	1910. June 30	<i>Feet.</i> 13	Dense.....	<i>No.</i> 0.0	<i>No.</i> 10.0	<i>No.</i> 77.8	<i>Bu.</i> 35	<i>Bu.</i> 778	<i>Bu.</i> 813

FLOGGER BED.

Flogger bed lies along Joe Flogger Shoal, which separates Blake Channel from the ship channel. As developed by the survey, it is the largest bed in Delaware, having a length of over 3 miles, an average width of about one-third mile, and an area of about 660 acres. Owing to its exposed situation and the depth of water, as well as to the contradictory information received as to its approximate location and extent, it was the most troublesome bed encountered by the survey. Lines were run across Joe Flogger Shoal from its extreme southern end, but no indications of shells or oysters were encountered until within about one-half mile of east line. From this point scattering shells were found, but when the bottom was examined with the tongs these were discovered to be more or less submerged in the sand.

The bed as outlined on the chart was located almost entirely by means of the chain. At its upper end it lies on the eastern or ship-channel side of Joe Flogger Shoal, but about a mile from its upper end it expands to the westward over an area of somewhat deeper water, and thence, to its southern end, continues on the western or Blake Channel side of the shoal. It was at this point of expansion only that oysters were found, in one small patch of very scattering growth and two or three areas of depleted bottom. The results were not of sufficient importance to exhibit in detail on the chart. It is possible that oysters are to be found in limited quantities in some of the deeper water, but the chain readings did not indicate patches of sufficient importance to warrant the expense of making dredgings. It is reported that there are oysters in some of the deep water of the ship channel, but no indications were found in such places as were examined.

It is understood that Flogger bed has not been dredged for several years, and the survey indicated that while formerly it may have been of importance commensurate with its area, it has become covered with sand throughout practically its whole extent. It may again become productive, but there is no present indication of this probability.

Oysters were reported around the buoy at the head of Flogger Shoal and at another buoy on the opposite side of Blake Channel, but a careful examination, especially in the latter place, failed to disclose them.

THE BEDS IN SUMMARY.

The oyster bottoms of Delaware all lie between Woodland Beach and the vicinity of Bowers Beach, covering an area about 21 miles long and with an average width of about 3 miles. South and west of a line running east from the old Mahon River Lighthouse and thence approximately southeasterly along Blake Channel, the bottoms are excluded from the common oyster fishery and a considerable proportion of the area is leased to private persons and firms for purposes of oyster culture.

With this area this report will not deal, as it was examined by the writer in but the most cursory manner and the survey of the private beds was being made solely as a State undertaking. It may be stated, however, that the private beds are planted partly with shells, mostly brought from points on Chesapeake Bay, but generally with seed oysters taken from the natural beds. The grounds are in large part leased or controlled by residents of Philadelphia and New Jersey, and the product is consumed principally in Philadelphia, being marketed through Maurice River Cove in New Jersey.

The natural rocks, with which alone this report is concerned, lie in a narrow strip between Blake Channel and the main ship channel on what is known as Joe Flogger Shoal, and between these channels and the Delaware shore in a belt which stretches from the east line above mentioned to about abreast of the upper pier at Woodland Beach, a distance of about 13 miles.

At its southeastern end, where it adjoins the planted area, this zone is about 3 miles in width, but it gradually narrows to the northward until at its upper extremity it is hardly one-half mile wide. The most extensive beds lie in the lower half of the zone and the most intense fishery is carried on in that region. During the time of the survey this was practically the only place in which the dredgers were operating, and we were informed that but little had been done elsewhere earlier in the season.

The following tables summarize the data of the extent, condition, and general distribution of oyster growth on the several beds previously discussed in more detail:

AREAS OF OYSTER BEDS.

Name of bed.	Character of oyster growth.					Total. Acres.
	Dense.	Scatter- ing.	Very scatter- ing.	Depleted.	Not deter- mined.	
	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.
Bombay.....	111	12	6	26		155
Thrum-cap.....	6	14	55	3		78
Over-the-Bar.....	109		15	39		163
Between Over-the-Bar and Sand.....			21			21
Sand.....	16		11	27		54
Leipsic Rock.....	4					4
North of Silver.....		8	17			25
Between Silver and Simons Creek.....		5	7	5		17
Silver.....	65	20	45	140		270
Between Silver and Ridge.....	3		12		21	36
Drum.....	16	21	19	12		68
Ridge.....	49	86	65	171		371
Northeast of Ridge.....			7			7
Old.....			20	17		37
Outside of Old.....	16					16
Between Ridge and Southwest.....	11				12	23
Southwest.....	11	8	31	56		106
Stone.....			33			33
East Line.....	(¹)					(¹)
Flogger.....					2 660	660
Total.....	417	174	364	496	693	2,144

¹ Less than 1 acre.² Practically all depleted.

ESTIMATED OYSTER CONTENT OF NATURAL BEDS, JULY 1, 1910.

Name of bed.	Character of oyster growth.					Total.
	Dense.	Scatter- ing.	Very scatter- ing.	Depleted.	Not deter- mined.	
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Bombay.....	40,515	2,512	162			43,189
Thrum-cap.....	1,164	1,106	1,925			4,195
Over-the-Bar.....	29,975		615			30,590
Between Over-the-Bar and Sand.....			1,200			1,200
Sand.....	3,700		700	200		4,600
Leipsic Rock.....	3,000					3,000
North of Silver.....		900	750			1,650
Between Sand and Simons Creek.....		375	375			750
Silver.....	15,925	2,180	2,070	1,400		21,575
Between Silver and Ridge.....	2,500		420		1,300	5,920
Drum.....	3,552	1,995	950	84		6,581
Ridge.....	8,967	10,406	3,705	855		23,933
Patch northeast of Ridge.....			200			200
Old.....			840	221		1,061
Outside of Old.....	6,800					6,800
Between Ridge and Southwest.....	5,300				1,500	20,300
Southwest.....	8,624	1,376	961	280		11,241
Stone.....			1,750			1,750
East Line.....	1,500					1,500
Flogger.....					(?)	(?)
Total.....	130,522	20,850	16,623	3,040	18,000	189,035

¹ Estimated from chain indications.

² Practically all depleted.

Combining the foregoing data, an interesting comparison may be instituted between the beds sustaining a heavy fishery with dredges and those which recently have been worked but little. According to the best information, supported by our own observations in the latter part of the season, practically all of the dredging in 1910 was on the beds south of Over-the-Bar, although a few vessels were observed apparently working on Thrum-cap. These beds, excluding Flogger, had a total area of 1,088 acres and a total estimated oyster content of 111,061 bushels, or an average of 102 bushels per acre, at the end of the season. On the beds which were reported or observed to be most severely worked the oyster content averaged considerably less than this. On the Ridge the average for the whole bed was about 60 bushels per acre, on Drum bed about 97 bushels, on Silver bed about 80 bushels, on Old bed 30 bushels, and on Southwest bed about 106 bushels, and for the five beds taken as a whole the average was about 75 bushels per acre.

The beds above and including Over-the Bar have an area of 396 acres and a total estimated content of 77,984 bushels of oysters of all sizes, or an average of 197 bushels per acre. These beds, owing to their position, are probably more subject than the lower beds to damage from freshets and are probably naturally less productive, yet they had at the time of examination an oyster growth over $2\frac{1}{2}$ times as dense. If we consider the various small patches surrounding the five beds enumerated above, which are in general too small to dredge or which, if large enough, have been overlooked during the season

recently closed, the disparity is still greater. Those which were examined by tonging had an area of 46 acres and an estimated content of 18,000 bushels of oysters, an average of nearly 390 bushels per acre, over five times the density of growth on the large beds in the vicinity.

The number of bushels taken from the beds of Delaware during the past season is not known but it was probably several hundred thousand bushels, and from the conditions found in the survey and the data just deduced it probably can be safely assumed that oysters were from three to five times as abundant at the beginning of the season as they were in its closing days when the survey was made.

This heavy draft on the beds would be less serious were it not accompanied by an abuse for which there is no excuse. In a region devoted mainly to planting and where a comparatively small quantity of oysters is marketed directly from the natural beds it is economically advisable to permit the taking of small oysters as well as large. So long as there is an abundance of shells on the bottom and a reasonable quantity of oysters is left to furnish spawn there will be, under favorable conditions of water and temperature, a more or less regular set of spat and the oyster population of the beds will be fairly maintained, although, of course, the proportion of oysters of marketable size will diminish. When, however, the beds are stripped of shells, as appears to be the case in Delaware, they will surely become depleted.

During the survey, although a number of vessels were actively dredging, no member of the party observed a boat engaged in culling. Inquiry among the oystermen elicited the information that while the boats catching seed oysters for sale generally cull their catch because the planters will not pay oyster prices for shells, the vessels owned or operated by planters when dredging on the public beds rarely do so. They are charged with carrying away everything which the dredge picks up, the shells being valuable for hardening the bottoms on their planting grounds and as cultch for catching a set of spat.

That some vessels are guilty of such behavior is within the knowledge of the writer, and moreover the charge is supported by the condition of the beds. One of the most noteworthy of the facts disclosed by the tong examinations was the small quantity of shells found as compared with similar examinations of beds in other States. On the five important beds in the vicinity of the Ridge there are less than 2 per cent as many shells per square yard as are found on the seed beds of James River, Va., where culling is strictly enforced. In places the deep pavement of shells which must have existed formerly has been completely removed and the underlying mud now shows itself in patches in the midst of the beds. A hard-worked bed to be in a healthy condition should contain an abundance of shells. The ultimate result of the continuance of this state of affairs is not difficult

to foresee. Oysters can not set on the mud. They must have some hard, clean object to which to attach when they settle down from their infantile free-swimming habit, and on the beds the old shells and the oysters themselves offer the only possibilities. If there be few or no shells the recuperation of exhausted beds is correspondingly retarded. If both shells and oysters are persistently removed, the most productive bed eventually will be hopelessly depleted.

PHYSICAL AND BIOLOGICAL CONDITIONS.

TIDES AND CURRENTS.

A staff tide gauge was established at the wharf at Mahon River Light-house and readings were taken hourly from 8 a. m. until 5 p. m. during the period of the survey. This does not furnish a very accurate plane of reference, but as the location of the gauge was central with respect to the more important beds it is sufficiently accurate for the purposes of this report. The average rise and fall of the tide between June 19 and July 10 was 5.4 feet, the minimum being 4.5 feet on July 10 and the maximum 6.3 on July 2.

No measurements of the velocity of currents were made, but in general it may be stated that they are strong throughout the region embraced in this report.

SALINITY OF THE WATER.

The salinity of the water exhibited a very considerable range within the limits covered by the survey. From June 18 to July 10 observations were made three times daily at the anchorage of the *Fish Hawk* and several times each day on the oyster beds undergoing examination. Most of the observations on the *Fish Hawk* were made at a point about 1 mile south of the east line and about 3 miles offshore, but others, fewer in number, were made near the southern limit of the planted beds, near the middle of the north and south extent of the public beds, and at the upper limit of oyster growth opposite Woodland Beach.

The data obtained are shown in the following table:

SALINITY OF WATER OVER OYSTER BEDS, JUNE 18 TO JULY 10.

Locality.	Number of observations.	Specific gravity of water corrected.			Average temperature of water.
		Maximum.	Minimum.	Average.	
Opposite Woodland Beach.....	3	1.0074	1.0032	1.0057	°F. 79
Midway between Ship John and Elbow Light-house.....	3	1.0121	1.0100	1.0107	77
3½ miles southeast by east of Mahon River Light.....	33	1.0149	1.0103	1.0136	77
6 miles east-northeast of Bowers Beach.....	6	1.0178	1.0158	1.0164	68

At the upper limit of oyster growth the salinity of the water was low at a time when there had been comparatively little rainfall, and it is probable that it may become practically fresh at this point during periods of freshet. This is without much doubt the cause inhibiting the growth of oysters at places higher up the river.

At the southern end of the planting grounds the salinity is comparatively high and in consequence it is to be expected that the drill or borer would be destructive. On the more important of the public beds, those lying between the east line and the mouth of Leipsic Creek, the density is favorable for the welfare of the oysters. It probably never falls so low as seriously to threaten the beds, and, on the other hand, it is hardly high enough, excepting close to the east line, to favor an abundance of drills.

ENEMIES OF THE OYSTER.


It is stated that schools of drumfish occasionally appear on the oyster beds of Delaware Bay and cause much damage, but none were observed during the survey. This enemy of the oyster is usually more destructive on planted beds than on the public rocks, probably because the single-culled oysters on the former are easier to crush than are the clustered, sharp-edged specimens more common on the natural beds. The inroads of the drumfish are sporadic and unexpected in most places, although on the coasts of some of the Southern States they are frequent enough to warrant the inclosure of the planted beds with wire fences. This appears to be the only really adequate protection, though if the presence of a school on the beds or in their vicinity is discovered in time it can often be driven from the neighborhood by the use of explosives.


The principal enemy to the oyster on the Delaware beds is the drill or borer, a small marine snail which drills a hole through the oyster's shell and thus gains access to the contents, which it consumes. The perforation is made by actual drilling with a rasplike organ protruded from the mouth, and so far as is known no acid or other solvent is employed to soften the shell. The drill breeds during late spring and summer, laying its eggs in vase-shaped, leathery capsules attached in clusters to shells and other hard bodies on the bottom. These capsules, each containing several eggs, are readily recognizable, being about one-fourth inch long and usually yellow in color.


In the few places examined on the planted beds there were considerable numbers of drills and many small oysters killed by them. On the public beds near the east line some drills and killed oysters were found, but over most of the area surveyed the salinity of the water is somewhat too low to permit these pests ever to become a serious factor. Below a salinity represented approximately by a mixture of equal parts of salt and fresh water, having a specific gravity of about 1.012 or 1.013, the drill will not thrive.

Although in the absence of other food the drill will attach and sometimes kill oysters of marketable size, it invariably attacks smaller ones by preference. Seed oysters 2 or $2\frac{1}{2}$ inches in diameter are comparatively immune, and in places where the drills are particularly troublesome such seed should be planted in preference to smaller. Although such is not known to be the case in Delaware, there are localities in which it is useless to plant shells or other cultch, as the spat is drilled before its shell has lost its first paperlike thinness.

The drill is a difficult enemy to combat. Where it is sufficiently abundant to be a menace on private beds the oysters are usually dredged up and the drills removed by hand and destroyed, after which the oysters are again laid down. Much can be done by destroying the drills and their egg capsules wherever found. The common practice of some Delaware planters of depositing rough seed on their beds undoubtedly helps to maintain the abundance of the drill.








Ship John Hunt Knight

Note
 Abandoned Oyster Beds shown in red
 Area of Oyster growth of species:
 - - - - - *ostrea*
 - - - - - *edulis*
 - - - - - *irradiata*
 - - - - - *edulis*
 Soundings depth at each bar unless expressed in feet
 Bottom symbols
 h. for hard s. for soft

NATURAL OYSTER BEDS
DELAWARE
 SURVEYED BY
 UNITED STATES BUREAU OF FISHERIES
 AND
 DELAWARE OYSTER SURVEY COMMISSION
 Under the direction of
 H. H. Bennett and J. C. Bennett
 June and July 1908



Sheet of Chart No. 11



THE FISHERIES OF ALASKA IN 1910

By MILLARD C. MARSH


Agent at the Salmon Fisheries of Alaska

and

JOHN N. COBB

Assistant Agent

Bureau of Fisheries Document No. 746



CONTENTS.

	Page.
Summarized statistics.....	5
Persons engaged.....	5
Investment.....	6
Products.....	7
The salmon industry.....	11
Hatcheries.....	11
Statistics.....	12
Catch.....	12
Canning.....	15
Canneries in operation.....	17
Pickling.....	21
Mild curing.....	23
Fresh salmon.....	25
Minor preserving processes.....	25
Return of marked salmon.....	26
Observations in Wood River region.....	27
Count of breeding run in Wood River.....	28
Significance of Wood River data.....	33
Explorations of Lake Aleknagik.....	37
The cod fishery.....	37
Shore stations.....	38
Statistics for central Alaska.....	39
Persons engaged.....	39
Investment.....	39
Products.....	39
Vessel fishing.....	40
The halibut fishery.....	40
Fishing grounds.....	40
Methods and conditions.....	42
Statistics.....	44
Persons engaged in the southeast Alaska halibut fisheries.....	44
Investment in the southeast Alaska halibut fisheries.....	44
Products of the southeast Alaska halibut fisheries.....	45
Puget Sound fishing fleet.....	45
The herring fishery.....	45
Abundance of fish.....	45
Uses for food and bait.....	46
The fertilizer question.....	47
Statistics.....	50
Persons engaged.....	50
Investment.....	50
Products.....	51
Fertilizer and oils.....	51

	Page
The crab fishery.....	51
The whale fishery.....	52
Furs.....	53
Aquatic furs.....	54
Beaver.....	54
Muskrat.....	55
Land otter.....	55
Sea otter.....	55
Fur seal.....	56
Miscellaneous aquatic mammals.....	58
Hair seals.....	58
Walruses.....	59
License taxes and hatchery rebates.....	60
Complaints and prosecutions.....	61
Proposed legislation.....	63
Recommendations.....	65
Appendix.—Fishery laws and regulations.....	67
An act for the protection and regulation of the fisheries of Alaska.....	67
An act to prohibit aliens from fishing in the waters of Alaska.....	70
Fishery regulations.....	71
Regulations for protection of fur-bearing animals.....	71

THE FISHERIES OF ALASKA IN 1910.

By MILLARD C. MARSH, *Agent at the Salmon Fisheries of Alaska,*
and
JOHN N. COBB, *Assistant Agent.*

SUMMARIZED STATISTICS OF THE FISHERIES.

As in the similar reports for previous years, the District of Alaska is considered in the four geographic sections generally recognized, as follows: Southeast Alaska, embracing all that narrow strip of mainland and the numerous islands adjacent, from Portland Canal northwestward to and including Yakutat Bay; central Alaska, the region on the Pacific, or south side, from Yakutat Bay westward, including the Aleutian chain; western Alaska, the shores of Bering Sea and islands in this sea; and arctic Alaska, from Bering Strait to the Canadian border.

With the exception of arctic Alaska and a portion of central and western Alaska, practically all of the fishing localities were visited by one or the other of the two agents engaged in the inspection work this year. Considerable commercial fishing is carried on in the Yukon River and its tributaries, where fish wheels, nets, and spears are employed, but unfortunately it has been found impossible so far, owing to the short time available for the agents each season, to extend the inspection work over this large area, or to secure data showing the extent of the fisheries there. Owing to the impossibility of the agents visiting arctic Alaska in the limited open season, the data for this section are incomplete, but what have been secured are shown.

It has been found an impossibility to secure even approximate data as to the persons engaged or the investment in the hunting of aquatic animals (except fur seals and sea otters), which is general among the natives.

PERSONS ENGAGED.

The number of persons engaged in the fisheries of Alaska in 1910 was 15,620, an increase of 3,032 over 1909. Of these 6,836 were whites, 4,147 Indians, 2,411 Chinese, 2,206 Japanese, 4 Koreans, and

16 Filipinos, as compared with 5,608 whites, 2,823 Indians, 1,998 Chinese, and 2,159 Japanese, in 1909, showing an increase in 1910 of 1,228 whites, 1,324 Indians, 413 Chinese, and 47 Japanese. The most gratifying feature is the large increase in the number of whites and Indians employed, as all of the Indians and many of the whites are permanent residents of the District. The fact that the fishermen act as sailors on the transporting vessels to and from the salmon canneries and salteries explains the small number of transporters shown in the table as compared with the large number of transporting vessels.

PERSONS ENGAGED IN THE ALASKA FISHERIES IN 1910.

Occupation and race.	Southeast Alaska.	Central Alaska.	Western Alaska.	Arctic Alaska.	Total.
Fishermen:					
Vessel—					
Whites.....	402	6			408
Indians.....	35	48			86
Japanese.....	4				4
Total.....	444	54			498
Shore—					
Whites.....	1,149	737	1,589		3,475
Indians.....	1,710	196	72	438	2,416
Chinese.....	10		9		19
Japanese.....	3				3
Total.....	2,872	933	1,670	438	5,913
Total fishermen.....	3,316	987	1,670	438	6,411
Shoresmen:					
Whites.....	731	396	1,232	10	2,369
Indians.....	1,103	132	331		1,566
Chinese.....	705	468	1,218		2,391
Japanese.....	472	393	1,323		2,188
Koreans.....		4			4
Filipinos.....			16		16
Total.....	3,011	1,393	4,120	10	8,534
Transporters:					
Whites.....	264	115	205		584
Indians.....	69	10			79
Chinese.....	1				1
Japanese.....	8	3			11
Total.....	341	129	205		675
Grand total.....	6,668	2,509	5,995	448	15,620

INVESTMENT.

The total investment in the fisheries is \$20,711,422, an increase of \$10,829,740, as compared with 1909. A considerable proportion of this increase is due to the showing of cash capital once more, this item having been eliminated for the first time in 1906. Nearly all forms of apparatus show increases as compared with 1909.

INVESTMENT IN THE ALASKA FISHERIES IN 1910.

Items.	Southeast Alaska.		Central Alaska.		Western Alaska.		Arctic Alaska.		Total.	
	Num-ber.	Value.	Num-ber.	Value.	Num-ber.	Value.	Num-ber.	Value.	Num-ber.	Value.
Fishing vessels:										
Steamers and launches.	71	\$501,100							71	\$501,100
Tonnage	1,024								1,024	
Outfit		177,049								177,049
Sailing	3	3,800	2	\$4,000					5	7,800
Tonnage	35		87						122	
Outfit		875		1,300						2,175
Transporting vessels:										
Steamers and launches.	135	366,850	28	252,050	44	\$650,950			207	1,269,850
Tonnage	1,378		1,195		2,662				5,235	
Outfit		210,800		77,900		109,000				398,300
Sailing	20	180,150	13	385,500	32	711,000			65	1,276,650
Tonnage	6,753		17,395		41,748				65,896	
Outfit		33,200		22,000		48,000				103,200
Steamers and launches (under 5 tons).	240	401,030	12	26,225	7	18,200			259	^a 445,455
Boats, sail and row	1,090	59,648	570	33,880	845	186,840	82	\$13,360	2,587	293,668
Scows and lighters	142	67,183	111	58,300	130	107,529			383	233,012
Pile drivers	22	45,197	21	46,300	17	38,300			60	129,797
Apparatus, vessel fish-eries:										
Purse seines	10	3,995							^b 10	3,995
Lines, trawl		22,080								22,080
Shotguns			48	476					48	476
Whaling gear		1,015								1,015
Apparatus, shore fish-eries:										
Haul seines	48	9,797	56	21,285					^c 104	31,082
Purse seines	152	43,079							^d 152	43,079
Gill nets	416	58,659	132	17,295	903	90,682			^e 1,451	166,636
Dip nets	13	123	18		9				31	132
Lines, hand		521		1,245						1,766
Lines, trawl		15,870								15,870
Traps, stake	41	109,550	38	51,162	14	19,500			93	180,212
Traps, floating	13	22,728	1	1,500					14	24,228
Crab pots	366	1,082							366	1,082
Spears	120	115							120	115
Hoes	14	10	5	4					19	14
Shotguns	40	1,200							40	1,200
Whaling gear								18,450		18,450
Cash capital		3,544,333		1,593,444		3,456,660		10,000		8,604,437
Shore and accessory prop-erty		2,376,584		1,346,405		3,030,008		4,500		6,757,497
Total		8,257,623		3,940,280		8,485,706		46,250		20,711,422

^a Includes outfit.^b Aggregate length of 3,280 yards.^c Aggregate length of 36,190 yards.^d Aggregate length of 59,030 yards.^e Aggregate length of 412,176 yards.

PRODUCTS.

The total quantity of products was 214,536,433 pounds, valued at \$13,259,859, an increase of 12,553,195 pounds and \$2,078,471 over 1909. Except for salmon bellies and backs, fertilizer, oil, furs, and hides, the weights are round weights, or the weights of products when first taken from the water; for weights of prepared products the reader is referred to the subsidiary tables of the report. As the packing establishments almost invariably catch their own fish, it has been found practically impossible to show the value of the products as they leave the fishermen's hands, hence the values shown are for the prepared products.

PRODUCTS OF ALASKA FISHERIES IN 1910.

Products.	Southeast Alaska.		Central Alaska.		Western Alaska.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Black cod:						
Fresh.....	13,800	\$572				
Frozen.....	10,172	326				
Pickled.....	72,673	1,934				
Cod:						
Fresh.....	6,000	300	16,000	\$560		
Pickled.....			125,806	3,320		
Dry-salted.....			2,877,157	59,433		
Tongues, pickled.....			3,600	130		
Eulachon:						
Fresh.....	2,600	104				
Pickled.....	40,000	1,200				
Smoked.....	600	36				
Flounders, or sole:	5,000	150				
Halibut:						
Fresh.....	19,038,001	731,914	51,000	2,040		
Frozen.....	2,467,125	73,548				
Fletched.....	73,893	2,534				
Pickled.....	270	14				
Herring:						
Fresh.....	574,359	5,203	10,000	300		
Frozen.....	522,500	5,225				
Pickled.....	731,560	12,255	60,480	1,728		
Dry-salted.....	45,600	954				
Eggs, dried.....	1,000	100				
Pollock:						
Fresh.....			1,800	90		
Redfish, or black bass:	19,100	960	8,000	400		
Rock cod:						
Fresh.....	22,000	1,080	11,000	440		
Pickled.....	160	7				
Salmon:						
Fresh—						
Coho, or silver.....	52,588	2,419	7,500	225		
Humpback, or pink.....	24,000	300				
King, or spring.....	977,348	45,770				
Red, or sockeye.....	77,577	4,378	28,000	840		
Frozen—						
Coho, or silver.....	97,529	3,889				
Dog, or chum.....	17,337	695				
King, or spring.....	38,576	1,235				
Canned—						
Coho, or silver.....	5,841,990	404,907	1,394,960	99,103	814,870	\$55,656
Dog, or chum.....	16,221,450	703,555	9,170	403	1,564,640	69,451
Humpback, or pink.....	34,382,285	1,565,358	2,225,790	101,380	2,194,360	97,317
King, or spring.....	24,360	1,998	1,105,020	85,235	1,686,090	127,569
Red, or sockeye.....	18,247,740	1,466,918	25,541,250	1,959,539	57,729,700	4,347,933
Mild-cured—						
King, or spring.....	3,824,900	218,441	35,650	2,232		
Pickled—						
Coho, or silver.....	9,450	296	33,750	1,208		
Humpback, or pink.....	84,780	1,905	3,510	78	810	15
King, or spring.....					95,040	3,399
King, or spring, fins.....	400	24				
Red, or sockeye.....	540	20	400,950	12,278	2,819,880	92,351
Red, or sockeye, tips.....					800	60
Dry-salted—						
Coho, or silver, backs.....			10,000	290		
Dog, or chum.....	29,570	554				
King, or spring.....	9,600	288				
Humpback, or pink, backs.....	21,800	278	1,500	25		
Red, or sockeye, backs.....			17,000	410		
Smoked—						
Coho, or silver, backs.....			2,000	200		
Dog, or chum.....	440	60				
Humpback, or pink, backs.....	100	5				
Red, or sockeye, backs.....			16,058	1,608		
Salmon bellies, pickled:						
Coho, or silver.....			25,200	1,135		
Dog, or chum.....	14,000	770				
Humpback, or pink.....	84,200	4,410	39,000	1,725		
King, or spring.....	1,200	128				
Red, or sockeye.....	600	24	161,000	10,815		
Smelt:	4,085	205				
Tomcod:	800	32				

PRODUCTS OF ALASKA FISHERIES IN 1910—Continued.

Products.	Southeast Alaska.		Central Alaska.		Western Alaska.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Trout:						
Cutthroat.....	1,000	\$50				
Dolly Varden, or salmon trout—						
Fresh.....	50,000	2,000	15,000	\$750		
Canned.....			13,510	618		
Pickled.....	1,000	50				
Rainbow.....	7,100	284				
Steelhead—						
Fresh.....	3,800	168				
Frozen.....	19,215	1,153				
Fertilizer:						
Herring.....	2,617,000	40,000				
Whale.....	869,141	16,456				
Oil:						
Herring.....	2,077,500	55,000				
Shark.....	165	10				
Whale.....	2,744,480	117,270				
Abalone shells.....	70	30				
Clams.....	6,880	430	3,200	120		
Crabs.....	116,904	4,902	32,000	2,400		
Seaweed.....	2,000	300				
Aquatic furs and skins:						
Beaver.....	368	1,922	608	2,763	1,026	\$5,883
Castoreum.....			11	59	52	160
Muskrat.....	1,592	5,086	560	917	25,834	69,245
Otter—						
Land.....	1,232	5,213	1,117	4,493	2,302	8,843
Sea.....	15	600	120	5,900	20	670
Sea, pups.....			3	5	6	32
Seal—						
Fur.....	828	4,207			85,476	468,042
Fur, unborn.....					242	12
Hair.....	2,790	796			871	150
Walrus ivory.....	80	85				
Whale products:						
Bones, unground.....	400,000	4,500				
Bones, ground.....	395,000	4,789				
Stearin.....	114,711	5,249				
Whalebone, or baleen.....	55,025	4,805				
Total.....	113,223,554	5,542,633	34,288,340	2,365,195	67,022,019	5,346,788

Products.	Arctic Alaska.		Total.	
	Pounds.	Value.	Pounds.	Value.
Black cod:				
Fresh.....			13,800	\$572
Frozen.....			10,172	326
Pickled.....			72,673	1,934
Cod:				
Fresh.....			22,000	860
Pickled.....			125,866	3,320
Dry-salted.....			2,877,157	59,433
Tongues, pickled.....			3,600	130
Eulachon:				
Fresh.....			2,600	104
Pickled.....			40,000	1,200
Smoked.....			600	36
Flounders, or sole.....			5,000	150
Haiibut:				
Fresh.....			19,089,001	733,954
Frozen.....			2,467,125	73,548
Fletched.....			73,893	2,534
Pickled.....			270	14
Herring:				
Fresh.....			584,359	5,503
Frozen.....			522,500	5,225
Pickled.....			792,040	13,983
Dry-salted.....			45,600	954
Eggs, dried.....			1,000	100
Pollock.....			1,800	90
Redfish, or black bass.....			27,100	1,360

PRODUCTS OF ALASKA FISHERIES IN 1910—Continued.

Products.	Arctic Alaska.		Total.	
	Pounds.	Value.	Pounds.	Value.
Rock cod:				
Fresh.....			33,000	\$1,520
Pickled.....			160	7
Salmon:				
Fresh—				
Coho, or silver.....			60,088	2,644
Humpback, or pink.....			24,000	300
King, or spring.....			977,348	45,770
Red, or sockeye.....			105,577	5,218
Frozen—				
Coho, or silver.....			97,529	3,889
Dog, or chum.....			17,337	695
King, or spring.....			38,576	1,235
Canned—				
Coho, or silver.....			8,051,820	559,666
Dog, or chum.....			17,795,260	773,409
Humpback, or pink.....			38,802,435	1,764,055
King, or spring.....			2,815,470	214,802
Red, or sockeye.....			101,518,690	7,774,390
Mild-cured—				
King, or spring.....			3,860,550	220,673
Pickled—				
Coho, or silver.....			43,200	1,504
Humpback, or pink.....			89,100	1,998
King, or spring.....			95,040	3,399
King, or spring, fins.....			400	24
Red, or sockeye.....			3,221,370	104,649
Red, or sockeye, tips.....			800	60
Dry-salted—				
Coho, or silver, backs.....			10,000	290
Dog, or chum.....			29,570	554
King, or spring.....			9,600	288
Humpback, or pink, backs.....			23,300	303
Red, or sockeye, backs.....			17,000	410
Smoked—				
Coho, or silver, backs.....			2,000	200
Dog, or chum.....			440	60
Humpback, or pink, backs.....			100	5
Red, or sockeye, backs.....			16,058	1,608
Salmon bellies, pickled:				
Coho, or silver.....			25,200	1,135
Dog, or chum.....			14,000	770
Humpback, or pink.....			123,200	6,135
King, or spring.....			1,200	128
Red, or sockeye.....			161,600	10,839
Smelt.....			4,085	205
Tomcod.....			800	32
Trout:				
Cutthroat.....			1,000	50
Dolly Varden, or salmon trout—				
Fresh.....			65,000	2,750
Canned.....			13,510	618
Pickled.....			1,000	50
Rainbow.....			7,100	284
Steelhead—				
Fresh.....			3,800	168
Frozen.....			19,215	1,153
Fertilizer:				
Herring.....			2,617,000	40,000
Whale.....			869,141	16,456
Oil:				
Herring.....			a 2,077,500	55,000
Shark.....			b 165	10
Whale.....			c 2,744,480	117,270
Abalone shells.....			70	30
Clams.....			d 10,080	550
Crabs.....			e 148,904	7,302
Seaweed.....			2,000	300
Aquatic furs and skins:				
Beaver.....			f 2,002	10,568
Castoreum.....			63	219
Muskrat.....			g 27,986	75,248

a Represents 277,000 gallons.

b Represents 22 gallons.

c Represents 369,930 gallons.

d Represents 1,260 bushels.

e Represents 70,452 crabs.

f Represents 2,002 skins.

g Represents 223,863 skins.

PRODUCTS OF ALASKA FISHERIES IN 1910—Continued.

Products.	Arctic Alaska.		Total.	
	Pounds.	Value.	Pounds.	Value.
Aquatic furs and skins—Continued.				
Otter—				
Land.....			a 4,651	\$18,549
Sea.....			b 155	7,170
Sea, pups.....			c 9	37
Seal—				
Fur.....			d 86,304	472,249
Fur, unborn.....			e 242	12
Hair.....			f 3,661	946
Walrus ivory.....	186	\$186	266	271
Whale products:				
Bones, unground.....			400,000	4,500
Bones, ground.....			395,000	4,789
Stearin.....			114,711	5,249
Whalebone, or baleen.....	2,334	5,057	57,359	9,862
Total.....	2,520	5,243	214,536,433	13,259,859

a Represents 1,861 skins.

b Represents 31 skins.

c Represents 3 skins.

d Represents 14,384 skins (of these, 660 skins were from a seized Japanese schooner).

e Represents 121 skins (these were from a seized Japanese schooner).

f Represents 1,221 skins.

THE SALMON INDUSTRY.

The run of salmon was very good in all sections except western Alaska. For a time the outlook was bad in southeast Alaska owing to the excessive rains which prevailed during the first half of the season, causing the salmon to rush up the streams, but an exceptionally dry spell lasting six weeks followed, which made the streams quite low and kept the fish from going up too rapidly. As a result the fisherman were enabled to make large catches during this period.

HATCHERIES.

Seven salmon hatcheries were operated during the season of 1909-10, as follows:

SALMON HATCHERIES OPERATED IN 1910.

Name.	Location.	Owner and operator.
Yes Lake.....	Yes Lake.....	United States Bureau of Fisheries.
Afognak.....	Afognak Island.....	Do.
Fortmann.....	Naha Stream.....	Alaska Packers Association.
Karluk.....	Karluk River.....	Do.
Klawak.....	Klawak Lake.....	North Pacific Trading and Packing Co., and North Alaska Salmon Co.
Hetta.....	Hetta Lake.....	Northwestern Fisheries Co.
Quadra.....	Quadra Lake.....	Do.

The Alaska Packers Association reports as follows on a subject of interest to fish culturists:

We have been quite successful in retaining the fry in our nursery ponds [at Fortmann hatchery] for a definite period and feeding them on fresh steelheads macerated

to a pulp. In two ponds containing about 10,000,000 fry, from 30 to 40 pounds of this food was fed each day, and they appeared to thrive wonderfully well upon it, as scarcely any dead fry were found.

The rainfall was 160.80 inches and the snowfall 289 inches for the year ended June 30, 1910, at Fortmann hatchery; which record will give a slight idea of the weather conditions with which the superintendents of hatcheries in Alaska have to contend. Despite the adverse weather conditions, however, all of the hatcheries except Fortmann and Afognak operated at full capacity, and taken as a whole the season was a fairly successful one.

The Klawak Lake hatchery of the North Pacific Trading & Packing Co. was enlarged the present summer so that it is now able to handle 10,000,000 eggs.

At the dam on the stream leading to Capt. John C. Callbreath's hatchery on McHenry Inlet a man has been stationed each year since the hatchery was shut down, for the purpose of lifting the salmon over the structure. In 1908, 1,022 males and 876 females were put over, and in 1909, 516 males and 434 females.

OUTPUT OF THE SALMON HATCHERIES OF ALASKA.

Hatcheries.	Year ended June 30, 1910. ^a				Eggs taken 1910-11.	
	Red, or sockeye.		Humpback, or pink.		Red, or sockeye.	Humpback, or pink.
	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.		
Yes Lake.....	^b 72,005,000	69,879,600	72,000,000	114,000
Afognak.....	76,020,000	68,422,170	499,400	363,740	30,725,000	405,000
Fortmann.....	53,340,000	50,725,000	34,920,000
Karluk.....	45,228,000	40,620,000	49,626,000
Klawak.....	(c)	5,300,000	(c)
Hetta.....	10,313,000	9,000,000	9,141,000
Quadra.....	10,863,000	9,850,000	11,200,000
Total.....	253,796,770	499,400	363,740	9,141,000	519,000

^a In three instances fry were held until July, 1910, and in order to make the record for the season complete these have been included.

^b Of these, 5,000 were reported as coho eggs.

^c No report.

STATISTICS.

CATCH IN 1907, 1908, 1909, AND 1910.

Following is a table showing, for the geographic sections, by apparatus and species and by species alone, the number of salmon caught in the years 1907, 1908, 1909, and 1910. All species, except red salmon, show increases over 1909. The total catch in 1910 is smaller than in any of the other years shown.

CATCH OF SALMON IN ALASKA IN 1907, 1908, 1909, AND 1910, BY SECTIONS,
SPECIES, AND APPARATUS.

Apparatus and species.	1907	1908	1909	1910
SOUTHEAST ALASKA.				
Seines:				
Coho, or silver.....	302,963	273,993	165,177	322,521
Dog, or chum.....	1,101,822	1,378,339	387,774	1,566,221
Humpback, or pink.....	8,614,551	8,900,467	5,572,005	6,228,732
King, or spring.....	259	1,812	293	152
Red, or sockeye.....	1,419,221	1,691,149	1,285,265	1,481,898
Total.....	11,438,816	12,245,760	7,410,514	9,599,522
Traps:				
Coho, or silver.....	139,783	119,034	112,213	165,023
Dog, or chum.....	158,170	368,709	337,395	437,726
Humpback, or pink.....	3,438,335	5,102,843	3,628,940	3,151,684
King, or spring.....	26,835	3,448	5,107	2,546
Red, or sockeye.....	615,084	486,646	893,816	860,737
Total.....	4,378,807	6,080,680	4,977,471	4,617,716
Gill nets:				
Coho, or silver.....	83,943	84,176	78,845	164,990
Dog, or chum.....	74,298	56,431	9,041	28,802
Humpback, or pink.....	18,029	59,582	127,422	32,357
King, or spring.....	70,358	64,148	68,659	51,667
Red, or sockeye.....	214,442	378,834	478,398	574,251
Total.....	461,100	643,171	762,365	852,067
Lines:				
Coho, or silver.....	1,052	1,329	8,000	6,000
King, or spring.....	23,082	61,633	134,606	204,823
Total.....	24,134	62,962	142,606	210,823
Spears:				
Red, or sockeye.....	20,000	4,000	45,400	70,000
Wheels:				
King, or spring.....		27		
Total:				
Coho, or silver.....	527,741	478,532	364,235	658,534
Dog, or chum.....	1,334,290	1,803,479	734,210	2,032,749
Humpback, or pink.....	12,070,915	14,062,892	9,328,367	9,412,773
King, or spring.....	120,564	131,068	208,665	259,188
Red, or sockeye.....	2,269,347	2,560,629	2,702,879	2,986,886
Grand total.....	16,322,857	19,036,600	13,338,356	15,350,130
CENTRAL ALASKA.				
Seines:				
Coho, or silver.....	48,759	60,847	52,258	64,202
Humpback, or pink.....	252,373	268,466	127,549	375,041
King, or spring.....	4,015	3,028	3,907	1,598
Red, or sockeye.....	3,568,069	2,709,750	2,038,833	2,227,803
Total.....	3,873,216	3,042,091	2,222,547	2,668,644
Traps:				
Coho, or silver.....	163,076	90,616	89,918	115,922
Dog, or chum.....				1,318
Humpback, or pink.....	6,420	375,140	3,740	273,023
King, or spring.....	36,791	17,216	44,632	34,007
Red, or sockeye.....	2,711,142	2,285,401	2,152,555	2,095,563
Total.....	2,917,429	2,768,373	2,290,845	2,519,833
Gill nets:				
Coho, or silver.....	15,000			18,826
King, or spring.....	27,022	18,351	18,059	15,995
Red, or sockeye.....	358,649	512,464	487,984	298,915
Total.....	400,671	530,815	506,043	333,736
Total:				
Coho, or silver.....	226,835	151,463	142,176	198,950
Dog, or chum.....				1,318
Humpback, or pink.....	258,793	643,606	131,289	648,064
King, or spring.....	67,828	38,595	66,598	51,600
Red, or sockeye.....	6,637,860	5,507,615	4,679,372	4,622,281
Grand total.....	7,191,316	6,341,279	5,019,435	5,522,213

CATCH OF SALMON IN ALASKA IN 1907, 1908, 1909, AND 1910, BY SECTIONS, SPECIES, AND APPARATUS—Continued.

Apparatus and species.	1907	1908	1909	1910
WESTERN ALASKA.				
Traps:				
Coho, or silver.....	29,199	20,000	9,930	6,340
Dog, or chum.....	36,141	114,534	101,456	58,039
Humpback, or pink.....	1,500	261,519	15	513,072
King, or spring.....	5,011	4,856	3,096	4,382
Red, or sockeye.....	1,078,869	860,516	508,011	326,833
Total.....	1,150,720	1,261,425	622,508	908,666
Gill nets:				
Coho, or silver.....	109,650	86,088	71,393	132,860
Dog, or chum.....	472,586	340,309	346,340	252,179
Humpback, or pink.....	337,514	138,138	31,811	149,057
King, or spring.....	134,391	87,174	128,893	97,373
Red, or sockeye.....	9,181,034	16,013,966	15,133,872	11,266,776
Total.....	10,235,175	16,665,675	15,712,309	11,898,245
Total:				
Coho, or silver.....	138,849	106,088	81,323	139,200
Dog, or chum.....	508,727	454,843	447,796	310,218
Humpback, or pink.....	339,014	399,657	31,826	662,129
King, or spring.....	139,402	92,030	131,989	101,755
Red, or sockeye.....	10,259,903	16,874,482	15,641,883	11,593,609
Grand total.....	11,385,895	17,927,100	16,334,817	12,806,911
TOTAL.				
Seines:				
Coho, or silver.....	351,722	334,840	217,435	386,723
Dog, or chum.....	1,101,822	1,378,339	387,774	1,566,221
Humpback, or pink.....	8,896,924	9,168,933	5,699,554	6,603,773
King, or spring.....	4,274	4,840	4,200	1,750
Red, or sockeye.....	4,987,290	4,400,899	3,324,098	3,709,701
Total.....	15,312,032	15,287,851	9,633,061	12,268,168
Traps:				
Coho, or silver.....	332,058	229,650	212,061	287,285
Dog, or chum.....	194,311	483,243	438,851	497,083
Humpback, or pink.....	3,446,255	5,739,502	3,632,695	3,937,779
King, or spring.....	68,637	25,520	52,835	40,935
Red, or sockeye.....	4,405,695	3,632,563	3,584,382	3,283,133
Total.....	8,446,956	10,110,478	7,920,824	8,046,215
Gill nets:				
Coho, or silver.....	208,593	170,264	150,238	316,676
Dog, or chum.....	546,884	396,740	355,381	280,981
Humpback, or pink.....	355,543	197,720	159,233	181,414
King, or spring.....	231,801	169,673	215,611	165,035
Red, or sockeye.....	9,754,125	16,905,264	16,070,254	12,139,942
Total.....	11,096,946	17,839,661	16,950,717	13,084,048
Lines:				
Coho, or silver.....	1,052	1,329	8,000	6,000
King, or spring.....	23,082	61,633	134,666	204,823
Total.....	24,134	62,962	142,666	210,823
Spears:				
Red, or sockeye.....	20,000	4,000	45,400	70,000
Wheels:				
King, or spring.....		27		
Total:				
Coho, or silver.....	893,425	736,083	587,734	996,684
Dog, or chum.....	1,843,017	2,258,322	1,182,006	2,344,285
Humpback, or pink.....	12,668,722	15,106,155	9,491,482	10,722,966
King, or spring.....	327,794	261,693	407,252	412,543
Red, or sockeye.....	19,167,110	24,942,726	23,024,134	19,202,776
Grand total.....	34,900,068	43,304,979	34,692,608	33,679,257

NUMBER AND GROSS WEIGHT OF EACH SPECIES OF SALMON CAUGHT IN 1907, 1908, 1909, AND 1910.

Species.	1907		1908	
	<i>Number.</i>	<i>Pounds.</i>	<i>Number.</i>	<i>Pounds.</i>
Coho, or silver	893, 425	5, 360, 550	736, 083	4, 416, 498
Dog, or chum	1, 843, 017	14, 744, 136	2, 258, 322	18, 066, 576
Humpback, or pink	12, 668, 722	50, 674, 888	15, 106, 155	60, 424, 620
King, or spring	327, 794	7, 211, 468	261, 693	5, 757, 246
Red, or sockeye	19, 167, 110	95, 835, 550	24, 942, 726	124, 713, 630
Total	34, 900, 068	173, 826, 592	43, 304, 979	213, 378, 570

Species.	1909		1910	
	<i>Number.</i>	<i>Pounds.</i>	<i>Number.</i>	<i>Pounds.</i>
Coho, or silver	587, 734	3, 526, 404	996, 684	5, 980, 104
Dog, or chum	1, 182, 006	9, 456, 048	2, 344, 285	18, 754, 280
Humpback, or pink	9, 491, 482	37, 965, 928	10, 722, 966	42, 891, 864
King, or spring	407, 252	8, 959, 544	412, 543	9, 075, 946
Red, or sockeye	23, 024, 134	115, 120, 670	19, 202, 776	96, 013, 880
Total	34, 692, 608	175, 028, 594	33, 679, 254	172, 716, 074

CANNING.

When the season of 1909 opened, all grades of salmon, except pinks and chums, were commanding remunerative prices. The prices of these two grades began to crumble in 1908 and kept on dropping through 1909, until finally they reached bottom at \$2.40 per case for pinks (a drop of \$1.05 per case from the 1907 prices) and \$2.28 per case for chums (a drop of 96 cents per case from the 1907 prices). The demand for pink and chum salmon began to fall off in 1907, despite which the packers kept on piling up stock during the next two years, with the result that they became a drug on the market, and for a time it was difficult to move them, even at the above unremunerative prices. Late in 1909 the demand began to improve, and when the season of 1910 opened but few pinks and chums were left in first hands.

Early in the season rumors began to circulate that prices on all grades would be advanced, and the buyers, who had been content to buy only for immediate necessities as long as prices seemed to be crumbling, now came into the market with orders for large stocks. As a result, the packers soon were obliged to prorate the orders, as the pack did not begin to equal the demand. The expected high prices were realized, and before the pack had come out of Alaska it was all sold at the most remunerative figures realized by the packers in years.

In 1909, owing to the expected quadrennial heavy run of sockeye salmon on Puget Sound, the canneries of Gorman & Co., at Kasaan, of the Astoria & Puget Sound Packing Co., in Excursion Inlet, and of the Fidalgo Island Packing Co., at Ketchikan, all in southeast

Alaska, were shut down, as these companies felt it would be more profitable to devote all their energies to their Puget Sound plants. In 1910 all were operated. In addition new canneries were opened by the St. Elias Packing Co., at Alsek, in southeast Alaska, by the Northwestern Fisheries Co., at Kenai, on Cook Inlet (succeeding the mild-curing plant formerly operated by the San Juan Fishing & Packing Co.), and by the Columbia River Packers Association, at Chignik, in central Alaska. The cannery of the Alaska Salmon Co., on Wood River, western Alaska, which was closed down in 1909, owing to the loss of its supply ship, was operated this year.

New canneries which will likely be finished in time to operate in 1911 are the Hawk Fishing Co., at Hawk Inlet, Tee Harbor Packing Company, at Tee Harbor, southeast Alaska, and the Alaska Packers Association, at Naknek, western Alaska. For some years the Alaska Packers Association has operated two canneries at Karluk, on Kodiak Island. Karluk has no harbor, except for boats drawing less than 4 feet of water, and the association, fearing a repetition of the disaster of 1907, when the bark *Servia*, with a full cargo of salmon, was driven ashore in a gale and totally destroyed, began in 1909 the erection of a new cannery at Larsen Bay, a well-sheltered spot near by. This establishment will operate in 1911, the two Karluk canneries being held in reserve. Fishing will be carried on as usual at Karluk, the fish being transported to the new cannery. C. A. Burckhardt & Co., who now operate two canneries in southeast Alaska, have bought the saltery formerly owned by Mrs. A. E. King, at Sunny Point, southeast Alaska, and will convert this into a one-line cannery. The Alaska Fishermen's Packing Co. have purchased the Nelson, Olsen & Co. saltery in Kvichak Bay, western Alaska, and will replace the old plant by a one-line cannery. Several canneries are also engaged in making, or are contemplating, extensive changes to and enlargements of their present plants.

On August 10 the cannery of the Alaska-Portland Packers' Association, at Snag Point, Nushagak Bay, was completely destroyed by fire. The warehouse alongside, with much of the gill-netting and all of the trap web, together with part of the season's pack, was also consumed. The bunk houses, store, office, and residence, and the floating property, were saved. The property loss was about \$200,000, partly covered by insurance. The company will rebuild next spring and hopes to have the cannery completed in time to operate that season.

On the night of September 12 fire broke out in the cannery of Gorman & Co., at Kasaan, in southeast Alaska, and resulted in the total destruction of the cannery, warehouse, store, hotel, and part of the season's pack. The company will erect a new cannery in time to operate next season.

Several canneries packed some thousands of cases of salmon in the new seamless or sanitary can with such success that it is probably a question of but a few seasons until this will be the only form of can in use in Alaska.

The two cannery fires resulted in the loss of the following cases of salmon:

	Cases.
Cohos, 1-pound tall.....	1, 552
Chums, 1-pound tall.....	4, 896
Pinks, ½-pound flat.....	141
Pinks, 1-pound tall.....	11, 956
Reds, 1-pound tall.....	22, 178
Total.....	40, 723

These have been included in the statistical tables, as they had passed through all the stages of packing and were eventually paid for by the insurance companies.

CANNERIES IN OPERATION.

Following is a list of the canneries operated during the season of 1910:

Name.	Location.
Southeast Alaska:	
John L. Carlson & Co.....	Taku Harbor.
George T. Myers & Co.....	Sitkoh Bay.
Yakutat & Southern Railway Co.....	Yakutat.
Astoria & Puget Sound Canning Co.....	Excursion Inlet.
Pacific American Fisheries.....	Do.
Northwestern Fisheries Co.....	Dundas Bay, Quadra Bay, Santa Ana, Hunter Bay.
North Pacific Trading & Packing Co.....	Klawak.
Fidalgo Island Packing Co.....	Ketchikan.
Shakan Salmon Co.....	Shakan.
Gorman & Co.....	Kasaan.
F. C. Barnes Co. (Inc.).....	Lake Bay.
Thlinket Packing Co.....	Funter Bay.
Alaska Packers Association.....	Loring and Wrangell.
St. Elias Packing Co.....	Aleek River.
Pillar Bay Packing Co.....	Point Ellis.
Metlakahla Industrial Co.....	Metlakahla.
Pacific Coast & Norway Packing Co.....	Petersburg.
Yes Bay Canning Co.....	Yes Bay.
Chilkoot Fisheries Co.....	Chilkoot Inlet.
Central Alaska:	
Northwestern Fisheries Co.....	Chignik, Uyak, Kenai, and Orca.
Alaska Packers Association.....	Kasilof, Karluk (2), Alitak, and Chignik.
Columbia River Packers' Association.....	Chignik.
Western Alaska:	
Alaska Packers Association.....	Nushagak Bay (2), Kvichak Bay (2), Naknek River (2), and Ugagak River.
North Alaska Salmon Co.....	Kvichak Bay, Nushagak Bay, Ugagak River, and Lockanok.
Northwestern Fisheries Co.....	Nushagak Bay.
Naknek Packing Co.....	Naknek River.
Red Salmon Canning Co.....	Ugashik River.
Alaska-Portland Packers Association.....	Nushagak Bay.
Bristol Bay Packing Co.....	Kvichak Bay.
Alaska Fishermen's Packing Co.....	Nushagak Bay.
Columbia River Packers Association.....	Do.
Alaska Salmon Co.....	Wood River.

Persons engaged.—The fishermen engaged this year numbered 3,722, of whom slightly more than one-half were white. The cannery employees numbered 8,194, of whom all nationalities show increases as compared with 1909. The transporters numbered 515, an increase over 1909. All branches of the industry show increases as compared with 1909. In all, 12,431 persons were employed, an increase of 1,909 over 1909.

PERSONS ENGAGED IN THE SALMON-CANNING INDUSTRY IN 1910.

Occupation and race.	Southeast Alaska.	Central Alaska.	Western Alaska.	Total.
Fishermen:				
Whites.....	444	485	1,541	2,470
Indians.....	1,153	80	1,233
Japanese.....	10	9	19
Total.....	1,607	565	1,550	3,722
Shoresmen:				
Whites.....	529	359	1,203	2,091
Indians.....	1,060	121	326	1,507
Chinese.....	705	467	1,216	2,388
Japanese.....	472	393	1,323	2,188
Koreans.....	4	4
Filipinos.....	16	16
Total.....	2,766	1,344	4,084	8,194
Transporters:				
Whites.....	184	111	189	484
Indians.....	23	2	25
Chinese.....	1	1
Japanese.....	2	3	5
Total.....	209	117	189	515
Grand total:				
Whites.....	1,157	955	2,933	5,045
Indians.....	2,236	203	326	2,765
Chinese.....	705	468	1,216	2,389
Japanese.....	484	396	1,332	2,212
Koreans.....	4	4
Filipinos.....	16	16
Total.....	4,582	2,026	5,823	12,431

Investments, wages, etc.—There were 52 canneries in operation—23 in southeast Alaska, an increase of 4 over 1909; 10 in central Alaska, an increase of 2 over 1909; and 19 in western Alaska, an increase of 1 over 1909; a total increase for all Alaska of 7.

There were 176 steamers and launches over 5 tons, 55 under 5 tons, and 59 sailing vessels engaged in transporting supplies and the pack, and doing general work for the canneries. This is a large increase over 1909.

All forms of apparatus except floating traps show increases over 1909. The increases are especially noticeable in purse seines and stake traps, which increased in number 43 and 27 respectively.

Included in this table for the first time are the items of cash capital, materials used, and wages paid. Considerable misapprehension

seems to have arisen among readers of this report as to the profits of the cannerymen, which have appeared to them enormous. Such an erroneous conclusion is apparently based on the assumption that the price received for the canned product represents practically net profits. For eight years prior to the 1910 season but few of the cannerymen received an adequate return upon the capital invested, while many of them sustained heavy losses during certain years. It has been found difficult to secure accurate data showing the cost of operation, and several items, such as insurance, taxes outside of Alaska, commissions paid the brokers, etc., have not been taken into account, but it is hoped in time to include these.

INVESTMENT IN THE SALMON-CANNING INDUSTRY IN 1910.

Items.	Southeast Alaska.		Central Alaska.		Western Alaska.		Total.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Canneries.....	23	10	19	52
Transporting vessels:								
Steamers and launches								
over 5 tons.....	110	\$310,450	24	\$212,050	42	\$605,950	176	\$1,128,450
Tonnage.....	1,186	1,077	2,507	4,770
Outfit.....		175,000		72,000		104,000		351,000
Sailing.....	16	160,250	11	348,000	32	711,000	59	1,219,250
Tonnage.....	6,332	17,160	41,748	65,240
Outfit.....		30,000		20,000		48,000		98,000
Steamers and launches								
under 5 tons.....	39	86,300	10	24,025	6	13,700	55	124,025
Boats, sail and row.....	541	36,163	263	23,990	822	178,148	1,626	238,293
Lighters and scows.....	108	46,983	108	57,500	130	107,529	346	212,312
Pile drivers.....	22	45,197	21	46,300	17	38,300	60	129,797
Apparatus:								
Haul seines.....	45	9,372	24	18,100	69	27,472
Purse seines.....	133	38,784	133	38,784
Gill nets.....	271	31,134	127	16,545	880	88,957	1,278	136,636
Traps, stake.....	41	109,550	38	51,162	14	19,500	93	180,212
Traps, floating.....	13	22,728	1	1,500	14	24,228
Spears.....	75	75	75	75
Cash on hand.....		230,000		100,000		190,000		520,000
Shore and accessory prop-								
erty.....		2,016,144		1,291,405		2,913,008		6,220,557
Materials used.....		1,964,493		778,531		1,646,775		4,389,799
Wages paid.....		1,100,678		638,886		1,562,295		3,301,859
Total.....		6,413,301		3,700,294		8,227,154		18,340,749

Output.—The table of products shows the quantity and value of each species packed, with size and style of cans. As usual, western Alaska leads in value of the pack, but southeast Alaska leads in quantity packed. Red, or sockeye, salmon predominate in central and western Alaska, while humpback, or pink, salmon predominate in southeast Alaska.

OUTPUT OF SALMON FROM THE CANNERIES IN 1910, BY SPECIES AND SIZE OF CANS.^a

Products.	Southeast Alaska.		Central Alaska.		Western Alaska.		Total.	
	Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
Coho, or silver:								
½-pound flat.....	326	\$1,299					326	\$1,299
1-pound flat.....	2,249	12,357					2,249	12,357
1-pound tall.....	80,045	391,251	19,928	\$99,103	11,641	\$55,656	111,614	546,010
Total.....	82,620	404,907	19,928	99,103	11,641	55,656	114,189	559,666
Dog, or chum:								
1-pound tall.....	231,735	703,555	131	403	22,352	69,451	254,218	773,409
Humpback, or pink:								
½-pound flat.....	6,375	15,871					6,375	15,871
1-pound flat.....	7,900	35,550					7,900	35,550
1-pound tall.....	480,088	1,513,937	31,797	101,380	31,348	97,317	543,233	1,712,634
Total.....	494,363	1,565,358	31,797	101,380	31,348	97,317	557,508	1,764,055
King, or spring:								
½-pound flat.....	108	432					108	432
1-pound tall.....	294	1,566	15,786	85,235	24,087	127,569	40,167	214,370
Total.....	402	1,998	15,786	85,235	24,087	127,569	40,275	214,802
Red, or sockeye:								
½-pound flat.....	43,166	170,489			1,474	5,896	44,640	176,385
1-pound flat.....	39,941	236,453					39,941	236,453
1-pound tall.....	199,158	1,059,976	364,875	1,959,539	823,973	4,342,037	1,388,006	7,361,552
Total.....	282,265	1,466,918	364,875	1,959,539	825,447	4,347,933	1,472,587	7,774,390
Grand total.....	1,091,385	4,142,736	432,517	2,245,660	914,875	4,697,926	2,438,777	11,086,322

^a All pound cases contain 48 1-pound cans; the ½-pound cases contain 48 ¼-pound cans. Reduced to a common basis of cases containing 48 1-pound cans, the pack is 2,413,052½ cases.

Comparison of pack of 1907, 1908, 1909, and 1910.—With the exception of 1908, the pack of 1910 exceeds in quantity that of any of the four years, and it exceeds in value any of them, being the most valuable pack ever put up in Alaska.

COMPARISON OF THE OUTPUT OF THE SALMON CANNERIES IN 1907, 1908, 1909, AND 1910.^a

Products.	1907		1908		1909		1910	
	Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
Coho, or silver:								
½-pound flat.....	969	\$4,273	209	\$627			326	\$1,299
1-pound flat.....	3,933	17,292	2,414	9,903	1,206	\$5,543	2,249	12,357
1-pound tall.....	80,772	315,819	66,309	263,559	55,350	225,486	111,614	546,010
Total.....	85,674	337,384	68,932	274,089	56,556	231,029	114,189	559,666
Dog, or chum:								
½-pound flat.....	491	1,228						
1-pound flat.....	664	2,125	107	321				
1-pound tall.....	183,262	544,404	218,406	553,876	120,712	274,110	254,218	773,409
Total.....	184,417	547,757	218,513	554,197	120,712	274,110	254,218	773,409
Humpback, or pink:								
½-pound flat.....	17,589	46,093					6,375	15,871
1-pound flat.....	7,406	26,662	569	1,590			7,900	35,550
1-pound tall.....	545,772	1,726,525	643,564	1,731,789	464,873	1,114,839	543,233	1,712,634
Total.....	570,767	1,799,280	644,133	1,733,379	464,873	1,114,839	557,508	1,764,055
King, or spring:								
½-pound flat.....	28	98	125	425			108	432
1-pound tall.....	43,410	181,620	23,667	99,442	48,034	207,624	40,167	214,370
Total.....	43,438	181,718	23,792	99,867	48,034	207,624	40,275	214,802
Red, or sockeye:								
½-pound flat.....	45,383	160,731	21,817	68,083	16,385	63,888	44,640	176,385
1-pound flat.....	29,821	154,646	26,950	138,120	85,193	236,609	39,941	236,453
1-pound tall.....	1,242,600	5,599,850	1,613,911	7,318,048	1,611,916	7,310,053	1,388,006	7,361,552
Total.....	1,317,804	5,915,227	1,662,678	7,524,251	1,713,494	7,610,550	1,472,587	7,774,390
Grand total.....	2,202,100	8,781,366	2,618,048	10,185,783	2,403,669	9,438,152	2,438,777	11,086,322

^a All pound cases contain 48 1-pound cans; the ½-pound cases contain 48 ¼-pound cans.

The following table shows, by species, the average price received by the packer per case of 1-pound talls for a series of years. The 1-pound tall cases are used because they form the vast majority of the pack and are the ones in common use by the consumer, the flat cans being packed for a special trade.

AVERAGE ANNUAL PRICE PER CASE OF 48 1-POUND TALL CANS OF SALMON, 1905-1910.

Products.	1905	1906	1907	1908	1909	1910
Coho, or silver.....	\$3.20	\$3.63	\$3.91	\$3.98	\$4.07	\$4.89
Dog, or chum.....	2.69	2.87	2.97	2.53	2.28	3.04
Humpback, or pink.....	2.95	3.00	3.16	2.69	2.40	3.15
King, or spring.....	3.28	3.78	4.18	4.20	4.32	5.34
Red, or sockeye.....	3.38	3.77	4.59	4.52	4.53	5.30

PICKLING.

Owing to the low prices which have prevailed during several seasons for whole pickled salmon, there was but little incentive for the salteries to engage in this business very heavily this year. Some shut down altogether, while others very materially curtailed operations. Prices improved during the latter part of the season, but it was then too late.

The action of the Department in forbidding the packing of salmon bellies without making some economic use of the backs contributed to the depression in the pickled trade, as bellies were the most remunerative product prepared. Nearly all of the salters are now agreed, however, that this action was wise and necessary. Under the old wasteful method from one-half to two-thirds of the edible portion of the fish was thrown away and the belly only was pickled.

Persons engaged.—This year 261 persons (196 fishermen, 51 shoresmen, and 14 transporters) were employed, a decrease of 135 as compared with 1909.

PERSONS ENGAGED IN THE SALMON-PICKLING INDUSTRY IN 1910.

How engaged.	Southeast Alaska.	Central Alaska.	Western Alaska.	Total.
Fishermen:				
Whites.....	29	3	46	78
Indians.....	13	105	118
Total.....	42	108	46	196
Shoresmen:				
Whites.....	5	7	20	32
Indians.....	16	3	19
Total.....	21	10	20	51
Transporters:				
Whites.....	2	2	4	8
Indians.....	6	6
Total.....	2	8	4	14
Grand total.....	65	126	70	261

Investment.—There were 12 salteries (6 in southeast Alaska, 4 in central Alaska, and 2 in western Alaska) in operation, a decrease of 4 as compared with 1909. In addition, a few of the canneries and mild-curing plants also pickled their surplus catch, and while the product has been included in the present table, the men and investment could not be separated from the statistics of the other branches of the industry.

INVESTMENT IN THE SALMON-PICKLING INDUSTRY IN 1910.

Items.	Southeast Alaska.		Central Alaska.		Western Alaska.		Total.	
	No.	Value.	No.	Value.	No.	Value.	No.	Value.
Salteries.....	6		4		2		12	
Transporting vessels:								
Steamers and launches.....	1	\$2,500	1	\$12,000	1	\$5,000	3	\$19,500
Tonnage.....	7		40		9		56	
Outfit.....		500		2,400		1,600		4,500
Sailing.....	1	900					1	900
Tonnage.....	16						16	
Outfit.....		200						200
Launches under 5 tons.....	5	6,550	1	1,000	1	4,500	7	12,050
Boats, row and sail.....	16	870	39	1,160	23	8,700	78	10,730
Lighters and scows.....	5	400	2	200			7	600
Apparatus:								
Haul seines.....	2	350	22	2,230			24	2,580
Purse seines.....	10	2,800					10	2,800
Gill nets.....	6	800			23	1,725	29	2,525
Cash capital.....		8,200		11,250		35,000		54,450
Shore and accessory property.....		15,300		9,500		27,000		51,800
Wages paid.....		5,925		16,577		22,590		45,092
Total.....		45,295		56,317		106,115		207,727

Output.—The output in 1910 amounted to 14,405 barrels, valued at \$130,641, as compared with 26,915 barrels and 6,997 half barrels, valued at \$208,758, in 1909. A small part of this output is composed of salmon bellies. A few of the backs were pickled and appear in this table, while the rest were either dried, dry-salted, or smoked, and appear under their proper headings in this report.

BARRELS OF SALMON PICKLED IN 1910, BY SPECIES.

Products.	Southeast Alaska.		Central Alaska.		Western Alaska.		Total.	
	No.	Value.	No.	Value.	No.	Value.	No.	Value.
Coho, or silver.....	35	\$296	125	\$1,208			160	\$1,504
Coho bellies.....			126	1,135			126	1,135
Dog, or chum, bellies.....	70	770					70	770
Humpback, or pink.....	314	1,905	13	75	3	\$15	330	1,998
Humpback bellies.....	421	4,410	195	1,725			616	6,135
King, or spring.....					352	3,399	352	3,399
King fins.....	2	24					2	24
King bellies.....	6	128					6	128
Red, or sockeye.....	2	20	1,485	12,278	10,444	92,351	11,931	104,649
Red tips.....					4	60	4	60
Red bellies.....	3	24	805	10,815			808	10,839
Total.....	853	7,577	2,749	27,239	10,803	95,825	14,405	130,641

MILD CURING.

At the opening of the present season the mild-curing industry was in better condition than for several years previous, as the pack of 1909 had been disposed of and prices for the new pack were ruling fairly high. Owing to this the packers extended their operations as much as possible, and as a result the pack this year is the largest ever put up in Alaska.

With the exception of a small quantity put up in Cook Inlet, central Alaska, the packing of mild-cured salmon was confined to southeast Alaska, although it is more than probable that the packers will soon extend their operations into western Alaska and parts of central Alaska not now worked.

As in previous years the principal trouble the packers experience is in getting rid of the white-meated king salmon with the least possible loss. These fish average about one-fourth of the total catch, and the fishermen insist that the dealers shall take them along with the others, which they do at a considerably lower price. A few of the larger of these white-meated kings are mild-cured. Early in the season many of them, together with the small red-meated fish, are shipped fresh to the Puget Sound ports, but after the kings begin to run in the Sound this is unprofitable.

Persons engaged.—This year 656 persons (560 fishermen, 68 shoresmen, and 28 transporters) were engaged in the mild-curing industry, as compared with 521 in 1909, a gain of 135. A number of others also were engaged for limited periods, but as their work in connection with other branches of the salmon business was more important they have been included there.

PERSONS ENGAGED IN THE SALMON MILD-CURING INDUSTRY IN 1910.

Occupation and race.	Southeast Alaska.	Central Alaska.	Total.
Fishermen:			
Whites.....	354	10	364
Indians.....	196		196
Total.....	550	10	560
Shoresmen:			
Whites.....	65		65
Indians.....	3		3
Total.....	68		68
Transporters:			
Whites.....	15		15
Indians.....	13		13
Total.....	28		28
Grand total.....	646	10	656

Investment.—There were 14 fixed plants (13 in southeast Alaska and 1 in central Alaska)—i. e., plants with permanent buildings and a chief business of mild-curing salmon—operated in Alaska this year. A considerable part of this industry is done by schooners and launches, the crews of which catch the fish in small boats and pack them aboard the vessels, moving from place to place with the schools of salmon.

INVESTMENT IN THE SALMON MILD-CURING INDUSTRY IN 1910.

Items.	Southeast Alaska.		Central Alaska.		Total.	
	No.	Value.	No.	Value.	No.	Value.
Fixed plants.....	13		1		14	
Transporting vessels:						
Steamers and launches (over 5 tons).....	23	\$51,500			23	\$51,500
Tonnage.....	179				179	
Outfit.....	2	35,000			2	35,000
Sailing vessels.....	2	4,000			2	4,000
Tonnage.....	67				67	
Outfit.....		3,000				3,000
Steamers and launches (under 5 tons).....	35	42,750			35	42,750
Boats, sail and row.....	402	14,365	5	\$1,000	407	15,365
Scows.....	20	10,100			20	10,100
Apparatus, shore fisheries:						
Gill nets.....	138	26,225	5	750	143	26,975
Lines, trolling.....		471				471
Shore and accessory property.....		40,920				40,920
Cash capital.....		86,000				86,000
Wages paid.....		46,537		1,200		47,737
Total.....		360,868		2,950		363,818

^a Includes outfit.

Catch, by apparatus and products.—All told, 164,520 red-meated and 22,525 white-meated king salmon were required in preparing the pack. The greater part of these fish were caught with trolling lines. The pack of 3,357 tierces, which sold for \$220,673, is an increase of 1,065 tierces and \$71,373 over 1909.

CATCH OF SALMON FOR MILD-CURING, 1910, BY APPARATUS AND SPECIES.

Apparatus and species.	Southeast Alaska.	Central Alaska.	Total.
Gill nets:	<i>Number.</i>	<i>Number.</i>	<i>Number.</i>
Red king salmon.....	20,864	1,767	22,631
White king salmon.....	2,656		2,656
Total.....	23,520	1,767	25,287
Lines:			
Red king salmon.....	141,889		141,889
White king salmon.....	19,869		19,869
Total.....	161,758		161,758
Grand total.....	185,278	1,767	187,045

PRODUCTS OF THE SALMON MILD-CURING INDUSTRY IN 1910.

Products.	Tierces.	Round weight of fish.	Dressed weight of fish.	Value.
Southeast Alaska:		<i>Pounds.</i>	<i>Pounds.</i>	
Red king salmon.....	3,022	3,475,300	2,468,198	\$209,826
White king salmon.....	304	349,600	246,700	8,615
Total.....	3,326	3,824,900	2,714,898	218,441
Central Alaska:				
Red king salmon.....	31	35,650	24,800	2,232
Total:				
Red king salmon.....	3,053	3,510,950	2,492,998	212,058
White king salmon.....	304	349,600	246,700	8,615
Grand total.....	3,357	3,860,550	2,739,698	220,673

FRESH SALMON.

As in previous years large quantities of king salmon (mainly white-meated and small red-meated fish) were shipped fresh to Puget Sound ports, where they brought very good prices up to the time king salmon began to run in the Sound waters.

Shortly after the canning season opened certain fishermen with headquarters at Petersburg and Wrangell became dissatisfied with the prices offered by neighboring canneries, and failing to come to an agreement began shipping their catches of red and coho salmon fresh to Puget Sound ports, where they received fair prices.

MINOR PRESERVING PROCESSES.

Dry salting and drying.—At a few places in central Alaska the bellies of red and coho salmon are cut out and pickled, after which the backs are dried in the sun, and the resulting product, called "ukalu," used for fox food at the fox ranches and for dog food.

The dry salting of dog salmon for food has almost ceased, but 22,178 pounds, valued at \$554, being prepared this year.

Smoking.—A delicious smoked product, known locally as "beleke," is put up at Kodiak and several other places, the backs of red, coho, and humpback salmon being used. A considerable quantity of white-meated king salmon, cut into steaks, was smoked in southeast Alaska this year.

Freezing.—The only establishments engaged in freezing salmon are at Taku Harbor and Ketchikan, in southeast Alaska. Only a small business is done in the freezing of salmon, halibut being the principal product of these plants. Black bass, black cod, and steel-head trout are among the miscellaneous products prepared.

RETURN OF MARKED SALMON.

A number of salmon bearing mutilations of certain fins, apparent brands, or with missing fins, were observed during the summer, as occurs every season. So far as these concern single fins they are not to be referred to any known artificial marks placed upon fish as a means of identification. Twelve of them, however, were red salmon lacking both ventral fins and are identified as returns from a definite marking experiment which has yielded annual results since 1906. This continued return of marked red salmon to southeast Alaska is of particular interest. These fish were marked by Mr. F. M. Chamberlain as fingerlings about three months old, in August, 1903, at Fortmann hatchery, and liberated in Naha Stream above Heckman Lake. The mark consisted of the complete excision of both ventral fins. The number of marked fish liberated was 1,600. The returns which are considered to have been satisfactorily identified are shown, by the year and locality, in the following table:

MARKED SALMON IDENTIFIED UPON RETURN TO STREAMS, 1906-1910.

Years.	Naha.	Yes Bay.	Karluk.	Total.	Age of fish.
1906.....	2	2	Years. 3½
1907.....	13	13	4½
1908.....	5	3	8	5½
1909.....	4	1	5	6½
1910.....	1	10	1	12	7½
Total.....	21	17	2	40

One of the 10 fish credited to Yes Bay in 1910 was caught in the bay by commercial fishermen and preserved by freezing at Ketchikan, where it was examined by the assistant agent on July 23. It was a male 20.5 inches in length and weighed 3¾ pounds. All the other marked fish assigned to Yes Bay for any year were taken at the Government hatchery at the head of Yes Lake.

These 40 fish are 2½ per cent of the 1,600 marked. The observed return is certainly somewhat larger and possibly greatly surpasses these figures. An indeterminate number, estimated at between 50 and 100, were reported to have been seen at Yes Lake hatchery in 1906, but of these no specimens were saved. No account has been taken of these in the above table, since there is no basis for determining how many of the presumed marks were certainly of the same nature as those accepted as representing actual returns. Salmon lacking a single ventral fin are frequently seen in the runs, and some mutilations of this pair of fins are to be distinguished from the results of artificial marking. While the table shows but one marked fish

taken at Karluk in 1909, several were reported, the exact number being unknown. The one of which account has been taken is based upon examination of a preserved specimen. The few taken at Karluk are the only specimens known to have returned outside of south-east Alaska.

The relation of the return to the parent stream and adjoining streams of southeast Alaska, in which most of the marked fish were retaken, is of importance. Excluding the uncertain return to Yes Bay in 1906, over half the returning fish succeeded in reaching the parent stream, and even with these Yes Bay fish included, a considerable proportion still belongs to the parent stream, while by far the larger part of the known return is confined to the region within 40 miles of the parent stream. It is obviously indicated that red salmon return to the general region in which they were hatched, rather than to remote regions, and that a considerable number reach the particular region of their origin, or their parent stream.

The return from the original plant of marked fish has now covered five successive seasons, indicating a variation of at least five years in the life period of a single hatch of red salmon. The known return had been diminishing in numbers since 1907 up to the current year, when it considerably increased. This is a somewhat anomalous result, and inconsistent with that gradual dwindling in numbers and disappearance from the runs of fish bearing this mark which was expected to occur. While the acceptance of these fish as conclusively identical with the marked salmon of 1903 depends on the cessation of their occurrence within a reasonable time, there is at present no sufficient reason for doubting that they are the same.

OBSERVATIONS IN WOOD RIVER REGION.

Mr. H. C. Fassett, inspector of fisheries in Alaska, represented the Bureau in western Alaska, with headquarters on Nushagak Bay, and had charge of the investigations in the Nushagak region. The order closing both Wood and Nushagak Rivers was uniformly observed, and without its restrictive effect a considerable proportion of the reduced quota escaping to the spawning grounds through Wood River would have been taken. Eight fish traps were operated on the bay and two in Igushik River, the latter yielding but few fish. The total take of traps was about 596,000, of which about 29 per cent were red salmon. These traps took 11.2 per cent of the whole catch of the Nushagak region, and 3.9 per cent of the whole red salmon catch.

The following table shows the total Nushagak catch (including 85,000 red salmon from Igushik River) and its content as to the five species of salmon. The red salmon catch is 83.5 per cent of the total number of salmon taken.

CATCH OF DIFFERENT SPECIES OF SALMON IN NUSHAGAK REGION, 1910.

Species.	Catch.	Species.	Catch.
King.....	86,433	Pink.....	440,369
Red.....	4,427,626	Dog.....	206,220
Coho.....	139,200	Total.....	5,299,848

COUNT OF THE BREEDING RUN IN WOOD RIVER.

The count of salmon escaping from the fishermen and ascending to the spawning grounds by way of Wood River was again made as in the two past years. The actual daily tally made at the rack at the foot of Lake Aleknagik is as follows:

DAILY TALLY OF REDFISH INTO LAKE ALEKNAGIK DURING THE SEASON OF 1910.

Date.	Number.	Date.	Number.	Date.	Number.
July 4.....	167	July 15.....	125,621	July 26.....	1,162
5.....	1,042	16.....	64,026	27.....	927
6.....	2,717	17.....	29,964	28.....	715
7.....	12,036	18.....	31,628	29.....	873
8.....	13,131	19.....	13,642	30.....	708
9.....	72,073	20.....	10,928	31.....	385
10.....	105,835	21.....	10,000	Aug. 1.....	361
11.....	70,252	22.....	4,881	2.....	139
12.....	26,772	23.....	3,618	Total.....	670,104
13.....	24,223	24.....	2,747		
14.....	37,612	25.....	1,919		

The run came into Nushagak Bay about July 3. The rack at the lake was completed and made tight on July 3, but no fish were seen until the 4th. The tally of July 7 probably represents the advance of the main run. As in the preceding year, there were two distinct impulses in the run at the lake, the height of the run or largest tally occurring on the 15th, or one day later than in the two preceding seasons.

RECORD OF METEOROLOGICAL OBSERVATIONS AT THE SALMON RACK AT LAKE ALEKNAGIK, ALASKA, DURING SEASON OF 1910.

Date.	Hour.	Temperatures.			Barom-eter read- ing.	Weather conditions.			Lake conditions.			Remarks.
		Air.		Lake at 6 feet depth.		Clouds in sky (amount 1 to 10).	Wind (force and direction).	Rainfall (heavy, moderate, light, trace).	Depth at rack-gate.	Current at rack (per min- ute).	Drift (much, little, none).	
		At read- ing.	Max- imum.									
June 25	12 m.	53	58	51	30.16	10	SE, light.				Much.	
	6 p. m.	50	52	50	30.19	10	S, light.				Much.	
	12 p. m.	42	49	42	30.17	10	SW, light.	Trace.			Much.	
	6 a. m.	40	42	37	30.21	10	SW, light.				Much.	
	12 m.	43	43.5	39.5	30.20	10	N, light.				Much.	
June 27	6 p. m.	43.5	46.5	40.0	39.5	10	Calm.	Trace.			Little	Lake like glass.
	12 p. m.	39.5	43.0	38.5	39.6	10	S, light.	Trace.			Little	
	6 a. m.	42.0	42.5	39.0	40.0	10	NW, moderate.	Trace.			Little	
	12 m.	45.2	44.8	40.8	40.0	10	SW, moderate.	Light			Little	
	6 p. m.	44.6	44.8	43.5	39.6	10	SSE, moderate.	Light			Little	
June 28	12 p. m.	41.5	44.0	41.0	40.0	10	S, moderate.	Light			Little	
	6 a. m.	40.2	41.0	38.5	39.6	10	SW, moderate.	Light			Little	
	12 m.	44.2	43.2	39.0	39.6	10	W, light.	Light			Little	
	6 p. m.	45.0	42.2	41.8	39.8	10	W, light.	None.			Little	
	12 p. m.	42.0	42.2	41.8	39.8	10	WSW, light.	Mist.			Little	
June 29	6 a. m.	50.0	49.3	37.6	40.5	10	NNE, moderate.	None.			Little	
	12 m.	44.0	49.7	44.0	40.0	10	SE, light.	None.			Little	
	6 p. m.	43.5	43.8	42.0	40.0	10	S, light.	Light			Little	
	12 p. m.	42.2	43.8	39.8	40.0	10	SW, light.	Light			Little	
	6 a. m.	43.0	43.0	40.0	39.8	10	SE, light.	Light			Little	
July 1	6 p. m.	45.5	45.0	43.0	40.0	10	ESE, light.	Light			Little	
	12 p. m.	43.0	45.0	42.0	40.0	10	S, moderate.	None.			Little	
	6 a. m.	42.8	45.0	39.8	40.0	10	E, light.	Light			Little	
	12 m.	47.0	48.0	42.6	39.8	10	W, light.	Light			Little	
	6 p. m.	45.2	48.5	43.0	40.0	10	E, light.	Light			Little	
July 2	12 p. m.	43.0	45.8	42.8	40.0	10	NW, light.	Light			Little	
	6 a. m.	43.5	44.0	41.2	39.81	10	NW, light.	Light			Little	
	12 m.	50.2	50.0	43.0	40.2	10	NW, light.	Light			Little	
	6 p. m.	54.5	53.8	43.0	40.6	10	W, light.	None.			Little	
	12 p. m.	49.0	51.2	49.2	40.0	10	NW, light.	Light			Much.	
July 3	6 a. m.	42.6	44.8	41.2	39.75	10	NW, light.	None.			Much.	
	12 m.	40.2	49.5	42.7	40.8	10	N, light.	None.			Much.	
	6 p. m.	46.0	52.8	43.5	39.75	10	S, light.	Light			Much.	
	12 p. m.	46.0	45.0	43.0	40.0	10	S, light.	None.			Much.	
	6 a. m.	53.0	51.0	43.0	39.75	9	N, light.	Light			Much.	
July 4	6 a. m.	63.2	63.2	42.0	29.84	9	SE, light.	None.			Much.	
	12 m.	63.0	63.2	42.0	29.87	9	SE, light.	None.			Much.	

RECORD OF METEOROLOGICAL OBSERVATIONS AT THE SALMON RACK AT LAKE ALEKNAGIE, ALASKA, DURING SEASON OF 1910—Continued.

Date.	Hour.	Temperatures.			Barometer reading.	Weather conditions.		Lake conditions.			Remarks.		
		Air.		Lake at 6 feet depth.		Clouds in sky (amount 1 to 10).	Wind (force and direction).	Rainfall (heavy, moderate, light, trace).	Depth at rack-gate. (per minute).	Current at rack (per minute).		Drift (much, little, none).	
		At reading.	Maxi-mum.										Mini-mum.
July 4	6 p. m.	59.8	65.1	56.0	41.0	10	NW, light.	Light.	11	7.2	Much.	Showers in p. m.	
	12 p. m.	46.2	61.0	46.0	41.0	10	N.E., light.	None.	11	7.9	Little.		
	6 a. m.	50.2	69.8	41.0	40.8	30.03	9	N., light.	None.	11	7.7		Much.
5	12 p. m.	61.8	63.5	41.6	41.0	8	SE, moderate.	None.	11	7.7	Much.	Hot sunshine.	
	6 p. m.	64.2	63.5	55.5	41.5	8	S., light.	None.	11	7.5	Much.		
	12 p. m.	44.6	67.0	44.6	41.0	30.12	2	N., light.	None.	11	7.5		Much.
6	6 a. m.	50.7	51.0	37.5	41.0	3	W., light.	None.	11	7.6	Much.	More drift than usual.	
	12 m.	76.2	59.2	52.5	41.0	30.24	2	S., light.	None.	11	7.5		Much.
	6 p. m.	60.2	78.0	59.2	41.0	30.25	3	SE, moderate.	None.	11	7.0		Much.
7	12 p. m.	50.0	60.8	50.0	41.0	30.30	8	SE, light.	None.	11	7.0	Much.	Gloomy and threatening.
	6 a. m.	44.8	44.8	37.2	41.0	30.32	9	FSE, moderate.	None.	11	7.0	Much.	
	12 m.	57.2	57.0	42.1	41.8	30.33	7	SE, light.	None.	11	6.7	Much.	
8	6 p. m.	55.1	62.8	51.2	42.0	30.33	8	W., light.	None.	11	6.7	Much.	Gloomy and threatening.
	12 p. m.	43.2	68.0	43.2	42.0	30.33	7	ESE, light.	None.	11	6.6	Much.	
	6 a. m.	42.2	43.0	40.7	41.0	30.33	9	E., light.	None.	11	6.6	Much.	
9	12 p. m.	62.5	60.0	40.2	41.2	30.33	2	S., light.	None.	11	5.8	Little.	Gloomy and threatening.
	6 p. m.	69.1	67.8	59.1	42.0	30.27	8	W., light.	None.	11	5.7	Much.	
	12 p. m.	52.6	66.2	48.5	42.0	30.24	7	N.E., light.	None.	11	5.6	Much.	
10	6 a. m.	60.2	63.0	42.5	42.0	30.19	8	N.E., fresh.	None.	11	5.2	Little.	Gloomy and threatening.
	12 p. m.	60.2	65.2	52.5	42.0	30.16	9	N.E., light.	None.	11	5.0	Little.	
	6 p. m.	52.2	61.0	51.0	41.0	30.15	10	SW, light.	None.	11	5.0	Little.	
11	12 p. m.	48.9	48.8	48.8	41.0	30.18	10	N., light.	None.	11	4.2	Little.	Gloomy and threatening.
	6 a. m.	49.9	46.0	41.0	41.0	30.21	10	SW, light.	None.	11	4.0	Little.	
	12 p. m.	59.2	49.5	42.5	42.5	30.22	8	WSW, light.	None.	11	3.8	Little.	
12	6 p. m.	57.7	68.8	51.2	43.5	30.15	7	WSW, light.	None.	11	3.1	Little.	Gloomy and threatening.
	12 p. m.	58.1	57.6	53.6	43.5	30.15	4	NW, light.	None.	11	2.7	Little.	
	6 a. m.	50.2	55.2	45.0	42.0	30.11	7	NW, light.	None.	11	2.4	Little.	
13	12 m.	59.7	72.2	50.0	44.8	30.11	10	S., light.	None.	11	2.0	Little.	Gloomy and threatening.
	6 p. m.	61.0	71.2	56.8	43.0	30.09	8	W., fresh.	None.	11	1.8	Little.	
	12 p. m.	48.4	59.2	47.5	42.0	30.09	9	WSW, light.	None.	11	1.0	Much.	
14	6 a. m.	46.2	49.8	40.2	43.2	30.15	6	W., light.	None.	11	0.8	Much.	Gloomy and threatening.
	12 p. m.	76.1	77.2	46.4	44.0	30.15	2	SW, fresh.	None.	11	0.5	Much.	
	6 p. m.	51.3	77.2	50.0	42.0	30.21	6	N., light.	None.	11	0.2	Much.	
15	12 p. m.	44.2	50.8	43.0	42.0	30.21	1	NW, light.	None.	10	11.5	Much.	Hard blow from west.
	6 a. m.	48.0	48.8	41.4	42.0	30.21	1	NW, light.	None.	10	11.0	Much.	
	12 m.	57.8	57.2	42.7	45.0	30.22	4	W., heavy.	None.	10	10.5	Much.	
6 p. m.	58.2	49.3	49.3	45.0	30.22	3	N., light.	None.	10	10.5	Much.		

14	12 p. m.	53.0	51.1	45.5	42.5	30.27	0	N., light.	None.	10	10.3	Much.
	6 a. m.	51.6	40.4	42.0	42.0	30.33	8	NW., light.	None.	10	9.4	Much.
	12 p. m.	57.9	59.0	44.5	44.5	30.34	4	W., light.	None.	10	9.0	Much.
	6 p. m.	62.4	69.2	50.2	45.0	30.29	4	S., light.	None.	10	8.5	Much.
15	12 p. m.	41.8	60.2	41.8	45.0	30.24	10	S., light.	Fog.	10	8.0	Much.
	6 a. m.	39.4	42.4	38.4	47.0	30.42	10	NW., light.	Fog.	10	7.5	Do.
	12 p. m.	49.8	50.6	39.0	44.6	30.11	10	NW., light.	None.	10	7.0	Do.
	6 p. m.	50.8	53.2	49.0	44.0	29.99	10	Calm.	Light.	10	6.5	Do.
16	12 p. m.	49.0	52.0	47.5	44.0	29.98	10	Calm.	Light.	10	6.0	Do.
	6 a. m.	45.8	51.0	43.6	43.0	30.00	10	Calm.	None.	10	5.0	Warm.
	12 p. m.	65.0	65.0	41.5	45.0	30.07	9	NW., light.	None.	10	4.5	Warm.
	6 p. m.	58.5	66.3	50.9	43.0	30.08	9	Calm.	None.	10	5.0	Warm.
	12 p. m.	50.2	58.5	50.2	44.0	30.14	9	Calm.	None.	10	4.5	Warm.
17	12 p. m.	46.3	50.8	45.1	42.5	30.18	10	Calm.	None.	10	4.0	Warm.
	6 a. m.	48.0	44.8	44.8	43.2	30.20	10	Calm.	Fog.	10	3.5	Foggy.
	12 p. m.	47.3	40.7	40.7	43.2	30.20	10	SSE., light.	Mist.	10	3.0	Foggy.
	6 p. m.	40.3	40.7	43.0	43.0	30.24	10	SSE., light.	None.	10	3.5	Disagreeable.
18	12 p. m.	42.0	40.7	40.7	43.0	30.27	10	SSE., light.	None.	10	2.5	Disagreeable.
	6 a. m.	46.8	40.8	43.5	43.0	30.35	10	SSE., light.	None.	10	2.0	Boisterous.
	12 p. m.	47.8	40.8	44.5	43.5	30.36	8	SSE., light.	None.	10	1.3	Misty.
	6 p. m.	43.5	40.8	44.0	44.0	30.38	10	SSE., light.	None.	10	0.7	Gloomy.
19	12 p. m.	48.2	42.8	43.5	44.0	30.34	10	SSE., light.	None.	10	0.3	Sun at times.
	6 a. m.	48.2	41.7	43.0	43.0	30.27	10	SSE., moderate	Light.	10	0.0	Threatening.
	12 p. m.	47.0	51.3	46.5	43.0	30.27	10	SSE., moderate	Moderate	9	11.5	Stormy.
	6 p. m.	45.5	51.4	44.8	43.0	30.25	10	SSE., light.	Light.	9	11.0	Stormy.
20	12 p. m.	45.5	52.8	44.7	42.5	30.14	10	SSE., light.	Light.	9	10.5	Gloomy.
	6 a. m.	46.5	52.8	44.7	42.5	30.14	10	Calm.	Mist.	9	10.3	Gloomy.
	12 p. m.	49.5	53.0	44.7	42.5	30.09	10	SE., light.	Mist.	9	10.0	Thick; misty.
	6 p. m.	49.5	53.0	44.8	43.0	30.07	8	S., light.	None.	9	9.5	Do.
	12 p. m.	48.3	53.7	47.2	43.0	30.08	10	SSW., moderate	Moderate	9	9.3	Bright moonlight.
21	6 a. m.	46.5	51.7	44.3	43.5	30.06	10	SW., moderate	Heavy	9	9.0	Clearing.
	12 p. m.	46.0	50.0	42.0	44.0	30.03	6	SW., moderate	None.	9	8.7	

RECORD OF METEOROLOGICAL OBSERVATIONS AT THE SALMON RACK AT LAKE ALEKNAGIK, ALASKA, DURING SEASON OF 1910—Continued.

Date.	Hour.	Temperatures.			Barom-eter read-ing.	Weather conditions.			Lake conditions.			Remarks.	
		Air.				Clouds in sky (amount 1 to 10).	Wind (force and direction).	Rainfall (heavy, moderate, light, trace).	Depth at rack-gate.	Current (per minute).	Drift (much, little, none).		
		At read-ing.	Maxi-mum.	Mini-mum.									Lake at 6 feet depth.
July 26	6 p. m.	54.1	60.2	48.3	Inches. 30.55	0	ESE., light.	None.	8	11.5	None.	Pleasant.	
	12 p. m.	41.8	60.0	41.8		4	E., light.	None.	8	11.9	None.	None.	Misty and raw.
	6 a. m.	42.0	45.7	40.5		4	E., light.	None.	8	10.5	None.	None.	Cloudy and cool.
27	12 m.	52.5	56.7	41.8	4	ESE., light.	None.	8	9.5	144	None.	Pleasant.	
	6 p. m.	57.2	63.6	52.0	4	E., light.	None.	8	9.0	None.	None.	Cloudy and cool.	
	12 p. m.	46.4	60.3	45.6	8	Calm.	None.	8	8.5	150	None.	Overcast.	
28	6 a. m.	44.3	49.8	43.3	10	Calm.	None.	8	8.3	None.	None.	Pleasant.	
	12 m.	60.3	75.1	44.3	9	S., light.	None.	8	8.0	None.	None.	Do.	
	6 p. m.	56.8	75.0	56.4	9	S., light.	None.	8	7.5	None.	None.	Overcast.	
29	12 p. m.	46.9	59.2	46.9	10	S., light.	None.	8	7.0	None.	None.	Rainy.	
	6 a. m.	44.7	50.7	44.2	10	Calm.	None.	8	6.8	160	None.	Pleasant.	
	12 m.	53.0	85.6	44.2	2	Calm.	None.	8	6.5	None.	None.	Do.	
30	6 p. m.	83.0	86.8	58.7	1	SW., light.	None.	8	6.0	None.	None.	Do.	
	12 p. m.	43.0	64.0	43.0	5	Calm.	SSE., moderate.	None.	8	5.5	None.	Unsettled.	
	6 a. m.	40.8	47.6	38.4	10	Calm.	None.	8	5.0	150	None.	Stormy.	
31	12 p. m.	41.4	49.5	40.8	10	Calm.	None.	8	4.8	None.	None.	Misty.	
	6 p. m.	47.3	51.1	40.3	10	Calm.	None.	8	4.5	None.	None.	Pleasant.	
	12 p. m.	44.4	50.7	42.4	2	W., light.	None.	8	4.0	None.	None.	Do.	
Aug 1	6 p. m.	47.7	50.8	40.0	2	WSW., light.	None.	8	3.5	140	None.	Unsettled.	
	12 m.	52.6	69.8	47.8	10	SSW., light.	Moderate	8	3.0	None.	None.	Stormy and cold.	
	6 a. m.	48.8	57.9	44.8	10	S., light.	None.	8	2.5	None.	None.	Unsettled.	
2	12 p. m.	43.2	48.0	43.0	10	Calm.	None.	8	2.3	None.	None.	Stormy.	
	6 a. m.	44.5	47.8	43.0	10	WSW., light.	None.	8	2.0	130	None.	Unsettled.	
	12 m.	56.1	64.7	44.1	10	Calm.	None.	8	2.3	None.	None.	Do.	
3	6 p. m.	54.5	72.4	53.8	10	SSE., light.	None.	8	2.0	None.	None.	Stormy.	
	12 p. m.	42.0	57.4	42.0	10	NE., moderate.	None.	8	2.0	None.	None.	Do.	
	6 a. m.	44.7	47.8	41.7	10	SSE., moderate.	None.	8	2.0	100	None.	Do.	
3	6 p. m.	44.2	50.3	42.8	10	SSE., moderate.	None.	8	1.5	None.	None.	Do.	
	12 p. m.	41.5	46.0	41.5	10	SSE., light.	Trace.	8	1.3	None.	None.	Boisterous.	
	6 a. m.	42.3	46.4	40.5	10	Calm.	None.	8	1.0	None.	None.	Moderating.	
12 m.	50.0	53.5	42.5	9	SW., light.	None.	8	1.0	(Demolition of rack com- menced.)	None.	Unsettled.		
6 p. m.	49.2	54.6	48.8	8	SW., light.	Trace.	8	1.0	None.	None.	Do.		
12 p. m.	42.0	52.2	42.0	10	Calm.	None.	8	1.0	None.	None.	Do.		

SIGNIFICANCE OF WOOD RIVER DATA.

The spawning run up Wood River again shows a loss in comparison with the preceding season. The total was 670,000 in 1910, as against 893,000 in 1909. The commercial catch of Nushagak Bay also fell off, being 4,400,000 in 1910 as against 4,900,000 in 1909. The Wood River run in 1910 was 75 per cent of the 1909 run; the Nushagak Bay catch in 1910 was 89.8 per cent of the 1909 catch. Thus in each of these years the Wood River spawning run has declined much more rapidly than the catch in the bay has declined. The following table shows the numerical results in round numbers for the three years of Wood River investigations. The last column gives the sum of the bay catch and the Wood River run, this total constituting far the greater part of the whole run into Nushagak Bay.

SPAWNING RUN IN WOOD RIVER, 1908, 1909, AND 1910.

Years.	Nushagak Bay catch.	Wood River tally.	Total.
1908.....	6,400,000	2,600,000	9,000,000
1909.....	4,900,000	893,000	5,793,000
1910.....	4,400,000	670,000	5,070,000

The commercial catch for the whole bay has fallen off since 1908 by two annual losses of $1\frac{1}{2}$ millions and $\frac{1}{2}$ million, respectively. The corresponding loss to the Wood River tally was in 1909 numerically even greater than the loss on the catch, while in both 1909 and 1910 the percentage loss in Wood River was greater than on the catch.

According to observations in the river and the head of the bay, and the reports of the packers, the run up the main river was unusually large this season, evidently greater than the Wood River run. By taking the latter as a minimum and twice the number as a maximum for the main river run, and estimating otherwise on the same basis as in previous seasons, about 6,400,000 is obtained as the estimated run for the whole bay in 1910, which in view of the maximum error probable may be accepted as within one-half million of the actual run. Of this estimate over 79 per cent, or more than 5 million fish, are fish actually counted in Wood River by the observers and in Nushagak by the commercial fishermen.

The total escape to the spawning grounds for the whole Nushagak region during the current season lies between 25 per cent and 36 per cent of the total run, with 31 per cent probable. In other words, the industry took between 64 per cent and 75 per cent of the whole run, and probably took about 69 per cent.

As bearing on the rate of increase the figures for the season corroborate broadly the conclusions reached the year previously and tend

to narrow the limits between which this rate is indicated to lie. From such a slender basis of facts as are available, a rate of increase of from 200 per cent to 250 per cent is to be inferred if there is neither under nor overfishing. If these figures are too high the Nushagak industry is overfishing. If they are too low, fish are being uselessly wasted to the spawning grounds. The latter of these alternatives would hardly be maintained by anyone, and can hardly hold over a course of years, yet it may possibly be true of an occasional season, such as that of 1908.

Value of a census of salmon runs.—If the establishment of the increment percentage, rate of increase, or measure of the tendency of red salmon to multiply by their own natural and unaided reproductive powers is of any importance to the fisheries, then the Wood River investigations or their counterpart ought to be continued and made to include a complete salmon catchment basin, the larger and more isolated the better. It can hardly be maintained that the factors of temperature, wind, chance, etc., affect so erratically the movements of the great schools that the annual run to a given basin is little or not at all related to the preceding spawning runs which escaped capture therein. Salmon of course do not all return to the region where they were hatched. Some go elsewhere and a continuous flux or ebb and flow of interchange results.

But the number of the spawners inevitably measures the reproductivity. If this number could be ascertained for all Alaska, it would soon be known how prolific the salmon are. Since this is impossible it remains to make the determination on as large a section of the spawning grounds as can be handled. A somewhat longer time is required in order that the annual variations affecting the particular fragment of the fishery under observation shall reach an average making it representative of the whole. It matters little whether the adult salmon return to their parent waters, or whether they interchange freely, even to the extent of none returning to their birthplaces. The essential point is to determine how large are the runs which succeed year after year to a series of known spawning escapes.

As a matter of fact, there is much difference of opinion among fishermen respecting the controlling effect of winds on the movements of salmon. In Bering Sea few days pass without strong blows, and it is easy to relate the suddenly arriving salmon run to some particular wind, just as the so-called equinoctial storm is supposed to have some essential connection with the autumnal equinox. But whatever resultant physical influences have, they do not prevent an unfailling annual rush of hordes of red salmon into Nushagak Bay, their advent predictable almost to the day and their numbers expected with perfect certainty to be measured in millions. During the countless years in which this has occurred before the commercial fishery

existed the uniformity was presumably greater than at present. The variations in size of the run known to have occurred since man disturbed the balance of nature in these fisheries are reasonably due mainly to the exigencies of the commercial industry, which has been unable to make any correlation between its take and the quota necessary for spawning. Even with these variations, no such thing as a failure in the run is known to history or tradition. Even at the lowest ebbs of the commercial fishery the salmon had still to be counted by millions. As fisheries go, the Nushagak region and most of the Bristol Bay streams are constant and perennial sources of salmon.

That the determination of the rate of increase of red salmon, or the limits within which it varies, is a matter of high importance is self-evident. Of course a high rate has already been implied by the great productivity of salmon fisheries and their failure in Alaska to deplete rapidly under enormous drains. Presumably it has been known to many that the fishermen have been, in many fisheries, taking almost every year more than half the run. The lesser portion must therefore have reproduced the whole run, which placed the annual increment at over 100 per cent. Just how small this escaping portion may be and still reproduce a maximum run has been and is yet the vital and crucial question. But three long steps in answer have been taken by the three years of Wood River investigations.

There is no other way to obtain this increment percentage than by continued counting of the breeders, which, with the commercial catch, amounts to a census of the run. The three annual counts already made in Wood River, coupled with general knowledge of the other rivers of the bay, already show roughly what proportion of the Nushagak Bay run has reached the spawning grounds in these years, and since the Bering Sea fisheries are not rapidly declining this is probably not much below the proportion which should reach the spawning grounds.

This showing is definite enough to be safely used in a practical way as a basis for dividing the whole run into a commercial and a breeding quota. At the beginning the tentative figures might be 70 per cent for the former and 30 per cent for the latter. Seventy per cent is not far from representing the proportion of the run the industry has been taking from Nushagak Bay in each of the past two years. By the use of racks in the rivers the run could be divided as it came into alternate daily portions, one to escape, the other for the packers. Thus a definite proportion of the run would be insured to the spawning grounds, and the actual number of fish of which it consisted would be known. Even if a considerable inaccuracy existed in the tentative fixing of 30 per cent for the breeding quota, no injury would result, for the annual counts would constantly

correct the figures. It is only necessary to begin such a system of catching and releasing at proportions just to the industry and reasonably safe for the fisheries. It may be assumed for this purpose that a 30 per cent escape will approximately maintain the Nushagak fisheries. This implies a rate of increase of 233 per cent, which means that for three salmon which reach the spawning grounds, spawn, and die, ten adult salmon return during the next few years, and that if no more than seven of these are taken by the fishermen the process can continue indefinitely.

The Pacific salmon, and particularly the red salmon, alone among commercial fishes, are surprisingly adapted to the control of man for the purpose of perpetuation and exploitation as a commercial asset. They leave the sea regularly at a certain season and make their way en masse to the narrow channels of the fresh and more or less clear waters, where they may be confined, held, captured, or counted and released to the spawning grounds without injury—all with comparative ease and convenience. Spawning is definitely confined to the single season of sexual maturity and is soon followed by the death of the adult, so that breeding salmon never themselves become a part of subsequent runs. These facts make it possible not only to measure their reproductive power, but to put into effect a system of fishing whereby from a minimum reservation of breeding salmon the fishery may be maintained perpetually at a maximum. At the same time the industry may obtain its fish for packing easily and cheaply. The pack may be made in a perfectly fresh condition. The canneries can operate uniformly throughout the season, instead of with the present alternations of scarcity and abundance. Runs of more uniform size would finally succeed upon a more uniform release of breeders, and would therefore be more accurately predictable.

There is a certain quantity of seed represented by spawning salmon, a more or less definite fraction of the whole run, varying within presumably narrow limits, which nicely produces without waste from the spawning fields and the feeding grounds of the seas a maximum crop of fish. Any greater quantity is an excess, being a total waste of nonproductive seed, while any lesser quantity is a more serious loss, the waste of a multiplied return from potential seed which should have been used as such. No system of fishing can possibly make this measured sowing of the spawning grounds without actually counting the whole run. This the present system does not do. It counts the catch alone, and therefore it almost always wastes fish, either as nonproductive breeders or as the multiplied (by about $2\frac{1}{2}$) return from fish which should have been allowed to breed. The tendency is toward the latter or greater loss. Only occasionally and by chance will both forms of waste be avoided.

These opportunities which the peculiar specialized habits of the red salmon afford for perpetually exploiting them commercially without depleting their abundance should be utilized. The packing industry would greatly profit in the end and the Alaska fisheries would enhance in value as a national asset. At present the law does not provide power to establish such a system of fishing, but it would permit a trial in a suitable region by mutual agreement between the packers concerned and Federal authority.

EXPLORATIONS OF LAKE ALEKNAGIK.

During the summers of 1908 and 1909 every stream tributary to Lake Aleknagik, which gives rise to Wood River, was examined by the agent. During the current summer Mr. W. T. Bower, of the Division of Fish Culture of the Bureau, spent the period from July 17 to July 27 in explorations of the lake and streams. By means of these observations the streams have been thoroughly prospected with reference to spawning salmon and hatchery possibilities. Two suitable and feasible hatchery sites have been selected, and on either a properly equipped expedition, arriving as soon as navigation opened, could erect a hatchery in time to obtain a portion at least of the same season's spawn.

Such a hatchery could be located on the lake shore and be accessible directly from tidewater for light-draft boats. No single stream of the lake would afford eggs enough to fill a large hatchery, and collections would have to be made over the whole lake in some seasons. There is, however, no more suitable location in the Bristol Bay region for accessibility and proximity to large spawning grounds. The second lake could be drawn upon for eggs if necessary. There is no hatchery in western Alaska, a region which furnishes some 63 per cent of the total pack of Alaska red.

THE COD FISHERY.

All but one of the firms and individuals [John H. Nelson, of Squaw Harbor] operating in the district for cod exclusively have their headquarters at San Francisco, Cal., or Seattle, Anacortes, or Tacoma, Wash., at which places, or in their immediate vicinity, the cured fish are received and prepared for marketing. About half of the operators have shore stations located at favorable places in central Alaska, on the Shumagin and Sannak Islands, and Unimak Island. From thence the dory fishermen carry on their operations, bringing in their catch daily, and when they have accumulated enough to form a cargo a vessel is dispatched from the home port or else a fishing vessel completes its fare from the station catch and carries the fish to the curing establishments in the States.

The industry has suffered severely in the past from the spreading broadcast of exaggerated ideas as to its possible profits. As a result of this persons totally unfamiliar with the work have engaged in it, and instead of building up a trade by the preparation of a good product at a living price have prepared goods in a slipshod manner and then disposed of them by cutting below the prices of more reputable dealers.

When the present season opened the trade was in a demoralized condition, owing to excessive cutting of prices. During the summer certain changes in ownership took place. A new company, the Western Codfish Co., took over the plants, vessels, etc., of King & Winge Co. and the Seattle-Alaska Fish Co. The Union Fish Co., of San Francisco, bought and had delivered to it the catches of the vessels owned and operated this year by the Robinson Fisheries Co., of Anacortes, Wash., and the Blom Codfish Co., of Tacoma, Wash.

Through this centralizing of the industry, price cutting was eliminated, temporarily at least, and when this report closed the market was in excellent condition. A considerable surplus is on hand, but the dealers are content to hold this for their own price, which, owing to the shortage of cod on the Atlantic coast, they are reasonably sure of getting.

Mr. J. A. Matheson, of Anacortes, Wash., has incorporated his plant, and it is now known as the Matheson Fisheries Co. The Pacific States Trading Co., of San Francisco, which did not operate this year, will probably resume operations in 1911.

The winter of 1909-10 was severe, and the cod fishermen were very much hampered as a result. Up to June 1 heavy winds prevailed, and after that, while winds were light, heavy fogs were frequent. Owing to the severe weather practically no fish were caught in Dublin Bay.

On March 28 the codfish schooner *Stanley*, owned by the Union Fish Co., of San Francisco, Cal., when approaching Pavlof Harbor, on San-nak Islands, in central Alaska, grounded on a reef and immediately began to go to pieces. In the heavy seas continually breaking over her one man was washed overboard and drowned and three men, including the master, died from exposure before rescuing parties from the shore could reach the ship. The rest of the crew, five men, were saved. The vessel was carrying supplies to the company's shore stations in Alaska, and her loss seriously hampered the operation of these for several months.

SHORE STATIONS.

During 1910 the following shore stations were operated: Alaska Codfish Co.: Unga, Baralof (Squaw Harbor), and Kelleys Rock (Winchester), on Unga Island; and Companys Harbor and Moffats Cove,

on Sannak Island. John H. Nelson: Squaw Harbor, Unga Island. Union Fish Co.: Pirate Cove, Popof Island; Northwest Harbor, Little Koniuji Island; Pavlof Harbor and Johnson Harbor, on Sannak Island; Sanborn Harbor, on Nagai Island; and Unga, on Unga Island. Several which were shut down this year will be operated in 1911.

STATISTICS FOR CENTRAL ALASKA.

During the year 197 fishermen, 22 shoresmen, and 37 transporters were employed. The total investment amounted to \$162,655. The catch amounted to 3,019,023 pounds of fish as taken from the water. When cured this weighed 2,269,914 pounds and sold for \$63,443, a very large decrease from 1909.

PERSONS ENGAGED IN THE CENTRAL ALASKA COD FISHERIES IN 1910.

Occupation and race.	Number.
Fishermen (shore fisheries):	
Whites.....	197
Shoresmen:	
Whites.....	18
Indians.....	3
Chinese.....	1
Total.....	22
Transporters:	
Whites.....	37
Grand total.....	256

INVESTMENT IN THE CENTRAL ALASKA COD FISHERIES IN 1910.

Items.	Number.	Value.	Items.	Number.	Value.
Transporting vessels:			Boats, sail and row.....	197	\$5,950
Steamers and launches.....	3	\$28,000	Apparatus: Hand lines.....		1,205
Tonnage.....	78		Cash capital.....		45,000
Outfit.....		3,500	Stations, with accessory prop- erty.....		39,500
Sailing.....	2	37,500	Total.....		162,655
Tonnage.....	235				
Outfit.....		2,000			

PRODUCTS OF THE CENTRAL ALASKA COD FISHERIES IN 1910.

Products.	Round weight.	Dressed weight.	Value.
	<i>Pounds.</i>	<i>Pounds.</i>	
Cod, fresh.....	16,000	14,000	\$560
Cod, salted.....	2,877,157	2,157,914	59,433
Cod, pickled.....	125,806	94,400	3,320
Cod tongues, salted.....		3,600	130
Total.....	3,019,023	2,269,914	63,443

VESSEL FISHING.

The following fleet ^a of 11 vessels, with headquarters in California and Washington, operated in Alaskan waters this year, several of them spending the winter of 1909-10 in the north.

COD-FISHING FLEET IN ALASKAN WATERS, WINTER OF 1909-10.

Name.	Class.	Net tonnage.	Owner.
Fanny Dutard.....	Schooner.....	252	Matheson Fisheries Co., Anacortes, Wash.
Alice.....	do.....	220	Robinson Fisheries Co., Anacortes, Wash.
Joseph Russ.....	do.....	235	Do.
Maid of Orleans.....	do.....	171	Seattle-Alaska Fish Co., Seattle, Wash.
Vega.....	do.....	233	King & Winge Codfish Co., Seattle, Wash.
Fortuna.....	do.....	138	Blom Codfish Co., Tacoma, Wash.
W. H. Dimond.....	do.....	376	Alaska Codfish Co., San Francisco, Cal.
City of Papeete.....	Barkentine.....	370	Do.
John D. Spreckles.....	Schooner.....	253	Do.
Fremont.....	do.....	328	Union Fish Co., San Francisco, Cal.
Stanley ^a	do.....	253	Do.

^a Lost at sea.

The vessels from Washington operating in Alaskan waters caught 911,500 fish, with a cured weight of 3,563,000 pounds, which sold for \$97,983, while those from California caught 498,399 fish, with a cured weight of 1,992,000 pounds, valued at \$54,780.

THE HALIBUT FISHERY.

FISHING GROUNDS.

The fishery for this very choice food fish occupies second place in the commercial fisheries of Alaska. At present the industry is practically restricted to southeast Alaska, the few fish taken in central Alaska being consumed in the towns in that section. This is due almost wholly to the fact that the present steamship facilities to this section of Alaska are inadequate for the handling of this species as expeditiously as is required. Halibut are reported from various places in Cook Inlet, from all along the Alaska Peninsula and the adjacent islands, and in Prince William Sound.

In western Alaska the fish is reported from a number of places, the natives usually catching and using it for food. The natives of the Pribilof Islands, when fishing off the islands, catch numbers of halibut and these are usually very choice specimens.

In southeast Alaska halibut appear to be most abundant in the numerous sounds and straits during the winter months. Icy, Chatham, Peril, and Sumner Straits, and Frederick Sound are the chief centers of abundance. The best grounds are to be found in Frederick Sound, especially around the Five Finger Islands. Good banks are to be found scattered all over Icy Straits. The waters of

^a None of the data relating to this fleet appear in the statistical tables.

Chatham Strait are too deep for general fishing, but off Point Gardiner and at several spots off Baranof Island, are to be found good fishing banks, while Kootznahoo Inlet, on Admiralty Island, yields good fishing in summer. In Sumner Strait are to be found very good deep-water winter fishing grounds. During the winter of 1909-10 some of the fishermen fished here in water as deep as 250 fathoms. The vicinity of the Eye Opener is the best ground to be found in the strait. Indians fish considerably in Boca de Quadra and the vicinity of Kah Shakes Cove, Mary's Island, and the mouths of Kasaan Bay and Cholmondeley Sound. In Stephens Passage considerable fishing is done in and just off the mouth of Seymour Canal. Most of the fishing in the protected waters of southeast Alaska has heretofore been done in winter, as the fish were then most abundant and the prices realized were better than in summer when the Puget Sound fleet operates on the Flattery Banks, off the Washington coast, and brings the fish in in such abundance that the Alaska-caught fish, which have to be shipped on the steamers plying between Seattle and southeast Alaska ports, at considerable expense, can not compete. This summer, however, the New England Fish Co. bought and froze all halibut brought to its Ketchikan plant and as a result a number of fishermen continued halibut fishing throughout the year.

For many years the Puget Sound steamers and large power vessels fished in Hecate Strait and off the chain of islands lying outside the British Columbia mainland. During the last few years these banks have been growing less and less productive, and as the Canadian fishery protection boats have very much harassed our fishermen who were operating in these waters, or who were driven into its harbors by stress of weather or for wood and water, they have been gradually extending their operations northward into Alaska waters, where they would be free from molestation. It has been known for some years that halibut were abundant at certain regions in the ocean off the outer fringe of islands in southeast Alaska, more particularly off Baranof Island and the mainland between Cape Spencer and Yakutat Bay, and it was surmised that other and possibly more extensive banks would be found if looked for. During the winter of 1909-10 several of the vessels prospected the open waters between Cape Muzon and Sitka, with the result that halibut were found in great abundance throughout the greater part of this area. Off Forrester Island seemed to be the center of greatest abundance. Here an average depth of 80 fathoms is found for about 4 miles from shore; a little farther out it deepens to 150 fathoms. The first few cargoes from here averaged 15 pounds to the fish, but the average soon dropped to 14 pounds. One steamer early in July caught about 250,000 pounds of halibut on the Forrester Island banks during one trip.

Halibut frequent the sandy banks on which coral and a small shellfish known to the fishermen as "sea cocks" abound. The latter is sought by the halibut as a choice morsel of food. The fish is a very voracious and promiscuous feeder. The stomach of one opened at the Ketchikan plant of the New England Fish Co. contained an octopus, a crab, a salmon, and a dogfish. Sand lance and fish eggs of a large size appear to be its favorite food at certain seasons. One dealer reports finding a 6-inch section of a tree branch in the stomach of one. The fishermen say that frequently when pulling up a hooked halibut, other halibut will follow the hooked one to the surface, biting at its tail and body.

A few female halibut with roe reach the dealers, but the fish are usually dressed on the banks, and the roe, when present, is thrown away. Several fish with roe were received by the New England Co. in August and September.

METHODS AND CONDITIONS.

Within the protected area in summer the fish are scattered considerably, but during the winter they school on banks in the waters noted above. During this season the greater part of the year's catch is made by the smaller vessels, which are unable to stand the rough weather usually encountered on the banks in the open ocean.

Dealers located at Hoonah, Juneau, Douglas, Scow Bay, Petersburg, Wrangell, and Ketchikan handle the fish from the fishing boats. Scow Bay, which is on Wrangell Narrows, about 5 miles from its head, is the principal shipping point. Here are moored several large house scows, floats, and barges, alongside of which the fishing boats tie up and deliver their catch, to be boxed in ice for shipment and put aboard the regular steamers for Seattle, which pass through the narrows every few days. The fish are cleaned and packed in ice in bins aboard the vessel on the banks. The fishermen furnish their own ice, which is frequently secured from icebergs which have broken off from nearby glaciers and are floating around in the bays, sounds, and straits. The dealer furnishes the shooks for making the boxes, which hold about 500 pounds. Where glacier ice is not available the fishermen buy from the artificial ice plants, paying from \$3 to \$5 per ton.

A few years ago halibut weighing over 50 pounds were usually fletched aboard the vessel, but the demand for fletched halibut is so small, and the price realized is so inadequate to the work involved, that but few are now prepared in this manner, and these usually on shore. In fletching the sides are taken off in two complete pieces, which are then put into bins and buried in salt so that the brine will run off. It usually requires about three weeks for the fish to strike properly. Half-ground California salt is used in curing.

In shipping fresh, the best fish are from 25 to 30 pounds in weight. A 10-pound fish is quite a small one. Those smaller are known as "chickens." Most of the Alaska halibut are of good grade. But few logy halibut are found; that is, with watery flesh which clings to the knife when cut and does not have the blue tint of the first-class fish.

Sometimes the dealer makes a contract with a vessel owner at a certain fixed figure, but when the fish are received on consignment the commission charged is generally 5 per cent. The dealers usually purchase outright, at the current rates, the fish landed by the small boats.

Large halibut are occasionally taken, one being delivered at Juneau in 1904 which weighed 365 pounds. According to the fishermen the females appear to have well developed eggs at any season of the year.

Shooks for making a halibut box cost from 65 to 70 cents for each box, depending upon the quantity ordered. The only other expense is for nails and the labor required in making the box. The fishermen deliver the halibut at the scows in an eviscerated condition. When being packed for shipment the head is removed and the fish thrown into the box with the tail toward the middle. Under ordinary conditions 1 ton of ice is required for 6 tons of fish, which is quite reasonable when it is taken into consideration that the fish must be carried a distance of over 700 miles by steamer. The freight rate to Seattle varies from \$7 to \$7.50 per cubic ton, depending upon the distance of the shipping point from Seattle. For shipments of less than 6 boxes the rate is somewhat higher. In addition wharfage has to be paid in Alaska (usually about \$1 per ton) and in Seattle (40 cents per ton). Six boxes of fish are considered to weigh 2½ tons.

The greater portion of the Pacific coast halibut is shipped to points east of the Mississippi River, Chicago, New York, and Boston being the principal distributing centers. The demand from the Pacific coast and adjacent States, however, is showing a healthy growth, and will eventually absorb the greater part of the catch.

Heretofore the vessels of the New England Fish Co. have operated from the company's plant in Vancouver, British Columbia, the fish landed from the vessels with American register having been shipped through to places in the United States in bond, free of duty. Since the establishment of the company's station at Ketchikan these steamers have virtually made this place their headquarters and have been so credited in this year's report.

On December 29, 1909 (too late to be included in the report for that year), as the gasoline schooner *Capella* was being towed from Wrangell to Petersburg by the gasoline boat *Neptune*, the latter broke down and both vessels drifted onto the northeast shore of

Vanks Island. The *Capella* became a total wreck, and her master and a sailor lost their lives from exposure and exhaustion after reaching land.

On November 13 the gasoline schooner *Sea Light*, of Ketchikan, while on a halibut fishing cruise, was wrecked at Larch Bay, near Cape Ommaney, in southeast Alaska, during a severe gale. After suffering much hardship the crew of 8 men managed to reach safety in their dories. Later the vessel was found on the beach by another fishing vessel which worked her off and towed her into Petersburg.

STATISTICS.

During the year 1910 there were 829 persons employed in all branches of the halibut industry. The number of steamers and launches increased enormously over 1909, because of the highly remunerative prices realized for halibut the previous year. The catch as reported in 1910 amounted to 21,579,289 pounds, valued at \$808,010, as compared with 5,189,924 pounds, valued at \$195,529 in 1909. Part of this great increase in showing is due to the changing of the headquarters of the New England Co.'s fleet of steamers from Vancouver, British Columbia, to Ketchikan, thus bringing them within the scope of this report.

PERSONS ENGAGED IN THE SOUTHEAST ALASKA HALIBUT FISHERIES IN 1910.

Occupation and race.	Number.	Occupation and race.	Number.
Fishermen:		Shoresmen:	
Vessel fisheries—		Whites.....	29
Whites.....	343	Indians.....	2
Indians.....	34	Total.....	31
Total.....	377	Transporters:	
Shore fisheries—		Whites.....	1
Whites.....	240	Grand total.....	829
Indians.....	180		
Total.....	420		

INVESTMENT IN THE SOUTHEAST ALASKA HALIBUT FISHERIES IN 1910.

Items.	Number.	Value.	Items.	Number.	Value.
Fishing vessels:			Scows.....	5	\$7,600
Steamers and launches....	66	\$468,800	Apparatus:		
Tonnage.....	842		Vessel fisheries, trawl		
Outfit.....		165,049	lines.....		22,080
Sailing.....	3	3,800	Shore fisheries, trawl		
Tonnage.....	35		lines.....		15,870
Outfit.....		875	Cash capital.....		52,500
Packing barges.....	1	15,000	Shore and accessory property.		252,200
Tonnage.....	338		Total.....		1,258,004
Launches under 5 tons.....	151	\$253,330			
Boats, sail and row.....	20	600			

a Outfit included.

PRODUCTS OF THE SOUTHEAST ALASKA HALIBUT FISHERIES IN 1910.

Products.	Round weights.	Dressed weights.	Value.
Vessel catch:	<i>Pounds.</i>	<i>Pounds.</i>	
Halibut, fresh.....	18,251,519	14,601,215	\$702,245
Halibut, frozen.....	2,343,644	1,876,915	69,871
Halibut, fletched.....	66,560	49,920	2,259
Total.....	20,661,723	16,528,050	774,375
Shore catch:			
Halibut, fresh.....	786,482	645,186	29,669
Halibut, frozen.....	123,481	98,785	3,677
Halibut, fletched.....	7,333	5,500	275
Halibut, pickled.....	270	200	14
Total.....	917,566	749,671	33,635
Grand total.....	21,579,289	17,277,721	\$08,010

In Central Alaska 51,000 pounds, valued at \$2,040, was marketed in addition to above.

PUGET SOUND FISHING FLEET.

A fleet of Puget Sound power vessels visits southeast Alaska during the months from October to March, when, owing to stormy weather and a scarcity of fish, it is not safe nor profitable to visit the banks near the home ports. This fleet makes its headquarters mainly at Petersburg, at the head of Wrangell Narrows, shipping the catch home from Scow Bay, near by, via the regular steamship lines. A few rendezvous at Ketchikan and Juneau. This fleet was composed of 60 vessels, valued at \$782,230, employed 1,800 men, and used trawls valued at \$70,850. As a result of its operations in Alaska the fleet (with the exception of the steamers) caught and shipped 3,531,644 dressed pounds (the round weight of this catch or the weight of the fishes taken from the water was approximately 4,414,555 pounds), valued at \$158,260. The steamers carry their own catches to the Sound ports and these have not been included in the above amount. During the summer months most of this fleet fishes on the Flattery Banks off the State of Washington, or else off the British Columbia coast.

THE HERRING FISHERY.

ABUNDANCE OF FISH.

At times herring are quite abundant along the coasts of southeast, central, and western Alaska. At Captains Harbor, on Unalaska Island, they appear twice each year, in July and September. Residents of Port Heiden, in Bering Sea, report that large schools visit that bay in the spring and fall, and there is said to be a large annual run at Atka Island. Herring are quite abundant in Port Clarence also, and some fishermen located at Grantley Harbor, near the head of

this bay, have been salting on a small scale during the past three or four years and selling the fish at Nome and the various settlements in that section of Alaska. The schools generally visit Cook Inlet, in central Alaska, from July to October, and these fish are the largest and finest found in Alaskan waters. In southeast Alaska herring are found in varying abundance in almost every bay, strait, and sound.

According to the best information obtainable, the herring in southeast Alaska begin to spawn during April or May and continue in some localities as late as July 1. Immediately after spawning the fish school in great abundance out in deep water, especially in Frederick Sound and the southern end of Stephens Passage, and then reenter the bays for the purpose of feeding. During July and August they are filled with red feed (certain species of small crustaceans) which makes them very difficult to cure. In September and October apparently they change their food, for the red feed is not then noticeable in their stomachs, and at this time they are in their prime. The runs are usually composed of mixed sizes, although in early summer there are said to be numerous bays where all the herring will be of small size. In western Alaska, according to Nelson, the herring spawn in the neighborhood of St. Michael in June.

At this time these fish form a continuous line along the beach, passing from south to north in unbroken succession, spawning on the seaweeds and rocks from above low-tide mark to a fathom below it. They enter all the inner bays and swarm about every reef and rocky point. The water boils with them along shore as they struggle about in a dense mass among the short seaweed in spawning, and they can be easily caught in one's hands. The females move slowly among the weeds, and press in the midst of them, depositing their eggs, which adhere to whatever they come in contact with, by means of a gummy secretion with which they are coated. Thrusting my hand under water for a half minute was sufficient for it to be covered with eggs.^a

In southeast Alaska during the spawning season, the natives place spruce boughs in the water, and after the eggs have adhered, remove the boughs and dry the eggs in the sun, using them later as food. In this way many thousands of eggs are destroyed each season. This practice should be prohibited by law.

USES FOR FOOD AND BAIT.

Unfortunately, but little commercial use is made of herring as a food fish in central, western, and arctic Alaska. In 1907 a herring saltery was established on Simeonof Island, one of the Shumagin group, in central Alaska. Owing to the low prices realized for the prepared product, and the high cost of transportation, the plant was closed down in 1908 and 1909, but it was reopened this year. A small quantity is marketed fresh, but the great bulk of the catch is made by the Indians, who consume the fish, either fresh or after being dried.

^a Report upon Natural History Collections made in Alaska between the years 1877 and 1881, by Edward W. Nelson, p. 320-21 (1887).

In southeast Alaska the fishery has attained to considerable prominence. Here herring are sold fresh and salted for food; but the principal use is as bait in the halibut and king salmon fisheries and as fertilizer and oil. In baiting, fresh herring are used whenever possible; but when the fisherman has to hold them for a few days the herring are usually dumped round into a barrel with enough salt to preserve them until needed. There is also a demand from the States for the larger herring for smoking purposes, and each season a few dressed and rolled in salt are packed in halibut boxes holding about 500 pounds, and shipped.

Several inquiries were received this year from Seattle and San Francisco brokers and commission men in regard to supplying salted herring for the China trade, and it is to be hoped that some business in this line will eventuate.

Each season there are many complaints from the halibut fishermen as to the scarcity of herring and the heavy loss sustained through the boats being tied up for days at a time owing to the lack of bait. The question of a constant and abundant supply of bait is, in fact, the most serious problem confronting the halibut fishermen. During the summer months halibut fishing is carried on in a desultory manner; but about the middle of September the fleet from Puget Sound arrives, and this, joined with the local fleets, soon causes a tremendous demand for herring, which is the only bait used in the fishery to any extent. The matter is still further complicated by the erratic behavior of the herring itself, which may appear in countless numbers in a certain bay one year, while the next year there may not be one.

The most feasible method for overcoming this handicap would be by the establishment of small freezers at Wrangell, Scow Bay or Petersburg; Juneau, and Hoonah, where herring could be received from the fishermen during the summer and early fall, when most abundant, and frozen and stored away until needed in the late fall and winter. The New England Fish Co., at its Ketchikan plant, freezes a large quantity of herring each year, which it supplies to its own steamers and to the smaller vessels which deliver their catches of halibut at its plant.

THE FERTILIZER QUESTION.

The use of herring in the manufacture of fertilizer and oil as conflicting with its use by man directly as a food and bait fish, and indirectly through the dependence of the valuable king salmon fishery upon it as food material, gives rise to a somewhat puzzling question of right and administrative policy. The present fisheries law does

not prohibit such use of food fishes, and there is now one plant—that of the Alaska Oil & Guano Co., at Killisnoo, in southeast Alaska—engaged in the industry. This year this plant caught 59,000 barrels of herring, with an aggregate weight, roughly, of 11,800,000 pounds. Of these all but 130 barrels, which were pickled for use as bait, were converted into fertilizer and oil.

It is easy to conceive of commercial uses to which fishes are put which take precedence over other uses with respect to public advantage. Thus the manufacture of fertilizer and oil from fishes is a lower use, inferior to the business of preparing food products from fishes, or even to their use as bait for food fishes. Thus the menhaden ranks lower than the herring. Such a view in part grows out of the fact that these fertilizer and oil products, quite legitimate in themselves, do not depend entirely on fishes for their raw material. Furthermore even fish fertilizer and fish oil do not depend upon the herring, for various nonedible fishes, as the menhaden, are available. The general view of a higher use denoted by the appropriation of fishes for human food has widely obtained and is evidenced by various legislation prohibiting the lower use where it has conflicted with the higher. The dependence of a highly prized food fish and a correspondingly valuable fishery upon another fish as food for the former, as in the case of the king salmon upon the herring, may be classed with the higher uses. This in fact is one of the most important aspects of the value of the herring fishery, if not its chief use. An important food of the king salmon is herring, and as the catching of king salmon by trolling now forms one of the most important and profitable of the fisheries of southeast Alaska, no condition that adversely affects it in a material degree should exist unless by the justification of a paramount right and importance.

In the absence of a material higher use the manufacture of the lower products is to be commended, in so far as it causes no depletion, as making a legitimate use of fishes which would otherwise go to waste. Certainly were there no other demand for the herring, such a use should be encouraged. The king salmon of course makes a continual demand upon it, and the king salmon fishery is a permanent one. Even the satisfaction of this demand might perhaps leave a margin of the natural increase of herring for other uses.

Other things being equal it is of course the operation of the law of supply and demand which will determine what use shall be made of commercial fishes, the product being prepared for sale in the highest market. Under such circumstances the matter of use might be left to competition which would exploit the fishery for its most profitable end. Perhaps no such legitimate use could be regarded as indefensible, though lower from some standpoints, but without

discussing this question it may suffice to point out that equality of conditions in practice soon ceases to exist, as is the case with the present herring fishery in Alaska. An established industry with plants and special machinery might continue a less profitable use on account of its possession of facilities and the loss involved in change or abandonment, and make thereby serious inroads upon a supply which would otherwise actually be taken for food uses. It would then seem the part of justice to prohibit the lower use after such time or under such conditions as would secure the interdicted industry from serious loss.

The practice evidently has been, with the approval of public sentiment concerned, to make legislative choice as between material conflicting uses on the general grounds of higher and lower uses, as already discussed. In the concrete instance of the Alaska herring fishery, although some demand an immediate ban on its manufacture into fertilizer and oil, it is not clear that a material conflict of interests exists. As a matter of fact, owing to distance from market, high freights, and the necessity for competing with the British Columbia and Puget Sound packers, the Alaskan herring has not made its way to any great extent as a food fish. As bait for the halibut fishery it is in great demand, but when most needed the herring run is usually small, and the salted herring, while used, is inferior as bait. Both the food and bait uses combined consumed only about 20 per cent of the take in 1910, a season of abundance of herring. The rest was manufactured into fertilizer and oil. Certainly an exigent demand for herring for other purposes could have been met to a larger extent from the large run of the current season.

It is for the future rather than the present that it is desirable to take action looking toward the end of the use of herring as the raw material for fertilizer and oil. It is safe to assume that all the uses of the herring are destined to increase, and therefore at some future time a conflict of uses is probably inevitable. There is but one establishment engaged in the fertilizer and oil industry in Alaska. To prevent extensions of the business and provide for its termination without injury to existing interests it is only necessary to prohibit it by legislation effective at a future date, allowing ample time for the present concern to wind up its affairs. The Bureau has already through the Department recommended to Congress an early tentative date, in part for the sake of eliciting the facts on which to base a reasonable interim. Evidence has been taken on both sides of the question and a common ground reached for a settlement of the question which is believed to be just for all concerned. It is maintained and conceded that the continuance of the herring fertilizer and oil industry is likely to become inconsistent with public policy

respecting the fisheries. The Department on the other hand is inclined to allow a liberal term before any prohibition upon the industry shall become effective, and upon the fixing of this term the question may be said to pend. A few years' delay in the inauguration of this change, intended to hold indefinitely, is a matter of little moment to the fisheries, but of imminent importance to the industry.

STATISTICS.

The following tables show the condition of the herring industry in 1910:

PERSONS ENGAGED IN THE ALASKA HERRING FISHERIES IN 1910.

Occupation and race.	Southeast Alaska.	Central Alaska.	Total.
Fishermen:			
Vessel fisheries—			
Whites.....	59	59
Indians.....	4	4
Japanese.....	4	4
Total.....	67	67
Shore fisheries—			
Whites.....	30	9	39
Indians.....	5	5
Total.....	35	9	44
Shoresmen:			
Whites.....	35	2	37
Indians.....	31	2	33
Japanese.....	6	6
Total.....	72	4	76
Grand total.....	174	13	187

INVESTMENT IN THE ALASKA HERRING FISHERIES IN 1910.

Items.	Southeast Alaska.		Central Alaska.		Total.	
	No.	Value.	No.	Value.	No.	Value.
Fishing vessels:						
Steamers and launches.....	5	\$32,300	5	\$32,300
Tonnage.....	182	182
Outfit.....	12,000	12,000
Launches, under 5 tons.....	6	10,000	1	\$1,200	7	11,200
Boats, sail and row.....	42	2,470	4	400	46	2,870
Scows.....	4	2,100	1	300	5	2,400
Apparatus:						
Vessel fisheries—						
Purse seines.....	10	3,995	10	3,995
Shore fisheries—						
Haul seines.....	1	75	3	400	4	475
Purse seines.....	9	1,495	9	1,495
Gill nets.....	1	500	1	500
Cash capital.....	80,000	2,000	82,000
Shore and accessory property.....	50,800	5,000	55,800
Total.....	195,735	9,300	205,035

* Includes outfit.

PRODUCTS OF THE ALASKA HERRING FISHERIES IN 1910.

Products.	Southeast Alaska.		Central Alaska.		Total.	
	Quantity.	Value.	Quantity.	Value.	Quantity.	Value.
Herring, fresh, for food.....pounds..	10,000	\$300	10,000	\$300
Herring, fresh, for bait.....do.....	574,359	\$5,203	574,359	5,203
Herring, frozen, for bait.....do.....	522,500	5,225	522,500	5,225
Herring, pickled, for food...barrels..	979	9,056	216	1,728	1,195	10,784
Herring, pickled, for bait.....do.....	1,906	3,199	1,906	3,199
Herring, salted, for food....pounds..	45,600	954	45,600	954
Herring eggs, dried, for food....do.....	1,000	100	1,000	100
Herring fertilizer.....do.....	2,617,090	40,000	2,617,090	40,000
Herring oil.....do.....gallons..	277,000	55,000	277,000	50,000
Total.....do.....	113,737	2,028	115,765

FERTILIZER AND OILS.

The only plant operated this year for the preparation of fertilizer and oil from fish was that of the Alaska Oil & Guano Co. at Killisnoo, in southeast Alaska. During the fishing season the company's vessels caught 59,000 barrels of herring, as compared with 52,000 barrels of herring and 3,846 barrels of salmon in 1909.

The Revilla Reduction Works have constructed a plant for the treatment of dogfish and mud shark livers at Ketchikan, in southeast Alaska. While the plant is primarily for the extraction of oil from the livers, it is also hoped by the owners to be able to dry-salt the flesh for shipment as food to China and Japan, and to dry the skins for sale. Unfortunately the flesh so far treated has turned yellow and brown, and until this fault can be corrected it will be of little value. The plant was completed so late in the season that practically nothing was done this year.

THE CRAB FISHERY.

As stated in previous reports, crabs are exceedingly abundant in nearly every section of Alaska, but it is only in southeast Alaska that they are put to any considerable commercial use, many being consumed locally, while large numbers are shipped to the Puget Sound markets, and a few to points in the Northwest Territory, Canada.

The principal shipping places are Petersburg and Wrangell, and the fishermen from here crab on the flats in Dry Straits, opposite Ideal Cove, and at Scow Bay, in Wrangell Narrows. They use a rectangular pot of wooden framework, about 40 inches long, 18 inches high, and 30 inches wide, with 3½-inch stretch mesh net covering. The tunnels, of which there is one at each end, are 7 inches in width and 5 inches in height. These pots cost about \$3 each.

The pots are set on trawls, about 25 or 30 to a trawl. Each is attached to a gangion about 5 fathoms long, thus permitting the raising and emptying of the pot without bringing to the surface the trawl itself. The trawls are marked by buoys and held by anchors.

On some of the trawls baited hooks are placed between the gangions for the purpose of catching bait for the pots. All sorts of fish, clams, etc., are used as bait.

When fishing the pots the fishermen throw back into the water all crabs under 6 inches in width, measured the broad way of the back, all females, and the soft-shell ones, the latter because there is usually very little meat in them.

At first the crabs shipped out of the district were packed alive in seaweed, but so many died on the way or arrived in bad condition that now all are boiled before being shipped. The shippers classify them as follows: Large, 7 inches and over; medium, $6\frac{1}{2}$ to 7 inches; and small, 6 to $6\frac{1}{2}$ inches. The prepared crabs are packed in boxes holding between 12 and 14 dozen each, and are set on their bottoms in three tiers with layers of ice at the bottom, between each tier, and at the top. The freight to Seattle is \$7.50 per measured ton, which would include 35 dozens of crabs.

There is ample room for a large development of this industry, both in canning and marketing fresh, and it is probable this will take place as soon as knowledge of the abundant supplies to be had in Alaska becomes more general.

THE WHALE FISHERY.

The only shore whaling station in the United States where all the parts of a whale are utilized is at Tyee, at the lower end of Admiralty Island, in southeast Alaska, and this plant was operated more vigorously than ever this year. In addition to the steamer *Tyee, Junior*, and the gasoline schooner *Lizzie S. Sorrenson*, which composed the fleet in 1909, the steamer *Fearless* (85 net tons) was fitted out this year. In order to permit the fleet to operate more freely in the open ocean, where most of the whales are now killed, the bark *Diamond Head*, loaded with supplies of coal, gasoline, provisions, etc., was anchored in a convenient bay, to which the fleet could resort when in need and thus save the long trip to the station except when necessary to tow the catch there.

The *Lizzie S. Sorrenson* early in the season met a most unusual fate. As she was cruising around in the ocean about 8 miles southwest of Cape Addington the evening of May 10 a whale was sighted. She was cautiously worked to within gunshot and a harpoon driven into the animal. The weapon failed to reach a vital spot, and the whale made off at a terrific rate, but finding its progress checked it suddenly turned and charged directly at the vessel. Unavailing efforts were made by the crew to work the ship out of the way of the infuriated creature, and the whale, striking her a terrific blow in the stern, knocked out a portion of the bottom. Efforts made to plug the hole were without success, and as the pumps did not suffice, the crew took

to their boats and the vessel soon sank. Two days later the shipwrecked crew was picked up by the whaler *Fearless*.

The station fleet secured 146 whales, of which 6 were sperm whales and one a right whale. As the sperm and right whales produce more valuable by-products than the ordinary whales secured here, the financial return this year was better than in previous seasons. Since the fleet began fishing in the open ocean, moreover, a greater number of sulphur-bottom whales, which are the largest, have been secured, thus adding materially to the output of the station with but slight addition to the cost of operating in the interior waters. It is probable that the plant will be removed to a spot nearer the present scene of operations in order to eliminate the time and expense now necessary in order to get the killed whales from the grounds to the station.

There are a number of shore whaling stations along the Arctic shores of Alaska, at Cape Smythe, Point Hope, and Point Barrow. These stations are quite different affairs from the shore whaling station at Tyee, in southeast Alaska, being virtually trading stations which, in addition to their regular mercantile business, furnish the capital to outfit Eskimos who wish to hunt whales in the ocean close to shore. When a whale is killed the whalebone is removed and sold to the trader, while the natives eat or preserve as food as much of the blubber and flesh as they feel will be required to support them through the long winter. At Cape Smythe there are about 19 boats whaling, at Point Hope about 22, and at Point Barrow about 36 boats. The crews average about 8 men to a boat and the darting gun is quite generally used. The season lasts about 2 months, and comprises a part of April, all of May, and a part of June. The bone shipped out from these stations appears in the statistical tables.

Owing to the glut in the whalebone market, but few of the Arctic fleet operated this year. The fleet comprised the following: Steamer *Herman* (229 net tons), steamer *Karluk* (247 net tons), brigantine *Jeanette* (217 net tons), schooner *Rosie H.* (69 net tons) which went north in 1908, gasoline schooner *Confianza* (84 net tons), and the schooner *Lettitia* (233 net tons). The gasoline schooner *Olga* (43 net tons) sailed north in 1908 and was wrecked in the Arctic late in 1909, the news not coming out until this year. While whales were plentiful they were excessively shy and hard to approach. The fleet secured 27 whales, the *Karluk* alone taking 21, which however, represents two seasons' work on the part of the *Karluk*, she having spent the winter of 1909-10 in the North.

FURS.

Except in the case of fur seals and sea otters, no effort has heretofore been made to conserve the supply of fur-bearing animals of the district, but "An act to protect the seal fisheries of Alaska, and for

other purposes," approved April 21, 1910, consigns these resources to the charge of the Department of Commerce and Labor.

In accordance with section 4 of this law a set of regulations have been promulgated by the Secretary of Commerce and Labor, as given in full in the appendix to this report (p. 71).

The following table shows the number and value of furs of all kinds shipped from Alaska in 1910:

SHIPMENT OF FURS FROM ALASKA IN 1910.

Products.	Southeast Alaska.		Central Alaska.		Western Alaska.		Total.	
	No.	Value.	No.	Value.	No.	Value.	No.	Value.
Bear, black.....	478	\$4,935	326	\$3,065	532	\$3,821	1,336	\$11,841
Bear, black, stuffed.....	1	20					1	20
Bear cubs, black, alive.....			4	125	2	10	6	135
Bear, black, skulls.....					1	10	1	10
Bear, blue.....			2	50			2	50
Bear, brown.....	4	75	27	1,285	2	200	33	1,560
Bear, brown, skulls.....			4	20	1	15	5	35
Bear, glacier.....	3	105	1	20			4	125
Bear, grizzly.....	3	30			3	115	6	145
Bear, polar.....	3	150			53	2,648	56	2,798
Bear castors.....						65		65
Bear galls.....			8	2			8	2
Beaver.....	368	1,922	608	2,763	1,026	5,883	2,002	10,568
Beaver castors.....				59		160		219
Coyote.....					11	6	11	6
Ermine.....	694	447	1,221	997	1,682	1,477	3,597	2,921
Fox, black.....	1	450			1	250	2	700
Fox, blue.....	2	60	492	14,730	660	5,636	1,154	20,426
Fox, blue, live.....			5	175			5	175
Fox, cross.....	2	20	156	1,007	199	1,822	357	2,849
Fox, grey.....			1	100			1	100
Fox, red.....	38	370	3,714	30,084	5,618	38,688	9,370	69,142
Fox, silver.....			50	8,650	3	390	53	9,040
Fox, silver grey.....			56	3,680	57	4,019	113	7,699
Fox, white.....			13	120	1,989	20,443	2,002	20,563
Hares, arctic.....	4	4					4	4
Lynx.....	182	3,541	85	1,856	782	18,685	1,040	24,082
Marten.....	493	4,294	462	3,738	4,702	41,319	5,567	49,351
Mink.....	4,290	22,081	2,534	10,138	16,974	76,369	23,738	108,588
Muskrat.....	12,738	5,086	4,479	917	206,676	69,245	223,893	75,248
Otter, land.....	493	5,213	447	4,493	921	8,843	1,861	18,549
Otter, sea.....	3	600	24	5,900	4	720	31	7,170
Otter pups, sea.....			1	5	2	32	3	37
Rabbit.....					4	4	4	4
Seal, fur.....	138	4,207			14,246	468,042	14,384	472,249
Seal, unborn pup fur.....					121	12	121	12
Squirrel.....	20	5	180	39	9	2	209	46
Weasel.....	36	24	62	31	11	15	109	70
Wolf.....	57	281	5	40	16	86	78	407
Wolverine.....	28	175	75	397	7	42	110	614
Total.....		54,095		94,506		760,024		917,625

^a This table does not take into account the shipments of furs by mail nor of those carried out among the personal effects of passengers.

^b Of these 660 skins were from seized Japanese schooners and were sold by the United States marshal for \$23,100.

^c These were also from the above seized Japanese schooners and were sold by the United States marshal.

AQUATIC FURS.

BEAVER.

This is the most valuable fur-bearing aquatic animal found in the interior waters of Alaska, and has been hunted with such vigor that its ultimate extinction seems to be now but a question of a few years. The range of this animal covers all of the mainland of Alaska, except-

ing only the belt of barren-coast country bordering the Arctic Ocean from Point Hope north and east to the Canadian line. It is also found on a few of the islands in southeast Alaska, and generally in the lakes and streams of the interior, avoiding the large rivers, owing to the great change in level likely to occur at different seasons. During the last three years a considerable proportion of the supply has come from the Kuskokwim and Yukon Valleys. The natives catch beavers in steel traps set at a frequented spot or shoot them from a concealed place near the beaver house or dam.

Castoreum, an oily odorous compound secreted by the preputial glands of the animal, also the dried preputial follicles and their contents, are sometimes prepared and find a sale in China, where they occupy a place in the pharmacopœia.

In 1905, 1,935 skins; in 1906, 1,536; 1907, 1,159; 1908, 1,280; 1909, 2,323, and in 1910, 2,002 skins were secured.

MUSKRAT.

This animal is found on the mainland, except along the extreme northern coast line, wherever bogs and ponds or running water occur; it is also found upon Nunivak and St. Michaels Islands. The Kuskokwim and Yukon Valleys, especially the former, furnish the vast majority of the output. The natives also use a large number each year for clothing and in barter with other native tribes. The value of muskrat has been steadily increasing during the last three years and as a result the animal has been hunted more vigorously each season. In 1905, 12,599 skins, valued at \$1,192; in 1906, 3,611 skins, valued at \$302; in 1907, 6,481 skins, valued at \$498; in 1908, 31,712 skins, valued at \$6,257; in 1909, 121,568 skins, valued at \$34,074, while in 1910, 223,893 skins, valued at \$75,248, were secured and shipped from the district. This takes no account of the local trade in skins between the different tribes.

LAND OTTER.

This species is widely distributed in Alaska, being found on nearly every part of the mainland. It also occurs on many of the islands. A steel trap is generally used in capturing the animals. The supply of land otter skins is fairly constant from year to year.

SEA OTTER.

But two vessels, the schooner *Everett Hays*, owned by Mr. Samuel Applegate, of Unalaska, and the schooner *Elvira* (formerly the Japanese sealing schooner *Kinsei Maru*), owned by Mr. Fred Schroeder of Dutch Harbor, fitted out for sea-otter hunting in 1910. The hunting is generally carried on between Chirikof and Tugidak Islands (the

latter one of the Trinity Islands) in central Alaska, and the season is from about May 15 to September 1, depending largely upon the state of the weather. This year the weather was very rough and as a result there were only about four days of actual hunting throughout the whole season. The *Everett Hays* secured 4 skins, while the *Elvira* took 12, a total of 16.

A few natives living at Kayak this year hunted for sea otter off Cape St. Elias and on June 7 shot two and on June 15 one. These skins were sold at the near-by town of Katalla.

Mr. Nils Christensen, of Cold Bay, on the Alaska Peninsula, hunts sea otters in winter along the reefs offshore, but secured nothing last winter. The same was true of Mr. Charles Rosenberg, who patrols a stretch of some 30 miles of beach on the Bering Sea side of Unimak Island on the lookout for dead sea otter which may be washed ashore.

This summer a native killed a sea otter near the Naknek River in Bristol Bay, where they are very rarely to be found. One was also killed in the neighborhood of Unga Island in central Alaska.

The Canadian sealing fleet again devoted a considerable part of its energies to the hunting of sea otter off Chirikof Island. The schooner *Thos. F. Bayard* secured two, while the *Pescawha* secured seven.

Several vessels from the Japanese sealing fleet also engaged in sea otter hunting, but with what success we are unable to state, owing to their secretiveness in such matters.

FUR SEAL.

The only place on the coast of Alaska which maintains a fur-seal fishery is Sitka. In April and May the herd passes Baranof Island, on which Sitka is located, on its way to the Pribilof Islands in Bering Sea, to breed. About the middle of April the native hunters, who are the only persons permitted to engage in the work, with their families, leave for the hunting grounds and establish their camps on Tava, Wrangell, and Biorca Islands, small islands a few miles from Sitka.

This year 10 boat parties had their headquarters on Biorca Island, four on Wrangell Island, and 18 on Tava Island. Each boat party is composed of from 3 to 5 men, and these use sailboats costing about \$130 each. Repeating shotguns, costing from \$25 to \$35 each, are the only weapons used. The hunting is done in the open ocean, and the boats from the various camps cover an area of from 35 to 50 miles directly out from shore and about the same distance up and down the coast. Good weather is essential, and in 1910 the natives were unfortunate, bad weather being frequent, with the result that the catch was very small.

This year 135 skins were taken and sold at a price aggregating \$4,117 (price paid the hunters and not the London price). In numbers this is a big decrease from last year, when the natives secured 396 skins. Prices received for the skins averaged much higher than in 1909, when \$18.60 was received per skin, as compared with \$30.50 this year.

The Biorka Island parties secured 50 skins, the Wrangell Island parties 13, and the Tava Island parties 72. The largest number secured by any one boat was 8.

In outfitting these boats the hunter, who is head man, furnishes the boat and gun, while the rowers furnish the ammunition and food. The gross proceeds arising from the sale of the skins taken are divided equally among the crew, with the exception of the hunter, who gets \$3 or \$4 more than the others.

The hunting parties return to Sitka the latter part of May. A committee of two is then appointed to supervise the sale of the skins, which usually takes place on a date between June 1 and 5, when the buyers from the States have reached Sitka. On sale day the skins are all brought to one house, where they are sorted into three sizes—"small," "medium," and "large"—care being taken to keep each boat's catch separate from the others. The "small" skins are those of the pups born during the previous two years. The "medium" skins are said to have the best fur, but the buyers prefer the "large" ones on account of their size. The buyers are not allowed to pick out the choice skins and bid on these alone, but must take them as they run, the subdivision in the beginning being made merely in order that the buyers may see what they are bidding on.

These skins are usually much sought after by the dealers, because, being taken by the natives, and a certificate from the collector of customs to this effect being attached to the catch, they can, under the law, be sent abroad to be cleaned and dyed and brought back to be sold in our markets. The possession of such a certificate is considered to add about \$10 to the value of the skin.

The Japanese schooners were again troublesome. During bad weather, when the natives could not go out with their small boats, the schooners came in close, and then when the good weather came they would work out just ahead of the native boats and pick up most of the seals.

The Japanese sealing schooner *Kaise Maru*, which was seized on May 3, 1909, by the deputy marshal at Sitka, is still at that place. The crew were charged with killing seals within the 3-mile limit, and also landing on certain islands near by. They were tried at Juneau in September of the same year and acquitted, but the owners failed to resume possession of their vessel after their release.

In 1909 revenue cutters seized the Japanese sealing schooners *Saikai Maru* and *Kinsei Maru*, and charged them with sealing within the 3-mile limit of the Pribilof Islands. The captured vessels were taken to Unalaska and later the officers and men were carried to Valdez, where all were tried and convicted at the November term of court. Condemnation proceedings against the vessels were instituted, and on April 18 of this year the deputy marshal at Unalaska sold the vessels with their stores and equipment, the *Kinsei Maru* bringing \$4,600 and the *Saikai Maru* \$321.50. When seized the schooners had 660 seal skins, and these sold for \$21,780. The vessels were purchased by Mr. Fred Shroeder, of Dutch Harbor, who renamed the *Kinsei Maru* the *Elvira*, and outfitted and sent her out this year on a sea-otter cruise. The skins sold have been included in the statistical tables of this report.

This year the Treasury Department adopted the policy of permitting sealing vessels to take on merely enough water to carry them to the nearest United States port, or if homeward bound, to take them home. Heretofore the vessels have taken aboard water whenever and wherever they pleased, thus being enabled to extend their cruise indefinitely. Several sealing vessels which visited ports in southeast and central Alaska were affected by this rule. Under the law no resident of the United States is permitted to furnish supplies to a sealer at any time.

The lease of the North American Commercial Co. of the Pribilof Islands expired this year, and the Government, through this Department, took possession of the islands. From St. Paul Island 10,754 skins were shipped, while St. George shipped 2,834, a total of 13,586.

MISCELLANEOUS AQUATIC MAMMALS.

HAIR SEALS.

These animals are to be found all along the coast of Alaska, occurring in places in almost countless numbers. While they form a very insignificant part of the commerce in which the white traders participate, owing to the fact that their fur is worthless, they are of immense value to the natives, for from the flesh and oil is secured a considerable part of the winter food, while the skins are highly prized for covering the kayaks and umiaks, and for boot soles, trousers, mittens, clothing bags, and caps, and when cut into strips make a very strong and durable cord. The coast natives also barter the flesh, oil, and skins with the interior tribes for reindeer hides and furs, thus creating a very important branch of trade of which it is impossible to form an accurate idea, owing to the inaccessibility of most of the tribes and the secrecy they observe when discussing such matters with white men.

WALRUSES.

This animal, which is not found south of the Bering Sea shore of the Aleutian chain, was at one time very numerous north of there, and the hunting of it and the seal formed the principal occupation of the Eskimos during the summer. It goes north as the ice breaks up in the spring and returns again in the fall, stopping but a short time at any spot and keeping close to the ice pack all this time.

While the hunting was carried on solely by the natives the herd suffered no appreciable diminution, but in 1868 the whalers began to turn their attention to walrus catching with serious results to the natives, as set forth in a former report.^a

To many of the Eskimos, especially on the Arctic shore, the walrus is almost a necessity of life, and the devastation wrought amongst the herds by the whalers has been, and is yet, the cause of fearful suffering and death to many of the natives. The flesh is food for man and dogs; the oil is used for food and for lighting and heating the houses; the skin, when tanned and oiled, makes a durable cover for the large skin boats; the intestines make waterproof clothing, window covers, and floats; the tusks are used for lance or spear points or are carved into a great variety of useful and ornamental objects, and the bones are used to make heads for spears and for other purposes.

During the first part of every season there is but little opportunity to capture whales, they being within the limits of the icy barrier. As a result much of the whalers' time during July and August was devoted to capturing walruses. Men would be landed on the shore in June and left to watch for the animals to haul up on the beach at certain points. The walrus must either come ashore or get on the ice, and when a herd is well ashore one or two old bulls are generally left on watch. The best shot among the hunters now creeps up, and by a successful rifle shot or two kills the guard. Owing to their very defective hearing the noise made by the rifle does not awaken them. The gun is then put aside and each hunter, armed with a sharp ax, approaches the sleeping animals and cuts the spines of as many of them as possible before the others become alarmed and stampede for the water and escape.

The natives hunt the walrus in kayaks, with ivory-pointed spears and sealskin line and floats. When the animal is exhausted by its efforts to escape, the hunters draw near and give the death stroke with a lance.

In 1908 Congress passed an act for the protection of game in Alaska, and in this the killing of walrus north of latitude 62° was permitted only from August 1 to December 10, both inclusive, while no one person was permitted to kill more than one.

This year new regulations were promulgated by the Department of Agriculture, and in these the open season for walruses in Bering Sea and Strait north of the Kuskokwim River is from May 1 to July 1, while all killing in Bristol Bay and Bering Sea south of the Kuskokwim River is prohibited until 1912.

As the natives are permitted to kill the walrus for food and clothing at any time when in need of food, the object of the law, which is

^a The Commercial Fisheries of Alaska in 1905. By John N. Cobb, Bureau of Fisheries Document 603, p. 35, 1906.

to prevent the indiscriminate killing by whites, is accomplished, and very few of the animals are now killed except by the few sportsmen who visit the Bering Sea district in summer. This year's reports indicate that walrus are increasing. The inspector of fisheries for Alaska saw a large number on the ice in Bristol Bay in May, while the master of the trading schooner *Helen Johnston* claims to have encountered in Bering Strait, near the Diomed Islands, on July 5 a large herd of swimming walrus which covered several acres of water. Capt. S. F. Cottle, of the steam whaler *Karluk*, reports having seen large pods of walrus this year.

LICENSE TAXES AND HATCHERY REBATES.

Under the provisions of the act for the protection and regulation of the fisheries of Alaska (approved June 26, 1906) the packers in Alaska are compelled to pay license fees or taxes on their season's output, as noted in the table following. The collection of these license fees or taxes is in the hands of the clerk of the court of the judicial district in which the packer is operating. The law literally requires the packer to pay the license fee in advance, but as the fee is based upon the pack he makes and it would be impossible in such an uncertain industry as fishing to estimate in advance exactly the quantity that will be packed, it is the custom to require the operator to apply for a license before beginning operations and then at the end of the season make return of the amount due the district.

The following table shows the quantity of taxable fishery products prepared, the stated license tax on the product, and the total amount of tax due on each. The last item is approximate, being based upon returns on file at this Bureau, some of which are sworn to and some estimated, and therefore perhaps varying somewhat from those sent to the clerk of the court. It is not probable, however, that the amount given will vary much either way from the correct amount as shown by the returns of the clerks:

LICENSE TAXES ON PREPARED FISHERY PRODUCTS.

Items.	Unit of quantity.	Quantity prepared.	License tax per unit of quantity.	Estimated amount of tax due.
Canned salmon.....	Cases.....	2,413,052½	\$0.04	\$100,522.08
Pickled salmon.....	Barrels.....	14,405	.10	1,440.50
Mild-cured salmon.....	Tierces α	3,357	.40	1,342.80
Dry-salted salmon in bulk.....	100 pounds	77,478	.05	37.70
Fish oil.....	Barrels.....	578	.10	57.80
Fertilizer, from fish.....	Tons.....	1,308½	.20	261.70
Fertilizer, from whales.....	do.....	435	.20	87.00
Total.....				103,749.58

α As the net weight of a tierce of fish is 800 pounds, this item is figured on a basis of 4 barrels to the tierce in working out the amount of tax.

The following table shows the name of the owner, location of each private salmon hatchery operated during the year ending June 30, 1910, the number of salmon (red) liberated, and the amount of rebate certificates due each hatchery:

REBATES CREDITED TO PRIVATE SALMON HATCHERIES IN 1910.^a

Owners.	Location.	Red salmon fry liberated.	Rebate due.
Alaska Packers Association	Naha Stream.....	40,725,000	\$16,290
Northwestern Fisheries Co.	Karluk Stream.....	35,875,000	14,350
	Quadra Lake.....	9,850,000	3,940
North Pacific Trading & Packing Co.	Hetta Lake.....	8,000,000	3,200
	Klawak Lake.....	5,300,000	2,120
Total.....		99,750,000	39,900

^a Some of the hatcheries did not complete their distribution of fry before July 1; those remaining will be counted next year.

COMPLAINTS AND PROSECUTIONS.

On Sunday, May 22, in Taku Inlet, southeast Alaska, the assistant agent discovered Henry Hoeke, S. Nelson, John Hanula, Tom Carvo, Abraham Lahti, Oscar Lustig, Van Oleson, and Ole Oleson fishing during the weekly closed season. All were brought before the United States commissioner at Juneau for preliminary hearing and bound over to the next grand jury. On October 24 all were indicted by the grand jury held at Ketchikan, and on the 29th of the same month all but Van and Ole Oleson pleaded guilty. S. Nelson and Henry Hoeke were fined \$50 each, while the others were fined \$25 each. The Oleson brothers elected to be tried in Juneau, and on December 10 they appeared in court there and pleaded guilty; sentence was deferred for six months.

In October a man named Mitchell was reported by other fishermen as violating the weekly closed season in the Taku River. He was indicted by the December grand jury, but was acquitted upon his trial the same month.

A visit to Tamgas Stream, a tributary of Tamgas Harbor, on the south end of Annette Island, in southeast Alaska, on July 25, developed the fact that a trap was being fished in the creek in violation of the law. Tamgas Stream is a short and narrow stream draining a lake, and a run of red and other salmon annually ascends the stream. About 300 yards from its mouth are a succession of cascades and falls. In the narrowest part of the cascades a rack had been constructed of poles driven into the bottom and covered with wire netting in such way as almost wholly to prevent salmon from passing up, the portion uncovered being too steep for any but the strongest to surmount. Just below and running parallel to the rack, and at right angles to the shore, was constructed a flume, with a flaring

mouth at the outer end; at the shore end a sharp turn of the flume led into a square box with slat bottom and covered over with boughs. The fish in ascending the stream would be stopped by the rack and in swimming around at the outer end many of them would be carried by the current into and down the flume, eventually landing in the receiving box at the end.

Inquiry among the few Indians camped near the mouth of the stream developed the fact that a native named James, of Metlakahtla, who died last winter, had first constructed the trap several years ago. This spring his two sons, boys under 18 years of age, rebuilt the trap. They were ordered to remove it and did so at once. Owing to the youth of the offenders and other extenuating circumstances, the matter was not presented to the United States attorney for action.

On July 6 Mr. Nels Moen, of Wrangell, complained in regard to the location of the Alaska Packers Association trap in Humpback Bay, Bradfield Canal, and also said his partner in the operation of a rival trap in the same bay, Mr. Oscar Williamson, could prove that the association's trap had been fishing on Sunday, July 3. As soon as possible thereafter a visit was made to Humpback Bay, where an inspection of the trap showed that it was constructed and placed in conformity with the law. As Mr. Williamson was confident of having evidence enough to justify his charge that the trap had been operated during the weekly closed season, the matter was brought before the United States commissioner at Wrangell, Mr. Williamson making the sworn complaint. Mr. H. A. Oleson, the trap foreman, was arrested and brought to Wrangell for preliminary hearing. The evidence, however, clearly showed no intent at violating the law and the defendant was discharged.

On the occasion of a visit to Sarkar Stream, on the west coast of Prince of Wales Island, southeast Alaska, on August 26, Mr. Fred Brockman was discovered fishing a gill net which had been stretched from bank to bank. The net had 13 coho salmon in it at the time. Brockman was arraigned before the United States commissioner at Wrangell on September 3 and by him was bound over to the next grand jury, which began its sessions at Ketchikan on October 24 and indicted the defendant on the same date. On October 24 he appeared in court and pleaded guilty. Owing to the defendant's physical condition the court imposed the small fine of \$25, but gave an impressive warning that the next offender appearing in court charged with this serious offense would be severely dealt with.

In the latter part of July several natives reported to the deputy marshal at Sitka that native fishermen were fishing within the prohibited area around the mouth of Necker Stream, which empties into Necker Bay, on the west coast of Baranof Island, in southeast Alaska. Twenty-five natives were brought in by the deputy marshal and

given a hearing before the United States commissioner at Sitka, who discharged all of the defendants, however, for lack of evidence.

Several complaints were made in regard to alleged illegal fishing by gill netters operating in Karta Bay, Prince of Wales Island, south-east Alaska, but diligent search failed to substantiate any of these, and as the nets were soon withdrawn the complaints, which had come from purse seiners, ceased.

On June 25 the deputy marshal and deputy collector of customs at Cordova visited Eyak River and found Perry and Causa Sabella, fishermen employed by the Northwestern Fisheries Co. at Orca, with a gill net stretched from shore to shore. The net held at the time of the visit some 40 or 50 fish. The men were brought before the United States commissioner at Cordova and fined \$1 and costs, amounting in all to \$50 each.

An evil which at present is slight, but will grow more and more serious as the district becomes more settled and the superabundant water power, which at present largely goes to waste, is harnessed and made to serve the purposes of the manufacturer, prospector, lumberman, etc., is the building of dams in streams which the salmon frequent. By the terms of the law it is—

unlawful to erect or maintain any dam, barricade, fence, trap, fish wheel, or other fixed or stationary obstruction, except for purposes of fish culture, in any of the waters of Alaska at any point where the distance from shore to shore is less than five hundred feet, * * * with the purpose or result of capturing salmon or preventing or impeding their ascent to their spawning grounds, and the Secretary of Commerce and Labor is hereby authorized and directed to have any and all such unlawful obstructions removed or destroyed.

In the past, builders of such obstructions have been very negligent in consulting the salmon agents in regard to the legality of their structures, and as a result considerable expense has been caused to them by their failure to observe the plain provisions of the law. Where some municipal or commercial benefit is to result the agents have been willing to meet the parties more than half way and to supply all needful plans for the placing of fishways in such dams where feasible.

PROPOSED LEGISLATION.

At the hearings held between April 19 and May 25, before the Committee on the Territories of the House of Representatives, on H. R. 22579, Sixty-first Congress, second session, known as the Wickersham bill, in amendment of the Alaska fisheries law of June 26, 1906, representatives of the Bureau furnished statements and testimony bearing on the provisions of this bill in their relation to the fisheries. At the close of these hearings the following letter was transmitted by the Secretary of Commerce and Labor in response to a request for the opinion of the Department with respect to changes or additions

desirable in the law. The proposals for legislation increase the taxes somewhat, and aim to extend and increase the power of the Department over all Alaskan fisheries save the fur seal.

DEPARTMENT OF COMMERCE AND LABOR,
OFFICE OF THE SECRETARY,
Washington, May 25, 1910.

HON. E. L. HAMILTON,
Chairman Committee on the Territories,
House of Representatives, Washington, D. C.

SIR: In reply to your letter of the 20th instant, in which you request the opinion of the Department with respect to changes advisable in the present Alaska fisheries law, after consideration of the act of June 26, 1906, section by section, the following recommendations are submitted:

1. Sections 5, 7, 8, 10, 11, 12, 13, 14, 15, and 16 are satisfactory.

2. Section 1 should be modified in accordance with the schedule already submitted at the hearing of May 3. This schedule is along the lines indicated by Judge Wickersham in H. R. 22579.

3. Section 2 should remain until more adequate facilities are provided for fish-cultural work by the Federal Government. All fish-cultural work in Alaska should eventually be carried on by the Federal Government. This can be brought about by the abolition of the present exemption system, the taking over of such private hatcheries as the owners may desire to turn over to the Government, and the establishment of additional Federal hatcheries.

4. In section 3, line 2, strike out the words "for purposes of fish-culture" and insert in lieu thereof "by direction of the Secretary of Commerce and Labor"; and in lines 4 and 5 strike out the words "where the same is less than five hundred feet in width."

5. In section 4, line 2, strike out the words "for purposes of fish culture" and insert in lieu thereof "by direction of the Secretary of Commerce and Labor."

6. In section 6, lines 6 and 7, strike out the words "five hundred yards of the mouth thereof" and insert in lieu thereof "such distance from the mouth thereof as in his judgment is necessary."

7. The matter covered by section 9 is now fully covered by the pure food and drugs act, food inspection decision No. 105, and this section may therefore be omitted.

8. The following additional sections are now recommended:

"SECTION —. That for the purposes of this act the Secretary of Commerce and Labor is authorized to determine and indicate by suitable markers the mouth of any creek, stream, or river in Alaska which salmon enter for spawning purposes.

"SEC. —. That the Secretary of Commerce and Labor is authorized and directed to establish such regulations, not inconsistent with existing law, as may in his judgment be necessary for the proper protection and conservation of shellfish and other aquatic animals not otherwise mentioned in this act.

"SEC. —. That it shall be unlawful to erect, maintain, or operate in Alaska any new establishment for canning or otherwise preserving for commercial use any salmon or other fish or fishery product, or to increase the capacity of any such existing establishment, or to reopen and operate any such establishment which has remained closed for the period of three years immediately preceding the passage of this act, without first obtaining the approval in writing of the Secretary of Commerce and Labor.

"Provided, however, That in the case of salmon-packing establishments approval shall be withheld only when in his judgment the fishing operations and investigations in the region adjacent to the proposed location indicate that the number of salmon taken is larger than the reproductive increase of salmon from adjacent spawning grounds: *And provided further*, That in case approval is withheld the applicant interested shall upon demand be given a hearing, of which he shall be notified at least thirty days previously.

"Sec. —. That it shall be unlawful, after January first, nineteen hundred and eleven, to utilize any part of any food fish save the offal and refuse thereof in the manufacture of fertilizer or fish oil.

"Sec. —. That the provisions of sections thirteen and sixteen of chapter four hundred and twenty-five of an act entitled 'An act making appropriations for the construction, repair, and preservation of certain public works on rivers and harbors, and for other purposes,' approved March third, eighteen hundred and ninety-nine, shall be applied to the protection of the fisheries of Alaska, and the Secretary of Commerce and Labor and his agents for the protection of the salmon fisheries of Alaska, and any officer or employee of the Department of Commerce and Labor designated by him, shall be charged with the enforcement of said section thirteen and shall have the same power and authority in all respects to swear out process and arrest as the several officials named in section seventeen of chapter four hundred and twenty-five of the above act."

Respectfully,


CHARLES NAGEL, *Secretary.*

There is pending before Congress a measure for reorganization and expansion of the Alaska work of the Bureau of Fisheries, under the one head of Alaska Fisheries Service. This division will include, if the law is enacted, the salmon-inspection service and the fur-seal service, together with supervision of all other fisheries and fur resources of Alaska.

RECOMMENDATIONS.

1. That vessels be provided for the inspection service as recommended in the report for 1909, and that immediate provision be made for the two smaller launches requested, as the most urgent needs of the service, for use during the 1911 season.

2. That in addition to the recommendations contained in departmental letter of May 25, 1910, printed on page 64 of this report, for the amendment of the present fisheries act of June 26, 1906, the weekly close season for salmon, as expressed in section 5 of the existing law, be extended over all Alaskan waters except Bering Sea and its arms; and that in sections 3 and 4 the word "salmon" be substituted for red salmon.



APPENDIX—FISHERY LAWS AND REGULATIONS.

The following laws relating to the fisheries and fur-bearing animals of Alaska, and the regulations established thereunder, which are now in force in the District, are published herewith for the guidance of those interested:

AN ACT for the protection and regulation of the fisheries of Alaska.

Be it enacted by the Senate and House of Representatives of the United States of America in Congress assembled, That every person, company, or corporation carrying on the business of canning, curing, or preserving fish or manufacturing fish products within the territory known as Alaska, ceded to the United States by Russia by the treaty of March thirtieth, eighteen hundred and sixty-seven, or in any of the waters of Alaska over which the United States has jurisdiction, shall, in lieu of all other license fees and taxes therefor and thereon, pay license taxes on their said business and output as follows: Canned salmon, four cents per case; pickled salmon, ten cents per barrel; salt salmon in bulk, five cents per one hundred pounds; fish oil, ten cents per barrel; fertilizer, twenty cents per ton. The payment and collection of such license taxes shall be under and in accordance with the provisions of the Act of March third, eighteen hundred and ninety-nine, entitled "An Act to define and punish crimes in the district of Alaska, and to provide a code of criminal procedure for the district," and amendments thereto.

SEC. 2. That the catch and pack of salmon made in Alaska by the owners of private salmon hatcheries operated in Alaska shall be exempt from all license fees and taxation of every nature at the rate of ten cases of canned salmon to every one thousand red or king salmon fry liberated, upon the following conditions:

That the Secretary of Commerce and Labor may from time to time, and on the application of the hatchery owner shall, within a reasonable time thereafter, cause such private hatcheries to be inspected for the purpose of determining the character of their operations, efficiency, and productiveness, and if he approve the same shall cause notice of such approval to be filed in the office of the clerk or deputy clerk of the United States district court of the division of the district of Alaska wherein any such hatchery is located, and shall also notify the owners of such hatchery of the action taken by him. The owner, agent, officer, or superintendent of any hatchery the effectiveness and productiveness of which has been approved as above provided shall, between the thirtieth day of June and the thirty-first day of December of each year, make proof of the number of salmon fry liberated during the twelve months immediately preceding the thirtieth day of June, by a written statement under oath. Such proof shall be filed in the office of the clerk or deputy clerk of the United States district court of the division of the district of Alaska wherein such hatchery is located, and when so filed shall entitle the respective hatchery owners to the exemption as herein provided; and a false oath as to the number of salmon fry liberated shall be deemed perjury and subject the offender to all the pains and penalties thereof. Duplicates of such statements shall also be filed with the Secretary of Commerce and Labor. It shall be the duty of such clerk or deputy clerk in whose office the approval and

proof heretofore provided for are filed to forthwith issue to the hatchery owner, causing such proofs to be filed, certificates which shall not be transferable and of such denominations as said owner may request (no certificate to cover fewer than one thousand fry), covering in the aggregate the number of fry so proved to have been liberated; and such certificates may be used at any time by the person, company, corporation, or association to whom issued for the payment pro tanto of any license fees or taxes upon or against or on account of any catch or pack of salmon made by them in Alaska; and it shall be the duty of all public officials charged with the duty of collecting or receiving such license fees or taxes to accept such certificates in lieu of money in payment of all license fees or taxes upon or against the pack of canned salmon at the ratio of one thousand fry for each ten cases of salmon. No hatchery owner shall obtain the rebates from the output of any hatchery to which he might otherwise be entitled under this Act unless the efficiency of said hatchery has first been approved by the Secretary of Commerce and Labor in the manner herein provided for.

SEC. 3. That it shall be unlawful to erect or maintain any dam, barricade, fence, trap, fish wheel, or other fixed or stationary obstruction, except for purposes of fish culture, in any of the waters of Alaska at any point where the distance from shore to shore is less than five hundred feet, or within five hundred yards of the mouth of any red-salmon stream where the same is less than five hundred feet in width, with the purpose or result of capturing salmon or preventing or impeding their ascent to their spawning grounds, and the Secretary of Commerce and Labor is hereby authorized and directed to have any and all such unlawful obstructions removed or destroyed.

SEC. 4. That it shall be unlawful to lay or set any drift net, seine, set net, pound net, trap, or any other fishing appliance for any purpose except for purposes of fish culture, across or above the tide waters of any creek, stream, river, estuary, or lagoon, for a distance greater than one-third the width of such creek, stream, river, estuary, or lagoon, or within one hundred yards outside of the mouth of any red-salmon stream where the same is less than five hundred feet in width. It shall be unlawful to lay or set any seine or net of any kind within one hundred yards of any other seine, net, or other fishing appliance which is being or which has been laid or set in any of the waters of Alaska, or to drive or construct any trap or any other fixed fishing appliance within six hundred yards laterally or within one hundred yards endwise of any other trap or fixed fishing appliance.

SEC. 5. That it shall be unlawful to fish for, take, or kill any salmon of any species in any manner or by any means except by rod, spear, or gaff, in any of the waters of Alaska over which the United States has jurisdiction, except Cook Inlet, the Delta of Copper River, Bering Sea, and the waters tributary thereto, from six o'clock post-meridian of Saturday of each week until six o'clock antemeridian of the Monday following, or to fish for, or catch, or kill in any manner or by any appliances except by rod, spear, or gaff, any salmon in any stream of less than one hundred yards in width in Alaska between the hours of six o'clock in the evening and six o'clock in the morning of the following day of each and every day of the week. Throughout the weekly close season herein prescribed the gate, mouth, or tunnel of all stationary and floating traps shall be closed, and twenty-five feet of the webbing or net of the "heart" of such traps on each side next to the "pot" shall be lifted or lowered in such manner as to permit the free passage of salmon and other fishes.

SEC. 6. That the Secretary of Commerce and Labor may, in his discretion, set aside any streams or lakes as preserves for spawning grounds, in which fishing may be limited or entirely prohibited; and when, in his judgment, the results of fishing operations in any stream, or off the mouth thereof, indicate that the number of salmon taken is larger than the natural production of salmon in such stream, he is authorized to establish close seasons or to limit or prohibit fishing entirely for one year or more within such stream or within five hundred yards of the mouth thereof, so as to permit salmon to increase: *Provided, however,* That such power shall be exercised only after

all persons interested shall be given a hearing, of which due notice must be given by publication; and where the interested parties are known to the Department they shall be personally notified by a notice mailed not less than thirty days previous to such hearing. No order made under this section shall be effective before the next calendar year after same is made: *And provided further*, That such limitations and prohibitions shall not apply to those engaged in catching salmon who keep such streams fully stocked with salmon by artificial propagation.

SEC. 7. That it shall be unlawful to can or salt for food any salmon more than forty-eight hours after it has been killed.

SEC. 8. That it shall be unlawful for any person, company, or corporation wantonly to waste or destroy salmon or other food fishes taken or caught in any of the waters of Alaska.

SEC. 9. That it shall be unlawful for any person, company, or corporation canning, salting, or curing fish of any species in Alaska to use any label, brand, or trade-mark which shall tend to misrepresent the contents of any package of fish offered for sale, *Provided*, That the use of the terms "red," "medium red," "pink," "chum," and so forth, as applied to the various species of Pacific salmon under present trade usages shall not be deemed in conflict with the provisions of this Act when used to designate salmon of those known species.

SEC. 10. That every person, company, and corporation engaged in catching, curing, or in any manner utilizing fishery products, or in operating fish hatcheries in Alaska, shall make detailed annual reports thereof to the Secretary of Commerce and Labor, on blanks furnished by him, covering all such facts as may be required with respect thereto for the information of the Department. Such reports shall be sworn to by the superintendent, manager, or other person having knowledge of the facts, a separate blank form being used for each establishment in cases where more than one cannery, saltery, or other establishment is conducted by a person, company, or corporation, and the same shall be forwarded to the Department at the close of the fishing season and not later than December fifteenth of each year.

SEC. 11. That the catching or killing, except with rod, spear, or gaff, of any fish of any kind or species whatsoever in any of the waters of Alaska over which the United States has jurisdiction, shall be subject to the provisions of this Act, and the Secretary of Commerce and Labor is hereby authorized to make and establish such rules and regulations not inconsistent with law as may be necessary to carry into effect the provisions of this Act.

SEC. 12. That to enforce the provisions of this Act and such regulations as he may establish in pursuance thereof, the Secretary of Commerce and Labor is authorized and directed to depute, in addition to the agent and assistant agent of salmon fisheries now provided by law, from the officers and employees of the Department of Commerce and Labor, a force adequate to the performance of all work required for the proper investigation, inspection, and regulation of the Alaskan fisheries and hatcheries, and he shall annually submit to Congress estimates to cover the cost of the establishment and maintenance of fish hatcheries in Alaska, the salaries and actual traveling expenses of such officials, and for such other expenditures as may be necessary to carry out the provisions of this Act.

SEC. 13. That any person, company, corporation, or association violating any provision of this Act or any regulation established in pursuance thereof shall, upon conviction thereof, be punished by a fine not exceeding one thousand dollars or imprisonment at hard labor for a term of not more than ninety days, or by both such fine and imprisonment, at the discretion of the court; and in case of the violation of any of the provisions of section four of this Act and conviction thereof a further fine of not more than two hundred and fifty dollars per diem may, at the discretion of the court, be imposed for each day such obstruction is maintained. And every vessel or other apparatus or equipment used or employed in violation of any provision of this Act, or

of any regulation made thereunder, may be seized by order of the Secretary of Commerce and Labor, and shall be held subject to the payment of such fine or fines as may be imposed.

SEC. 14. That the violation of any provision of this Act may be prosecuted in any district court of Alaska or any district court of the United States in the States of California, Oregon, or Washington. And it shall be the duty of the Secretary of Commerce and Labor to enforce the provisions of this Act and the rules and regulations made thereunder. And it shall be the duty of the district attorney to whom any violation is reported by any agent or representative of the Department of Commerce and Labor to institute proceedings necessary to carry out the provisions of this Act.

SEC. 15. That all Acts or parts of Acts inconsistent with the provisions of this Act are, so far as inconsistent, hereby repealed.

SEC. 16. That this Act shall take effect and be in force from and after its passage.

Approved, June 26, 1906.

AN ACT To prohibit aliens from fishing in the waters of Alaska.

Be it enacted by the Senate and House of Representatives of the United States of America in Congress assembled, That it shall be unlawful for any person not a citizen of the United States, or who has declared his intention to become a citizen of the United States, and is not a bona fide resident therein, or for any company, corporation, or association not organized or authorized to transact business under the laws of the United States or under the laws of any State, Territory, or district thereof, or for any person not a native of Alaska, to catch or kill, or attempt to catch or kill, except with rod, spear, or gaff, any fish of any kind or species whatsoever in any of the waters of Alaska under the jurisdiction of the United States: *Provided, however,* That nothing contained in this Act shall prevent those lawfully taking fish in the said waters from selling the same, fresh or cured, in Alaska or in Alaskan waters, to any alien person, company, or vessel then being lawfully in said waters: *And provided further,* That nothing contained in this Act shall prevent any person, firm, corporation, or association lawfully entitled to fish in the waters of Alaska from employing as laborers any aliens who can now be lawfully employed under the existing laws of the United States, either at stated wages or by piecework, or both, in connection with Alaskan fisheries, or with the canning, salting or otherwise preserving of fish.

SEC. 2. That every person, company, corporation, or association found guilty of a violation of any provision of this Act or of any regulation made thereunder shall, for each offense, be fined not less than one hundred dollars nor more than five hundred dollars, which fine shall be a lien against any vessel or other property of the offending party or which was used in the commission of such unlawful act. Every vessel used or employed in violation of any provision of this Act or of any regulation made thereunder shall be liable to a fine of not less than one hundred dollars nor more than five hundred dollars, and may be seized and proceeded against by way of libel in any court having jurisdiction of the offense.

SEC. 3. That the violation of any provision of this Act or of any regulation made thereunder may be prosecuted in any United States district court of Alaska, California, Oregon, or Washington.

SEC. 4. That the collector of customs of the district of Alaska is hereby authorized to search and seize every foreign vessel and arrest every person violating any provision of this Act or any regulation made thereunder, and the Secretary of Commerce and Labor shall have power to authorize officers of the Navy and of the Revenue-Cutter Service and agents of the Department of Commerce and Labor to likewise make such searches, seizures, and arrests. If any foreign vessel shall be found within the waters to which this Act applies, having on board fresh or cured fish and apparatus or imple-

ments suitable for killing or taking fish, it shall be presumed that the vessel and apparatus were used in violation of this Act until it is otherwise sufficiently proved. And every vessel, its tackle, apparatus, or implements so seized shall be given into the custody of the United States marshal of either of the districts mentioned in section three of this Act, and shall be held by him subject to the proceedings provided for in section two of this Act. The facts in connection with such seizure shall be at once reported to the United States district attorney for the district to which the vessel so seized shall be taken, whose duty it shall be to institute the proper proceedings.

SEC. 5. That the Secretary of Commerce and Labor shall have power to make rules and regulations not inconsistent with law to carry into effect the provisions of this Act. And it shall be the duty of the Secretary of Commerce and Labor to enforce the provisions of this Act and the rules and regulations made thereunder, and for that purpose he may employ, through the Secretary of the Treasury and the Secretary of the Navy, the vessels of the United States Revenue-Cutter Service and of the Navy: *Provided, however,* That nothing contained in this Act shall be construed as affecting any existing treaty or convention between the United States and any foreign power.

Approved, June 14, 1906.

FISHERY REGULATIONS.

1. During the inspection of the salmon fisheries by the agents and representatives of this Department, they shall have at all times free and unobstructed access to all canneries, salteries, and other fishing establishments, and to all hatcheries.

2. All persons, companies, or corporations owning, operating, or using any trap-net, pound-net, or fish-wheel for taking salmon or other fishes shall cause to be placed in a conspicuous place on said trap-net, pound-net, or fish-wheel the name of the person, company, or corporation owning, operating, or using same, together with a distinctive number, letter, or name which shall identify each particular trap-net, pound-net, or fish-wheel, said lettering and numbering to consist of black figures and letters, not less than six inches in length, painted on white ground.

3. All persons, companies, or corporations engaged in canning salmon shall forward to the Bureau of Fisheries, Department of Commerce and Labor, Washington, D. C., three copies of each and every different can label which it is designed to place upon the canned product.

CHARLES NAGEL, *Secretary.*

REGULATIONS FOR THE PROTECTION OF FUR-BEARING ANIMALS IN ALASKA.

[Alaska Fisheries Service—Circular No. 1.a]

MARCH 8, 1911.

To whom it may concern:

Section 4 of "An act to protect the seal fisheries of Alaska, and for other purposes," approved April 21, 1910, provides that—

No person shall kill any otter, mink, marten, sable, or fur seal, or other fur-bearing animal, within the limits of Alaska Territory or in the waters thereof; and every person guilty thereof shall, for each offense, be fined not less than two hundred nor more than one thousand dollars or imprisoned not more than six months, or both; and all vessels, their tackle, apparel, furniture, and cargo found engaged in violation of this section shall be forfeited; but the Secretary of Commerce and Labor shall have power to authorize the killing of any such mink, marten, sable, fur seal, or other

a The sundry civil bill passed by Congress March 4, 1911, provided for a reorganization and expansion of the Alaska service of the Bureau of Fisheries, as referred to on page 65 of this report. This circular, while by its date not strictly within the scope of the report for 1910, is printed here for its usefulness in connection with the other laws now administered by the Alaska fisheries service.

fur-bearing animal under such regulations as he may prescribe; and it shall be the duty of the Secretary of Commerce and Labor to prevent the killing of any fur seal except as authorized by law and to provide for the execution of the provisions of this section until it is otherwise provided by law.

Fur-bearing animals enumerated below may, subject to the provisions of regulation No. 12, be hunted and killed in the Territory of Alaska, except during the seasons specified with respect to each of the several animals mentioned.

1. *Sea otter*.—The hunting or killing of sea otter is prohibited until November 1, 1920.

2. *Beaver*.—The hunting or killing of beaver is prohibited prior to November 1, 1915.

3. *Land otter and mink*.—The hunting or killing of land otter or mink is prohibited throughout the season from April 1 to November 15, both days inclusive, of each year.

4. *Marten, fisher, sable, ermine, and weasel*.—The hunting or killing of marten, fisher, sable, ermine, or weasel is prohibited throughout the season from April 1 to November 15, both days inclusive, of each year.

5. *Muskrat*.—The hunting or killing of muskrat is prohibited throughout the season from May 16 to November 30, both days inclusive, of each year.

6. *Black bear*.—The hunting or killing of black bear is prohibited throughout the season from June 1 to August 31, both days inclusive, of each year.

7. *Fox, lynx, and wildcat*.—The hunting or killing of fox, lynx, or wildcat is prohibited throughout the season from March 1 to November 15, both days inclusive, of each year.

8. *Wolf, wolverine, spermophile, and rabbit or hare*.—The killing of wolves, wolverines, spermophiles (ground squirrels), and rabbits or hares is not prohibited.

9. The killing of any fur-bearing animal by means of strychnine or any other poison is prohibited at all times.

10. Permits or licenses may be issued by the Secretary of Commerce and Labor for the taking of fur-bearing animals for scientific purposes, for shipment to zoological parks, or for breeding purposes.

11. The penalties and forfeitures imposed by the act will be strictly enforced against all persons who take, capture, or kill, or attempt to take, capture, or kill, any fur-bearing animal in the Territory of Alaska during the prohibited seasons herein established, or who barter or have in their possession the skin or pelt of any fur-bearing animal taken in the close or prohibited season.

12. Shipments of furs, which may be made at any time, will be reported to the Bureau of Fisheries, Department of Commerce and Labor, on appropriate blanks which will be supplied for that purpose.

These regulations supersede all others previously in force.

Approved:

CHARLES NAGEL, *Secretary*.

**SPECIAL INVESTIGATION OF THE ALASKA
FUR-SEAL ROOKERIES, 1910**

By **HAROLD HEATH**

Professor of Invertebrate Zoology, Stanford University

Bureau of Fisheries Document No. 748

CONTENTS.

	Page.
Brief sketch of natural history of the seal.....	3
The rookeries.....	4
Rookery development.....	4
Harem counts.....	5
Active bulls.....	6
Idle bulls.....	7
Young bulls.....	7
Counts of idle and young bulls.....	8
Breeding cows.....	8
Counts of pups.....	9
Estimates of cows and pups.....	10
Yearlings and 2-year-olds.....	11
The reserve.....	11
Estimate of all classes.....	12
The quota.....	13
Conservation and some involved problems.....	13
The question of an equilibrium of the herd.....	15
The patrol and pelagic sealing.....	16
The pelagic catch.....	18
Cows in drives.....	19
Causes of death.....	19
Ages of seals.....	20
Appendix—Extract from field notes.....	21

SPECIAL INVESTIGATION OF THE ALASKA FUR-SEAL ROOKERIES, 1910.

By HAROLD HEATH,
Professor of Invertebrate Zoology, Stanford University.

Under the act of Congress of April 21, 1910, involving various changes in the administration of the Pribilof Islands and the seal fisheries and providing for the appointment of additional officers and employees, it was decided that a naturalist should be designated to study and report upon the condition of the seal herd. Pending the selection of a permanent occupant of this position, to take effect July 1 under the law, the writer was sent to the islands as a special investigator to perform the naturalist's duties for the season which was already beginning. Observations were made on St. Paul Island, beginning June 29, the date of arrival on the island, and continuing until July 15, then for a week on St. George Island, and again on St. Paul until August 29. A report of these observations is contained in the following pages.

I am indebted to the Government agents on the islands and to the officers of the revenue fleet for valuable data and many courtesies in connection with my work.

BRIEF SKETCH OF NATURAL HISTORY OF THE SEAL.

As popularly applied the term "seal" includes a fairly large group of aquatic mammals, such as the sea lion and the fur and hair seals, all of which bear a superficial resemblance to each other. Strictly speaking, the last named are the only ones deserving of the name. Unlike the hair seal, the fur seal, or sea bear, is able to progress readily on land, is able to hold its head erect, and its fore limbs, finlike in form, are used in swimming. Concerning its life at sea, we know that the seals of the Pribilof Islands spend their winter months along the western coast of North America, the adult females extending their migrations as far as southern California. Early in May the adult males or bulls begin to appear on the rookeries, where each is subsequently joined by 30 females on the average, the height of the

breeding season occurring about the 15th of July. Shortly after her arrival each cow gives birth to a pup, and after a sojourn of perhaps two weeks, during which time she is served by the bull, she puts out to sea on the first of several journeys in search of food.

During this time the young males or bachelors are arriving, and are usually found in groups on the outskirts of the rookeries. It is from these young males that the land catch of skins is made.

Early in August disorganization of the harems commences. The greater number of cows have been served, the active bulls accordingly relax their vigilance, the idle bulls and those less mature wander about without serious molestation, the pups congregate at various points on shore or in the shallows, where they learn to swim, and as autumn advances the roving instinct becomes more and more apparent in all classes, finally leading to the abandonment of the shore early in November.

THE ROOKERIES.

In position and extent the rookeries have undergone but few changes since last year. The number of active bulls and the attendant harems have decreased slightly, but whether this indicates an actual decrease in the number of cows is doubtful, since the count of pups, as noted in a succeeding section, was made on one rookery only and the data derived therefrom are not perfectly trustworthy. The decline in the number of harems on St. Paul is most apparent on Gorbatch, the Zapadnis, and Tolstoi, where there are 55 less than in 1909. On the other hand, there are 47 more on the Reef, Kitovi, Polovina, and Vostochni. On St. George the very slight increase noted on Staraya Artel and Zapadni is almost exactly counterbalanced by a decline on North and East rookeries.

This year the fleet operated chiefly about Northeast Point and to the south and east between St. Paul and St. George, but the results of their operations do not appear to be so distinctly reflected in a corresponding decline of adjacent rookeries as in 1909. Such a definite effect requires that the seals put out to sea along radii centering in either one of the islands, but on numerous occasions I have watched cows, and especially bachelors, leaving the rookeries, and their course is far from being either direct or uniform. The problem, however, is of interest chiefly to the naturalist as matters rest at present, and is without any very practical bearing on the conservation of the herd.

ROOKERY DEVELOPMENT.

At present there appear to be no very definite problems associated with the development of the rookery, but following the custom observed for several years past counts of harems and cows were made whenever and wherever it was possible. Kitovi especially received

attention and as far as practicable was examined at intervals of about three days with the following results:

DEVELOPMENT OF KITOVI ROOKERY, SEASON OF 1910, AS SHOWN BY COUNTS OF SEALS ON DIFFERENT DATES.

Date.	Harems.	Cows.	Reserve bulls.	Half bulls.
June 30.....	16	27	37	6
July 2.....	32	107	24	14
July 6.....	43	326	19	7
July 9.....	47	500	14	10
July 13.....	62	929	9	10

The past winter was unusually severe and long continued, delaying the breaking up of the drift ice, the melting of the snow, and the appearance of flowering plants for upward of three weeks. It is interesting to note, however, that this delay did not affect the summer resident birds, which put in an appearance at the customary time, though compelled in numerous instances to deposit their eggs on the snow. Nor did it hinder the migration of the seals, though several cows likewise took up positions on snow drifts, where they and the pups appeared to be unmindful of their unusual habitat.

HAREM COUNTS.

In accordance with the custom pursued in past years, the counts of harems were made as nearly as possible at the "height of the season," occurring July 12-16. Owing to stress of weather Sivutch, or Sea Lion Rock, rookery was not counted, but was estimated as containing 61 harems, the number found last season.

SUMMARY OF HAREM COUNTS, 1910, AND COMPARISON WITH 1897 AND 1909.^a

Rookery.	1897	1909 ^a	1910	Rookery.	1897	1909	1910
St. Paul Island:				St. George Island:			
Gorbatch.....	308	120	112	Little East.....	46	4	4
Ardiguen.....	33	11	11	East.....	128	65	59
Reef.....	454	184	206	Zapadni.....	133	43	47
Sea Lion Rock.....	102	61	61 ^b	Staraya Artel.....	57	42	48
Kitovi.....	179	55	62	North.....	196	106	103
Lukanin.....	139	39	41	Total.....	560	260	261
Polovina.....	143	42	50	Grand total.....	4,418	1,387	1,381
Polovina Cliffs.....	61	23	20				
Little Polovina.....	40	19	12				
Morjovi.....	233	45	47				
Vostochni.....	910	184	204				
Zapadni.....	458	147	118				
Little Zapadni.....	176	62	54				
Zapadni Reef.....	114	11	7				
Tolstoi.....	295	87	77				
Tolstoi Cliffs.....	98	25	29				
Lagoon.....	115	12	9				
Total.....	3,858	1,127	1,120				

^a Figures for 1909 are those of Mr. George A. Clark.

^b Estimated.

Assuming that Sea Lion Rock is occupied by the same number of harems as in 1909 or neglecting it for both seasons, there are 7 fewer harems on St. Paul this year than last.

Comparing the number of harems on St. George during the years 1909 and 1910 there is 1 more, and when both islands are considered 6 fewer. As there is one bull to a harem, this is another way only of stating that there are 6 fewer bulls this year than last; and obviously such an estimate affords no indication whatever of the actual number of breeding cows.

ACTIVE BULLS.

The number of active bulls, each in control of a harem, is as just noted, somewhat smaller this year than last (as 1,387 to 1,381); but it is the universal verdict that as a class they have lost none of those characteristics that make them successful masters. As usual there was considerable skirmishing among them as the harems were forming, but the wounds inflicted were comparatively insignificant and no deaths were recorded. Early in the season one dead female was found on Gorbach whose wounds may have been caused by a bull, and later six cows were seen on various rookeries that had been severely though not fatally slashed.

In a few cases young bulls or "quitters" were found with harems on various rookeries, but usually they held sway on the outskirts of the community and joined the females in the mad rush to the sea whenever they were approached. It was the rare exception that they held a position in the more crowded portions of the rookery, where they would be called upon to defend their cows against the attempted inroads on the part of more seasoned harem masters.

By some authorities it has been urged that this infusion of young male life into the general herd is beneficial, but in all probability its value is overestimated. It is not disclaimed that some animals are born with more vigorous constitutions than others, and that in all probability their offspring will be more hardy in consequence. And furthermore, it is a truism that in the struggle for existence it is a gain that the feeble are weeded out; but this is an entirely different question from the one relating to the effects of age. In the case of the female a long existence may lessen the production of milk or alter its composition, and consequently inhibit the proper nourishment of the offspring, but with the male no such argument may be brought forward. In the case of the race horse, which has been studied as much as any other mammal, attempts have been made to show that it is desirable to breed young males, and again, with essentially the same data, such a position has been attacked. To-day we know far less about the seal, but it is a safe proposition to argue in favor of perpetuating, as far as possible, those fully developed males that are able to protect their harems.

IDLE BULLS.

These animals are victims of circumstances. Owing usually to an unfavorable location, they have failed to secure harems, though they are as physically able to control them as any of their class. Furthermore, the term "idle" is a misnomer, for no one who has watched them on the rookeries would ever accuse them of being sluggish. On the other hand, they are aggressive in the extreme, and especially during the height of the season engage in frequent quarrels with the harem masters, from whom they usually pilfer a small number of cows before the close of the season.

It can not well be doubted that an excess of this class of animals is more or less of a menace to the normal, or at all events what appears to be the most successful, type of seal existence. Claims have been made to the effect that for untold ages the seal has fought the battle of life successfully and that in the present time the hand of man is not required to control his destinies. The first part of this statement is undeniably correct, but the last is open to criticism, for it assumes that the seal is to-day leading a normal existence. Unfortunately this is not true, for we know that the number of breeding cows is becoming alarmingly reduced. In the open Pacific the number of captured males and females may be approximately equal, but the Bering Sea catch, as past records show, contains from 70 to 80 per cent of females. Since, on the average, there is 1 male to every 30 cows in the harem, there must inevitably result an excess of males, an unnatural state of affairs, and the belief that in cutting down this excess we are conferring a benefit appears to rest on a firm foundation.

This season the number of idle bulls was 221, not so great a number but that they were kept at bay until the disintegration of the harems had commenced, when they usually became the possessors of a small number of cows.

YOUNG BULLS.

Young bulls, otherwise known on the islands as "quitters", are usually 6 or 7 years old, and at the approach of man retire. They frequently haul out with the bachelors or form a shifting fringe about the group of breeding seals. In rare cases they controlled harems, usually on the margins of the rookeries, and in a few cases were seen in the act of copulation.

An accurate count of these animals was not made, unfortunately, since a considerable number had hauled out with the bachelors and could not be numbered without interfering with subsequent drives. At the height of the season the number on the rookeries was 184, and at various times 386 in all were included in the drives. Some were doubtless driven more than once, but it seems certain that the

actual number was at least 200, giving a total of 384. As the average life of the male is 13 years, of which 5 are spent as harem master, the decrease annually of the present active list is 276. It is apparent therefore that killing in the past has not been too close, and that there is a sufficient reserve at the present time.

COUNTS OF IDLE AND YOUNG BULLS.

The following count of idle and young bulls was made at the time the census of harems was taken. It was not possible without causing undue disturbance to enumerate members of the latter class that had hauled out with the bachelors on four important rookeries—Northeast Point, Gorbatch, the Reef, and Tolstoi.

COUNTS OF IDLE AND YOUNG BULLS ON ST. PAUL AND ST. GEORGE ISLANDS, 1910.

Rookery.	Idle bulls.	Young bulls.	Rookery.	Idle bulls.	Young bulls.
St. Paul Island:			St. Paul Island—Continued.		
Gorbatch.....	12	17	Tolstoi Cliffs.....	5	1
Ardiguen.....	1		Lagoon.....	3	
Reef.....	28	17	Total.....	144	136
Kitovi.....	9	9			
Lukanin.....	5	11	St. George Island:		
Polovina.....	5	12	East.....	20	
Polovina Cliffs.....	5	5	Zapadni.....	19	17
Little Polovina.....	2	7	Staraya Artel.....	17	21
Morjovi.....	1	1	North.....	21	10
Vostochni.....	29	26	Total.....	77	48
Zapadni.....	22	13			
Little Zapadni.....	10	8	Grand total.....	221	184
Zapadni Reef.....		3			
Tolstoi.....	7	6			

BREEDING COWS.

While there is a steady increase in the number of cows hauling out on any rookery for a month after the middle of June, a seagoing stream soon makes its appearance, consisting of cows en route to the feeding grounds after their pups are born. Hence at the "height of the season," about the middle of July, the number of cows on the beach is no true indication of the total number, nor does it always bear a constant ratio to the whole. Under certain circumstances, possibly due to climatic conditions, nearly the full complement may be present at the height of the season, and again in other years not over 30 per cent of the community may be on the rookery. It thus becomes apparent that such counts, of varying character from season to season, must be used with extreme caution, if at all, in estimating the entire number of females on any rookery or the annual decline or increase. As has been pointed out by others, we may arrive at an approximate estimate only by a count of the pups, and under that heading an attempt has been made to show that even here we must use the results with the greatest care in making a census of the herd.

During the height of the season counts were made on the following rookeries:

COUNTS OF COWS ON SOME ST. PAUL ROOKERIES DURING HEIGHT OF SEASON,
1897, 1909, AND 1910.

Rookery.	1897	1909 ^a	1910
Lagoon.....	1,319	281	229
Tolstoi Cliffs.....	1,286	698	646
Zapadni Reef.....	1,049	137	78
Ardiguen.....	470	207	218
Kitovi.....	2,436	892	837
Kitovi Amphitheater.....	654	127	92
Lukanin.....			820
Polovina Cliffs.....			426
Little Polovina.....			421
	7,214	2,342	3,767

^a Counts of Mr. George A. Clark.

COUNTS OF PUPS.

Owing to the fact that all the cows are never present on the rookeries at a given time, it is obvious that the only approach to an accurate census of the breeding females is to be made by counting all the pups on all the rookeries. Such a procedure is not only arduous but wasteful, since the cows in early August, when the counting is usually done, are readily driven into the sea and a portion must inevitably fall a prey to the pelagic sealer. Accordingly it was the custom, for several years prior to 1906, to count the pups on a number of rookeries, and with such data estimate the entire herd. In more recent times the number of such pup counts has become gradually lessened until this year Kitovi was the only rookery examined, with the following result: Total number of pups, 1,966; dead, 62.

The implication that Kitovi is a typical average rookery must rest upon the assumption that it stands between those in which the decline is great and those in which it is at a minimum. As a matter of fact, an examination of the counts of Kitovi during the past four years shows that in reality it has been remarkably constant so far as the cows are concerned. Commencing with 1907 the number of pups each year is 1,959, 1,960, 1,979, and this year there are 1,966.

Last year there were 55 active bulls on Kitovi and 1,979 pups; this year there are 62 bulls and 1,966 pups. The average harem last year was 36; this year, 31.7; a difference due almost wholly to the increased number of active bulls. And, furthermore, this slight difference is of far-reaching importance when we come to consider the application of these data to the estimate of the entire herd. With 1,381 harems, each numbering 36 cows, the estimate would be 49,716; if each comprised 31.7 cows there are then 43,777 in the breeding herd, a difference of 5,939, or 11,878 when the pups are included in the count, due solely to the presence of 7 active, extra bulls.

Then, again, on the other rookeries an increase or decrease in the number of active males produces a corresponding rise or fall in the estimated number of cows. For example, on Vostochni there may be 6,500 cows and 200 active bulls. If 20 idle bulls, before the height of the season, secure 1 cow apiece, they enter the active list, and there are then 220 harems. As the average harem is 31.7, this increase affects the estimate to the extent of a gain of 634 cows, though in reality the number of cows has remained constant. At present this gain or loss in the active bull list outside of Kitovi is of relative unimportance, but it is conceivable that under certain circumstances it may assume a more prominent rôle.

I have in mind the fact that in treating this phase of the problem we are, after all, dealing in generalities, but the results may become so general that they have little actual value. In my opinion it is highly desirable that a pup count on all of the rookeries be made during August, or even early in September, in stress of weather, or possibly after the sealing fleet has left Bering Sea; and again a similar survey should be made five years later, when the typical rookery could be determined and questions relating to the increase or decrease of the herd be settled beyond a reasonable doubt.

ESTIMATES OF COWS AND PUPS.

Assuming that the average harem comprises 31.7 cows, the total number in the entire seal herd is computed in the following table:

COMPUTATION OF COWS AND PUPS ON ST. PAUL AND ST. GEORGE ISLANDS, 1897, 1909, AND 1910.

Rookery.	1897	1909 ^a	1910	Rookery.	1897	1909 ^a	1910
St. Paul Island:				St. Paul Island—Contd.			
Gorbatch.....	9,086	4,320	3,551	Tolstoi Cliffs.....	2,891	1,452	888
Ardguen.....	736	355	349	Lagoon.....	2,598	693	285
Reef.....	13,393	6,624	6,530	Total.....	112,023	41,266	35,502
Sea Lion Rock.....	3,009	2,196	^b 1,934	St. George Island:			
Kitovi.....	5,289	1,979	1,966	Little East.....	1,190	144	127
Lukanin.....	4,100	1,404	1,299	East.....	3,776	2,340	1,870
Polovina.....	4,218	1,512	1,585	Zapadni.....	3,923	1,548	1,490
Polovina Cliffs.....	2,200	828	634	Staraya Artel.....	1,681	1,512	1,522
Little Polovina.....	1,180	684	380	North.....	5,782	3,816	3,266
Morjovi.....	6,873	1,620	1,490	Total.....	16,342	9,360	8,275
Zapadni.....	13,511	5,292	3,740	Grand total.....	128,365	50,626	43,777
Vostochni.....	26,845	6,624	6,467				
Little Zapadni.....	5,192	2,232	1,711				
Zapadni Reef.....	3,041	319	222				
Tolstoi.....	8,702	3,132	2,471				

^a Estimates of Mr. George A. Clark.^b Estimated.

In the above census it is to be remembered that the totals apply to cows and pups and that both together number 87,554 in 1910.

YEARLINGS AND 2-YEAR-OLDS.

Of the various computations necessary to arrive at an estimate of the entire seal herd those concerned with the 2-year-olds and yearlings are the least satisfactory. And yet by restricting the quota of skins taken to 3-year-olds we could in a relatively short period arrive at a fairly close approximation, and at the same time settle other vexed questions that are in need of solution. At the present time we are compelled to base our estimates largely on the quota and those males dismissed from the killing grounds.

In the quota this year 10,210 skins weighed less than 7 pounds each, and 2,603 males were dismissed from the drives because they were undersized. Some of the latter were doubtless driven more than once, but even so it is probable that the number was not less than 1,800. Besides these, 337 2-year-olds were branded early in the season. This accounts for 12,347. That there are yet others is evidenced by the fact that fully 700 bachelors of killable size appeared on the hauling grounds of both islands in early August after the killing season, in addition to which there were probably other young animals in considerable numbers, though how many is uncertain. And it is probable, also, that some were at sea, but here again we have no exact information. A conservative estimate of 2-year-old males is therefore 13,000, which is also the number of virgin 2-year-old females that during the late summer arrived at the rookeries.

It appears to be the general belief that in 1909 there were 12,000 yearlings of each sex, and judging from estimates based on pup counts and the quota, the herd appears to have been stationary for the past three or four years. Hence we might suppose that the number of yearlings for this year is approximately the same as last. However, it is possible that the estimates based largely on Kitovi are misleading and that the quota was maintained by closer and closer killing. Future observations alone will settle this question. In order to be on the safe side we may assume that a shrinkage of 10 per cent has taken place and that accordingly the number of yearlings of each sex for the year 1910 is 10,800.

THE RESERVE.

For six years prior to 1910 two thousand 2 and 3 year old males were reserved annually, but as the brand, made by clipping the hair on the head, was not permanent, we have no means of knowing how many of these were subsequently killed. If 1,000 were actually exempted each year and there is an annual mortality of 10 per cent there should be between 500 and 600 this year remaining of the reserve of 1905. And if the decline of the present number of active bulls is approximately 300 there should this year be an increase of

over 200. As a matter of fact there is a slight decline, so that it appears that males exempted one year were killed the next. In reality, if we may judge from the records of past years, there is no necessity of reserving annually a number greater than one-half of the total number of active bulls, but these should be chosen from the class that will be wiggged next year, or branded with a permanent mark.

This year 1,271 males were set aside as a reserve. Very nearly 1,000 4-year-olds and older were dismissed from the drives. Some of these were doubtless driven more than once, but it is assuredly safe to conclude that 600 were actually present. In addition there were others on the water front and in the water to the number of at least 100, and finally there were 605 idle and half bulls. This gives a total of 2,576, a number considerably in excess of the requirements.

ESTIMATE OF ALL CLASSES.

The following is an itemized estimated census of the seals forming the herd in 1910:

ESTIMATED CENSUS OF SEAL HERD IN 1910.

Class.	1910
Active bulls.....	1,381
Breeding cows.....	43,777
Pups.....	43,777
Idle bulls.....	221
Young bulls.....	384
Bachelor reserve.....	1,971
2-year males.....	5,500
2-year females.....	13,000
Yearling males.....	10,800
Yearling females.....	10,800
Quota killed.....	13,584
Total.....	145,195

According to this estimate and Mr. Clark's estimate of 158,488 for 1909, the herd has diminished by 13,293 within the past year. Whether this is a just conclusion must be decided by computations to be made during the next few years. Accuracy is impossible so long as the present methods are employed. During late years it has been assumed that the error is not greater than 12 per cent, and this is probably a fair conclusion. Last year the herd numbered between 150,000 and 160,000; this year it seems to fall between 140,000 and 150,000.

THE QUOTA.

In 1897 it was estimated that the ratio of bachelors to the entire herd was 1:20; this year it is approximately 1:10. The conditions that have brought about this change are matters largely of conjecture, for our knowledge of the seal is too imperfect to warrant a satisfactory explanation. It is reasonably certain that the mortality among pups is less than formerly and, as Mr. Lembkey states in his report of 1909, this would insure a proportionately larger return of yearlings, males and females, and subsequently of breeding cows, both of which are factors tending to the increase of bachelors. Then again the death rate of the young, estimated to be 50 per cent during the first year, may have been excessive and the proportion of bachelors to the the entire herd may have been greater than was estimated in 1897. But even if these problems were solved to our complete satisfaction they do not bear directly on the question of the conservation of the herd. As noted in another paragraph, the essential point to be settled is regarding the reserve. If it is sufficient to supply the requisite number of males, as the active ones disappear, then it appears to be the best policy to kill those remaining. The herd is declining or at best stationary. The pelagic sealer is hovering about the islands and close killing diminishes his catch. That the quota should consist of the skins of 3-year-olds is obviously the most economical plan, but from a purely zoological standpoint this is a matter of detail and relatively unimportant.

This year 10,749 skins were taken on St. Paul and 2,834 on St. George, a total of 13,583, or 785 less than in 1909. The weights of these, together with data relating to the drives and numbers dismissed, are given in the report of the agent in charge.

CONSERVATION AND SOME INVOLVED PROBLEMS.

It has been seen from the foregoing paragraphs that the number of males for breeding purposes is sufficient, and this has been so for many years. On the other hand the number of females has been decreasing steadily, and there is no question but that the pelagic sealer is, and has been, an important factor in producing this decline. Furthermore, another fact is evident, that with the conservation of the females on land and the setting aside annually of a sufficient male reserve no additional care will add one jot or tittle to the number of cows. It is perfectly true that the elements involved in the problem of the male reserve are intricate and some are not clearly understood, but in the last analysis the important question to be answered is this: Is there a sufficient number of males to take the place of those active on the rookeries? and every year the answer has been in the affirmative. On land, killing may be close, and skins below the

maximum value may be taken, but if the females are protected and the male reserve be adequate other questions sink into a position of relative unimportance as the seal problem now presents itself.

The foregoing paragraph is written from a purely biological standpoint, having in mind only the conservation of the herd, but there are other questions of a more practical bearing that should be settled before the sealing business can be conducted on the most economical basis. In the first place it is highly desirable that the number of pups born annually be more accurately determined, reducing the possible error below 10,000, where it stands at present. In 1896 the error was estimated to be about 6 per cent, but last year and this it is probably twice as great. With the herd approaching the vanishing point accuracy is more than ever a desideratum and should be had even at the cost of an unusual amount of labor.

Again, we have no information, within narrow limits, of the number of males or females returning at the close of the first year, or if this be beyond computation, then of the number returning the second or even the third year. This, as the sexes are of approximately equal numbers, will give more nearly than any other practicable method the number of females taking their places on the rookeries. Beyond this time observations should be made to determine the number of reserved 3-year-olds that appear the next year, and finally the percentage that ultimately becomes active on the rookeries. From such observations the reserve of males may ultimately be made with an accurate knowledge of facts, and not with such hazy ideas as we have at present.

It is highly desirable that the quota be taken from the males in prime condition, and I heartily agree with Mr. Lembkey and Mr. G. A. Clark, who argue in their reports of 1909 for the killing of 3-year-olds. I am by no means convinced that even by the branding of every pup, and so destroying the fur to some extent, we can, by this means alone, reduce the value of the skin to such a degree that the pelagic sealer will be forced out of business. It may indeed be a fact, but the brands made in the past were in some cases fatal and are supposedly about all that the young seal is able to survive, and yet not over one-tenth or at most one-eighth of the fur is destroyed. The resulting depreciation of value will probably not amount to more than \$10, and two San Francisco furriers place it as low as \$5. The price of skins is gradually advancing and on the other hand we do not know what returns will pay the schooner owners to keep a ship in the sea. The crew, averaging 35, receives \$5 per man each month (Captain Quinan of the revenue cutter *Tahoma* says \$2.50) and 12½ cents goes to each man for every skin taken by his particular row-boat. Let us suppose each schooner is out six months, and, judging from past records, 8,000 skins will be taken this year, or 320 per

schooner. If the price per skin were only \$15 (\$30 was the price they received last year) \$4,800 would certainly be a paying investment.

On the other hand there is another factor making toward the reduction of the sealing fleet which, together with the partial destruction of the skins through branding, may possibly put the pelagic sealer out of business or, more probably, so limit the number of vessels that an equilibrium of the seal herd may become a fixed feature. This element is competition. With 25 schooners in the sea, rivalry must this year have been very keen, and with a diminishing herd some competitors must sooner or later leave the field. Any depreciation in the value of skins must hasten the desirable result, provided—and here an unknown factor enters—that the price of skins does not advance. But with the decline of the number of skins it is probable that prices will advance, and it appears very questionable whether branding and competition will drive away all of the pelagic fleet for many years to come. It may, however, make it possible for the herd to remain practically stationary until some form of treaty insures more perfect conservation.

The branding process may be made to include the male pups, but as the pelagic sealer secures but few bachelors this would greatly destroy the value of the land catch without giving adequate returns. It is possible that the males dismissed from the drives might be penned up for a month or so, but unfortunately I can not speak with authority regarding this plan, that was once put into execution several years ago. Some advocates claim that it is entirely possible; that after a few days the captives show no signs of restlessness in their unnatural surroundings. Others are equally certain that the experiment was not a success, as several of the larger animals broke through the barriers and some less fortunate became restless in the extreme and finally died of exhaustion. Furthermore, it is reported the bachelors ordinarily put to sea from time to time in search of food, and it is difficult to see how food would be forthcoming even if they desisted from their attempts to escape. The fact that placing animals in captivity would prevent redriving does not appear in itself to be sufficient reason for carrying out the plan. If by these schemes we hope to drive the pelagic sealer from his elected calling then it seems to me they will not succeed, but that they may increase the value of the land catch is possible.

THE QUESTION OF AN EQUILIBRIUM OF THE HERD.

The question of an equilibrium of the herd is one of very high importance. In 1897 the Fur Seal Commission agreed that such a state of affairs would ultimately occur, and in 1909 Mr. G. A. Clark argues in favor of the possibility that there is now an equilibrium.

Unfortunately, in the present year a sufficiently large pup count was not made whereby to settle the question. The estimated decline may be approximately correct or it may be due to the methods of taking the census. If an equilibrium does exist it means that if the number of guards stationed on the islands is sufficient to prevent poaching the entire land catch may amount annually to something in the neighborhood of 10,000 skins and the herd would be in no danger of extinction. If instead of allowing matters to rest as they are the Government orders the branding of female pups, then some of the pelagic sealers may be compelled to abandon their calling, and the herd would probably increase, but there is nothing to prevent the return of the entire sealing fleet when the herd is larger and a profitable catch may be made even though each skin is much reduced in value.

As matters appear there is one way only whereby the pelagic sealer may be driven away entirely, and that is by the further reduction of the seal herd. This is at best a cold-blooded proposition and will probably not meet with general approval, but there seems to be no other way to destroy the activity of the fleet.

The question now stands, Shall the pelagic sealer be driven from the sea and the financial gain from the then highly diminished herd be reduced to a minimum, or is it better policy to place the business more nearly on a paying basis though the pelagic sealer share in the returns? Until pelagic sealing is discontinued by an agreement with the countries concerned the revenue fleet must be kept about the islands, under any circumstances the natives must be cared for, and in various ways a heavy financial outlay must be made annually. Personally I favor the latter plan, reaping as large a harvest as is compatible with the conservation of the herd and at the same time leaving as little as possible to those on the high seas.

THE PATROL AND PELAGIC SEALING.

The revenue fleet maintained throughout the season of 1910 a most thoroughgoing and careful patrol about the islands, where reefs, and shifty currents, and impenetrable fogs are of the most treacherous character. Three cutters, the *Tahoma*, Capt. Quinan, commanding; the *Manning*, Capt. Cardin; and the *Perry*, Capt. Haake, constituted the fleet, with Capt. Foley at Unalaska in command. Prior to July 26 each vessel remained 12 days in the vicinity of St. Paul, and after 5 days returned from coaling at Unalaska. On the date named the *Perry*, during a dense fog, went ashore at Rocky Point on St. Paul and was never floated. The duties of the remaining vessels became correspondingly increased, but so far as known no schooner pushed inside of the 3-mile zone after this accident, and

generally speaking the infractions of the law throughout the season were of minor importance.

Pelagic sealing, on the part of the Japanese, continued with unabated vigor. During this season 25 vessels were reported, 7 more than in 1909, and the reports in Capt. Foley's office in Unalaska show that each schooner carried approximately 25 to 40 men and from 5 to 10 boats. Furthermore, several of these ships cleared from Japan early in the year, and, arriving at various points from California to Sitka, followed the herd to the breeding grounds in Bering Sea. In the vicinity of St. Paul Island, none ventured, so far as known, within the 3-mile zone, but in one or two instances violations were reported by the natives on St. George, where the revenue-cutter patrol is far less vigilant. On June 28 the *Tokai Maru* was seized and fined for violation of the alien fishing law, and on July 25 the *Toro Maru* was seized and fined for violation of the custom laws (section 2773 of the Revised Statutes). On July 18 two row boats were sighted in the vicinity of Zapadni, on St. George, so close to shore that one was seen to contain at least one unskinned seal. And again during foggy weather on July 30 two boats' crews from the schooner *Hoko Maru* landed at Northeast Point and Lukanin, respectively, and the next day 4 sailors from the *Toro Maru* were captured en route to Zapadni. Though pleading stress of weather, all were taken into custody and were subsequently tried in Unalaska.

Generally speaking, the fleet operated to the east and north of St. Paul, presumably in the path of the seals leaving the Reef, Kitovi, Lukanin, the Polovinas, and Northeast Point. On July 10 the steamer *Homer* reported at least a dozen schooners with their attendant boats, which had formed a great circle between St. Paul and St. George and were slaughtering the seals compelled to cross the line of fire at two points. Although the nearest of these vessels was at least 8 miles from the shores of St. Paul, the reports of the shotguns could be heard distinctly on land, and a count I made on that day from 11.20 to 11.50 a. m. showed that 228 shots were fired, an average of 7.6 per minute.

In this connection it may be mentioned that on certain days, owing to meteorological conditions, sounds travel amazing distances. According to Capt. Quinan, shots were heard one day in July seemingly well within the 3-mile zone, but with the lifting of the fog the nearest boat was fully 7 miles distant. Somewhat later in the month a fusilade was distinctly heard on St. Paul, but with the clearing away of the mists not a single boat could be detected even with powerful glasses used from the top of a 70-foot hill. It thus becomes apparent that alleged transgressions, based on this species of evidence alone, are far from being trustworthy.

To an outsider the practice of having Japanese stewards aboard the cutters is not above criticism. They must inevitably come into possession of valuable information that may be of service to Japanese prisoners, for whom they act as interpreters, if I am informed correctly. Furthermore, the Japanese detained for 10 days on St. Paul this year were in constant communication with the natives of the village, and it was no fault of theirs if they did not learn more of the island than is disclosed by the chart. One has a certain amount of sympathy for the pelagic sealer, who receives a mere pittance for his services and is the only sufferer when his boat is captured; but his imprisonment is not a serious hardship, especially if he be allowed to work on the coal pile at \$2 per day and is ultimately sent back to Japan.

These are, after all, matters of comparative unimportance. The arrest, and even the severe punishment, of such offenders do not seriously interfere with the activities of the schooners and their owners. Such devices as branding to partially destroy the value of the skins, and of penning up male seals released from the drives, are not complete preventives, so that until an agreement is consummated the international struggle between watcher and watched must forever go on with all of the attendant aggravating features. It is possible that the herd is not in a state of equilibrium, but is actually diminishing. If this continue the hunter on the high seas must ultimately vanish from the scene of his pernicious activity; but is the Government of the United States compelled to place the seal herd on the altar of sacrifice in order to bring about this desired result?

If this, indeed, be true then we must decide, and that right early, whether this be a lesser evil than the other, hypothetical to a certain degree, of branding the females, which form the greater portion of the pelagic catch, and by the depreciation of their skins, making it necessary for a greater number than at present to be taken with profit by the pelagic sealer. At the same time this would render it possible for an increased number of cows to escape and breed on the rookeries, and so add materially to the bachelor herd and consequently to the land catch.

THE PELAGIC CATCH.

Regarding the pelagic catch of this year, our evidence must rest upon a very slender reed—the reports of the Japanese themselves. According to these, 4,213 skins were taken prior to August 15, of which 2,098 came from Bering Sea. Last year the reported Japanese catch up to August 15 was 4,954 skins. As a matter of fact, it was then probably twice as large, for the entire season's catch, as reported from the London market, was 10,561 skins. This year it is safe to predict that there will be at least 8,000.

COWS IN DRIVES.

During the killing season proper, closing August 1, the discipline maintained by the active bulls on the rookeries was very strict, and accordingly a very insignificant number of cows made their way into the neighborhood of the bachelors and were driven to the sealing grounds. Such as did so, of course, were subsequently released. During a food drive on August 10, when the harems had commenced to disintegrate, several cows appeared in the drive, but I was unable to find a single one among the dead on the killing grounds. Doubtless females may occasionally be clubbed accidentally, but this year I can testify that the greatest care was exercised, and I know of no occurrences of the kind.

FEEDING OF PUPS.

For various reasons, up to the time of my departure from the islands, no attempt was made to raise pups. The pair handled successfully by Boatswain Thurber had shed the first coat and were fully 3 months old; he was unsuccessful with the young, black pups. These last named may possibly be reared if food of the proper character be fed, but at the present time we are ignorant of the composition of seal's milk. In any event one must have not only a large store of patience but an abundance of time, and whatever may be said regarding the first requisite the latter is not forthcoming during the summer, when one is concerned with numerous other matters pertaining to the herd in general and must leave the islands in August.

CAUSES OF DEATH.

Under normal circumstances the life of the seal of either sex is probably from 12 to 13 years. Since the bulls are active for not more than five seasons, one-fifth of the active list dies each year, and as the cows are believed to breed during ten seasons one-tenth of their number disappears annually.

Judging from the reports of former years the season of 1910 was one of comparative quiet. No fatalities due to fighting were noted among the bulls, and only one cow was discovered whose death may be attributed to rough handling on the part of a bull.

On the killing grounds between 20 and 30 bachelors were found with from one to three buckshot imbedded in various parts of the body. Some of the resulting wounds were severe, but no deaths were directly traced to this cause.

In earlier times the ravages of the parasitic worm, *Uncinaria*, were especially noticeable on the Tolstoi sand flat and portions of Zapadni, but in recent years, due to the shrinkage of the herd, these areas have been abandoned. Very few cases were noted by Dr. Chichester

in 1909, and not one was detected this year. The dead pups dissected showed no lesions whatever, their emaciated appearance and empty alimentary canal indicating death from starvation.

AGES OF SEALS.

Last year 34 branded cows that had been marked as pups not later than 1902 were observed on the rookeries. This year 11 were seen prior to August 1, but during this time there is little opportunity to examine the cows critically, and later in the season such an examination would produce an unwarrantable disturbance on the rookeries. However, the fact is established that there are branded cows in existence, and the time of their disappearance and their possible age may be decided at a later date. It is interesting to note that two cows on St. George bore the T brand of 1899.

Practically every active bull on both islands was examined critically, but not a single brand was seen and none was reported by the government agents or the natives. The branded bull on Kitovi, which last year completed his fifth season, has disappeared. Another bull, blind in one eye, occupied a site on Kitovi for the third season. In other years bulls with scars or other distinguishable marks have been seen at various stations, but these have rarely continued on the active list for more than three or four seasons. It is therefore an established fact that under ordinary circumstances the male becomes active at 8 years of age and lives three or four years thereafter. The age of the female is not known with the same degree of certainty, but it is commonly believed that she lives to the same age.

APPENDIX—EXTRACT FROM FIELD NOTES.

Beginning early in August, the harems begin to show signs of disorganization; the majority of the cows have been served and are free to come or go without serious let or hindrance; the idle and half bulls roam about at will and the breeding season thus passes into its last stage. From this time on observations producing no unwonted disturbance are to be made only from some place of concealment, such as are supplied by the cliffs of Ardiguen or Lukanin. To these two spots I repaired practically every day in August, and for varying lengths of time watched the life of the seal herd. It is unnecessary to detail observations that have already been recorded by several students of the subject, but I may voice again the general verdict that such a show of mammalian life is to be met with nowhere else on the face of the earth, and from several points of view it would indeed be a calamity if the seal meets the fate of the manatee, the sea otter, or the buffalo.

Concerning other life on the islands, much has been said and much remains to be investigated. For many years the bird life has received the attention of the ornithologist and the more important phases of the problems involved have probably been settled; yet there are other matters of minor detail relating to stray migrants, nest materials, and construction and feeding that well deserve attention.

The insects of the islands are numerous and of all the animals or plants doubtless afford some of the most important and interesting problems, if not the very greatest, of purely scientific character remaining to be solved. Owing to the brevity of the summer season, some of the stages in the life history are completed in a surprisingly short space of time, and a comparison of the life histories of related insects in adjoining regions would be interesting to say the least. Furthermore, the conditions under which they survive the winter will also be an interesting chapter in the life of the island organisms.

The flowering plants have been the subject of much study, and it is doubtful if many novelties will be recorded in the future. To a less extent this is true of the lichens, but there are unquestionably small species that have escaped detection; and again there are modifications due to habitat that make it altogether possible that superficially similar forms may in reality be distinct species. Among the fungi there are certainly new forms. On some of the upland slopes in the early season I have found species that do not correspond to any described in the reports of the region.

It is highly desirable that a museum be installed on the islands, containing, so far as is practicable, specimens of all the animals and plants. And equally desirable is a library, comprising all works that in any way are concerned with the biology of the country.

Finally, one word relating to the natives. Considering their antecedents, and especially their former mode of life and lack of advantages, these people have made truly remarkable strides, and yet there is obviously room for improvement. By nature conservative, they are somewhat nonplastic, but at heart they are anxious to better their condition, and they do respond with comparative readiness to all uplifting influences. In matters relating to personal hygiene there is much to be desired, and, improved, their span of life will doubtless be lengthened to a very noticeable degree. And, again, it is highly desirable that during the long and confining winter both the men and women have something to occupy their time—something profitable and yet agreeable, and if possible with a resulting value in some larger community. It is difficult to decide what is best. Numerous plans have suggested themselves, but none of them are free from certain inherent difficulties, and I earnestly hope that those more competent may give the subject their serious consideration, for certainly this species of missionary work carries a rich reward.

In addition to the questions here outlined are others of deep import. Years ago Darwin called attention to the remarkable similarity of the animals on the Galapagos Islands to those on the western slope of South America, and on the basis of this likeness formulated his theory of evolution. Doubtless on the Pribilof Islands the same conditions exist when compared with others of the mainland. Extensive breeding experiments are being carried on in several sections of our country, but it is by no means certain that new species are created in the period measured by a man's life or even in a hundred years. On the islands, however, in a normal habitat, evolutionary agencies have doubtless made their influence felt, even though the islands are geologically young. It seems therefore wise to make extensive collections of the island fauna and flora, to study these critically, and, finally, to compare them with related species on the mainland. These results might be very interesting when considered in connection with the newly formed island of Bogoslof. On this body of land, forced above the sea within the memory of man, we already find plants thriving, and there are doubtless animals on the land or along the shore. Even if there are no visible differences between organisms on this island and those of the Aleutian chain, we may gain some insight into the means whereby their transportation has been accomplished, and if collections and careful notes are kept in the near future the evolutionary side of the subjects may be studied sometime in the years to come.

THE FUR-SEAL FISHERIES OF ALASKA IN 1910

By WALTER I. LEMBKEY

Agent in Charge

Bureau of Fisheries Document No. 749

CONTENTS.

	Page.
The new administration.....	3
Increased scope of agents' duty.....	3
Hire of vessel and purchase and transportation of supplies.....	4
Employees.....	5
Transfer of lessee's property.....	6
Natives' affairs.....	8
Supply depot.....	9
Bank accounts.....	10
Resources of natives.....	11
Census of inhabitants.....	12
Management of seal herd.....	12
Marking of bachelors.....	12
Statistics of killing.....	14
Skins shipped.....	14
Record of drives.....	15
Weights of skins taken.....	19
Enumeration of breeding herd.....	21
Counts of harems.....	21
Counts of pups.....	23
Number of breeding cows.....	23
Census of entire seal herd.....	24
Estimate of half bulls.....	24
Estimate of 2-year-olds.....	24
Number of yearlings.....	25
Number of 3-year-olds.....	25
Summary of seal life in 1910.....	25
Pup-raising experiments.....	26
Pelagic sealing.....	28
Wreck of revenue-cutter Perry.....	29
Foxes.....	29
Conditions and trapping on St. George.....	30
Trapping on St. Paul.....	31
Recommendations.....	31
Killing of bachelor seals.....	31
Support of natives.....	33
Scheme of compensation of natives.....	36
Natives on the Aleutian Archipelago.....	36
Manual training for natives.....	39
Fire protection for Pribilof Islands.....	40

THE FUR-SEAL FISHERIES OF ALASKA IN 1910.

By WALTER I. LEMBKEY,
Agent in charge.

THE NEW ADMINISTRATION.

With the passage of the act of April 21, 1910, the leasing system, which since 1870 had required that the sealing right on the Pribilof Islands be let in 20-year periods to the highest bidder, was abrogated. This new law neither suspended the killing of seals on the islands nor required it to be curtailed, but provided that such killing should be done only by the authority of the Secretary of Commerce and Labor through officers, agents, or employees of that Department, the natives to be employed to perform the labor necessary to secure the sealskins and to receive fair compensation for their labor. So also the sealskins taken under the authority and by the persons already mentioned should be sold by the Secretary to the best advantage of the Government.

By this act the Secretary of Commerce and Labor was given authority also to appoint such additional officers, agents, and employees as may be necessary to carry out the provisions of the act; to purchase at a fair valuation the plant of the former lessee on the islands; to establish and maintain supply depots on the Pribilof Islands; to provide for the transportation of supplies by the charter of vessels; and, finally, to furnish food, fuel, clothing, and other necessaries of life to the natives of the Pribilof Islands, and to provide for their comfort, maintenance, education, and protection.

INCREASED SCOPE OF AGENTS' DUTY.

This act placed upon the Department heavy responsibilities which hitherto had been borne by the lessee. The business of killing seals and curing the skins, the mercantile business with a stock of approximately \$40,000 worth of goods, and, in short, all other practical affairs, were required to be actively managed by the Department agents, who previously had occupied the virtual status of inspectors of the lessee's operations, in addition to the duty of examination of the seal herd and the administration of the natives' affairs.

The act mentioned had not been approved by the President before those charged with the management of the seal fisheries were giving their attention to the working out of the details under the new con-

ditions. On May 9 the annual instructions to the agent in charge were signed; shortly afterwards \$2,000 in cash was advanced to the agent to pay for labor on the islands other than that of killing seals, bonds being given by himself and assistant agents to insure the proper handling of this fund and the faithful performance of duties in general. On May 17 the agent in charge left Washington to begin the preparations for carrying out the requirements of the act of April 21, 1910.

HIRE OF VESSEL AND PURCHASE AND TRANSPORTATION OF SUPPLIES.

On May 21 the agent arrived in San Francisco and on the 26th a charter for the steamer *Homer* at \$142.50 per diem was signed, subject to the approval of the Department of Commerce and Labor. This vessel was delivered under the charter June 1, and was sent first to the coal bunkers to receive her fuel and cargo coal and thence to the covered dock of the Cosmos Line to receive freight.

After the charter of the *Homer* was completed, the purchase of supplies for the natives and the islands in general was next to be taken up. It was found at once that the best prices on the goods required could not be obtained without inviting competitive bids; consequently, with the assistance of the North American Commercial Company, the retiring lessee, which placed its annual requisitions at the Department's disposal, schedules of the principal classes of merchandise were prepared in triplicate and presented to three of the largest mercantile firms in the several lines of business, with the request that each submit a bid in writing. All merchandise, with the exception of small articles of miscellaneous classification, was thus purchased from the lowest bidder, after a careful inspection of the goods to determine whether the quality as well as the price was satisfactory.

It was necessary to visit in person the place of business of each firm to solicit these bids; to go again to make purchases, and again to deliver the vouchers in payment of the articles purchased. With this and the attendant clerical work, it is considered that no time was wasted in the preparations incident to the sailing of the supply ship for the islands.

During the period from June 1 to 10, the supplies were purchased and the vessel loaded. On June 11 the *Homer* sailed from San Francisco, arriving at Dutch Harbor June 24. Coaling there, she proceeded to the islands, arriving at St. George June 27 and St. Paul June 29. Having discharged all freight, she left on July 1 for Dutch Harbor to load coal for the natives' use. Delivering this coal on July 7-11, she returned to San Francisco July 23.

Taking on another cargo of merchandise, together with coal enough for the round trip to the islands, the *Homer* again left San Francisco

August 6, arriving at Dutch Harbor August 21, at St. George the 23d, and St. Paul the 24th. Having received the sealskins aboard, she left St. Paul August 28 and arrived back at San Francisco September 12.

The sealskins were taken at once to Oakland Long Wharf, where, carefully packed in casks and placed in ventilated freight cars, they left on the night of September 14 for New York and thence were shipped to London to be sold at public auction.

EMPLOYEES.

It has already been stated that during the continuance of the leases of the two companies the Government agents on the islands were not concerned with the active management of business, but acted with regard to it merely as inspectors. This does not refer to the supervision of the natives' affairs, the management of which was never the subject of concern by any of the lessee's employees. Under these circumstances the services of the four agents were ample to oversee properly the operations of the lessee and to perform such duties as might be required of the Government's representatives. With the taking over, however, of the business which heretofore formed the exclusive concern of the lessee, an increase in the number of the Government employees on the islands became necessary.

Special biological study of the seal herd having been decided upon, a naturalist was appointed for this work, Dr. Harold Heath, of Stanford University, accepting the position until permanent arrangements could be made. The selection of the additional employees and the assignment of their duties were left to the agent. Of the force required, it was considered advisable to retain as many of the employees of the late lessee as could be used, as these men were efficient, skilled in their duties, and required no instruction other than that necessary to acquaint them with new conditions.

During the summer the force of employees on the islands, in addition to the agents and the naturalist, was as follows:

Name.	Position.	Period.	Annual salary.
On St. Paul Island:			
A. H. Proctor.....	Storekeeper and bookkeeper.....	Indefinite.....	\$1,800
S. Melovidof.....	School-teacher.....	do.....	1,200
H. C. Mills.....	Physician.....	Until fall.....	1,200
[Chinese].....	Cook.....	Indefinite.....	720
N. Bogdanof.....	Stockman.....	do.....	300
[Selected natives].....	Janitor Government house.....	do.....	240
Do.....	Janitor former company house.....	do.....	180
On St. George Island:			
James Murtha.....	Storekeeper.....	Until fall.....	1,200
C. M. Cunningham.....	Physician.....	do.....	1,200
Ned B. Campbell.....	School-teacher.....	Indefinite.....	900
[Chinese].....	Cook.....	Until fall.....	720
M. Iestenkof.....	Stockman.....	Indefinite.....	300
[Selected natives].....	Janitor Government house.....	do.....	240
Do.....	Janitor former company house.....	do.....	180

Mr. Proctor assumed his duties with the idea of serving during the winter on St. Paul. Subsequently, by an order of Secretary Nagel, made during the Secretary's visit to St. Paul, Mr. Proctor was transferred to St. George as acting assistant agent, in place of Assistant Agent Clark, who returned to the Department. Dr. Mills served only during the summer, returning to his home at his own request. The Chinese cooks on both islands were relieved at their own request by others brought up from San Francisco. Assistant Agents James Judge and E. W. Clark with Agent Lembkey returned to the Department on the *Homer*.

Messrs. Murtha and Cunningham served only during the summer, as was contemplated when they were first appointed. Dr. Pedro de Figanière was sent up by the Department to take the place of Dr. Cunningham. Mr. Campbell was appointed by the Department. All others were appointed provisionally from the force on the islands.

During the ensuing winter the force of employees on the two islands will be as follows:

St. Paul: H. D. Chichester, assistant agent in charge; Walter L. Hahn, naturalist; Norman P. Morgan, physician; S. Melovidof, school-teacher; a Chinese cook; and N. Bogadanof, stockman.

St. George: A. H. Proctor, acting assistant agent; P. de Figanière, physician; Ned B. Campbell, school-teacher; a Chinese cook; and M. Lestenkof, stockman.

The respective assistant agents are performing their usual duties in addition to those heretofore devolving upon the lessee's agents. When it is considered also that the office force of the lessee in San Francisco, with over \$20,000 in salaries, has been eliminated, it will be seen that the island service, while highly efficient, is conducted at a minimum of expenditure. No increase in administrative force has occurred. A bookkeeper, two physicians, and two school-teachers only have been added, in addition to cooks and miscellaneous native help.

TRANSFER OF LESSEE'S PROPERTY.

By a letter dated May 7, 1910, from the Commissioner of Fisheries, the agent was directed to confer with the North American Commercial Company and if possible to arrive at a fair and just valuation to be placed upon the property of that company on the Pribilof Islands, with a view to purchase by the Government.

Two days after arrival at San Francisco a conference was had with the company, at which a statement of the presumed value to the Government of the company's holdings on the islands was made. After consideration of the question the company several days later agreed to transfer the Pribilof Islands plant at the valuation proposed at the previous conference.

Upon arrival at the islands an inventory as of June 30 was taken. Later, the transaction having received the approval of Secretary

Nagel, who personally visited the islands and inspected the plant, vouchers were drawn to cover the various amounts shown on the inventory according to the basis of settlement proposed and accepted, and were transmitted to the Department for settlement.

A recapitulation of the inventories on the two islands, as taken on June 30, 1910, with a memorandum of the basis of settlement, follows:

ST. PAUL ISLAND.

Company's inventory.		Settlement price.	
Merchandise.....	\$5,154.33	San Francisco invoice cost.....	\$5,154.33
Tools and implements.....	3,522.83	50 per cent of inventory.....	1,761.41
Drugs and instruments.....	816.63	Inventory cost.....	816.63
Household furniture.....	2,957.22	25 per cent deducted from inventory.....	2,217.92
Dispensary furniture.....	159.97	Do.....	119.98
Boats and bidarras.....	3,835.40	Launch.....	\$2,000
		Boat.....	400
		Do.....	275
		3 bidarras, at \$175 each.....	525
		Lump sum.....	90.00
Telephone line.....	367.79	Do.....	257.00
School supplies.....	276.29	50 per cent of inventory.....	12,841.72
Company buildings.....	25,653.45	Do.....	8,634.55
Native dwellings.....	17,269.11	Inventory cost.....	138.00
Sea-lion skins.....	138.00	Do.....	967.62
Live stock.....	967.62	Do.....	1,260.02
Salt and twine.....	1,260.02	Do.....	61.88
Fox skins (traps, etc.).....	61.88	Lump sum.....	200.00
Library.....	1,012.86	50 per cent of inventory.....	402.31
Wharf ways and derrick.....	804.63	Same, at \$17.....	1,132.17
Coal, 66 tons 1,339 pounds, at \$20.....	1,331.97		
Total.....	65,620.00	Total.....	39,255.54

ST. GEORGE ISLAND.

Merchandise.....	\$6,352.03	San Francisco invoice cost.....	\$6,352.00
Coal, 38 tons, at \$20.....	760.00	Same, at \$17.....	646.00
Dispensary.....	718.97	50 per cent of inventory.....	359.48
Live-stock account.....	313.72	Inventory cost.....	313.72
Groceries, company house mess.....	227.73	San Francisco invoice cost.....	227.73
		San Francisco invoice cost after inspection.....	198.10
Salt and seal twine.....	198.10	Do.....	98.87
Old salt.....	98.87	Inventory cost.....	85.71
Sea-lion skins.....	85.71	Lump sum.....	700.00
Boats and bidarras.....	1,215.96	50 per cent.....	5,802.02
Company buildings.....	11,604.04	Do.....	868.61
Derrick and landing (including cars and track).....	1,737.23	25 per cent deduction.....	1,532.72
House and office furniture.....	2,043.63	Lump sum.....	100.00
Library.....	670.64	50 per cent.....	3,323.48
Native dwellings.....	6,646.96	Do.....	148.62
Telephone.....	297.25	Do.....	582.23
Tools and implements.....	1,164.47		
Total.....	34,135.31	Total.....	21,339.32

The foregoing lists represent a total valuation for both islands of \$60,568.17. Subsequent deductions because of errors in addition, computation, etc., reduced this amount by \$26.69. A final settlement was made by the Department for \$60,541.48 and checks for that amount were transmitted to the company.

With the exception of the buildings, practically everything on the inventory represents new stock, purchased by the company during its lease and not acquired from the former lessee. With regard to the buildings it may be said that, although erected by the former lessee, they have been kept from deterioration by constant repair and could not be replaced for anything approaching the price paid for them by the Government. On St. George the company's

dwelling house and warehouses were virtually rebuilt by the late lessee, when also several new native dwellings were added. On St. Paul constant repairs were made to all the buildings during the period of the lease, and the buildings not only are habitable but efficient. When it is considered furthermore that only 50 per cent of the inventory valuation was paid for these buildings, it may be seen that the price was not excessive.

NATIVES' AFFAIRS.

Upon the agents' arrival at the islands considerable anxiety was found to have existed in the minds of the natives and others as to the time of arrival of the supply ship and the arrangements which might be made for the conduct of affairs under the changed conditions. Through the revenue cutters which touched at the islands previous to the arrival of the *Homer*, information had been received of the assumption of active management by the Government, but no intimation as to what efforts were being put forth by the Department for taking charge of the practical affairs. This anxiety had been heightened by the fact that the supply of some articles of necessity, as food on St. Paul, had been almost consumed. In fact, to provide against an imminent shortage it had become necessary in the early part of June to obtain by the revenue cutter *Manning* a quantity of flour, biscuits, salt beef, and canned vegetables from Dutch Harbor. In addition to this fear of impending famine, the natives had received the impression that they would be obliged to labor for the Government without any compensation other than clothing and food, as had been actually the case under the Russian régime.

The agents' first effort, therefore, was to allay these impressions and to establish relations of confidence with the natives, though, as a matter of fact, the arrival of a shipload of supplies and of a gunny sack containing about 150 pounds of coin had the effect automatically of removing the greater portion of this uncertainty. In addition, conferences were had with individual natives and with the assembled communities, in which the changes which had occurred during the past season were explained and assurance was given that the intention of the Government was to improve the present condition of the natives wherever possible rather than to make it less favorable than under the late lessee.

It was necessary specifically to reassure them that cash payments for sundry labor would be continued under the new management. This has been the source of almost all the cash received by the natives, and the loss of it the occasion of their chief anxiety. The assurance of the continuation of these payments in cash, together with the increase in the rate of payment for taking sealskins, and the material reduction in the prices at which merchandise is to be sold to the natives out of the stores on the islands, all had the effect of

restoring confidence and obtaining a renewal of the natives' good will.

Supply depot.—Immediately upon the arrival of the *Homer* all hands not entirely occupied with sealing began taking an inventory of merchandise and other property belonging to the company, with a view to its being taken over by the Government, in accordance with instructions contained in the letter to the agent in charge dated May 7, 1910. This inventory was prepared in time to be transmitted on the return of the *Homer* on her first trip.

After completing the inventory the merchandise which arrived on the *Homer* was uncrated and checked with the invoices. The price was marked on the articles at the rate fixed in the instructions of the agent, namely, a flat rate of 33½ per cent advance over San Francisco wholesale prices. The prices of those articles of merchandise also which were taken over from the company were made to conform to the prices fixed for the new invoices of goods.

The application of this flat rate of 33½ per cent advance had the result of selling merchandise to the natives at lower prices than ever before in the history of the islands. Because of high market prices in San Francisco at the time the spring requisition was purchased the retail price of butter was increased from 35 cents to 42 cents; flour remained the same, at \$1.75 a quarter barrel; lard was raised from 18 cents to 21 cents a pound; rubber boots, from \$6 to \$6.35 a pair; canned beef from 30 cents to 35 cents each. Some few other articles were sold at the same rate as formerly; all other prices were reduced. A statement of some of these reduced prices follows:

Articles.	Former price.	Present price.	Articles.	Former price.	Present price.
Apples:			Needles.....	\$0.05	2 for \$0.05
Canned.....	\$0.25	\$0.20	Oil:		
Evaporated.....	2 for .30	3 for .25	Coal.....	.40	.26
Apricots, canned.....	.25	.20	Cottonseed.....	.35	.25
Arctic:			Onions.....	.06½	.05
Men's.....	2.25	1.90	Peaches, canned.....	.25	.20
Women's.....	1.50	1.35	Pears, canned.....	.25	.20
Beans, canned.....	.20	.15	Peas.....	.20	.15
Bedspreads.....	2.25	1.70	Potatoes.....	.03½	.02½
Beef, salt.....	.12½	.09	Baking powder.....	.20	.15
Blackberries, canned.....	.25	.20	Prunes.....	.15	3 for .25
Blankets.....	7.00	5.50	Raisins.....	.15	3 for .25
Calico.....	.10	3 for .25	Rice.....	3 for .25	3 for .20
Candies.....	.02½	.02	Worcestershire sauce, American.....	.25	.15
Candy, 2 pounds.....	.50	.25	Shoes:		
Chimneys, lamp.....	.15	2 for .15	Babies'.....	.75	.55
Coffee.....	.25	.20	Do.....	1.25	.90
Collars, white.....	.25	2 for .25	Boys'.....	3.00	2.00
Corn, canned.....	.20	.15	Children's.....	2.50	1.75
Crackers:			Do.....	2.00	1.40
Soda.....	.10	3 for .25	Men's.....	4.00	3.15
Sweet.....	.20	.15	Misses'.....	2.50	1.75
Cups and saucers.....set..	.20	.15	Women's.....	3.00	2.35
Dress goods.....	.60	.50	Do.....	4.50	2.60
Ewers and basins.....set..	2.00	1.25	Swiss, dotted.....	.25	.15
Gingham.....	.15	2 for .25	Soap.....	.06½	.05
Gloves, men's, wool.....	.50	.25	Socks.....	.50	.45
Knives, pocket.....	.40	.30	Tea.....	.50	.25
Jams.....	.25	.20	Tobacco, leaf.....	.50	.40
Jelly.....	.25	.20	Tomatoes, canned.....	.20	.15
Lining, cotton.....	.15	.12½	Trousers.....	5.00	4.00
Milk, condensed.....	.25	.20			

On every weekly order issued a saving of from 75 cents to \$1.50 was made by reason of these reduced prices. In addition the price of coal was reduced from \$20 a ton to \$12.75 plus a small charge for stevedorage at either end. While no accurate computation has yet been made, it is believed that by reason of the reduced prices of commodities sold the purchasing power of the natives will be increased by several thousands of dollars.

Bank accounts.—When the Alaska Commercial Company in 1870 began taking seals under its lease, in addition to providing comfortable dwellings for the native inhabitants, it also endeavored to encourage thrift among them by receiving deposits of money from such natives as desired to open savings accounts. On these accounts, which were subject to check at all times, the company paid interest at the rate of 4 per cent on balances found on May 31 of each year. During the period of this company's lease some natives had accumulated accounts of over \$2,000 each.

These accounts were taken over by the North American Commercial Company when it succeeded to the sealing privilege in 1890. While during the 20-year lease of the latter company these funds on deposit became smaller, due to the lessened amounts earned by the natives and to distribution to nonresident heirs upon death of the owner of the account, there still remained a few so-called bank accounts in the hands of the North American Commercial Company at the time of the expiration of its lease.

When the contract of the North American Commercial Company expired in 1910 these funds remained on deposit with it, and some action with reference thereto became necessary on the part of the Government, which then took over the active management of the business.

In the instructions dated May 9, 1910, it was directed that if the balance on the bank account of any native was small it should be paid by the company directly to the native; if, however, the native desired, it should be held by the company and deposited in a safe financial institution in San Francisco by the agent in charge as attorney in fact for the benefit of the native owning the account, the interest to be collected annually and paid directly to the native.

Upon arrival at the islands last spring the natives were informed of the situation and told that if they desired their money could be deposited in a bank in San Francisco previously selected, which would pay interest at the rate of $3\frac{1}{2}$ per cent per annum. They all assented to the redepositing of their funds in the manner stated.

Such small accounts as did not exceed \$25 were paid to the owner in cash by the company; the accounts of larger amount than that stated were closed by the company's presenting the respective owners with drafts for the several amounts.

Each native who possessed one of these drafts delivered the same to W. I. Lembkey and upon blanks previously provided signed a power of attorney to him authorizing him to deposit the drafts with a bank in San Francisco, to collect the amount of any interest due thereon and to give receipts for the same.

A list of the accounts and the persons to whom they belong follows:

St. George Island:		St. Paul Island—Continued.	
Fevronia Galanin.....	\$40. 00	Peter Bourdukofsky.....	\$130. 00
Dimitri Lestenkof.....	137. 00	Elizabeth Rookavishnikof.....	40. 00
Michael Lestenkof.....	240. 00	Agrifina Fratis.....	71. 00
Peter Prokopiof.....	83. 55	Agrifina S. Pankof.....	285. 00
Emanuel Zaharof.....	33. 20	Peter Oustigof.....	140. 00
Zoya Swetzofof.....	123. 00	Alexander Melovidofof.....	235. 00
Mary Galanin.....	245. 00	Julia B. Krukof.....	170. 00
Michael Shane.....	63. 55	Simeon Fratis.....	71. 00
Mary Philamonof.....	90. 05	Akalina Fratis.....	426. 00
		Alexai Emanof.....	230. 00
Total.....	<u>1,055. 35</u>	Tekan Volkof.....	966. 00
St. Paul Island:		Martha Fratis.....	71. 00
Alexander Mercuriof.....	170. 00	John Hansen.....	370. 00
Nekita Hopof.....	50. 00	Oulianna Fratis.....	71. 00
Agrifina Bbgadanof.....	161. 10		
Marina Stepetin.....	40. 00	Total.....	<u>4,050. 40</u>
Apollon Bourdukofsky.....	203. 30	Grand total.....	<u>5,105. 75</u>
Parascovia Kozlof.....	150. 00		

The St. Paul drafts were deposited to the credit of W. I. Lembkey, trustee for the various natives. Separate accounts were opened with each fund and pass books provided to be delivered to each native owning the account. In cases where the money was owned by a minor child, the account was opened in the name of its natural guardian—either one of its parents, or if an orphan, the person with whom it resides—with Agent Lembkey as trustee for the guardian.

Upon taking the St. George drafts to the bank it was discovered that by an oversight they had not been indorsed by the persons in whose favor they were drawn. Unfortunately, therefore, they could not be deposited. An arrangement was made with the North American Commercial Company, however, whereby the amount of these St. George drafts, \$1,055.35, was deposited by the company to protect the drafts which it will be necessary to take back to St. George Island for proper indorsement. After being so indorsed they will be paid by the bank and savings accounts opened with each of the persons named, in the same manner as the drafts from St. Paul.

The interest on these accounts will be collected annually and paid to the proper persons. The receipts for money so paid will be submitted with the annual report.

Resources of natives.—During the summer of 1910, from taking seals, and the previous winter from trappings foxes on St. George,

the natives of the islands earned the following amounts, to be applied to their support:

St. George:	
203 blue foxes, at \$5; 9 white, at \$1.....	\$1,024
2,834 sealskins, at \$1.....	2,834
St. Paul:	
664 sealskins, at 75 cents.....	498
10,088 sealskins, at \$1.....	10,088
Total.....	14,444

As the fox skins were delivered to the North American Commercial Company, that company paid directly to the agent on St. George for the natives the amount of \$1,024, due the natives on that account. The company also paid in cash to the agent on St. Paul the \$498 due the natives from the 664 sealskins which the Department authorized the company to take to complete its quota of 15,000 for 1909. The amounts of \$10,088 earned by the St. Paul natives and \$2,834 earned by the St. George natives for taking the sealskins shipped on Government account in 1910 were credited to the natives on the island books. Payments of cash therefrom were not made except of small sums in very rare instances. Each native sealer, however, was allowed to draw supplies against this fund at a fixed rate each week until the cost of such supplies equaled the amount of the native's credit from earnings; after this, supplies to be issued to him directly from the stores in sufficient quantity to support himself and family.

The various statements of the division of natives' earnings are filed in the Bureau of Fisheries at Washington.

Census of inhabitants.—On St. Paul, on June 30, 1910, there were 198 resident natives, including 98 males and 100 females, a net increase of 5 over the previous census. During the year 13 births, 1 arrival, and 9 deaths occurred.

On St. George, at the same date in 1910, 91 natives were present, of which 45 were males and 46 females. Six births and 2 deaths occurred during the year, leaving a net increase of 4 in the population.

Detailed censuses are filed in the Bureau of Fisheries at Washington.

MANAGEMENT OF SEAL HERD.

MARKING OF BACHELORS.

The general instructions to the agent, dated May 9, 1910, required that not any 2-year-old bachelors but only 500 3-year-old bachelors should be marked to form the breeding reserve. This was predicated upon the assumption that the 500 3-year-olds so reserved would be over 14 per cent of the whole number of such young males in the herd. Subsequently, by a telegram from the Secretary dated June 6, which, not having been delivered, presumably through the fault of the telegraph company, was repeated June 10, the number of

3-year-old males to be reserved by marking was increased from 500 to 1,000.

These were apportioned between the two islands, by assigning 800 to St. Paul and 200 to St. George, for the reason that there are in round numbers four times as many breeding seals on St. Paul as on St. George. Upon arrival at St. George Island a copy of the annual instructions was given to Assistant Agent Clark, and he was also informed that the quota of bachelors to be reserved on St. George was 200 3-year-olds. As the vessel remained at St. George only a few hours, and as numerous other matters required consideration, it was not possible to put into writing the various explanations of the instructions.

Upon my return to St. George Island two weeks later I was informed by Agent Clark that the quota of marked bachelors had been secured. No statement of the number so marked, however, was made, and at the close of the season among the data received detailing the season's work on St. George no mention was made of the number of bachelors branded. Upon meeting Agent Clark on the *Homer* after he had left St. George for San Francisco, upon specific inquiry I ascertained for the first time that the instructions were misapprehended by him and that he had sought to brand on St. George only 100 3-year-olds, and did actually brand only 108 of that class of young males. He had not the memoranda showing the dates on which drives were made for this purpose and the number secured from each drive. As the season then had been closed for three weeks it was useless to cause the marking of an additional number to make up the deficiency in the breeding reserve for that island.

On St. Paul, however, more young males were branded than the total number for both islands required by the instructions. Previous to my arrival on that island, on June 29, with the current instructions, Assistant Agent Judge, acting under the instructions for the previous year, had already marked 337 2-year olds in addition to 279 3-year-olds, 14 4-year-olds, and 5 5-year-olds. After my arrival additional 3-year-olds only were marked to complete the number of that class required for St. Paul. A record of the bachelors marked on St. Paul, showing also dates and rookeries driven from, follows:

RECORD OF BACHELORS MARKED ON ST. PAUL ISLAND FOR BREEDING PURPOSES,
SEASON OF 1910.

Date.	Rookery.	Two years.	Three years.	Four years.	Five years.
June 17	Reef.....	46	77
27	Zapadni.....	82	56
28	Reef and Gorbach.....	209	146	14	5
July 2	Northeast Point.....	246
4	Reef.....	191
5	Zapadni.....	91
	Total.....	337	807	14	5

The total number of bachelors marked on both islands, therefore, would be as follows: 2-year-olds, 337; 3-year-olds, 915; 4-year-olds, 14; 5-year-olds, 5; total, 1,271.

The report of London trade sales this year shows that 5,006 large pup and middling pup skins (which are accepted to be those of 3-year-old bachelors) appeared in the 1910 catch. Adding to these the 915 reserved 3-year-olds would make a total of 5,921 of that class which we might claim were in the herd in 1910. Of this whole number, the number reserved (915) is over 15 per cent.

Two-year-old males were not required by the current instructions to be reserved, for the reason that the number of 2-year-olds having skins of 5 pounds and under, together with those 2-year-olds which would not appear in the drives at all, of which there are always some, it was believed would be sufficient to supply the necessary number of 3-year-olds in 1911.

STATISTICS OF KILLING.

St. Paul.—From August 9, 1909, to June 17, 1910, 6 drives of seals on St. Paul and 2 on Sea Lion Rock were made to furnish food to the inhabitants of St. Paul. From these, 1,573 skins were obtained, including 1 from a seal found dead at Rocky Point. From July 3 to 31, 29 drives were made on St. Paul for skins, in which 8,683 skins were secured. On August 10, 1910, an additional drive was made to furnish food for the natives during the coming "stagey season," from which 496 skins were secured. From the sources enumerated a total of 10,752 skins were obtained during the season ended August 10, 1910.

St. George.—On St. George during the so-called food-killing season, from August to November, 1909, 18 seals were killed at various dates by the guard at Zapadni; 8 drives also were made, in which 482 seals were killed, filling the quota of 500 for food allowed for that island. During the season of killing for skins, 2,314 skins were secured in 10 drives, 16 were obtained from the seals killed at various times by watchmen for food, and 4 were left in salt from the previous season, a total of 2,334, in addition to the 500 taken during the food-killing season.

SKINS SHIPPED.

St. Paul.—Of the skins taken on St. Paul, 664 were delivered to the North American Commercial Company, under authority of the department's letter of January 5, 1910, to complete that company's quota of 15,000 skins for 1909. The remainder, 10,088 skins, were available for shipment on Government account. While this number

supposedly was shipped from St. Paul on the *Homer*, on August 28, word was received in October last from Assistant Agent H. D. Chichester, in charge on St. Paul, that after the departure of the *Homer* with the skins on board a bundle containing 2 sealskins was found wedged under the floor of the skin lighter or bidarra, in which crevice it had become obscured during the shipment of the skins. These two were placed in the salt house to apply on the shipment of the following year. The total number of skins, therefore, shipped from St. Paul in 1910 for Government account was 10,086.

St. George.—On August 23, 1910, the whole number of skins taken on St. George, from the sources enumerated (2,834), were placed on board the *Homer* to be shipped to San Francisco for Government account.

The whole number of skins from both islands, recapitulated from the data already given, is as follows:

From St. Paul:	
By North American Commercial Company	664
By Government.....	10,086
From St. George, by Government.....	2,834
Total.....	13,584

RECORD OF DRIVES.

On St. Paul, during the season of 1910, no record was kept of the seals dismissed from the food drive made on June 6 on Sea Lion Rock, as the configuration of the ground there is such that the seals can not be herded, but escape in every direction upon the landing of the clubbers, who kill such as they can while the seals are running off. So also no record was kept in the drive for "branding" on June 17, from which at the same time 145 seals were killed. The record of dismissals, therefore, begins on July 3, when the drive was made at Northeast Point for "branding," at which, at the same time, the 2-year-old bachelors in the drive, not being required to be marked, were killed.

In the 32 drives made on St. Paul from July 3 to August 10, a total of 12,434 seals appeared, of which 9,179, or 73 per cent, were killed and 3,255 dismissed. Those dismissed consisted of 1,581 small, 825 large, and 849 of those marked for the breeding reserve. This killing was 4 per cent closer than during the lessee's killing season of 1909, when 69 per cent of all seals driven were killed.

SEALS KILLED AND SEALS DISMISSED FROM DRIVES ON ST. PAUL ISLAND, SEASON OF 1910.

Date.	Rookery.	Killed.	Dismissed.			Total driven.	Per cent killed.
			Small.	Large.	Branded.		
July 3	Northeast Point.....	437	32	67	536	81
4	Reef.....	331	48	31	410	80
5	Zapadni.....	166	48	31	245	67
6	Tolstoi and Lukanin.....	142	6	39	28	215	66
7	Halfway Point.....	77	2	9	3	91	84
8	Northeast Point.....	293	37	47	85	462	63
9	Reef and Gorbach.....	437	21	28	116	602	72
9	Tolstoi and Lukanin.....	120	2	17	5	144	83
10	Zapadni.....	198	10	18	32	258	76
14	Northeast Point.....	407	16	35	15	473	86
14	Polovina.....	5	10	15	33
15	Reef and Gorbach.....	429	19	9	17	474	90
15	Tolstoi and Lukanin.....	131	17	8	2	158	82
16	Zapadni.....	339	77	22	24	462	73
20	Northeast Point.....	487	132	29	26	674	72
20	Halfway Point.....	5	1	6	83
21	Reef and Gorbach.....	548	56	33	42	679	80
21	Tolstoi and Lukanin.....	449	53	23	26	551	81
22	Zapadni.....	346	51	32	32	461	75
25	Northeast Point.....	465	48	65	38	616	75
25	Halfway Point.....	18	17	3	38	47
26	Reef and Gorbach.....	664	139	30	78	911	72
26	Tolstoi and Lukanin.....	336	32	35	37	440	76
28	Zapadni.....	318	55	14	44	431	73
28	Halfway Point.....	12	1	2	1	16	75
29	Northeast Point.....	589	64	68	23	744	79
30	Reef and Gorbach.....	575	86	37	55	753	76
30	Tolstoi and Lukanin.....	204	29	29	21	283	72
31	Zapadni.....	155	25	16	26	222	69
Aug. 10	Reef and Gorbach.....	496	475	24	69	1,064	46
	Total.....	9,179	1,581	825	849	12,434	73

CLASSIFICATION OF LARGE SEALS DISMISSED FROM DRIVES ON ST. PAUL ISLAND, SEASON OF 1910.

Date.	Rookery.	Four years.	Five years.	Six years.	Seven years.	Adult.
July 4	Reef.....	7	9	9	6
5	Zapadni.....	12	6	11	2
6	Tolstoi and Lukanin.....	11	8	11	6	3
7	Halfway Point.....	1	2	6
8	Northeast Point.....	10	9	14	14
9	Reef and Gorbach.....	8	9	2	9
9	Tolstoi and Lukanin.....	8	2	7
10	Zapadni.....	8	5	3	2
14	Northeast Point.....	12	6	10	7
14	Polovina.....	2	3	5
15	Reef and Gorbach.....	4	2	3
15	Tolstoi and Lukanin.....	4	2	2
16	Zapadni.....	10	4	3	3	2
20	Northeast Point.....	19	5	4	1
20	Halfway Point.....
21	Reef and Gorbach.....	2	9	12	10
21	Tolstoi and Lukanin.....	4	9	4	6
22	Zapadni.....	16	10	4	2
25	Northeast Point.....	24	21	18	2
25	Halfway Point.....	3	4	4	4	2
26	Reef and Gorbach.....	10	5	12	3
26	Tolstoi and Lukanin.....	13	16	5	1
28	Zapadni.....	8	2	2	2
28	Halfway Point.....	1	1
29	Northeast Point.....	17	9	3	5	4
30	Reef and Gorbach.....	14	16	4	1	2
30	Tolstoi and Lukanin.....	7	16	2	4
31	Zapadni.....	9	4	2	1
Aug. 10	Reef and Gorbach.....	12	1	2	6	3
	Total.....	255	195	148	111	19

On St. George the record of seals driven and dismissed covers the period from June 13 to July 31. In this time 3,065 seals were driven and 2,295 killed, while 240 small, 343 large, and 187 marked seals were released. The number killed represents 74 per cent of the whole number driven, an increase of 11 per cent over the killings of 1909, when 63 per cent of those driven were killed.

SEALS KILLED AND SEALS DISMISSED FROM DRIVES ON ST. GEORGE ISLAND, SEASON OF 1910.

Date.	Rookery.	Killed.	Dismissed.			Total driven.	Per cent killed.
			Small.	Large.	Branded.		
June 13	East.....	31	4	38	73	42
23	East and North.....	138	11	93	242	57
30	do.....	162	16	79	255	63
July 5	East, North, and Staraya Artel.....	171	55	30	58	314	54
12	do.....	313	26	14	21	374	83
16	North.....	258	18	5	5	286	90
21	North and East.....	376	48	15	27	466	80
26	East, North, and Staraya Artel.....	405	42	35	37	519	77
31	do.....	441	20	36	39	536	82
	Total.....	2,295	240	343	187	3,065	74

CLASSIFICATION OF LARGE SEALS DISMISSED FROM DRIVES ON ST. GEORGE ISLAND, SEASON OF 1910.

Date.	Rookery.	Four years.	Five years.	Six years.	Seven years.
June 13	East.....	17	9	9	3
23	East and North.....	25	43	18	7
30	do.....	39	7	21	10
July 5	East, North, and Staraya Artel.....	8	13	6	3
12	do.....	4	4	6
16	North.....	4	1
21	North and East.....	8	5	2
26	East, North, and Staraya Artel.....	13	6	11	5
31	do.....	13	11	6	6
	Total.....	131	98	78	36

It will doubtless be remarked that the percentage of seals killed in 1910 was greater than in the preceding year. The seals killed in 1910 were, however, neither larger nor smaller than those taken in 1909, but conformed at least as closely to the prescribed ages and weights as they did in 1909, the last year of the leasing system. Indeed, when doubt arose, as often it does arise, whether a seal was of the 3-year-old (or killable) age or whether it was of the 4-year-old (or prohibited) age, in 1910 the animal was allowed to escape, whereas in 1909 it would have been killed. In this respect it may be said that the killing in 1910 conformed even more closely to regulations than that of 1909.

Since the animals killed in 1910 were of the same class as those of the preceding year, and since the rejections from the drives were

fewer in proportion to those killed, it must be concluded that this condition is due not to closer killing, but to the absence, for some reason, of those animals which are not killable and which when they appear in drives make up the number of "rejected" seals. In other words, the bachelors driven were not culled more closely for killables, but fewer rejectable seals appeared in the drives, thereby making the rejection percentages smaller.

One certain reason for this increased percentage of killed in 1910 is to be found in the lessened number of "branded" or marked bachelors with which to deal during the killing. In previous years 2,000 of these marked bachelors were present during the killing season, while in 1910 only 1,000 of them were marked. Furthermore this missing thousand would have been composed of 2-year-olds which haul up on the bachelors' hauling-grounds much more frequently than do the 3-year-olds. With 1,000 2-year-olds marked for exemption from killing, it would have been certain that from 1,200 to 1,500 more rejections would have occurred during the season, the number of rejections of this class varying somewhat from year to year. On the other hand, rarely does the number of subsequent rejections of the 3-year-olds equal the number of that class actually marked.

Had 1,200 been added to the number of rejections obtained in 1910, the percentage of killed would have been 69, very nearly what it was in the year preceding.

Another presumed cause of the lack of small rejections last year is the probable fact that the smaller seals, i. e., those that had skins under 5 pounds in weight, failed to haul up on land proportionately in the same numbers as hitherto; that is to say, these small seals remained for longer periods in the water than usual. In respect to this matter we are met with the fact that we are wholly unable to state anything definite concerning the hauling habits of young bachelors. Some are always in the water and on inaccessible hauling grounds, for which reasons no definite idea of the whole number in existence can be obtained. Nevertheless, it is known that the hauling habits of seals vary from year to year; that these habits are altered by circumstances not incident to their natural environment, such as the action and movement of the pelagic fleet; that these bachelors haul in one year in greater numbers proportionately on one island than the other, or on one rookery than on other rookeries; that they return to their normal habits with the disappearance of the cause which forced them to abandon those habits temporarily.

For 1910 it can be shown that these small seals, which were yearlings the preceding year, were not killed, either as pups or yearlings. Yearlings are never killed on land except through unavoidable accident, and an analysis of London sales of skins shows that yearlings form but a small fraction of 1 per cent of the pelagic catch. Unless they

died from natural causes, of which there is no evidence, they must be in existence somewhere as 2-year-olds. Not having appeared on land during the summer, the natural inference must be that they were in the water and did not haul on land.

That there were in existence small seals which did not haul during the summer might be indicated by the fact that in the killing on August 10 the number of small seals turned away was entirely out of proportion to the usual number occurring in drives during the season. The absence of these small seals during the summer was a matter of remark, and their reappearance at the last drive of the season also was noted with interest.

In treating of this matter it is desired to show that notwithstanding the fact that of seals driven a greater percentage killed appears on the record for this year as compared with last, no smaller seals than usual were killed and not as large seals were taken as previously. The increased percentage is the result, first, of the absence of 2-year-old marked bachelors present in former years, and secondly, to a failure of young nonkillable seals to haul on land in their usual numbers during the summer.

WEIGHTS OF SKINS TAKEN.

Of the 10,752 skins taken on St. Paul, 10,749 were weighed. Of these 70 were under 5 pounds and 48 over $8\frac{1}{2}$ pounds. On St. George, 2,834 skins were weighed, of which 20 were under 5 pounds and 11 over $8\frac{1}{2}$. Of the overweight skins on St. Paul, nearly all were taken in a food killing on Sea Lion Rock, and before weighing were immersed in sea water until they were saturated. In this condition each carried several pounds of water, increasing their weight correspondingly. Had they been weighed dry, or even with the usual quantity of moisture, few of them would have been above the prescribed limit.

It is not possible to avoid wetting the seals taken on Sea Lion Rock, neither is it permissible to salt the skins without weighing. It is wholly undesirable also to alter the statistics of weights in such manner as to attempt to compensate for excess due to the presence of water or other foreign substances in the fur. The weights therefore have been recorded as taken, but due allowance must be made for conditions which change the weights and which have no relation to the size of the skins.

The skins that were underweight were likewise taken mainly in food drives, at a time when the natives were eager for fresh meat and when they were restricted to killing seals having skins under 7 pounds. With the necessity of rejecting all the females and all the larger males from the food drives, it can readily be appreciated that the tendency of the natives is to let few of the small males escape, even if the skins weigh a few ounces less than 5 pounds.

On the whole it can be seen that only a few skins of the whole catch were outside the weights prescribed and that these were taken unavoidably.

WEIGHTS OF SEALSKINS TAKEN ON THE PRIBILOF ISLANDS, ALASKA, DURING THE YEAR ENDED AUGUST 10, 1910.

Weight.	St. Paul Island. ^a	Weight.	St. George Island. ^b
<i>Pounds.</i>		<i>Pounds.</i>	
4.....	6	4.....	1
4½.....	4	4½.....	5
4¾.....	20	4¾.....	14
4¾.....	40	5.....	125
5.....	670	5¼.....	82
5¼.....	710	5½.....	406
5½.....	1,014	5¾.....	202
5¾.....	1,277	6.....	628
6.....	980	6¼.....	106
6¼.....	1,113	6½.....	524
6½.....	1,176	6¾.....	114
6¾.....	993	7.....	321
7.....	752	7¼.....	43
7¼.....	553	7½.....	168
7½.....	552	7¾.....	21
7¾.....	327	8.....	54
8.....	203	8¼.....	4
8¼.....	172	8½.....	5
8½.....	139	9.....	6
8¾.....	7	9¼.....	1
9.....	17	9½.....	2
9¼.....	4	10.....	1
9½.....	7	10½.....	1
9¾.....	4		
10.....	1	Total.....	2,834
10½.....	2		
11.....	1		
11¼.....	4		
11½.....	1		
12.....	1		
Total.....	10,749		

^a Nearly all the oversize skins listed from St. Paul Island were taken in a food killing from Sea Lion Rock, on which occasion the skins when weighed carried from 1 to 3 pounds of water each. Had they been dry when weighed, very few or none would have exceeded the prescribed weights. The major portion of skins underweight were taken in food drives for the natives, when large seals were released, and, consequently, the smaller seals were killed closely.

^b Of the skins from St. George over or under the limit of weight only 3 were taken during the sealing season proper. Four were taken by the company last year, and withheld from the quota; the others were taken during food killings, when the natives were particularly eager for fresh meat.

Following is a statement furnished by Messrs. C. M. Lampson & Co., of the sizes of the sealskins consigned to them by the United States Government for auction in London. This statement shows the classification of the 12,920 skins as weighed and assorted upon their receipt by the firm.

ASSORTMENT OF ALASKA SALTED FUR SEALSkins FOR ACCOUNT OF UNITED STATES GOVERNMENT, DEPARTMENT OF COMMERCE AND LABOR.

[London, 19th November, 1910, 64 Queen Street, E. C. Subject to recount.]

	Lbs. oz.		Lbs. oz.
78 smalls.....	7 15	195 middling pups, rubbed.....	6 6
713 large pups.....	7 2	290 small pups, rubbed.....	5 11
3,032 middling pups.....	6 7	75 ex. small pups, rubbed.....	5 3
4,899 small pups.....	5 12	36 faulty.....	
1,266 ex. small pups.....	5 5		
11 ex. ex. small pups.....	4 10	<u>12,732</u>	
33 smalls, low.....	7 11		
135 large pups, low.....	6 9	5 smalls.....	
498 middling pups, low.....	6 1	21 large pups.....	
501 small pups, low.....	5 9	48 middling pups.....	
88 ex small pups, low.....	5 0	94 small pups.....	
10 smalls, cut.....	7 2	18 ex. small pups.....	
71 large pups, cut.....	6 13	2 faulty.....	
238 middling pups, cut.....	6 2		
421 small pups, cut.....	5 6	<u>188</u>	
81 ex. small pups, cut.....	4 15	<i>a</i> 12,922	
6 smalls, rubbed.....	7 0		
55 large pups, rubbed.....	6 14		

a See p. 15. This number recorded as shipped, but two skins afterwards found wedged under floor of boat used for lightering skins to steamer *Homer*.

ENUMERATION OF BREEDING HERD.

COUNTS OF HAREMS.

The usual counting of harems and idle bulls at the height of the season of 1910 disclosed the following:

COUNT OF HAREMS AND IDLE BULLS ON ST. PAUL ISLAND, 1910.

Date.	Rookery.	Harems.	Idle bulls.	Quitters.	Water bulls.
July 12	Lagoon.....	9	3		
12	Tolstoi Cliffs.....	29	5	1	
12	Tolstoi.....	77	7	1	5
12	Zapadni Reef.....	7		3	
12	Little Zapadni.....	54	10	4	4
13	Kitovi.....	53	7	4	4
13	Amphitheater.....	9	2	1	
13	Lukanin.....	41	5	6	5
13	Ardignen.....	11	1		
13	Gorbatch Cliffs.....	2		2	
13-15	Gorbatch.....	110	12	15	
13-15	Polavina.....	50	5	2	10
13-15	Polavina Cliffs.....	20	5	5	
13-15	Little Polavina.....	12	2	7	
14	North East Point.....	251	30	17	10
15	Reef.....	206	28	4	13
16	Zapadni.....	118	22	9	4
	Total.....	1,059	144	81	55

The number of harems on Sea Lion Rock, which could not be visited at this season, is placed at 61, the number found last year.

COUNT OF HAREMS AND IDLE BULLS ON ST. GEORGE ISLAND, 1910.

Date.	Rookery.	Harems.	Idle bulls.	Hauling-ground bulls.	Quitters.
July 14	Little East.....	4			
	East Reef.....	22	6		
	East Cliffs.....	37	^a 14		
	North.....	103	21	10	
	Staraya Artel.....	48	17	21	
	Zapadni.....	47	19	16	1
	Total.....		261	77	47

^a Includes hauling-ground bulls.

A summary of the number of bulls on both islands, with a comparison of the number found in 1909, follows:

SUMMARY OF BULLS ON ST. PAUL AND ST. GEORGE ISLANDS, 1910.

	Harems.	Idle bulls.	Quitters.	Hauling-ground bulls.	Water bulls.
St. Paul.....	1,059	144	81		55
St. George.....	261	77	1	47	
Sea Lion Rock.....	^a 61				
Total, 1910.....	1,381	221	82	47	55
Total, 1909.....	1,399	172	139	98	13

^a Estimated.

Compared with 1909 the number of harems on both islands has decreased 18, or 1.3 per cent, an inappreciable decrease when contrasted with that which has occurred annually for years. This decrease in harems can not be laid to a scarcity of bulls, as can easily be proved, but to a lack of enough cows to provide other bulls with harems.

On the other hand the number of idle bulls—that is to say, those mature adult males stationed on rookeries waiting for cows—has been increased from 172 to 221, or a gain of 29 per cent. This is the result of the saving of young males by marking and of further restrictions upon killing, commenced in 1904.

The number of 7-year old males or "quitters," so termed because of their tendency while idle to desert their stations when approached by man, has decreased from 139 to 82; the number of water bulls has increased from 13 to 55, and of the hauling-ground bulls there has been a decrease from 98 to 47. As these latter classes are more or less unstable and as some of each class could have been in the water at the time these counts were made, it is not attempted to ascribe specific reasons for the fluctuations in them. The fact is demonstrated, however, that young bulls are present in fair numbers. The further fact that 13 per cent of the stationed bulls, excluding quitters,

are idle, indicates conclusively that the herd of breeding bulls is properly safeguarded from too close killing by existing regulations.

COUNTS OF PUPS.

Because of the presence of Japanese schooners in numbers close to the islands, counts of pups on St. Paul Island were limited to Kitovi rookery, including Amphitheater. On St. George Island, for the same reason, pups were not counted except on Little East rookery, which now embraces only a few seals. The St. Paul counts follow:

COUNTS OF PUPS ON ST. PAUL ISLAND, 1910.

	Live pups.	Dead pups.	Total pups.	Harems.	Average harem.
Kitovi.....	1,717	57	1,774	53	33.4
Amphitheater.....	187	5	192	9	21.3
Total, 1910.....	1,904	62	1,966	62	31.7
Total, 1909.....	1,915	64	1,979	58	34.1

From the comparisons which the foregoing data afford, it would appear that the breeding cows on this rookery have not decreased but have remained virtually stationery as regards numbers during this period. The harems thereon, however, are more numerous, thus giving fewer cows to each bull, or, technically speaking, lowering the average harem on this space from 34.1 in 1909 to 31.7 in 1910.

On St. George the count of pups on Little East, which, as stated, was the only count of pups made on that island, disclosed 75 pups in 4 harems, or an average of 18.7 cows per harem. The great decrease in this rookery (Little East) may be appreciated when it is noted that in 1897 the seal census made by the Jordan Commission gave to this rookery 46 harems and 1,190 cows. The number found there in 1910 represents a diminution in thirteen years on this small rookery alone of 42 harems and 1,115 cows.

NUMBER OF BREEDING COWS.

As it is highly impracticable to count the pups on all the rookeries, it has been customary to arrive at the whole number of breeding cows by estimation based upon an actual count of the whole number of harems on the islands and the average number of cows found to be in each of the harems of one rookery which is accepted as typical of all.

As the number of harems on all islands has been ascertained to be 1,381 and the average harem, as demonstrated by the count of Kitovi, to be 31.7, the whole number of breeding cows in 1910 would be 43,777. As 45,786 of such cows were shown by this method to

be present in 1909, the decrease between the years, 2,009, represents a loss of 4.3 per cent.

This for all practical purposes, is a fairly accurate measure of the number of breeding cows, which constitute the most important factor in the herd. While merely an estimate, the number is close enough to actual conditions to be approximately correct. A loss of only 4.3 per cent in the breeding cows from the pelagic sealing which has been practiced with such assiduity during 1910 would seem too small. However, the statistics of the seal herd for the last few years demonstrate that the rate of decrease during this period has not been large, and it is not out of the way to believe that it was small in 1910.

CENSUS OF ENTIRE SEAL HERD.

Beyond the breeding cows and pups, estimates of which contain much of accuracy, an estimate of the whole herd is very difficult to make, and is unsatisfactory in that it treats of elements which are not susceptible of ascertainment and must be approximated. There are also very few means of testing its accuracy at this or a future time. The methods used are, however, the best that can be devised and tend in the direction of accuracy rather than the opposite.

ESTIMATE OF HALF BULLS.

The record of rejections of seals from drives during the summer season of 1910 shows that 1,168 young males too large to be killed were released from the killing fields. It has been established that not by any means all of this class of animals haul in places where they can be enumerated and that the number of those actually turned away should be doubled at least to arrive at the whole number in existence.

By doubling the number found, 1,168, we would have 2,336 half bulls, from which we may look for recruits to the breeding bulls.

ESTIMATE OF 2-YEAR-OLDS.

In 1908 it was computed that 53,884 pups were born. Being equally divided as to sex, one half, or 26,942, were males and an equal number females.

In 1909, if we allow the diminution of 50 per cent for mortality at sea, which has been taken heretofore to occur among the pups during their first migration, one-half of these would return in 1909 as yearlings. There should have been then in 1909 by this method of computation 13,471 yearling males and an equal number of females. These, with a loss of something like 10 per cent, would return in 1910 as 2-year-olds to the number of approximately 12,124 of each sex.

We should have in 1910, therefore, by this computation, over 12,000 virgin or 2-year-old cows and an equal number of males.

From the latter, however, at least 7,500 were killed during the last summer, leaving approximately 4,500 2-year-old males in existence at the close of the season. The above computation would indicate that 12,124 2-year-old cows and 4,500 2-year-old males were present at the end of the killing season of 1910.

NUMBER OF YEARLINGS.

In 1909 it was estimated that 45,764 pups were born, half of which were males and half females. By applying a 50 per cent death rate during their initial migration we should have in 1910 11,441 yearling males and the same number of yearling females.

NUMBER OF 3-YEAR-OLDS.

Nine hundred and fifteen 3-year-olds were marked during the summer and released as breeders. An uncertain number in addition was not driven at all and still survive. It would be a moderate allowance to estimate the number of 3-year-olds remaining in the herd at 1,200.

SUMMARY OF SEAL LIFE IN 1910.

From the foregoing computations an approximate census of seal life present on the islands at the close of the sealing season of 1910 would be as follows:

Bulls, active with harems.....	1,381
Bulls, idle, and quitters.....	303
Half bulls.....	2,336
3-year-old bachelors.....	1,200
2-year-old bachelors.....	4,500
Yearling bachelors.....	11,441
Male pups.....	21,888
Breeding cows.....	43,777
2-year-old (virgin) cows.....	12,124
Yearling females.....	11,441
Female pups.....	21,888
Total.....	132,279

The foregoing "census," if we except the bulls with harems, and those idle, is nothing more than an estimate based upon such enumerations as could be made that were of value in determining the number of seals. While it shows over 2,000 seals less than a similar computation in 1909, it nevertheless exhibits apparent increases in certain classes of seals over the preceding census spoken of. For example, the 2-year-old bachelors estimated to be present in 1910 exceed in numbers by over 2,000 those stated to be in existence at the close of the season of 1909. The 2-year-old cows estimated in 1910 are 2,000 more than were assigned for the previous year.

This is the result solely of the method of estimation adopted alike for both years. Both are based upon the number of cows born two years previously. In 1907, 50,825 pups were estimated to have been born, and 10,165 of these were computed to have survived as 2-year-old males in 1909. On the other hand, in 1908, the same method of estimation would indicate that 53,884 pups were born in that year—3,000 more than in 1907—and that of these the number surviving as 2-year-olds in 1910 was 12,124.

It is believed that it is not the intention of anyone to claim that an increase in seal life has occurred at any time within the past few years in the face of the large catches of seals in the water, consisting mainly of breeding females. It is believed, on the other hand, that a marked decrease has occurred, a belief justified when the contracted space occupied by the breeding seals is viewed. But the measure of this decrease is ascertainable solely by estimation, the same methods being used from year to year. When using only a few seals in establishing a basis for computing the whole number, it is not difficult to realize that a few chance harems more or less on the space counted would have the effect of greatly increasing or decreasing the whole number computed to be in the herd. It would be easy to revise these calculations by adding to or subtracting from the estimated number to make it conform with one's idea of what number should or should not be found. But the idea one may have might be more incorrect than the result of the computation, so that in a revision it would not be possible to determine whether in increasing or decreasing the result one were moving in the direction of accuracy or away from it. It is much better to announce the number each year as it may appear from calculations made similarly, and to explain any apparent incongruity by the statement that the whole is an estimate and nothing else.

The result of the killing of 1910 has demonstrated that the number of 2-year-old bachelors estimated as remaining in the herd at the close of the season of 1909 was entirely too small. In the census of 1909 only 2,165 2-year-old bachelors were allowed. These of course would be 3-year-olds in 1910. As a matter of fact, the skins of 1910 when classified in London showed that perhaps 5,000 of the catch were 3-year-olds. In view of this fact it is believed that, in estimating the number of these as well as other immature seals, a smaller death rate should be allowed than hitherto.

PUP-RAISING EXPERIMENTS.

In accordance with the Bureau's instructions, attempts were made on both St. Paul and St. George Islands to feed starving pups and save their lives. On St. Paul Island the efforts were unsuccessful, but the St. George experiments yielded most interesting results.

ST. PAUL EXPERIMENTS.

Perhaps a dozen or more starving pups were gathered off the various rookeries and brought to the village. An inclosure was built at the end of the village pond and the pups were placed in this.

A bottle with an ordinary rubber nipple was used in a first attempt to induce the little animals to nurse. This method failing, however, milk was poured down the pups' throats from the bottle. But this, besides being difficult and tedious, was uncertain and wasteful, as most of the milk was ejected by the pups before being swallowed. To feed a dozen or more pups with a bottle, moreover, occupied the services of half a dozen men for nearly half a day. Afterwards a tube attached to a funnel was passed into the stomach of each pup and the feeding was accomplished by this means.

Owing to lack of proper material the inclosure in which the pups were placed could not be made tight enough to retain them. Some of the pups escaped to the sea; the others died. Feeding with solid food was not attempted.

Upon the departure of the *Bear* on her last trip from the islands, 10 healthy pups upon which no feeding experiments had been attempted were taken from St. Paul rookeries and placed aboard that vessel to be shipped to Seattle for the use of the Bureau. All of these arrived safely, having been schooled on the voyage to eat solid food.

ST. GEORGE EXPERIMENTS.

Fifteen starving pups were gathered on St. George Island at various times and different methods were tried to save their lives.

These starvelings readily ate all the small live fish that could be obtained and such other larger fish as the weather would permit the natives to capture offshore. In addition the pups ate salted salmon after it had been freshened in water. Had enough live or fresh dead fish been obtainable it is believed that at least some of the pups that were fed artificially could have been saved.

On September 10, 1910, four starving pups were secured and their frenums cut. All were fed by injections of milk into the stomach. One died that night from congestion of the lungs, probably because of the introduction into the pulmonary tract of milk while feeding. Upon autopsy of this animal, a piece of coal as large as a walnut was found lodged in the pylorus. Two of the others escaped the first night.

A corral, having a tank 4 feet by 8 feet and 1 foot deep, was then built and two more pups in addition to the one now remaining were placed in it on September 15. Into this tank filled with water were placed a number of small fish caught among the rocks (probably *Neoliparis*). The pups ate all of these at once and some sculpin cut

into small pieces. After this several attempts were made to provide sufficient fresh fish to feed the pups, but owing to rough weather only several days' supply could be obtained. After this salt salmon freshened in water was offered to the pups and eaten. When this latter was finally refused, milk and mutton broth were fed to sickly pups.

All but one of these pups, 15 in all, died on the island, and that one, after being placed aboard the *Bear*, died before reaching Seattle.

These experiments are of value, however, as demonstrating that by September 15 these pups have advanced to such a stage that they can eat and digest solid food even though they continue to nurse during October and November. The results also show, however, that on the seal islands these experiments can not be carried on with hope of success because fresh fish can not be obtained with regularity in sufficient quantity. Had these pups been taken to Unalaska, where small fish can be readily obtained, it is believed that much better results would have followed.

Of the 14 that died on St. George Island, the autopsies in 2 cases disclosed occlusion of the pylorus by stones taken through the mouth. The death of at least 1 of the pups was due to this condition.

PELAGIC SEALING.

During the season of 1910, 25 Japanese sealing schooners were boarded by revenue-cutter vessels on patrol in Bering Sea. Of these, 2 were seized by the cutters, 1 for a violation of the alien fishing laws and another for a violation of the customs law (section 2773, Revised Statutes). As a rule pelagic sealing vessels kept outside the 3-mile limit, and, so far as known, none of the men composing the crews landed upon the islands for the purpose of killing seals.

Eleven Japanese in 3 small boats landed on St. Paul Island on July 30 and 31. It was stated by them that they had been lost from their schooners and came to the islands as a place of refuge. They were quartered on the islands until August 8, when they were placed aboard the *Manning* and taken to Unalaska with 4 native witnesses, charged with having landed upon the islands without permission, in violation of the act of April 21, 1910.

Upon trial before the United States commissioner at Unalaska they were found guilty and each sentenced to a week's imprisonment. After serving this sentence they were placed aboard a Japanese sealing schooner with their boats, guns, and other property and sent home.

Unofficial reports indicate that 5 Canadian sealing vessels took seals last year in Bering Sea. Their catch from both the Pribilof

and Asiatic herds aggregated 3,775 skins. The total pelagic catch from the Pribilof herd, as shown by London trade sales, was in the neighborhood of 15,000 skins.

WRECK OF REVENUE CUTTER PERRY.

On the early morning of July 26, 1910, the revenue cutter *Perry* went ashore on Rocky Point Reef, St. Paul Island, in a thick fog. Shortly afterwards, by the action of the swell, her bottom was punctured on the rocks upon which she lay, and all efforts to get her off were futile. Such movable property (guns, stores, boats, etc.) as could be readily transported was brought ashore and stored in an empty warehouse at Rocky Point. The entire crew was quartered at the village for several days and was made as comfortable as circumstances permitted. The teams and native men on the islands were used for several days in rendering assistance. Later the *Perry's* men and stores were taken aboard the other cutters in the fleet and the wreck stripped and abandoned. On August 19 the hull was broken up by a strong southerly gale and scarcely anything was left to mark where she grounded.

FOXES.

The history of foxing on the Pribilof Islands is interesting. What number of fox skins were taken off these islands by the Russians will never be known. Petroff (1883) states that 34,767 were taken from 1842 to 1860, inclusive. From that date to 1867, the fox skins taken from the islands are not segregated from the returns of those taken from general Alaskan sources, which are given by Petroff as 27,731. From 1870 to 1890 fox skins to the number of 4,380 on St. Paul and 20,412 on St. George were taken and shipped by the Alaska Commercial Company. From 1890 to 1910, 2,963 fox skins were taken on St. Paul and 13,641 on St. George.

During the lease of the Alaska Commercial Company (1870-1889), there existed no contract with the Government for the right to purchase these skins, and the only expenditure by the company for the more than 24,000 skins it received was the 50 cents it paid the natives for each skin. The North American Commercial Company during the greater portion of its 20-year lease paid to the natives \$5 for each blue and \$1 for each white fox skin.

Foxes are trapped annually on St. George Island in house traps which do not injure the animal. The catch last year there was 227. On St. Paul Island, where these animals never have been as plentiful as they were on the other island, no trapping has been done since 1903 until last winter (1909-10), when 185 were secured. These were killed in steel traps. For the blues the natives received \$5 apiece; for the whites, \$1. This money was applied to the natives' support.

CONDITIONS AND TRAPPINGS ON ST. GEORGE ISLAND.

On St. George Island, during the winter of 1909-10, the feeding of foxes in the herd during the period from October 20 to June 1 was continued as in former years. Seal carcasses preserved from killings during the summer formed the greater portion of the material fed, together with about 3,000 pounds of salted codfish freshened in sea water.

For some reason, not ascertained exactly, a smaller number of foxes passed through the house and box traps during the winter in question than ever before since feeding the foxes and selective trapping began. Whether this is the result of an actual diminution in the herd, or whether other conditions, such as an abundance of food outside the traps or an instinctive fear of entering the traps, were the cause, can not be stated definitely.

During the winter of 1909-10 only 335 foxes passed through the traps on St. George Island. To show the smallness of this number as compared with former years, a table with the total number of foxes handled in the various years during which selective trapping has been followed is given below:

1898-99.....	842	1904-5.....	766
1899-1900.....	973	1905-6.....	1,061
1900-1901.....	1,335	1906-7.....	882
1901-2.....	1,104	1907-8.....	1,006
1902-3.....	1,011	1908-9.....	798
1903-4.....	1,061	1909-10.....	335

In trapping, the practice is to catch all animals alive, to release as breeders a certain number of pairs of the most vigorous, and to kill those that are not considered the best examples of the species. Those released are marked, so as to be thereafter recognizable, by clipping a ring out of the hair on the tail of the animal, the marks differing for the sexes. Such foxes as escape being trapped, not being marked of course, can be distinguished at sight.

No such number of foxes not marked was seen in the winter mentioned as to lead unquestionably to the conclusion that the herd has not diminished. There are, on the other hand, good grounds for believing that it has diminished. The causes of this probable fact, however, are obscure and conjectural. The very few found dead did not justify the belief that any epidemic had occurred.

A summary of the statistics of trapping on St. George Island during the winter of 1909-10 is appended:

Marked and released:

Blue males.....	51
Blue females.....	57

Killed for pelts:	
Blue males.....	126
Blue females.....	86
White males.....	5
White females.....	4
Skins from animals found dead, etc.....	6
Skins accepted by lessee, blue.....	203
Skins rejected by lessee, blue.....	6
Skins mangey, etc., thrown away.....	9
White fox skins accepted by lessee.....	9
Total number of animals handled.....	335

These pelts, having been taken during the period covered by the contract of the North American Commercial Company, were delivered to it upon payment at the stipulated rate of \$5 for each blue skin and \$1 for each white skin. The money thus derived was used exclusively for the support of natives.

TRAPPING ON ST. PAUL ISLAND.

During the winter of 1909-10, for the first time since 1904, there were considered to be foxes enough on St. Paul to justify trapping, which accordingly was carried on during a period of six days.

On this island, unlike St. George, notwithstanding repeated efforts to secure it, the foxes do not congregate in large groups, permitting systematic feeding and selective trapping. Any trapping therefore on St. Paul must be done with the spring steel trap, in the use of which the native trappers must scatter over the entire island.

In the 6 days of trapping mentioned the St. Paul natives secured on that island 130 blue and 35 white foxes. In addition, a boat load of native men went over to Otter Island, and there secured 19 blues and 1 white. Observations made during the past summer indicate that the fox herd on St. Paul Island has not diminished appreciably as the result of this trapping of the previous winter.

The skins taken on St. Paul and Otter Islands were delivered to the North American Commercial Company and payment made at the same rate as on St. George. This difference between the management of the two islands exists, however, that whereas the earnings on St. George from fox skins are formed into a community fund, on St. Paul each individual trapper is given the use of the money from such fox skins as he has been able to secure.

RECOMMENDATIONS.

KILLING OF BACHELOR SEALS.

The methods used in taking seals during the past season of 1910 were the same as those used by the two lessees in the preceding 40 years' tenure of the sealing right, and the same, in fact, in all

fundamental respects as those pursued by the Russians since 1840. They are the result of years of experience and are the best that can be devised to meet the conditions. No change in them should be made.

The practice of killing bachelor seals for skins as well as for natives' food should not be abandoned unless a cogent reason presents itself. No harm to the seal herd can result from the killing of surplus males. No benefit to the herd could accrue from the maturing of males unnecessary for purposes of reproduction, which, when of adult age, would have no female consorts, but which, by incessant and furious fighting, would destroy or cripple the breeding bulls and themselves as well.

It is true that a test to insure the survival of the fittest should be applied to the male fur seal, as in fact it should to all breeders. It is not true, however, that this test can only be made through trial of combat. With respect to some groups of animals, such as the Pinnipedia, conditions of their natural environment may be so severe as to eliminate weaklings as effectually or even more so, than would fighting amongst themselves, and nature provides an eliminative process in the case of the fur seal entirely apart from the struggling of bulls with each other for supremacy on land. This test begins almost with a seal's birth.

When the baby seal has scarcely learned to swim beyond the borders of the rookery on which it is born, while it is still a suckling and knows not how to seek other food, it is separated from its mother and driven off the land by the rigor of the climate. Weak and unskillful swimmer as the pup is, not only must it withstand the severe winter storms in the northern ocean but in the same unfavorable element pursue and capture its food and elude its natural enemies of the sea. As the result of this struggle with the natural conditions in which it is placed it is estimated that one-half of the pups die during the initial migration. Only the strongest and most wary can survive this trial.

This struggle for existence continues incessantly during the animal's life. From each migration it sends back to the breeding grounds only those animals hardy enough to withstand its severity. That animal leaving the rookeries with any physical imperfection does not return. It dies at sea. Those that do return are the most perfect examples of their class.

With this severe eliminative test occurring as the result of natural environment, to superimpose a violent struggle with his own kind after the animal has reached the breeding ground would be to subject him to further stress entirely unnecessary to prove his ability as a breeder. Having passed successfully through the winter's migration, the animal returns to the rookeries a perfect specimen of its kind. A severe trial by combat could not have the effect of increasing

its breeding efficiency, but on the other hand could only seriously impair if not wholly destroy it. It would be the same if two valuable stallions, each physically perfect, and matched in strength and courage, were allowed to fight with each other until one were killed. The survivor, if one did survive, would be so seriously injured by its opponent as to be rendered incapable of service for the time being, if not permanently.

To breed a large number of surplus male seals merely that they may fight amongst themselves and determine the strongest in combat is useless. By the time the strongest individuals have proved their superiority they have expended so much of their energy in fighting that physically weaker but fresh animals may overpower them and take their cows. Such is the history of the Pribilof rookeries during the time when thousands of idle bulls were present. Instances to substantiate this conclusion have been witnessed many times.

Since physical combat is not required to test the ability of a male fur seal, no reason is known for providing a number of males beyond that necessary to fertilize the females in the herd. Therefore the practice of killing surplus males at the time when their pelts have a considerable commercial value should be continued. Surely no purely sentimental reason should prevail over those of practical weight.

SUPPORT OF NATIVES.

The present system of supporting the natives on the Pribilof Islands should be changed. Under it the native receives enough food, fuel, and clothing to sustain life, but only a portion of the sum necessary for his maintenance comes to him as compensation for labor performed, the remainder being donated as a gratuity through an appropriation of Congress. This latter feature is the most objectionable of all and the one which it is sought to eliminate. Better to explain the situation the following brief summary is given of the manner in which the natives have been supported since they were first transported to these islands.

In 1787, the year following the discovery of St. George Island, the discoverer, Pribilof, brought to the islands a number of native families, principally from Unalaska, and landed them there to serve as laborers in taking skins from the animals with which the islands abounded. Several other adventurers also brought natives to these islands and founded small villages at several points thereon. In 1799, upon the taking over by the Russian-American Company of the administration of the whole of Alaska, the competing traders were sent away from the Pribilofs and the islands passed under the autocratic control of Baranof. A cessation of killing was ordered, and in 1806-7 nearly all the natives were removed to Unalaska.

In 1808 seal killing began again, with accessions of laborers mainly from Unalaska and adjacent villages. On St. Paul Island the natives were drawn together and huddled into one settlement at Halfway Point. About 1825, for convenience in handling cargo, the village was again changed to its present site.

On St. George Island several settlements existed originally, but were consolidated at the present site about 1830-1835.

Under the Russian régime, especially under the management of the Russian American Company, which provided the machinery of government for the territory during the tenure of its privilege, the natives were mere slaves. They had no redress for any injury or insolence which their masters might see fit to inflict upon them. Their habitations were large communal dwellings of earth, half underground, cold, and filthy. Here they lived and died unnoticed and uncared for. They subsisted on fish and the flesh of seals, with the addition of roots and a very little flour.

In 1835, Veniaminof states, the natives worked at whatever was found and whatever they were directed to do. Payment was not established by the day or year, but for each skin taken by them or for what was placed to their credit. They received no specific wage, though they were not all of equal ability, there being usually three or four classes. In these classes the sick and old workmen were counted, although they were only burdens, and therefore received the smallest shares, about 150 rubles, and the other and better classes 220 to 250 rubles a year. Those who were zealous were rewarded by a present of 50 to 100 rubles. The wives of the Aleuts, who worked only at seal killing, received from 25 to 35 rubles. These rubles were scrip currency, made of leather, equal in value to a franc, or about 20 cents.

In 1868, at the time of the purchase of Alaska by the United States, the natives were living in semisubterranean houses built of turf and such pieces of driftwood and whalebone as they were able to secure on the beach. Their food was seal meat and a few articles furnished in meager quantity by the Russian company. They had no fuel except driftwood and blubber, and depended for heat upon crowding together in the sod houses, sleeping upon the dried grasses secured upon the islands.

In 1870 the Alaska Commercial Company took charge of the islands under a lease. It at once built neat frame dwellings for the natives, and paid them 40 cents apiece for each sealskin taken. As 100,000 were taken annually this gave the natives about \$40,000 each year, enough to support them in qualified comfort. While this sum was divided on a communal basis, some natives by thrift and economy were able to save sums amounting to perhaps \$2,500 each. No interference with the expenditure of their earnings was made by the agents of the government.

When, however, after 1890, under the lease of the North American Commercial Company, the take of skins was reduced to a few thousands annually, the natives faced starvation. Their earnings at this time, at the rate of 50 cents for each skin, were entirely insufficient. To relieve this situation, the Government did not increase the wages of the natives for taking skins, but, as the reduction of the catch was due mainly to arbitrary restrictions by the Government, furnished an annual appropriation of \$19,500 to supplement the natives' earnings for their support.

This appropriation, while keeping the natives from starving, made an important change in their fiscal relations. Heretofore the native could expend his earnings as he pleased. After the appropriation, however, the earnings were sequestered by the agents, and the natives had no voice whatever in the expenditure of the money for which they toiled. Each native was allotted articles of necessity to a certain amount each week payable from his wages, and after the latter were expended the appropriation was drawn upon at the same rate until another sealing season intervened.

This practice exists to-day. The natives now receive \$1 for each skin taken, in addition to the annual appropriation of \$19,500. Their total income from taking seals and foxes, with the appropriation, was last year about \$34,000, or somewhat more than \$100 for each person.

The system of distribution of these earnings is one of pure communism. The native men are divided into about four classes, according to ability in taking seals. The members of each class receive a like sum, those in the first class being given more than those in the second, and so on to the fourth class, the lowest, which embraces apprentices. These sums, whatever they may be, are credited to each native and are drawn upon each week by orders on the store issued by the agent to the head of each family, the amount of the order varying with the size of the family. This plan of compensation, while assuring provision for the natives' immediate needs, is highly objectionable when considered from a sociological standpoint, its weakness being that it reduces all to a common level. It prevents that progress that accrues from the cultivation of superior skill or greater self-denial, and makes a virtual almshouse of the Pribilof reservation by dealing with the inhabitants as indigents. It requires willing service of the native, but takes from him his wage and expends it for his benefit without his consent. Incentive to increased individual efficiency is lacking because effort to that end is fruitless in bringing any greater benefit than if it had not been made.

It is reasonable to assume that the Government, while operating on the seal islands for its own profit, at the same time desires to better the condition of the native residents upon whose efforts it must depend

for successful conduct of its business. The first step in that direction is to do away with the appropriation of Congress for their support and to increase the wage earned through the taking of skins to a sum at least equal to the amount necessary for their maintenance. This would at once eliminate the objectionable element of charity in the present system and allow each man to support himself and family from his own earnings. Such a course is in my opinion not only an act of simple justice, the consummation of which would, moreover, involve no additional expense to the Government, but would go far toward increasing the moral tone of the native, by making him more self-reliant and self-respecting. It can be taken without additional legislation, the Secretary of Commerce and Labor now having the power under existing law to fix the natives' compensation for taking skins.

SCHEME OF COMPENSATION OF NATIVES.

The scheme of compensation embodied in the foregoing recommendations may be summarized as follows:

1. The appropriation for natives' support to be discontinued.
2. For natives' labor an allotment should be made of, say, \$3 for each sealskin.
3. The moneys thus derived should be formed into a general fund, which should be prorated among all the natives of both islands.
4. This fund, by agreement with the natives, to be used for their support at the rate of a certain weekly amount based upon the number of mouths in each family.
5. The balance or remainder of each native's account at the close of each year to be paid to the native in cash.

It must be understood that the native is restricted by his work to the seal islands and can not go forth to pursue any other vocation, be it more or less profitable. It is not fair to this laborer to deny him all progress in the world and to confine him in his necessarily restricted sphere to such compensation only as permits the bare necessities of life to him and his family. Whatever a corporation having a lease of the sealing privilege may have done, the United States Government ought not to put its laborer into the condition of constant and continuous vassalage with all progress denied him.

NATIVES ON THE ALEUTIAN ARCHIPELAGO.

The Aleut race is not found on the mainland, but inhabits the Aleutian Archipelago and several of the islands along the coast of the Alaskan Peninsula. It was never numerous and now embraces probably fewer than 1,000 souls, whose numbers are decreasing rapidly from disease and insufficient food. Some action should be taken to ameliorate their condition.

When discovered by the Russians in the eighteenth century, these Aleuts were a hardy race of fishermen and aquatic hunters. In their tiny bidarkis or skin boats they made long journeys and in them successfully weathered storms that would have sent the European rowboat to the bottom. They subsisted upon fish and the flesh of such warm-blooded animals as they could capture.

Being a tractable race, except when goaded to desperation, they were at once made use of by the Russians as hunters of the sea otter, which was the fur the white men most eagerly sought. Whole fleets of bidarkis with hundreds of native hunters would be transported hundreds of miles from their homes, and thence with a little food supplied them were put to sea to buffet with the storms of the northern ocean which withal were not so greatly feared by the natives as were their white masters. Thousands of them never returned.

Aleuts in numbers were taken to Sitka by the Russians as hunters and laborers, and kept there until they died. Entire fleets of bidarki hunters were loaned by the Russian company to foreign vessels to hunt sea otter, the profits of the venture being shared equally by the vessel and the company. The ship was required to pay the Russian company about 200 Mexican dollars for every Aleut lost at sea or killed by coast Indians. In 1805, 20 bidarkis were fitted out at Kodiak and with a colony of natives were taken to San Quinten bay in Lower California, where they were required to hunt for fur seals. This colony struggled on until 1841, when it was abandoned.

In the draft of the terms upon which the Russian-American Company should receive an extension of its charter, after its expiration in 1861, or thereabouts, the following paragraph is found:

10. The Aleuts and other peaceful natives within the colonies are relieved from compulsory labor on behalf of the Russian-American Company. They shall be allowed to settle in localities which they may find convenient, and shall be free to absent themselves from the places of their residence, subject only to such rules of police as may be established by the board of administration of the colonies.

This clause in the proposed charter was inserted to cure abuses in respect to the treatment of natives reported by Golovnin and the creole Kashevarof. In short, the Government would renew the charter only under such terms as the company would not accept.

When the Russian-American Company acquired control of Alaska the Aleuts were paid nothing for sea-otter skins, but in lieu of compensation received subsistence and "exemption from imperial taxes and dues." When this practice was forbidden by the Emperor Alexander I and the company instructed to pay the natives for every skin deposited by them with the company the natives received for every sea otter 10 rubles in leather scrip, the equivalent of \$2, but each hunter was required to furnish his own subsistence. The company sold the sea-otter skins for at least \$100 each.

Upon the occupation of the territory by Americans, the native from a condition of abject misery and want was plunged into a state of affluence of which he knew not how to take advantage. Rival trading companies established stations along the coast where sea otters abounded, and bid eagerly for the furs brought in by the native hunters. But while paying him liberally for the skins, the traders adroitly exposed for sale in the stores articles of sheer luxury to tempt the native's cupidity and encourage him to expend the money received for his skins. During the seventies and eighties the Aleut sea-otter hunter clothed his women in satins and silks of the gaudiest colors; his hut contained a brussels carpet and a parlor organ; his church received large donations; in short, a great deal of his earnings was expended at once for luxuries and he was forced to hypothecate his next year's catch of skins to obtain supplies to support his family during the winter.

With the commercial disappearance of the sea otter, however, the native again relapsed into a condition of penury bordering on starvation. Whereas in the days of plenty he lived on tinned meats and luxuries from the trader's store, now to sustain life he was driven again to fish and to hunt. Having contracted the vice of drunkenness, even in his poverty he would barter his skins for rum, or for sugar and flour with which to make the Russian strong beer. Disease sapped his vitality and decimated his villages.

Such practically is the condition to-day of the native on the Aleutian chain.— While formerly he had to subsist upon what he could wrest from nature, he was then as free from the vices of civilization as he is now of its saving benefits. His contact with the white race has encouraged appetites of which the native was previously ignorant and has taken away his self-reliance and ability to cope with his surroundings. In his state of poverty, the furs he is still able to gather are the object of desire of small traders, who visit his settlements annually and exchange trade goods for furs. The native has no resource but to part with his furs at such prices as the trader may wish to give.

Unless the Government takes active measures this interesting race of people will become extinct. And since the Government is trying to save species of the lower animals which are threatened with that calamity, it would seem proper that similar attention should be paid to a race of human beings which is rapidly disappearing. A simple and yet it is believed an effective plan to accomplish this end is offered and earnestly recommended to the attention of the Department:

1. The entire Archipelago to be made a special reservation. This can be accomplished without difficulty or friction. There are no vested rights in the entire range of islands, so far as known, except

at Dutch Harbor, a small portion of which has been surveyed and patented. The property of the Alaska Commercial Company at Unalaska is built on a Government reservation on which it has only squatter's rights. For its buildings it should receive compensation.

The islands in this chain are devoid of timber. Coal or minerals have appeared only in too small quantities to justify exploiting. Agriculture on any scale to support life is impossible. Grazing is impracticable. There are no good harbors except at Dutch Harbor and Unalaska. Fish are plentiful but the streams are so small that commercial fishing will not pay. In short, there exists no good reason why these islands should not be set apart for the use of those aboriginal inhabitants claiming them as their native land.

2. Trading by private persons or corporations to be prohibited.

3. The Government to maintain a station at each principal settlement, of which there are not over five. Each station to contain a store and a school, with a storekeeper and school-teacher, the whole to be under the supervision of a general agent.

4. The storekeepers should buy the natives' peltries and such other articles as they may have for sale, including baskets, at a fair price; the native should be encouraged to self-support and thrift.

5. Small fishing stations could be maintained, the product of which could be marketed for natives' account.

This plan can be worked out and operated with little trouble and expense. Without some provision of this nature the Aleuts on the Archipelago will be wiped out by disease and lack of food. With the Government willingly expending thousands of dollars to prevent extermination of the lower animals, surely no justification is needed for expenditure to prevent the extinction of a race of men who were hardy and self-reliant until brought into contact with European races.

MANUAL TRAINING FOR NATIVES.

During the Russian occupation certain native youths exhibiting special aptitude were trained in the useful arts, such as carpentry, boat building, iron and copper working, etc.

But those natives so educated in Russian times have nearly all died, and the new generation can not build its own houses or boats. No training of this character, although greatly needed, has been provided by our Government.

Some arrangement should be made to teach the Aleuts how to work at other employments than their natural one of hunting. A teacher of the useful trades should be provided on each of the Pribilof Islands. A small school could also be established at Unalaska, and the young men from the entire archipelago sent there for a course of instruction. I recommend this to careful consideration.

FIRE PROTECTION FOR PRIBILOF ISLANDS.

The villages of St. Paul and St. George are entirely without fire protection, and with the high winds that prevail are fortunate indeed in never having had a disastrous conflagration. Aside from the money loss entailed, such a contingency, should it occur in winter and destroy the food supply in the warehouses, would probably result in the starvation of the inhabitants. Native dwellings have been ignited by overturned kerosene lamps and in one case a whole native family while asleep was asphyxiated by fire in the interior of their house. In every case so far, however, the blaze has been discovered and extinguished before it could take serious hold upon the framework of the building.

I strongly urge the provision of adequate fire protection for both islands. The isolation of the locality demands that some means be supplied for the prevention of conflagration, which there would be a catastrophe. The investment of a small amount for this purpose would be sufficient to provide protection for years, and would be the cheapest fire insurance that could be obtained. Should these buildings burn, the business not only would be seriously interfered with, and the native and white inhabitants threatened with starvation, but the Government would lose the amount of its investment and be obliged to spend twice as much to replace the plant as was paid for it.

As to means, chemical apparatus could be used in summertime, but would be of little avail in winter because of the likelihood of freezing while not in use.

In winter, running water under pressure would be the only resort. Running water is not available at present, but could be supplied by any of the following means:

On St. Paul.—1. Sea water could be pumped through a small stand-pipe by a gasoline engine and distributed through mains in the village.

2. Fresh water from a lake a mile away could be piped to the village by pumping, and held in a large reservoir of sufficient capacity to furnish fresh water not only for fire protection but for natives' use.

3. Water from wells a half mile from the village could be pumped and used in the same manner as in suggestion 2.

On St. George.—1. Salt water could be pumped as in the preceding suggestion 1.


2. The water system already in use there, whereby water is brought by gravity and siphoning from a lake to the village, could be adapted to furnish a stream that would reach over any native dwelling and probably any larger warehouse or dwelling.

THE SALMON FISHERIES OF THE PACIFIC COAST

By JOHN N. COBB

Assistant Agent at the Salmon Fisheries of Alaska

Bureau of Fisheries Document No. 751



CONTENTS.

	Page.
Introduction	5
I. The species of salmon and the runs	6
Chinook, quinnat, or king salmon	6
Sockeye, blueback, or red salmon	8
Silver or coho salmon	9
Humpback or pink salmon	10
Dog or chum salmon	11
Steelhead trout	11
II. Fishing grounds and history of the fisheries	12
Washington	12
Columbia River	13
Oregon	15
California	18
Alaska	21
III. Apparatus and methods of the fisheries	23
Gill nets	23
Haul seines	24
Diver nets	25
Dip nets	25
Squaw nets	25
Purse seines	26
Traps, or pound nets	27
Indian traps	29
Wheels	30
Reef nets	32
Trolling	32
Bow and arrow	34
Spear and gaff	34
IV. Fishermen and other employees	35
V. Fishery regulations	37
Controversial forms of apparatus	37
Laws and their enforcement	40
VI. Methods of preparing salmon	46
Canning	46
Early days of the industry	46
Handling the salmon	49
Dressing	50
Cutting	50
Salting	51
Filling the cans	51
Washing the cans	52
Capping	53
Soldering	53
Testing	54
Cooking	54

VI. Methods of preparing salmon—Continued.	
Canning—Continued.	Page.
Sanitary, or solderless, cans	55
Repairing cans	56
Lacquering	57
Labeling	58
Brands	59
Boxing or casing	60
Can making	60
Mild curing	60
Pickling	62
Dry salting	62
Smoking	63
Freezing	64
Miscellaneous products	65
Oil and fertilizer	66
VII. Statistics of the Pacific salmon industry for 1909.	68
Persons employed	68
Investment	68
Products	70
Products canned	70
Miscellaneous products	73
Washington	74
Statistics by counties	75
Statistics by waters	83
Columbia River	90
Oregon	92
Statistics by counties	92
Statistics by waters	97
California	105
Statistics by counties	108
Statistics by waters	112
Alaska	115
British Columbia	119
VIII. Statistical data for other years.	121
Canning industry of Pacific coast of North America from 1864 to 1910.	121
Canning industry shown by species and waters	123
Pickling industry	136
Mild-curing industry	137
IX. Trade with outlying possessions.	138
X. Foreign trade in salmon.	140
Exports of canned salmon	140
Exports of fresh and cured salmon	149
Imports of fresh salmon	152
Imports of cured salmon	153
XI. Salmon culture.	154
California	154
Oregon	159
Columbia River and tributaries	164
Washington	168
British Columbia	172
Alaska	174

THE SALMON FISHERIES OF THE PACIFIC COAST

By JOHN N. COBB,

Assistant Agent at the Salmon Fisheries of Alaska.

INTRODUCTION.

The most valuable commercial fisheries in the world, excepting only the oyster and herring fisheries, are those supported by the salmons. Of these the most important by far are the salmon fisheries of the Pacific coast of North America, where California, Oregon, Washington, and Alaska, including also British Columbia, possess industries representing millions of dollars of investment and millions of output annually. No published reports contain data for the entire coast, or have pertained to the same year for both Alaska and the States. In the following pages, containing the returns from a canvass occupying several months, the data are complete for the United States coast and Alaska for the year 1909, and to make the report more comprehensive, historical and geographical aspects of the subject, as well as methods of the fisheries and allied industries, are discussed at some length. Figures for British Columbia have been included also, so far as possible, the official reports of the Dominion of Canada and of the Province itself having been drawn upon for this purpose. The statistics for Alaska are taken from the already printed (1909) report of Mr. Millard C. Marsh and the present writer.^a

^a The fisheries of Alaska in 1909. By M. C. Marsh and J. N. Cobb, agents at the salmon fisheries of Alaska. Bureau of Fisheries Document No. 730. 1910.

I. THE SPECIES OF SALMON AND THE RUNS.

The Pacific coast salmon are all included in the genus *Oncorhynchus*. With them the fishermen incorrectly class the steelhead trout, which really belongs to the closely related genus *Salmo*.

As long ago as 1731 the species of *Oncorhynchus* were first made known by Steller, who, almost simultaneously with Krascheninikov, another early investigator, distinguished them with perfect accuracy under their Russian vernacular names. In 1792 Walbaum adopted these vernacular names in a scientific nomenclature for these fishes.

Five species of salmon (*Oncorhynchus*) are found in the waters of the north Pacific, ranging northward from Monterey Bay on the American coast and Japan on the Asiatic, the extreme northern distribution of certain of the species having not yet been accurately determined. The five species are: (1) *Oncorhynchus tshawytscha*, quinnat, tyee, chinook, spring, or king salmon; (2) *Oncorhynchus nerka*, blueback, red, sukkegh, or sockeye salmon; (3) *Oncorhynchus kisutch*, silver, coho, or white salmon; (4) *Oncorhynchus keta*, dog or chum salmon; and (5) *Oncorhynchus gorbusha*, humpback or pink salmon.

CHINOOK, QUINNAT, OR KING SALMON.

The largest, best known, and most valuable of these is the chinook or king salmon (*O. tshawytscha*). It is found throughout the region from the Ventura River, Cal., to Norton Sound, Alaska, and on the Asiatic coast as far south as northern China. As knowledge extends, it will probably be recorded in the Arctic.

In the spring the body is silvery, the back, dorsal fin, and caudal fin having more or less of round black spots, and the sides of the head having a peculiar tin-colored metallic luster. In the fall the color is, in some places, black or dirty red. The fish has an average weight of about 22 pounds, but individuals weighing 70 to over 100 pounds are occasionally taken. One was caught near Klawak, Alaska, in 1909, which weighed 101 pounds without the head. The Yukon River is supposed to produce the finest examples, although this supposition is not based on very reliable observations. The southeast Alaska fish average as high as 23 pounds in certain seasons, followed by an average of about 22 pounds in the Columbia River, and about 16 pounds in the Sacramento.

In most places the flesh is of a deep salmon red, but in certain places, notably southeast Alaska, Bristol Bay, Puget Sound, and British Columbia, many of the fish, the proportion being sometimes as much as one-third of the catch, have white flesh. A few examples have been taken with one side of the body red and the other white, while some are found with mottled flesh. No reasonable explanation of this phenomenon has yet been given.

In its southern range the quinnat strikes in at Monterey Bay in sufficient numbers to justify commercial fishing about the middle of April, where it is seen feeding upon the inshore moving schools of herring and sardines, continuing until in August. There are two runs of spawning fish in the Sacramento, the first or "spring run" beginning in April and continuing throughout May and June, these fish spawning mainly in the cold tributaries of the Sacramento, such as the McCloud and Fall Rivers. The second or "fall run" occurs in August, September, and October, and these fish spawn in the riffles in the main river between Tehama and Redding, also entering the tributaries in that vicinity. The two runs merge into each other. It is also claimed that there is a third run which comes in December.

In former years the San Joaquin and the American and Feather Rivers of the Sacramento system had large runs of salmon, but excessive fishing and the operation of various mining and irrigation projects have practically depleted them.

The Eel and Mad Rivers of northern California have only a late or fall run, while the Klamath River has both a spring and a fall run, and Smith River has a spring run alone. Rogue River in Oregon has both a spring and a fall run, and the Umpqua and several other coast streams of Oregon have small early runs.

The Columbia River has three runs, the first entering during January, February, and March, and spawning mainly in the Clackamas and neighboring streams. The second, which is the best run, enters during May, June, and part of July, spawning mainly in the headwaters. The third run occurs during late July, August, September, and part of October, and spawns in the tributaries of the lower Columbia.

In Puget Sound chinook salmon are found throughout the year, although it is only during the spawning season that they are very abundant. In the Fraser River, a tributary of the Sound, the run occurs from March to August.

In the Skeena River, British Columbia, the run occurs from May to July, the same being approximately true of the Nass also.

In southeast Alaska they are found all months of the year. From March to the middle of June they are abundant and feeding in the numerous straits and sounds; in May and June the spawning fish enter the Unuk, Stikine, Taku, Chilkat, Alsek, and Copper Rivers

in large numbers, and in a few smaller streams in lesser abundance. In August, September, and October they are again to be found in large numbers feeding in the bays and sounds, while during the winter months a few have been taken on trawls set for halibut, showing that they are living in the lower depths at this time.

In Cook Inlet the run occurs during May and June and is composed wholly of red-meated fish; in the rivers of Bristol Bay the run comes in May and June, and the same is true of the Togiak, Kusko-kwim, and Yukon Rivers, although fish may be seen in the upper courses of the Yukon in July, the lateness here being due to the immense distance the fish have to cover.

On the Asiatic side the chinook is found in some of the rivers of Siberia.

SOCKEYE, BLUEBACK, OR RED SALMON.

The sockeye or blueback salmon (*O. nerka*), which forms the greatest part of the canned salmon of the world, when it first comes in from the sea is a clear bright blue above in color, silvery below. Soon after entering the river for the purpose of spawning the color of the head changes to a rich olive, the back and sides to crimson and finally to a dark blood red, and the belly to a dirty white. The maximum weight is about 12 pounds, and length 3 feet, with the average weight about 5 pounds, varying greatly, however, in different localities. Observations of Chamberlain^a in Alaska show that the average weight of a number of sockeyes taken from Yes Bay was 8.294 pounds, while the average weight of a number from Tamgas was only 3.934 pounds. Evermann and Goldsborough^b report as a result of the weighings of 1,390 red salmon, taken from as many different places in Alaska as possible, an average weight for the males of 7.43 pounds; for the females, 5.78 pounds; or an average weight for both sexes of 6.57 pounds. A run of small, or dwarf, males accompanies certain of the main runs, these being especially noticeable in the Chignik lagoon, Alaska, run. This species usually enters streams with accessible lakes in their courses.

A few specimens of the sockeye have been taken as far south as the Sacramento River. In Humboldt County, Cal., small runs are said to occur in Mad and Eel Rivers. Only an occasional specimen appears in the coastal streams of Oregon. The Columbia is the most southern river in which this species is known to run in any numbers, entering the river with the spring run of chinooks. From here south the species is called blueback exclusively. A considerable run enters the Quinnaault River, Wash., and there is also a small run in Ozette Lake, just south of Cape Flattery.

^a Some observations on salmon and trout in Alaska. By F. M. Chamberlain, naturalist, U. S. Fisheries Steamer Albatross. U. S. Bureau of Fisheries Document no. 627, p. 80.

^b The fishes of Alaska. By B. W. Evermann and E. L. Goldsborough. Bulletin Bureau of Fisheries, vol. xxvi, p. 257.

In the Puget Sound region, where it is known as the sockeye, this species ascends only the Skagit River in commercial numbers, although a small run appears in the Lake Washington system of lakes and, possibly, in the Snohomish, Stillaguamish, and Nooksack Rivers.

The greatest of all the sockeye streams is the Fraser River, British Columbia, and this stream has been famous from very early days for its enormous runs of this species, a peculiar feature of which is that there is a marked quadrennial periodicity in the run. The maximum run occurs the year following leap year, the minimum on the year following that. The greater part of the catch of the Puget Sound fishermen is made from this run as it is passing through Washington waters on its way to the Fraser. The fish strike in during July and August on the southwest coast of Vancouver Island, apparently coming from the open sea to the northwest. They pass the Straits of Juan de Fuca, Rosario, and Georgia, spending considerable time in the passage and about the mouth of the river. Small numbers run as early as May and as late as October, but the main body enters about the first week in August.

The sockeye occurs in most of the coastal streams of British Columbia, and is usually the most abundant species. The principal streams frequented are the Skeena, Rivers Inlet, Nass, Lowe Inlet, Dean Channel, Namu Harbor, Bella Coola, Smith Inlet, Alert Bay, and Alberni Canal.

In Alaska, where this fish is generally known as the red salmon, it is abundant and runs in great numbers in all suitable streams, of which, in southeast Alaska, the following are the most important: Boca de Quadra, Naha, Yes Bay, Thorne Bay, Karta Bay, Nowiskay, Peter Johnson, Hessa, Hetta, Hunter Bay, Klawak, Redfish Bay, Stikine, Taku, Chilkoot, Chilkat, Alsek, Seetuck, Ankow, etc.; in central Alaska, Copper, Knik, Kenai, Sushitna, Afognak, Karluk, Alitak, Chignik; in the Bristol Bay region, the Ugashik, Ugaguk, Naknek, Kvichak, Nushagak, and Wood. It is also supposed to occur in the Togiak, Kuskokwim, and Yukon Rivers, which debouch into Bering Sea, and probably occurs in the Arctic streams of Alaska. The run in Alaska begins usually in June and extends usually to the middle of August. It begins earlier in Prince William Sound, and sometimes extends into September in southeast Alaska.

On the Asiatic side the species is known to occur at Bering Island and in all suitable streams south to Japan, where it is found landlocked in Lake Akan, in northern Hokkaido.

SILVER OR COHO SALMON.

The silver or coho salmon (*O. kisutch*) is silvery in spring, greenish on the upper parts, where there are a few faint black spots. In

the fall the males are mostly of a dirty red. The flesh in this species is of excellent flavor, but paler in color than the red salmon, and hence less valued for canning purposes.

This species has a maximum weight of about 30 pounds, with a general average of about 6 pounds.

The silver salmon is found as far south as Monterey Bay, where it appears during the month of July and is taken by the trollers. From Eel River, in California, north, it is found in most of the coastal streams. It usually appears in July and runs as late as November, the time of appearance and disappearance varying somewhat in different sections. Owing to its late appearance comparatively few, and they usually in the early part of the season, are packed by the canneries, most of which shut down in July and August. This fish also tarries but a short time about the mouth of the stream it is to enter, and is wary of nets, which makes it rather unprofitable to fish for the latter part of the season when it is running alone.

On the Asiatic side the coho ranges down the coast to Japan.

HUMPBACK OR PINK SALMON.

The humpback or pink salmon (*O. gorbuscha*) is the smallest of the American species, weighing from 3 to 11 pounds, the average being about 4 pounds. In color it is bluish above, silvery below, the posterior and upper parts with many round black spots, the caudal fin always having a few large black spots, oblong in shape. The males in fall are dirty red and are very much distorted in shape, a decided hump appearing on the back, from which deformity the species acquires its name. The flesh is softer than in the other species; it is pale in color, hence its canned name, "pink" salmon.

The southern limit of the fish is the Sacramento River, but only occasional specimens are found here and in the rivers to the northward until Puget Sound is reached. Here a large run appears every other year, the only place on the coast where such is the case.

The humpback occurs in varying abundance in the waters of British Columbia, but it is in the waters of southeast Alaska that it appears in its greatest abundance. Many of the canneries in this region depend mainly upon the humpback for their season's pack, and the canned product now occupies an excellent position in the markets of the world. The fish spawn in nearly all of the small, short streams.

In central and western Alaska the runs are much smaller and the humpback is not much sought after by the cannery men, who are usually able to fill their cans with the more valuable species.

On the Asiatic side it is found in the rivers of Siberia (abundant in the Amur), but not in Japan.

In southeast Alaska the run begins in June and continues until September, or even later in some places. In western Alaska the period is somewhat shorter. In Puget Sound it continues until late in the fall.

DOG OR CHUM SALMON.

The dog or chum salmon (*O. keta*) reaches a maximum weight of 16 pounds, the average being about 8 pounds. When it first appears along the coast it is dirty silvery, immaculate or sprinkled with small black specks, the fins dusky, the sides with faint traces of grid-ironlike bars. Later in the season the male is brick red or blackish, and its jaws are greatly distorted. Its flesh is quite pale, especially when canned, when also it is mushy in texture. It is especially good for freezing, salting, and smoking.

This species has a wide distribution. It is found as far south as San Francisco, but is not utilized commercially in California except on Eel River. It is found in most of the coastal streams from here north, being especially abundant from Puget Sound northward to southeast Alaska, both inclusive. In this region it is being utilized in greater abundance each year, as the market for it widens.

In central, western, and arctic Alaska the species occurs in varying abundance, but is utilized sparingly, except by the natives, with whom it is the favorite species dried for winter food.

This is the most abundant species of salmon in Japan, where it is called sake, and large quantities are dry-salted each year. In Siberia the species is abundant and is known as kaita or kita.

The run of dog salmon comes later than that of any other species except the coho. In Alaska it begins in June, but the height of the season does not occur until late in August or early in September, and fish are found as late as November. In Puget Sound they run from about the middle of August till late in November, and practically the same is true in the Columbia River.

STEELHEAD TROUT.

The steelhead trout (*Salmo gairdneri*) is commonly classed as one of the salmons by the fishermen of the Pacific coast, and it has been included in this report on this account. In different localities the average weight is placed at from 8 to 15 pounds, while extreme sizes reach 45 pounds. The excellent quality of its flesh, causes it to be highly prized for the fresh market, but owing to its pale color only limited quantities are canned.

The principal center of abundance of this species is the Columbia River. It is found from Carmel River, Cal., north to central Alaska, and possibly has an even wider range in Alaska. It seems to be found in the rivers during the greater part of the year. In the Columbia River the spawning season is from February to May, in Puget Sound in the spring, and in southeast Alaska in May and June. The best commercial fishing is in January, February, and March. In California the catching of this species is restricted to hook and line fishing.

II. FISHING GROUNDS AND HISTORY OF THE FISHERIES.

WASHINGTON.

Puget Sound.—Strictly speaking, the name Puget Sound should be restricted to that long, narrow arm extending south from the Strait of Juan de Fuca, but a practice has developed, and is now common among fishermen and others, of designating all the great water area in the State of Washington comprising Puget Sound proper, Strait of Juan de Fuca, Canal de Haro, Rosario Strait, the Gulf of Georgia, and the smaller straits, bays, and sounds, as Puget Sound, and this practice, for convenience sake, has been followed in this report.

This great indentation in the coast, with its numerous islands and many fine harbors, has greatly aided the development of this portion of Washington and has been especially favorable to the prosecution of the salmon and other fisheries. Numerous rivers and creeks enter the Sound, the more important of these being on the eastern shore and comprising the Nooksack, Skagit, Stillaguamish, Snohomish, Duwamish, Puyallup, and Nisqually. On the southern and western shores the tributary streams are nearly all small, the more important being the Skohomish, Quilcene, Dungeness, and Elwha.

The first fishing operations by white men were begun soon after the settlement at what is now known as Seattle, about 1852. For many years the catch was sold either fresh or salted. The first salmon cannery on Puget Sound was erected in 1877, at Mukilteo, in Snohomish County. The first pack was of 5,000 cases, composed wholly of silver or coho salmon. Later this plant put up the first humpbacks ever canned. In 1880 the cannery was removed to West Seattle. In 1885 other canneries were erected at Mukilteo, Seattle, Tacoma, and Clallam Bay, most of them packing silver and humpback salmon alone. The first sockeye salmon cannery was established at Semiahmoo, in Whatcom County, in 1892, from which time on the industry fluctuated considerably, 15 canneries being operated in 1910.

Quillayute River.—This is a small stream, about 30 miles in length, which flows through the southwestern part of Clallam County and empties directly into the ocean. The Quillayute Indian Reservation is located here and the natives catch some salmon and market them on Puget Sound.

Quinault River.—This river, which enters the ocean in the northwestern part of Chehalis County, has a length from the ocean to Quinault Lake of about 40 miles, wholly within the boundaries of the Quinault Indian Reservation. Fishing is restricted to the Indians and the catch is generally shipped by rail to Hoquiam and Aberdeen, on Grays Harbor, and sold to the dealers at these places.

Grays Harbor.—This is the first important indentation on the coast of Washington south of Cape Flattery. It is about 40 miles long from east to west and about 20 miles wide in the widest part. The principal tributary is the Chehalis River, but there are a number of small streams which debouch into the harbor.

As early as 1878 there was a cannery on Grays Harbor, but from then until 1891 the data relating to this branch of the industry are very meager. In 1910 two canneries were in operation at Aberdeen and Hoquiam, respectively.

Willapa Harbor.—The entrance to this harbor, which also includes Shoalwater Bay, is about 27 miles south of Grays Harbor. The harbor runs east and west and is about 25 miles long. Shoalwater Bay extends south from it a distance of about 30 miles; its southern portion ending about a mile from the Columbia River, and on the western side being separated from the ocean by a spit varying in width from three-fourths to 1 mile. The bay is shallow, excepting in the main channel. The principal salmon streams entering the harbor are the Nasel and North Rivers, in which most of the pound or trap nets are located.

Data relating to the early history of the fisheries of this section are very meager. In 1887 there were four canneries in operation, probably the largest number ever operated. In 1910 there was but one—at South Bend.

COLUMBIA RIVER.

The Columbia, which is the largest river of the Pacific coast, rises in British Columbia, flows through Washington, reaching the northern border of Oregon about 75 miles west of the State's eastern boundary; from this point the river forms the dividing line between Oregon and Washington, its general course being westerly. It empties into the Pacific at Cape Disappointment. Its principal tributaries are the Snake, John Day, Deschutes, and Willamette Rivers, and through these the main river drains an enormous extent of territory.

This river, which has produced more salmon than any other river in the world, has had a most interesting history. Many years before the white man saw its waters the Indians visited its banks during the annual salmon runs and caught and cured their winter's supply of food. It was about the year 1833 that a small trading sloop, under the command of Capt. Lamont, came into the Columbia

River on one of her regular trips and dropped anchor near what is now known as St. Helens. While waiting several months for a return cargo the captain salted a number of barrels of chinook salmon, using old Jamaica rum kegs for the purpose. This is the first record of the export of this toothsome fish.

In 1861, H. N. Rice and Jotham Reed began packing salted salmon in barrels at Oak Point, 60 miles below Portland. The first season's pack amounted to 600 barrels. The venture proved fairly profitable and was soon participated in by others.

In the spring of 1866 William Hume, who had assisted in starting the first salmon cannery in the United States, on the Sacramento River, in 1864, finding the run of fish in the latter stream rather disappointing, started a cannery on the Columbia at Eagle Cliff, Wash., about 40 miles above Astoria. Then the river literally swarmed with salmon, and the cannery had no trouble in packing 4,000 cases, which it increased to 18,000 the next year and to 28,000 cases in 1868. In 1867 a crude cannery on a scow was started by S. W. Aldrich, who did all the work, from fishing to canning, himself. In 1868 a cannery was built near Eagle Cliff by one of the Humes, and from this time on for a number of years the industry grew by leaps and bounds.

The banner year in the canning industry was 1884, when 620,000 cases of chinook salmon were marketed. At this time the runs were so enormous that tons and tons of salmon were thrown overboard by the fishermen because the canneries were unable to handle them.

At the present time (1910) there are 10 canneries in operation on the river, while large quantities of salmon are also frozen, mild cured, pickled, smoked, and sold fresh in the markets of the world.

Commercial fishing is carried on mainly between the mouth of the Columbia and Celilo, a distance of about 200 miles, and in the Willamette River. The most of it is in the lower part of the river, within about 40 miles of its mouth. Bakers Bay, on the Washington or north side, and just within the river's mouth, is the favorite ground for pound-net fishing. The principal gill-net drifting ground is from the river's mouth to about 20 miles above Astoria, but drifting is done wherever convenient reaches are found much farther up the river. Most of the drag seines are hauled on the sandy bars in the river near Astoria, which are uncovered at low water. Wheels are operated in the upper river above the junction of the Willamette with the main river.

Astoria is the principal center for all branches of the industry, but more especially for canning. Other places in addition to Astoria at which canneries are located are Ilwaco, Eagle Cliff, Altoona, Brookfield, Pillar Rock, Cathlamet, on the Washington shore, and at Warrendale, Rooster Rock, and Seuferts, on the Oregon shore.

OREGON.

Necanicum Creek.—This short stream is in Clatsop County and enters the Pacific Ocean about 10 miles south of the Columbia River. Its fisheries are of small importance.

Nehalem River.—The Nehalem is a small coastal river that rises in the mountains of Clatsop and Columbia Counties, and flows into the Pacific Ocean in the northern part of Tillamook County. As early as 1887 there was a small cannery here, and the business has been followed ever since.

Tillamook Bay and River.—Tillamook River is a very short stream which enters Tillamook Bay, the latter being in Tillamook County and about 45 miles south of the mouth of the Columbia River.

Fishing is carried on mainly in the bay. The earliest record we have of canneries on this bay is of 1886, when two were in operation. Since 1891 but one has been operated.

Nestucca River.—This stream enters the ocean in the southwestern part of Tillamook County. A cannery operated here in 1887 and the business has been carried on intermittently since then.

Siletz River.—This river has its source in the mountains of Polk County, and enters the ocean in the northern part of Lincoln County. The commercial development of the fisheries was hampered for many years owing to the fact that the river was within the boundaries of what was then the Siletz Indian Reservation. The first cannery was established here in 1896.

Yaquina Bay and River.—The Yaquina ("crooked") River is about 60 miles long; its general course is nearly west through the county of Benton. The river is narrow throughout the greater part of its length. A few miles from its mouth it suddenly broadens out into an estuary from one-half to three-fourths of a mile wide which is commonly called Yaquina Bay. The river enters the Pacific about 100 miles south of the Columbia.

Salmon canning was begun on this river in 1887, when two small canneries were constructed. The next year an additional plant was erected. The business has fluctuated considerably since then and there is now but one cannery.

The fishing grounds are all in the bay and the lower section of the river. The fishermen of this section are fortunate in that they have railroad communication with the outside world, the only place on the ocean side of Oregon, except Tillamook, so situated.

Alsea Bay and River.—Alsea River rises in the southwestern part of Benton County, and flows in nearly a northwesterly direction to the Pacific, a distance of about 60 miles. Like the Yaquina, the "bay" is merely a broadening out of the river just inside its mouth.

The first cannery was established in 1886 and by 1888 there were three in operation. For many years past but one has been in operation.

The best fishing grounds are from the mouth of the river to about 5 miles inland.

Siuslaw River.—This river has its source in the mountains of Lane County, and its course lies first in a northwesterly direction and to the westward until the Pacific is reached. Through part of its course it is the dividing line between Lane and Douglas Counties.

As early as 1878 there were two canneries operated on this river, but from 1879 till 1888 there are no data available showing the extent of the fisheries. At present there are two canneries in operation.

The salmon fishing grounds extend from near the mouth of the river to about 20 miles upstream.

Umpqua River.—With the exception of the Columbia this is the largest and longest river in Oregon. It is formed by north and south forks, which unite about 9 miles northwest of Roseburg, and the river then flows northwestwardly and enters the Pacific. Practically all of this river is within the boundaries of Douglas County, one of the largest counties in the State. A railroad is now being built along this river and when this is completed there will doubtless be a large development of the fisheries of this region owing to the opportunities which will then be offered for shipping fresh fish.

As early as 1878 there were two canneries located on the Umpqua. The number has never been larger than this, and usually there has been but one operating. In 1910 there was but one, at Gardiner.

Coos Bay and River.—Coos Bay is a navigable semicircular inlet of the ocean with numerous arms or branches. There is much marshy ground in the bay, and a number of sloughs, or small creeks, which empty into the bay from both sides. Coos River proper is an unimportant stream, but a few miles in length. North Bend, Marshfield, and Empire are the principal towns on the bay. A branch railroad is being built to these points from the main line of the Southern Pacific Railway, and as soon as this is completed the fishing industry will receive a great impetus. Heretofore this region has depended upon steamers and sailing vessels plying to Portland and San Francisco for its communication with the outside world, and this slow and infrequent means of shipment has very seriously handicapped the fisheries.

Salmon canning began here in 1887, when two canneries opened for business. The business has fluctuated considerably since, most of the time but one cannery being operated, and such being the case in 1910.

Fishing is carried on mainly in the bay. A few set nets are operated in the river.

Coquille River.—This river is formed by three branches, called the North, Middle, and South Forks, which rise in the Umpqua Moun-

tains and unite near Myrtle Point, the head of tidewater, about 45 miles by river from the mouth of the stream. It is a deep and sluggish river, with no natural obstructions to hinder the free passage of fish. Its fisheries have been seriously hampered by the lack of railroad communication, but this will be remedied, as the railroad to Coos Bay will eventually connect with a short line now in existence between the Coquille and Coos Bay.

The principal towns on the Coquille River are Bandon, Prosper, Coquille, and Myrtle Point. Bandon is the shipping port.

Pickled salmon were cured and shipped from this river very early, the first recorded instance of any considerable quantity being in 1877, when 3,000 barrels of salmon were sent to San Francisco. The salt shipments were important until within recent years. The first salmon cannery was erected in 1883, at Parkersburg. In 1886 another was built at the same place, and the following year still another was erected close by. This was the largest number ever in operation in any one year. In 1910 two canneries were operated, both at Prosper.

The fishing grounds are from the mouth to Myrtle Point, about 45 miles inland.

Sixes River.—This small river is located in the northern part of Curry County, and is about 40 miles in length, entering the Pacific a very short distance above Cape Blanco. The salmon caught here are either salted or shipped fresh to the canneries on the Coquille River.

Elk River.—This is another small stream about 40 miles in length, which enters the Pacific just south of Cape Blanco. As on the Sixes River the salmon are either salted or sold fresh to the canneries on the Coquille River.

Rogue River.—This river has as its source Crater Lake in the Cascade Mountains, on the western border of Klamath County, flowing a distance of about 325 miles to the ocean, which it enters at Wedderburn. Its principal tributaries are the Illinois, Applegate, and Stewart Rivers. Owing to canyons and falls in the main river between the mouth of the Illinois River and Hellgate, the latter near Hogan Creek, which runs through the town of Merlin, navigation and fishing are impossible in that section. Except at the mouth of the river the population is very sparse until about the neighborhood of Hogan Creek, where the river approaches the railroad, and from here on for some miles there are numerous growing towns.

Owing to the fact of there being both a spring and a fall run of salmon in this river, the fisheries early became of importance, although sadly hampered because of being compelled to depend wholly on vessel communication with San Francisco, many miles away. In the early years the salmon were pickled and shipped to San Fran-

cisco. In 1877 Mr. R. D. Hume, who had been canning salmon on the Columbia River, removed to the Rogue River, and established near the mouth a cannery which he operated every season (except 1894, when the cannery burned down) until his death in November, 1908, since which date it has been operated by his heirs. Mr. Hume also operated a large cold-storage plant at Wedderburn for several years.

The development of the fisheries of the lower Rogue River was very much hampered by the monopoly which Mr. Hume acquired and maintained until his death. He bought both shores of the river for 12 miles from its mouth, and also owned an unbroken frontage on the ocean shore extending 7 miles north from the mouth of the river. As a result of this, independent fishermen could find no convenient places for landing, which was necessary in order to cure, handle, and ship the fish caught. Since Mr. Hume's death the property has been sold to various parties, but the people of Oregon, upon an initiative and referendum petition, voted in 1910 to close Rogue River to all commercial fishing.

In the upper river ranchers living along the banks have engaged in fishing for a number of years, the catch for the most part being sold fresh. In recent years, as the country has developed, this fishery has become fairly important.

Chetco and Windchuck Rivers.—These two unimportant streams empty into the Pacific in the lower part of Curry County, not far from the California line. The former is about 20 miles and the latter about 25 miles in length. Both have runs of salmon, and small fisheries have been maintained for some years, the catch being either pickled or sold to the California canneries.

CALIFORNIA.

Smith River.—This river, which is the most northerly one in the State, rises near the Siskiyou Mountains, and runs in a westerly direction to the Pacific Ocean.

The river has only a spring run of salmon, and the early recorded history of the fisheries is fragmentary. The pickling of salmon was the main business at first and has been important ever since, as the cannery, which was first established in 1878, operated irregularly, and seems to have shut down entirely in 1895.

Klamath River.—This is the most important river in California north of the Sacramento. It issues from the Lower Klamath Lake in Klamath County, Oreg., and runs southwesterly across Siskiyou County, passes through the southeastern section of Del Norte County, keeping its southerly course into Humboldt County, where it forms a junction with the Trinity River, and thence its course is directed to the northwest until it reaches the Pacific Ocean.

The Klamath River is important as a salmon stream because it has both a spring and fall run of salmon. In 1888 a cannery was established at Requa, at the mouth, and this has been operated occasionally ever since. The pickling of salmon has been done here for a number of years. Some years part of the catch has been shipped fresh to the cannery on Smith River, or to the Rogue River, Oreg., cannery.

Humboldt Bay and tributaries.—The shore line of Humboldt County is bold and high except in the vicinity of Humboldt Bay, where it is rather flat. The latter is the only harbor along the county shore, and it is quite difficult of access, owing to the bar at the entrance, upon which the sea breaks quite heavily. The bay is about 12 miles long and about 3 miles wide. Mad River, which has its rise in the lower part of Trinity County, runs in a northwesterly direction, then makes a sharp turn and enters the bay from the north side. Eel River, which has its rise in Lake County, far to the southeast, runs in a northwesterly direction and enters the bay at its southern extremity. Small railroads running south from Eureka traverse the shores of both rivers for some miles. A railroad to run from the north side of San Francisco Bay to Eureka is now nearing completion, and when in operation it will doubtless aid very materially in extending the market for salmon caught in these rivers.

Mattole River.—This is a small and unimportant river in the southern part of Humboldt County, and is said to have a good run of salmon each year, but no commercial fishing has as yet been carried on here.

Sacramento and San Joaquin Rivers.—These two rivers are the most important rivers in California. The Sacramento is quite crooked, the distance by river from Red Bluff to San Francisco being about 375 miles, while the distance by rail between these two places is only 225 miles. The river rises in several small lakes in the mountains about 20 miles west of Sisson, in Siskiyou County, and for nearly half its length flows through a narrow canyon. The upper portion is a typical mountain stream, with innumerable pools and rapids. A little above Redding the river emerges from the canyon and widens into a broad shallow stream. Below Sacramento it runs through a level country and is affected by tides. Sloughs are numerous in this stretch, some connecting it with the San Joaquin. The Sacramento and San Joaquin Rivers join as they empty into Suisun Bay.

The principal tributaries of the Sacramento which are frequented by salmon are the Pit and McCloud Rivers and Battle Creek. At one time salmon frequented the American and Feather Rivers, but mining and irrigation operations along these streams either killed them off or drove them away.

The San Joaquin River has its source in the Sierra Nevada Mountains. Flowing westerly and forming the boundary between Fresno and Madera Counties for a considerable distance, it then turns abruptly to the north just where it is joined by Fresno Slough, which drains Lake Tulare. From here its general course is northwesterly until it joins the Sacramento River, near the latter's mouth. The Chouchilla and Fresno Rivers are the principal tributaries of the San Joaquin.

The principal fishing grounds for salmon are Suisun Bay, the lower part of San Joaquin River, and the Sacramento River as high as the vicinity of Sacramento. Drift gill nets are used almost exclusively in this section. From Sacramento to Anderson there is considerable commercial fishing, more particularly with haul seines.

Owing to the early and excellent railroad facilities which the fisheries of the Sacramento River have enjoyed, they have not been handicapped so seriously as most of the other Pacific coast rivers in finding profitable outlets for the catch. Soon after the first trans-continental line was opened the shipping of fresh salmon to eastern points began and it has been an important feature of the industry ever since.

The chief event in the history of the salmon fisheries of this river is the fact that the canning of salmon on the Pacific coast had its inception here in 1864. The circumstances leading up to this event and its consummation are interestingly told by Mr. R. D. Hume in the following words:

The first salmon cannery of the United States was located at Washington, Yolo County, Cal. A part of the building was originally a cabin situated on the river bank outside of the levee just opposite the foot of K Street, Sacramento city. It was built in 1852 and occupied by James Booker, Percy Woodson, and William Hume. William Hume came to California in the spring of 1852, bringing with him a salmon gill net, which he had made before leaving his home at Augusta, Me. In company with James Booker and Percy Woodson, Mr. Hume began fishing for salmon in the Sacramento River just in front of the city of Sacramento. William Hume had been salmon fishing in the Kennebec River in the State of Maine with his father, where his father and grandfather had been engaged in the same business since 1780, and their ancestors in Scotland had for pleasure pursued the sportive salmon on the Tweed and Tay for centuries before. In 1856 William Hume went back to Maine, and on his return to California the same year was accompanied by his brothers, John and G. W. Hume, who also engaged in salmon fishing in the Sacramento River. Among the schoolmates of G. W. Hume was one Andrew S. Hapgood, who had learned the tinsmith's trade, and who a short time after G. W. Hume left for California went to Boston and entered the employ of J. B. Hamblen, a pioneer in the canning business, and was sent by him to Fox Island on the coast of Maine, to engage in canning lobsters. The canning of lobster was a new and growing industry, and Mr. Hamblen, to increase his business, a short time after sent Mr. Hapgood to the Bay of Chaleur, an arm of the sea which divides the Province of Quebec from that of New Brunswick,

where, in addition to the canning of lobster, they also canned a few salmon. I believe this was the first salmon canned on the American Continent, and I am informed that the business in a small way is still carried on in that section of the country. In 1863 G. W. Hume went back to Maine, and while there visited Mr. Hapgood at Fox Island, to which place he had been again sent by Mr. J. B. Hamblen to take charge of the works at that place. During the visit of Mr. G. W. Hume to his friend Hapgood a talk about salmon was had, and it was agreed that if salmon on the Pacific coast were as plentiful as represented by Mr. Hume much money could be made in a salmon-cannery business. The plan decided on was that Mr. G. W. Hume, on his return to California, should try and induce his brother William to engage in the business with them, and, if he succeeded in so doing, Mr. Hapgood should purchase the necessary machinery and come out to California in time for the spring season of 1864. Mr. William Hume being agreeable to take part in the enterprise, Mr. Hapgood set out on the journey and arrived at San Francisco on March 23, 1864, and a few days later at the location where the operations were afterwards conducted.^a

* * * * *

For a considerable time after the salmon-canning business was inaugurated the packers suspended operations in the early part of July of each year, as at that time the market would take only goods which showed a rich oil and the best food values.^b

The business languished after the firm established its cannery on the Columbia River, but in 1874 was renewed again by others and continued with varying success until 1905, when it ceased, owing to the smaller quantity of fish available and the difficulty of competing with the mild-cure packers and the fresh-fish dealers.

Monterey Bay.—The first harbor south of San Francisco is Monterey Bay, a large indentation cutting into Santa Cruz and Monterey Counties. Only a portion of it is well sheltered, however. For a number of years it had been known that salmon frequented the waters of this bay for the purpose of feeding on the young fishes which swarmed there. Sportsmen frequently caught them with rod and reel, but it was not until the early eighties that the industry was established on a commercial basis. It has since grown very rapidly. The catch has either been mild cured at Monterey or shipped fresh.

ALASKA.

Alaska is the most favored salmon-fishing region. Many rivers, some of great length and draining enormous areas, intersect the district in every direction, while the number of small creeks is countless. Almost every one of these have runs of salmon of varying abundance. The principal streams entering Bering Sea are the Yukon, Kuskokwim, Togiak, Nushagak, Kvichak, Naknek, Ugaguk, and Ugashik; in central Alaska the Chignik, Karluk, Alitak, Sushitna, and Copper

^a The description of the machinery used and the methods of canning have been quoted in full under "Canning" elsewhere in this report.

^b The first salmon cannery. By R. D. Hume. *Pacific Fisherman*, Seattle, Wash., vol. II, no. 1, January, 1904, p. 19-21.

Rivers are the main streams, while in southeast Alaska are found, among many others, the Anklow, Seetuck, Alsek, Chilkat, Chilkoot, Taku, Stikine, and Unuk Rivers. Most of the fishing in Alaska is carried on in the bays into which these rivers debouch. In southeast Alaska, which is composed largely of islands, the fishing is carried on mainly in the bays, sounds, and straits among these.

Even before the purchase of the District from Russia in 1867 our fishermen occasionally resorted to southeast Alaska and prepared salted salmon. The salmon fisheries did not become important, however, until canning was begun. The first two canneries in the District were built in the spring of 1878, both being located in southeast Alaska. One was built by the Cutting Packing Co. at the Redoubt, Old Sitka, on Baranof Island, while the other was constructed at Klawak, on Prince of Wales Island, by the North Pacific Trading & Packing Co., which latter company still operates at the same place.

The first cannery in central Alaska was built by Smith & Hirsch at Karluk, on Kodiak Island; in western Alaska the first was constructed on Nushagak Bay in 1884 by the Arctic Packing Co.

Owing to the increased demand for canned salmon and the inability of the coast States canneries to keep pace with it, the number of canneries in Alaska rapidly increased for some years until in 1890, when there were 38 in operation. The inevitable happened about this time, however, the production having far outstripped the demand, and canned salmon became a drug on the market.

Heretofore each cannery had operated without regard to the others, but with this condition of affairs prevailing it was soon perceived that steps to reduce the output would have to be taken, and a number of the companies pooled their packs, reduced the number of plants operated, and thus cut down the output nearly one-half. The first arrangement was only temporary, but in 1893 a number of the companies combined permanently and formed the Alaska Packers' Association, which was then, and is yet, the largest company operating in the District.

Since 1893 the industry has experienced periods of alternate prosperity and adversity. In 1910 there were in operation 23 canneries in southeast Alaska, 10 in central Alaska, and 19 in western Alaska, a total of 52. The high prices realized for salmon in 1910 have drawn more capital into the industry, and in 1911 13 new canneries will be constructed and operated.

III. APPARATUS AND METHODS OF THE FISHERY.

GILL NETS.

The gill net is the oldest and most popular form of apparatus in use in the salmon fisheries of the Pacific coast. There are two kinds, drift and set, these names clearly expressing the difference between them. Fine flax or linen twine is generally used in their manufacture, although in some places cotton twine is employed, and it has usually 12 threads and is laid slack. They are hung in the ordinary manner—to a rope with cork floats to support the upper portion of the gear, and to a line with lead sinkers attached, which keeps the net vertical in the water and all its meshes properly distended. The nets are tanned, usually several times each season.

Drift nets vary greatly in length and depth, depending upon the width of the fishing channels, the depth of water, etc. On the Sacramento River they average about 300 fathoms in length, are 45 meshes deep, and have a stretch mesh of from $7\frac{1}{2}$ to $9\frac{1}{2}$ inches. On the coastal rivers of Oregon these nets average about 125 fathoms in length, and are about 36 meshes in depth, the mesh varying with the species of salmon sought. On the Columbia River the nets average about 250 fathoms in length and have a stretch mesh for chinooks of 9 to $9\frac{1}{2}$ inches. On the Willamette River, the principal tributary of the Columbia, they average about 75 fathoms in length, with meshes of 8 and $9\frac{1}{2}$ inches. On Willapa Harbor drift gill nets run from 100 to 250 fathoms in length, are 30 meshes deep, with stretch meshes of 7 and $8\frac{1}{2}$ inches. On Grays Harbor they average 100 fathoms in length, the chinook nets run from 24 to 45 meshes in depth, with a stretch mesh of 9 inches, while the silver or coho nets are 35 meshes in depth, with a stretch mesh of 7 inches. In the Puget Sound region the nets average 300 fathoms in length, with meshes suitable for the particular species sought. In Alaskan waters the nets vary greatly in length and depth, depending upon the places where fished.

Drift gill netting is prosecuted chiefly in the estuaries of the rivers in and near the channels. If the water is clear the nets are set only at night, but should the water be muddy or discolored with glacial silt, fishing can be carried on either night or day. Night fishing is most common in the States, while day fishing is most common

in Alaska. When fishing in rivers it is necessary to work in a straight stretch of water of fairly uniform depth and free from snags or sharp ledges, these being called "reaches."

In setting the net the boat puller rows slowly across the stream while the other man pays out the apparatus, to the first end of which a buoy has been attached. When about two-thirds of the gear is out the boat is turned downstream at nearly right angles to her former course, so that the net, when set, approximates the shape of the letter L. The net is laid out at nearly right angles or diagonally to the river's course, so that it will intercept the salmon that are running in, and is usually put out about an hour before high water slack and taken in about an hour after the turn of the tide. In Alaska the fishermen usually fish on both the high and low slack. The nets are allowed to drift for the time specified, the fishermen drifting along at one end, then the net is hauled into the boat over a wooden roller fixed in the stern, and the fish, which have become gilled in the meshes, are removed and thrown into the bottom of the boat.

Set gill nets are made in the same way as drift nets, in many instances being fragments of the latter, and are usually operated in the upper reaches of the rivers. They vary in length from 10 to 100 fathoms, from 35 to 65 meshes in depth, and have the same sizes of meshes as the drift nets, the size varying, of course, with the species sought for. Sometimes these nets are staked, sometimes anchored, while occasionally only one end is tied to the shore or a stake set in the water.

On the flats off the mouth of the Stikine River, in southeast Alaska, a combination of the drift and set method is followed. A double set of stakes, about 6 feet apart, are set out from the shore for a distance of several hundred yards. An hour or two before slack water the fishermen pay out the net parallel to the line of stakes and about 50 feet from them. The tide drifts the net down until it is caught against the stakes, which retain it until slack water, when the fisherman takes it up and repeats from the opposite direction on the next turn of the tide.

HAUL SEINES.

On the Columbia River, where this form of apparatus plays a prominent part in the fisheries, the nets vary in length from 100 to 400 fathoms; the shallowest end is from 35 to 40 meshes deep, but it rapidly increases in width and is from 120 to 140 meshes deep at the other wing. The "bunt," or bag, in the central part of the net is about 50 fathoms long. These nets are usually hauled on the numerous sand bars which are a very noticeable feature of the river at low tide. Buildings are erected on piles on these sand flats, in which the

men and horses take refuge at high tide, when the bars are covered with water. Operations begin as soon as the beach or bar uncovers, so that the men can wade about. The net is placed in a large seine boat, with the shore end attached to a dory. At the signal the seine boat is headed offshore, while the dory heads toward the bar. As the seine boat circles around against the current the net is paid out in the shape of a semicircle. The dory men hurry to the bar with the shore end of the net, the idea being to get that in as soon as possible in order to prevent the escape of the salmon in that direction. As soon as this has been accomplished, the outer shore line is brought to the bar, when several horses are hitched to the line and begin to haul in the net, care being taken by the men to work it against the current as much as practicable, and to get it in as speedily as they can in order to prevent the escape of salmon either by jumping over the cork line or finding some outlet below the footrope or lead line.

The only other place on the coast where haul seines are important is at Karluk, on Kadiak Island, in Alaska. Here the seines are hauled upon the narrow sand pit dividing the lagoon from the strait, and practically the same method is followed as in the Columbia River.

DIVER NETS.

These are in use in the Columbia River, mainly throughout the middle and upper portions of the river. They vary from 100 to 200 fathoms in length and are used almost exclusively for chinook salmon. In construction they somewhat resemble a trammel net. Two nets are attached together side by side. The outer one, or the one toward the oncoming fish, has a larger mesh than the other, so that if the fish manages to pass through the first, it will be caught in the smaller meshes of the second.

DIP NETS.

These consist of an iron hoop secured to the end of a stout pole with a bag-shaped net fastened to the hoop. They are generally used at the cascades on the rivers, small platforms being erected upon which the operator stands while fishing. Indians formerly used them to a large extent, but, owing to the steady decline in the number of Indians, and the appropriation of favorable spots by the whites for other forms of apparatus, they are but little used now.

SQUAW NETS.

This type is virtually a set net. It consists of an oblong sheet of gill netting, about 12 feet long and 8 feet deep, its lower edge weighted to keep it down, and its upper edge attached to a pole that floats at the surface, and is held by a line or lines to another projecting pole which is securely fastened to the shore, so that it will not

swing around with the strain of the swift current on the net. A single block is attached to the pole, and through this passes a rope, thus making a tackle for the more convenient manipulation of the net. The dip-net fishermen of the Columbia River use this net, which derives its name from the fact that it used to be commonly operated by Indian squaws for taking salmon. But few are now in use, for the same reasons as given for the decline in the use of dip nets.

PURSE SEINES.

This form of apparatus is in quite general use in Puget Sound and southeast Alaska, and has proved highly effective in these deep, swift waters. These seines are about 200 fathoms long, 25 fathoms in the bunt, and 20 fathoms in the wings, all with a 3-inch mesh. The foot line is heavily leaded and the bridles are about 10 feet long. The purse line is made of 1½-inch hemp. The rings through which the purse line is rove measure about 5 inches in diameter and are made of galvanized iron.

On Puget Sound the purse seiners congregate mainly on what are known as the Salmon Banks, off the lower end of San Juan Island, during the run of sockeyes. After this run is over they go up the Sound and fish for dogs and cohos, and later go to the head of the Sound and fish for dogs, cohos, chinooks, and steelhead trout. In southeast Alaska they follow the fish all over the bays, straits, and sounds of that section. Purse seines are used in a few other places, but the fishery is secondary to those with other forms of apparatus.

On Puget Sound special power boats, which are fitted with a power winch for hauling in the net, are used almost exclusively in operating the purse seines. As soon as a school of fish is sighted one end of the seine is attached to a dory, and while this remains stationary the seine boat starts off, the crew paying out the net over a roller in the stern. A circle is made around the fish, the boat returning to the dory. The purse line is then attached to the winch, and the line slowly hauled in by power. As the net comes in, the slack is neatly coiled up on a platform in the stern of the boat, the cork line lying on one side and the lead line on the other. As the circle gradually narrows a man stands at the davit with a long pole which he continually plunges into the circle and between the purse lines for the purpose of frightening the fish away from the center of the net, which is open for about a third of the time required to purse it. The poleman in time becomes very expert and is able to plunge the pole into almost any part of the center and have it return unaided to his hands. After the net has been pursed, the bag is either rolled into the boat or the fish dipped or gaffed from the net into the boat.

This style of fishing is said to have been introduced on Puget Sound by the Chinese in 1886.

TRAPS OR POUND NETS.

A trap is stationary and consists of webbing, or part webbing and part wire netting, held in place and position by driven piles. This piling usually is held together above water by a continuous line of wood stringers, also used to fasten webbing to or to walk on if necessary.

In building, the "lead" is first constructed. This runs at right angles, or very nearly so, to the shore, and consists of a straight line of stakes, to which wire or net webbing is hung from top of high water, or a little higher, to the bottom, making a straight, solid wall.

At a little distance inshore of the outer end of the lead begin what are called the "hearts." These are V-shaped and turned toward the lead, beginning at a distance of 30 to 40 feet on either side of same and running in the same general direction, the "big heart" or outer heart first, the inner heart, supplementing the first, being smaller, and the end of the outer heart leading into it. The narrow end of the inner heart leads into the "pot" and forms what is known as the "tunnel." The tunnel ends in a long and narrow opening, running up and down the long way, and is held in position by ropes and rods. Below this is what is known as the "apron," a sheet of web stretched from the bottom of the heart upward to the "pot;" in order to lead the fish into the tunnel when swimming low in the water, and to obviate the necessity of building the pot clear to the bottom, which would be expensive, as the pots of the traps are usually in quite deep water.

Some traps have "jiggers" (a hook-shaped extension of the outer heart) on each side, which help to turn the fish in the required direction.

The "pot" is placed at right angles with the inner heart and immediately adjoining same. It is a square compartment, with web walls and bottom connected in the shape of a large square sack, fastened to piling on all sides. This pot is hauled up and down by means of ropes and tackles, either by hand or, as is most popular, by steam.

The "spiller" is another square compartment adjoining either end of the pot (sometimes there are two "spillers," one at each end), and is simply a container for fish. A small tunnel leads the fish from the pot into the spiller, from whence the fishermen lift them out. This is accomplished by closing the tunnel from the pot, after which the ropes holding the front of the spiller are loosened and the net wall allowed to drop almost to the level of the water. A steam tug then pushes a scow alongside the spiller and takes position on the outside of this scow. From the deck of the tug a derrick is rigged with a running line from the steam capstan through the block at the top of the derrick. This line is attached

to the far end of a net apron, called a "brailer," which is heavily weighted by having chains along each side and leaded cross-ways at several places. A small boat is run inside the spiller, and the men in this draw the brailer across the barge and let it sink in the spiller. The fish soon gather over it, when the steam capstan quickly reels it in, the net folding over as drawn in from its far side and spilling the fish out on the scow. Men on the scow pick out and throw overboard the undesirable fish. The apron is then drawn back across the pot and the operation repeated so long as any fish remain. In this manner a trap with many tons of salmon in it is quickly emptied.

Traps, like nearly all other fixed fishing appliances, are built on the theory that salmon, like most other fishes, have a tendency to follow a given course in the water, whether a natural shore line or an artificial obstruction resembling one; also that the fish very seldom turns in its own wake. The trap has taken advantage of these natural tendencies of the fish, and is arranged so that, although the salmon may turn, he will continually be led by the wall of net toward and into the trap.

If a trap is located in a place where fish play and where an eddy exists, and the fish run one way with the incoming tide and the opposite with the outgoing, it will fish from both directions; if located where the fish simply pass by, as, for instance, on a point or reef, it will fish from one side only.

A variation of the trap, to be used in places where piles can not be driven, is the floating trap. An experimental trap of this variety was used at Uganuk, on Kodiak Island, Alaska, as early as 1896. Its use was abandoned in 1897, not to be resumed until some years later. A number of floating traps (of the type invented by Mr. J. R. Heckman, of Ketchikan, Alaska) have been and are being used in southeast Alaska, the first having been installed in 1907. The design of this trap follows the shape of an ordinary Puget Sound driven trap. It is constructed of logs, 20 to 26 inches at the butt, bolted and braced together in one solid frame. Suspended from this frame through the logs are $2\frac{1}{2}$ -inch pipes extending down in the water 30 feet. Halfway down these pipes and also on the extreme lower ends are eyebolts, to which the web is drawn down and fastened. Thus the web is kept in place as well as if the pipes were driven piles. The lead is also a continuation of large piles or logs bolted firmly together with similarly suspended pipes and webbing.

The so-called wooden traps on the Columbia River are essentially weirs, being a modification of the brush weirs or traps used by the Indians for the capture of salmon long before the advent of the white men. They are built on shore, of piling and planks, the latter arranged like slats with spaces between. The bowl, or pot, is

provided with a movable trapdoor that can be opened during the closed season and on Sundays, so that the fish can pass through and run upstream. These weirs, after being built, are launched into the river, placed in proper position near the shore, and then ballasted so that they sink to the bottom.

According to Collins,^a "pound nets were introduced on the Columbia River in 1879. In May of that year Mr. O. P. Graham, formerly of Green Bay, Wis., built a pound net on the river similar to those used on the Great Lakes. The success of this venture led to the employment of more apparatus of this kind, and many fishermen went West to participate in the fishery."

According to the same authority^b Mr. H. B. Kirby, who had previously fished on the Great Lakes, set a pound net in Puget Sound about 1883, but it was a complete failure. On March 15, 1888, he again set a pound net, which he had designed to meet the new conditions, at Birch Bay Head, in the Gulf of Georgia. It proved a complete success, and was the forerunner of the present large number which are set annually in these waters.

In Alaska the first trap was set in Cook Inlet about 1885. British Columbia refused to permit the use of pound nets in its waters until 1904, when their use was allowed within certain limited regions.

Some of these trap nets, especially on Puget Sound, have proved extremely valuable. The years 1898 and 1899 covered practically the high-water mark, as several desirable locations changed hands in those years at prices ranging from \$20,000 to \$90,000 for single pounds, the original expense of which did not exceed \$5,000. But few have brought such high prices since, however, owing to the decline in the run of salmon.

The location of sites for these nets is regulated by law in Oregon, Washington, and British Columbia, but in Alaska the procedure is not well defined and has proved rather confusing to strangers. Some acquire the necessary shore line by mineral location or by the use of scrip, while still others have merely a squatter's right. Within the bounds of the forest reserve no land can be acquired except by lease, which may be secured from the United States forestry agent, Ketchikan, Alaska.

INDIAN TRAPS.

The natives, especially in Alaska, have various ingenious methods of catching salmon. In the Bering Sea rivers they catch them by means of wickerwork traps, made somewhat after the general style of a fyke net. These are composed of a series of cylindrical and conical baskets, fitting into each other, with a small opening in the

^a Report on the fisheries of the Pacific Coast of the United States, by J. W. Collins, Report of Commissioner of Fish and Fisheries for 1888, p. 210. 1891.

^b *Ibid.*, p. 257.

end connecting one with the other and the series terminating in a tube with a removable bottom, through which the captive fish are extracted. Some of the baskets are from 15 to 25 feet in length and are secured with stakes driven into the river bottom, while the leader, composed of square sections of wickerwork, is held in place by stakes.

During the summer of 1910 the author found and destroyed an ingenious native trap set in Tamgas stream, Annette Island, south-east Alaska. This stream is a short and narrow one, draining a lake, about midway of which are a succession of cascades. In the narrowest part of the latter, and in the part up which the fish swim, a rack had been constructed of poles driven into the bottom and covered with wire netting, so as almost wholly to prevent salmon from passing up. Just below, and running parallel to the rack and at right angles to the shore, was placed a box flume with a flaring mouth at the outer end. At the shore end the flume turned sharply at right angles and discharged into a square box with slat bottom and covered over with boughs. The fish in ascending the stream would be stopped by the rack and in swimming around many of them would be carried by the current into and down the flume, eventually landing in the receiving box alongside the shore.

WHEELS.

Fish wheels are of two kinds, the floating or scow wheel, which can be moved from point to point if need be, and the shore wheel, which is a fixed apparatus. They operate in exactly the same manner, however. The stationary wheel is located along the shore in a place where, experience has shown that the salmon pass. Here an abutment is built of wood and stone, high enough to protect it from an ordinary rise in the river. To this is attached the necessary framework for holding the wheel. The latter is composed of three large scoop-shaped dip nets made of galvanized-iron wire netting with a mesh of $3\frac{1}{2}$ to 4 inches. These nets are the buckets of the wheel, and they are so arranged on a horizontal axis that the wheel is kept in constant motion by the current, and thus picks up any fish which come within its sweep. The nets are fixed at such an angle that as they revolve their contents fall into a box chute through which the fish slide into a large bin on the shore. The wheels range in size from 9 to 32 feet in diameter and from 5 to 15 feet in width, and cost from \$1,500 to \$8,000, the average being about \$4,000. A number of them have long leaders of piling running out into the river, which aid in leading the salmon into the range of the wheel.

The scow wheel consists of a large square-ended scow that is usually decked at one end and open at the other. Several stanchions, some 8 to 10 feet high, support a framework upon which an awning

is spread to protect the fish from the sun's rays and the crew from the elements. To one end of the scow are fastened two upright posts, which are guyed by wooden supports, while projecting from the same end is the framework which supports the wheel, the latter being constructed in the same way, but on a smaller scale, than the stationary wheel. In operation the scow is anchored with the wheel end pointing downstream, and as the wheel is revolved by the current the fish caught fall from the net into a box-chute, through which they slide into the scow. As stationary wheels can be used only at certain stages of water, the scow wheel is a necessary substitute to be used at such times as the former can not be operated.

The above forms of wheels are used exclusively on the Columbia River.

An ingenious device is used by some of the wheelmen on the Columbia River in getting their catch to the canneries, a few miles farther down the river. The salmon are tied together in bunches and these attached to air-tight casks and sent down the stream. At the canneries small balconies have been constructed at the water end of the building. A man armed with a pair of field glasses is stationed here, and as soon as he sights one of these casks he notifies a boatman, who goes out and tows in the cask and salmon. About 800 pounds of salmon are attached to a keg, and a tag showing the wheel from which shipped is tied to the fish.

In 1908 the first fish wheel to be located in the coastal waters of Alaska was operated in the Taku River, in southeast Alaska. The wheel was set between two 4-foot scows, stationed parallel to each other, and each 40 feet in length. The wheel had two dips, each 22 feet in width and hung with netting. It could be moved from place to place, the same as the scow wheels on the Columbia River. It was operated throughout the king and red salmon runs, but caught almost no salmon, and was not set in the succeeding years.

For many years the natives of the interior of Alaska have been resorting to the banks of the Yukon River and its tributaries in order to secure a sufficient supply of salmon to sustain them through the succeeding winter. The favorite apparatus of these natives is a type of fish wheel of local invention, which has been in use by them for many years, probably long before the white man first saw the Yukon. A square framework of timbers is constructed in the water and moored to the bank by ropes. A wheel, composed of three dips, is placed in this, the axle resting upon the framework. The shape of the dip is such that the salmon caught roll off it into a trough, down which they slide into a boat moored between the wheel and the shore. Although crude in construction, it is very effective and a large number of them are set each season.

The Columbia River fish wheel is a patented device. It was first used by the patentees, Messrs. S. W. Williams & Brother, in 1879, and for several years they retained a monopoly in its use. A number are now operating on the river. The device was not new even when patented, as the natives of the Yukon River Basin had been using a precisely similar principle for an unknown number of years previously, while a similar "fishing machine," as it is called, had been in use prior to this time and is still used by white fishermen on the Roanoke River, in North Carolina.

REEF NETS.

As the name indicates, this device is used around the reefs. Under natural conditions the reef is covered with kelp throughout its length, the kelp floating at the top of the water. A channel is cut through this, and in it is placed a tunnel of rope and netting, which flares at the outer end, in deep water, and into which is thatched grass, kelp leaves, or any other article resembling submarine growth, to hide the construction sufficiently to avoid frightening the fish. Short leads of kelp are also arranged on the sides so as to draw the fish to the tunnel, which is held in place by anchors. On the reef itself two boats are anchored parallel to each other and some feet apart. An apron of netting is fastened to the rear of the two boats, while the other end extends under the small end of the tunnel and is kept in place by men in the forward ends of the boats, who have lines fastened so the apron can be raised by them. The device can only be used with the tide entering the tunnel at the large end. When the fish have entered and passed through the tunnel upon the apron, the men raise the floating end of the latter and dump them into the boats.

At one time this was a favorite device of the Puget Sound natives for catching sockeye salmon. They attribute its origin to one of the Hudson Bay Company's employees, who, they say, taught them a long time ago how to catch salmon in this way. Owing to the large number of men required to work them, and the fact that they can be worked only at certain stages of tide and in favorable weather, these nets have gradually been supplanted by other devices. In 1909 but five were used and these were operated off the shores of San Juan, Henry, Steuart, and Lummi Islands, and in the vicinity of Point Roberts.

TROLLING.

Each year the catching of salmon by trolling becomes of increasing importance commercially. For some years sportsmen had this exciting and delightful occupation to themselves, but eventually the mild curers created such a persistent and profitable demand for king, or chinook, salmon that the fishermen, who had previously restricted

their operations to the use of nets during the annual spawning runs, which last but a small portion of the year, began to follow up the fish both before and after the spawning run and soon discovered that they were to be found in certain regions throughout nearly every month in the year.

The Monterey Bay, Cal., trollers use 48 cotton line generally. A few inches below the main lead an additional line is added, with a small sinker on it. This gives two lines and hooks, and as the main line has but the one lead, and that above the junction with the branch line, it floats somewhat above the latter, which is weighted down with a sinker. The main stem is about 20 fathoms in length, while the branch lines are about 5 fathoms each. These lines cost about \$3.50 each. No spoon is used, but bait almost invariably. A few fishermen use a spread of stout steel wire, 4 feet long, with 5 or 6 feet of line on each end of the spread, two lines and hooks.

On the upper Sacramento River (mainly at Redding and Keswick) some fishing is done with hand lines. A small catch was made here in 1908, but none were so caught in 1909.

Even as early as 1895 trolling was carried on in the Siuslaw River, Oreg., for chinook and silver salmon. At Oregon City and other places on the Willamette River a number of chinook salmon are caught by means of trolling each year, mainly by sportsmen. A spoon is quite generally employed in place of bait. The fishermen claim that the salmon are not feeding at this time, as their stomachs are shriveled up.

For a number of years the Indians living at the reservation on Neah Bay, Wash., have annually caught large numbers of silver and chinook salmon in the Strait of Juan de Fuca. A few white fishermen also engage in this fishery at the present time in the same waters, while others troll for the same species, but more particularly silvers, in parts of Puget Sound proper. The ordinary trolling line, with a spoon instead of bait, is used.

The most remarkable trolling region is in southeast Alaska. For some years the Indians here had been catching king salmon for their own use during the spring months, and about the middle of January, 1905, king salmon were noticed in large numbers in the vicinity of Ketchikan. Observing the Indians catching these, several white fishermen decided to engage in the pursuit, shipping the product fresh to Puget Sound ports. They met with such success that 271,644 pounds, valued at \$15,600, were shipped. The next year several of the mild-cure dealers established plants in this region, thus furnishing a convenient and profitable market for the catch, and as a result the fishery has grown until, in 1910, 204,823 king salmon and 6,000 coho salmon were caught and marketed. The length of the fishing season has also lengthened until now the business is prosecuted vigorously during about seven months in the year,

and in a desultory manner for two or three months more, only the severe winter weather preventing operations the rest of the year.

In southeast Alaska the fishermen generally use either the Hendryx Seattle trout-bait spoon no. 5 or the Hendryx Puget Sound no. 8. The former comes in nickel or brass or nickel and brass, the full nickel preferred. The Siwash hook no. 9/0, known as the Victoria hook in British Columbia, is in quite general use. As a rule, but one hook is used, and this hangs from a ring attached to a swivel just above the spoon, while the point of the hook comes a little below the bottom of the spoon. Occasionally double or treble hooks are used. Some fishermen use bait, and when this is done the herring, the bait almost universally employed, is so hooked through the body as, when placed in the water, to stretch out almost straight and face forward as in life.

A small commercial fishery is carried on in this region for coho salmon, mainly in August and September, in the neighborhood of Turnabout Island, in Frederick Sound. A Stewart spoon with two hooks on one ring is used, baited with herring in such a way that the fish is straightened out and faced toward the spoon. The sportsmen of Ketchikan also fish with rod and reel for this species in the neighborhood of Gravina Island, using a Hendryx spoon (kidney bait no. 6), which is silvery in color on one side and red on the other. Although much smaller than the king, the coho salmon is more gamy.

Reports from the trollers of southeast Alaska prove that all species of salmon will take the hook at some time or other in the salt waters of this region, an examination of their stomachs generally showing that they are either feeding or in a condition to feed.

BOW AND ARROW.

On the Tanana River, a tributary of the Yukon River, in Alaska, the Indians hunt salmon in birch-bark canoes with bow and arrow. As the canoe is paddled along and the Indian sees the dorsal fin of the salmon cutting the surface of the muddy water he shoots it. The tip of the arrow fits into a socket, and when struck the tip, which when loose is attached to the stock by a long string, comes out of the socket and the arrow floats, easily locating the fish for the fisherman.

SPEAR AND GAFF.

Spears of varying shapes and styles have been in use by the Indians from time immemorial and are still employed on many rivers in which salmon run. With the exception of the Chilkoot and Chilkat Rivers of Alaska, practically all of the catch secured in this manner is consumed by the fishermen and their families. In the Chilkoot River the Indians have built numerous racks in the stream and on the banks, upon which they stand and hook the fish out with a gaff attached to a pole. The catch is sold to the cannery located on Chilkoot Inlet.

IV. FISHERMEN AND OTHER EMPLOYEES.

In the early days canning was a haphazard business, and workmen came and went as common laborers do in the wheat fields of the West. As the business increased in importance and the need of skilled labor became imperative, men were put to certain work and kept at it from season to season, with the result that in a few years a corps of highly skilled laborers had been evolved, and this had much to do with the rapid extension of the industry.

For many years Chinese formed the greater part of the cannery employees, the superintendent, foreman, clerks, machinists, and the watchmen alone being whites. No other laborers have ever been found to do the work as well or with as little trouble as the Chinese. In times of heavy runs, when the cannery would have to operate almost night and day in order to take advantage of what might be the last run for the season of the sometimes erratic salmon, the Chinese were always willing, even eager, to do their utmost to fill the cans, and if fed with the peculiar food they insisted upon having and due regard was had to certain racial susceptibilities, the cannery man could almost invariably depend upon the Chinese doing their full duty.

The Chinese-exclusion law cut off the supply of Chinese, and as the years went by and their ranks became decimated by death, disease, and the return of many to China, the contractors were compelled to fill up the rapidly depleting crews with Japanese, Filipinos, Mexicans, Porto Ricans, etc., with the result that to-day in many canneries special quarters have to be provided for certain of the races—more particularly the Chinese and Japanese—in order to prevent racial hatred from engendering brawls and disturbances.

The Japanese now compose about one-half of the cannery employees. While a few cannery men express themselves as well pleased with this class of labor, the majority find it troublesome.

In Alaska and at a few places in the States Indians are employed in the canneries. In Alaska more would be employed if they could be secured. They make fair workpeople, but are rather unreliable about remaining through the season.

The supplying of this kind of labor is done largely through the contract system. In the large cities along the coast are labor agencies, mainly owned by Chinese, which make a specialty of furnishing labor for this work. In the agreement between the canning

company and the contractor the company guarantees to pack a certain number of cases during the coming season and the latter agrees to do all the work from the time the fish are delivered on the wharf until they are ready to ship at the end of the season, for a certain fixed sum per case. Should the cannery pack more than the guaranteed number, which it usually does if possible, the excess has to be paid for at the rate per case already agreed upon, while if the pack, for any reason should fall below the contract amount the company must pay for the shortage the same as though they had been packed. The company transports the Chinese to the field of work and carries them to the home port at the end of the season. It provides them with a bunk house, and furnishes fuel, water, and salt. The contractor sends along with each crew a "boss," who has charge of the crew, and furnishes their food, the company transporting this free.

White men do the greater part of the fishing for salmon, many nationalities being represented, but Scandinavians and Italians predominating almost everywhere. A number of Greeks are to be found fishing in the Sacramento, while Slavonians do most of the purse-seining on Puget Sound. The native-born American is not often found actually engaged in fishing, but frequently is the owner of the gear or has a responsible position in the packing plants.

A number of Indians participate in the fisheries of Alaska, and a few fish in Washington. The only Chinese engaged in fishing are in Monterey Bay. A number of Japanese also fish in this bay, which is the only place in American territory where they fish for salmon, except in Alaska, where the small number of 13 were occupied in 1909. A number of Japanese engage in fishing in Canadian waters.

In many places on the coast, particularly in Alaska, fishing is a hazardous occupation. In Alaska most of it is done in the bays, sounds, and straits, where storms are frequent, and the annual loss of life is heavy. The records of the Alaska Fishermen's Union show for its members the following losses of life by drowning: 1905, 10 men; 1906, 5 men; 1907, 10 men; 1908, 17 men; and 1909, 17 men.

The fishermen early saw the advantages of organization, and nearly every river now has a union, which is subordinate to the general organization. One of the most typical of these is the Alaska Fishermen's Union, which has active jurisdiction over all sections of Alaska, except a portion of southeast Alaska. Early in the year this organization enters into contracts with the salmon canneries and salt-eries, by which the rates of wages, duties, etc., of the fishermen are fixed in advance. As a result of this mutual agreement upon terms, but little trouble is experienced with the fishermen, who generally conform scrupulously to the terms of the contract, and strikes and bickerings, which were very common a few years ago, are now almost entirely absent.

V. FISHERY REGULATIONS.

CONTROVERSIAL FORMS OF APPARATUS.

From time immemorial the users of certain forms of fishing apparatus have complained of and condemned the use of other forms, which, either through disinclination, through lack of financial means, or because it was not suitable for use in the section in which they fished, they themselves have not seen fit to employ. In some instances these complaints are well founded, but an unprejudiced observer is apt to view with suspicion charges advanced under conditions when personal interest may so easily cloud or color the individual judgment. In a court of equity it is a well-established principle that the plaintiff must appear with clean hands, and that is a difficult matter for the users of any form of apparatus in the salmon fisheries of the Pacific coast. If in one section the fishermen live strictly within the letter and spirit of the law, the users of the same apparatus in another section may be the most persistent and destructive violators. And, again, while the law may be strictly observed, the law itself may be inadequate or purposely deficient, and the apparatus therefore be doing incalculable damage to the fisheries.

While all forms of apparatus in use in the salmon fisheries of the Pacific coast have been objected to in some one section or another, the principal complaints have been against fish wheels and trap or pound nets. The wheels are used only in the Columbia River. The traps are found in the Columbia River and in the other waters of the State of Washington and in Alaska.

To the objections of other fishermen the owners of wheels and traps retaliate by charging prejudice and self-interest, and with some justification. It is unquestioned that these costly forms of apparatus are beyond the financial means of the ordinary fishermen, that their use reduces the number of persons employed in the fisheries, and that the owners, who are usually the packers or others closely affiliated with them, can, if they so desire, render themselves largely independent of other fishermen, such as the gill netters and seiners, and thus keep down the cost of the fish to the packers. Although not often advanced publicly, this is the real basis of the most of the complaints. Publicly the objections are based upon higher grounds,

such as the waste through catching and killing in wheels and traps of enormous quantities of salmon which can not be handled in the limited time available, or of species which the packers have no use for, and which they find it easier or less expensive to kill by much handling than to release and in so doing lose a few salmon.

One thing should never be lost sight of, however. Fishery apparatus is set for the purpose of catching fish, and its value is dependent upon the degree of effectiveness with which it accomplishes the object sought with the least expenditure of money and time for construction and operation.

It is a question whether, under present conditions, if the use of traps were abolished, the other forms of apparatus would be able to keep pace with the demand for fish. But the question of whether traps should be allowed or not in any one section should be settled by reference solely to the conditions prevailing in that section, and not to theoretical or general objections to traps as traps or to objections based upon trap fishing in some other and, possibly, vastly different section. There are some regions on the Pacific coast where if traps were permitted they would soon destroy the run of salmon, while there are many other sections where they would not injure the fisheries at all, unless possibly by use in too great numbers. The latter is especially true in many parts of Alaska, where the chief objection is that in a few places too many of them are grouped together.

A considerable part of the objection to the use of traps is doubtless due to the generally shameless disregard of the laws in the past, and in some sections also to-day. In Alaska up to 1908 the trap owners paid practically no attention to the laws, and the same is true to a large extent to-day on Puget Sound, and to a lesser extent, possibly, in the Columbia River. Since the enactment and rigid enforcement of the excellent trap law of 1906 in Alaska, the objections to trap nets have decreased very noticeably, though the traps have probably caught more fish than they did under the old conditions, the only difference being that the catch has been distributed more equally, and not, as in former times, caught chiefly in those traps situated nearest to the ocean, while those in the upper reaches took but few.

The Washington law prescribes minutely the method to be followed in closing traps during the weekly closed season and appears on its face to be an excellent plan. In practice it is quite otherwise, however, for one person can close or open the trap in one or two minutes' time, and all the watchman has to do in the weekly closed season is to let the apron down whenever he sees a boat approaching, raising it again as soon as he is sure the visitor is not a fish warden. Thus it is practically impossible to detect any but the boldest or most careless violations of the law.

The provision in the Alaska fisheries law regulating the manner of closing traps during the weekly closed season is without question the best in the country, and Washington could adopt it with much profit. It requires that "the gate, mouth, or tunnel of all stationary or floating traps shall be closed, and 25 feet of the webbing or net of the 'heart' of such traps on each side next to the 'pot' shall be lifted or lowered in such manner as to permit the free passage of salmon and other fishes." With two men stationed on the trap at least 15 or 20 minutes of most strenuous work is required to open or close the trap in this manner, and the fishery agent has ample time to reach the scene before the operation is completed. This fact has been found to be an excellent deterrent.

At first the owners advanced the plea that the lowering of 25 feet of the web of the heart next to the pot would so weaken the trap that it might be carried away by the very strong and high tides which prevail in Alaska, but three years' actual trial has proved this fear to be groundless, and now no objections are heard to this feature of the law.

Although not used to as great an extent, wheels have probably occasioned more controversy than traps. While the traps are usually set in either bays, straits, and sounds, where the water is salt or brackish, or in the lower reaches of all the rivers, the wheels are set in the upper courses of the Columbia River only. After the fish have run the gauntlet of the almost countless gill nets, seines, and trap nets in the lower and middle river, and are approaching their spawning beds, they meet with the runways leading to the wheels, which in some instances are set in natural channels in the cascades or falls, or in artificial channels through which the greater part of the run must of necessity pass. Nearly all of the salmon hatcheries on the Columbia are located either on the main river below Cascade Locks, or on one of the tributaries entering the river below there, while above this point there were operated in 1909 17 stationary wheels and 5 scow wheels.

It may be maintained that a salmon which has successfully evaded the nets in the section of the river below Cascade Locks is of vastly more importance to the preservation and perpetuation of the fisheries than a number which have not yet crossed the bar at the mouth of the river. Thus, it has been argued, while wheels have not done anything like the damage to the fisheries ascribed to them, a regard for the perpetuation of the fisheries of the Columbia River demands that their use, as well as that of all other forms of apparatus for the taking of fish commercially, should be prohibited above Cascade Locks.

This brings up the question of the justice of such an arrangement from the standpoint of the owners of the wheels. When they put

in these wheels their use was lawful, and the same is true to-day. They are expensive apparatus, and many thousands of dollars are invested in them. In addition there is an important salmon cannery located at Seuferts, just above The Dalles, which would be absolutely worthless if the above action were taken. It would be no more than just, if the States of Oregon and Washington decided to abolish all commercial fishing above Cascade Locks, that a fair valuation for losses be fixed by arbitration and paid to those affected.

There is also no question but what too many gill nets and trap nets are now being fished in the lower part of the river, and some scheme ought to be devised by which the number of licenses annually granted can be reduced very materially.

Strict regulations of the forms of apparatus used in the salmon fisheries and the curtailment of certain or all forms when they become too numerous will be of greater efficacy in the perpetuation of the industry than any other method which has been so far recommended or tried except that of closed seasons.

LAWS AND THEIR ENFORCEMENT.

The history of the enactment and enforcement of laws relating to the salmon fisheries of the Pacific coast (except possibly California) is not one that those earnestly and sincerely desirous of preserving and perpetuating the fisheries have reason to be proud of. In the first place, it has been and is yet exceedingly difficult to secure efficient laws, owing to the influence of the selfish interests which have no regard to the future. In the second place, it was and is yet difficult to secure the enforcement of even the laws that are on the statute books. In most States a change in the governorship almost invariably entails a change in fish commissioner, who is often more concerned with pleasing the interests that secured his appointment and retain him in office than in giving the affairs of his department the attention that they require. This condition, not peculiar to the Pacific Coast States alone, doubtless will eventually be removed to a great extent by divorcing the fisheries departments from politics. The Pacific Coast States have had in the past and still have some earnest men who have been and are doing good work, and this number can easily be increased by making the positions permanent. Under present conditions a fish commissioner scarcely has a comprehensive grasp of the intricate problems of his department and begins to be of value to the State before a change of administration occurs and he is compelled to give way to another man, who in turn must be taught all that his predecessor had learned.

The worst condition of affairs in regard to the making and enforcement of fishery laws is found to prevail in those waters which form the boundary between States or between Canada and the United States.

The Columbia River, which forms the boundary between Oregon and Washington, affords a typical example of the evils which can result from a division of responsibility between two States. For many years each State enacted laws regulating the fisheries of the river with very slight regard usually to laws already in force in the other State. As a result of this the fishermen transferred their residence for license purposes from State to State as the laws of one or the other best suited their particular purposes.

The fishermen and packers also were in apparently irreconcilable conflict as to the proper means to be taken to conserve the fisheries, and each session of the legislatures saw strong lobbies present to work for certain selfish ends, while the few earnest men who had the real welfare of the fisheries of the river at heart had difficulty in making the slightest headway against the influence of these lobbies.

To further complicate the matter, in 1894 Oregon claimed that under the provisions of the enabling act admitting it as a State it had jurisdiction to the Washington shore, and proceeded to arrest Washington men who were fishing in what was the open season according to Washington law but the closed season under Oregon law.

In June, 1908, the voters of the State of Oregon had presented for their consideration two bills radically affecting the waters of Columbia River. One closed the river, east of the mouth of the Sandy River, against all fishing of any kind except with hook and line, and was originated by gill-net fishermen of the lower river for the purpose of eliminating fish wheels in the upper waters. This bill was the first presented to the people, and when it appeared the upriver men retaliated by presenting a bill affecting the lower river to such an extent that it practically prohibited the net fishermen from operating.

Very much to the surprise of all concerned both bills were passed and became laws on July 1, to take effect, as provided, on August 25 and September 10, respectively. The Oregon master fish warden proceeded to enforce both laws, arresting all violators on both sides of the river, irrespective of whether or not they were operating under a Washington or Oregon license, and incidentally did the fisheries a great service by bringing prominently before the public the anomalous condition of affairs which were occasioned by the archaic system under which the fisheries of the Columbia were governed. The State of Washington appealed to the United States courts, which, after argument, issued an injunction preventing the warden from enforcing the laws so far as the Washington fishermen were concerned.

In the meantime the attention of the general Government had been drawn to the apparently irreconcilable conflict between the two States, and fearing that in the mêlée the interests of the fisheries would be lost sight of, President Roosevelt, in a message to Congress, after reciting briefly the lack of harmony in jurisdiction by the

States, recommended that the general Government take over the control of the fisheries of the Columbia, as well as other interstate rivers.

This had the effect of bringing matters to a head and negotiations were soon in progress looking to the preparation of a treaty between the two States by which uniform laws would be adopted, and thus each State have concurrent jurisdiction to the opposite shore of the river. The legislatures each appointed a committee of eight members to confer and frame joint legislation. The two committees met in Seattle, Wash., early in 1909, and agreed upon the following recommendations:

First. A spring closed season from March 1 to May 1.

Second. A fall closed season from August 25 to September 10.

Third. A Sunday closed season from 8 p. m. Saturday of each week to 6 p. m. the Sunday following between the 1st day of May and the 25th day of August.

Fourth. We suggest the mutual recognition by each State of the licenses issued to floating gear by the other State.

Fifth. That the State of Oregon repeal chapter 89 of the session laws of Oregon for the year 1907, relative to the operation of purse seines and other like gear on the Columbia River.

Sixth. We recommend the enactment of similar laws in both States carrying an appropriation of at least \$2,500 in each State and providing for the destruction of seals and sea lions and the granting of a bounty on the same, to be \$2.50 for seals and \$5 for sea lions.

Seventh. We recommend the repeal of both the fish bills passed under the provisions of the initiative and referendum in June, 1907, by the people of the State of Oregon, said bills being designated on the ballot as 318, 319 and 332, 333.

The recommendations were enacted into law by both States, and at the same time the State of Washington in its bill also prohibited fishing for salmon within 3 miles of the mouth of the Columbia between March 1 and May 1 and between August 25 and September 10, or salmon fishing on tributaries of the Columbia, except the Snake, between June 1 and September 15; and also prohibited fishing by any means for salmon save by hook and line in the Kalama, Lewis, Wind, Little White Salmon, Wenatchee, Methow, and Spokane Rivers and in the Columbia River 1 mile below the mouth of any of the rivers named. The agreement was subjected to a rather severe strain, however, when it was discovered that the Oregon Legislature had failed to provide the same closed periods for the tributaries that were enacted for the Columbia, thus leaving the Willamette, Clackamas, Lewis and Clark, and Youngs Rivers and Spikanon Creek open to fishing for 15 days in March and 15 days in April, while the Columbia was closed. The cry of bad faith was at once raised by the Washington fishermen, and for a short time it appeared that the agreement would be broken at the very beginning. The Oregon Board of Fish Commissioners took the matter up, however, and by

order closed these streams to all fishing during the times of closed season on the Columbia, and thus restored peace once more.

The conditions which prevail in Puget Sound adjacent to the boundary between Washington and British Columbia have also been the cause of serious anxiety to those interested in the perpetuation of the salmon fisheries. The great schools of sockeye salmon which are on their way from the ocean to the spawning beds in the Fraser River pass through this section, and it is here that the greater part of the fishing is done. The Province of British Columbia has made earnest efforts to preserve this run, but unfortunately the same can not be said of the State of Washington. The laws are fairly good, but owing partly to the small force and facilities available for executing them and partly to other reasons, they have not always been enforced as they should be.

This condition of affairs on Puget Sound and similar conditions in other boundary waters led the general Government to take up the matter, and on April 11, 1908, a convention was concluded between this country and Great Britain for the protection and preservation of the food fishes in international boundary waters of the United States and Canada. Both Governments appointed international commissioners—Dr. David Starr Jordan for the United States and Mr. S. T. Bastedo (who was succeeded later by Prof. Edward Ernest Prince) for Canada—whose duty it was to investigate conditions prevailing in these waters and to recommend a system of uniform and common international regulations. After an exhaustive investigation the commissioners submitted recommendations, which included the following affecting the boundary waters dividing the State of Washington and the Province of British Columbia, these waters being defined as the Strait of Juan de Fuca, and those parts of Washington Sound, the Gulf of Georgia, and Puget Sound lying between the parallels of $48^{\circ} 10'$ and $49^{\circ} 20'$:

GENERAL REGULATIONS.

3. *Disposition of prohibited catch.*—In case any fish is unintentionally captured contrary to the prohibitions or restrictions contained in any of the following regulations, such fish shall, if possible, be immediately returned alive and uninjured to the water.

4. *Dynamite, poisonous substances, etc.*—No person shall place or use quicklime, dynamite, explosive, or poisonous substances, or electric device in treaty waters for the purpose of capturing or killing fish.

5. *Pollution of waters.*—No person shall place or pass, or allow to pass, into treaty waters any substance offensive to fishes, injurious to fish life, or destructive to fish fry or to the food of fish fry, unless permitted so to do under any law passed by the legislative authority having jurisdiction.

No person shall deposit dead fish, fish offal, or gurry in treaty waters, or on ice formed thereon, except in gurry grounds established by the duly constituted authorities.

6. *Capture of fishes for propagation or for scientific purposes.*—Nothing contained in these regulations shall prohibit or interfere with the taking of any fishes at any time for propagation or hatchery purposes, and obtaining at any time or by any method specimens of fishes for scientific purposes under authority granted for Canadian treaty waters by the duly constituted authorities in Canada and for United States treaty waters by the duly constituted authorities in the United States.

12. *Capture of immature salmon prohibited.*—No salmon or steelhead of less than 3 pounds in weight shall be fished for, killed, or captured in treaty waters.

13. *Salmon weirs, etc., above tidal limits prohibited.*—No salmon and no steelhead shall be fished for, killed, or captured by means of a net of any sort, any weir or any fish wheel, above tidal limits in any river in treaty waters.

14. *Close season for sturgeon.*—During the term of four years next following the date of the promulgation of these regulations no sturgeon shall be fished for, killed, or captured in treaty waters.

15. *Capture of fish for fertilizer or oil prohibited.*—Fishes useful for human food shall not be fished for, killed, or captured in treaty waters for use in the manufacture of fertilizer, or of oil other than oil for food or medicinal purposes.

16. *Naked hooks and spears prohibited.*—No spear, grappling hook, or naked hook, and no artificial bait with more than three hooks, or more than one burr of three hooks attached thereto, shall be used for the capture of fish in treaty waters. This regulation shall not prohibit the use of a gaff in hook-and-line fishing.

17. *Torching prohibited.*—No torch, flambeau, or other artificial light shall be used as a lure for fish in treaty waters.

The following regulations relate specifically to the waters named:

STRAIT OF JUAN DE FUCA AND ADJACENT WATERS.

The following regulations (62 to 66, inclusive) shall apply to the Strait of Juan de Fuca, those parts of Washington Sound, the Gulf of Georgia, and Puget Sound lying between the parallels of 48° 10' and 49° 20' north latitude:

62. *Close season for salmon.*—From August 25 to September 15 in each year, both days inclusive, no salmon or steelhead shall be fished for, killed, or captured for commercial purposes in these treaty waters; provided, however, that in the waters to the westward of a line drawn southward from Gonzales Point to the shore of the State of Washington silver salmon, or coho salmon, may be fished for, killed, or captured from September 1 to September 15 in each year, both days inclusive.

63. *Weekly close season for salmon and steelhead.*—From 6 o'clock Saturday morning to 6 o'clock on the Monday morning next succeeding, no salmon or steelhead shall be fished for, killed, or captured in these treaty waters.

It is, however, provided that in the waters to the westward of a line drawn southward from Gonzales Point to the shore of the State of Washington the weekly close season shall begin 12 hours earlier, and shall end 12 hours earlier.

64. *Construction of pound nets.*—All pound nets or other stationary appliances for the capture of salmon or steelhead shall be so constructed that no fish whatever shall be taken during the weekly close season. The erection or addition to the pound net of a jigger is prohibited.

65. *Location of pound nets.*—All pound nets shall be limited to a length of 2,500 feet, with an end passageway of at least 600 feet between one pound net and the next in a linear series, such distance being measured in continuation

of the line of direction of the leader of such net, and a lateral passageway of at least 2,400 feet between one pound net and the next.

On and after January 1, 1911, the mesh in pound nets shall be 4 inches in extension in the leader and not less than 3 inches in other parts of the net.

66. *Nets other than pound nets.*—No purse net shall be used within 3 miles of the mouth of any river and no seine within 1 mile of the mouth of any river in these treaty waters.

No gill net of more than 900 feet in length or of a greater depth than 60 meshes shall be used in these treaty waters.

In Alaska previous to 1906 the conditions prevailing were very similar to those in Oregon and Washington, but in that year Congress enacted a comprehensive and excellent law regulating the fisheries, the enforcement of which was entrusted to the Bureau of Fisheries. The force of agents is still inadequate, although materially increased in 1911, and its facilities for covering the territory are very meager. Conditions approaching the ideal will not prevail until these defects have been remedied; but respect for the fishery laws in Alaska obtains very generally now as a result of their persistent enforcement during the past five years.

VI. METHODS OF PREPARING SALMON.

CANNING.

EARLY DAYS OF THE INDUSTRY.

In the salmon industry canning is, and has been almost from the time of the discovery of a feasible method of so preserving the fish, the principal branch. The first canning of salmon on the Pacific coast was on the Sacramento River in 1864, when Messrs. G. W. and William Hume and Andrew S. Hapgood, operating under the firm name of Hapgood, Hume & Co., started the work on a scow at Washington, Yolo County, Cal. The Hume brothers, who came from Maine originally, had been fishing for salmon in the Sacramento River for some years before the idea of canning the fish had entered their minds, while Mr. Hapgood had previously been engaged in canning lobsters in Maine, and was induced by the Humes to participate in order that they might have the benefit of his knowledge of canning methods. The late Mr. R. D. Hume, who worked in the original cannery and later became one of the best known cannery men on the coast, thus describes the plant and the methods employed:^a

Before the arrival of Mr. Hapgood [from Maine] the Hume brothers had purchased a large scow, on which they proposed to do the canning of salmon, and had added an extension to the cabin 18 by 24 feet in area, to be used as a can-making shop. This had a shed on the side next to the river for holding any cans that might be made in advance of the packing season. A few days after the arrival of Mr. Hapgood [March 23, 1864], the tools and machinery were packed and put in position. Mr. Hapgood made some stovepipe and two or three sheet-iron fire pots, and in a short time was ready for can making. The following list of tools and machinery will show how primitive our facilities were as compared with present methods: 1 screw hand press, 1 set cast-iron top dies, 1 set cast-iron bottom dies, 1 pair squaring shears, 1 pair rotary shears, 1 pair bench shears, 1 pair hand shears or snips, 1 pair 24-inch rolls, 1 anvil (weight 50 pounds), 1 forging hammer, 1 tinner's hammer, 1 set punches for making stovepipe, 1 rivet set, 1 grooving set, 2 iron slabs grooved on one side to mold strips of solder, 1 iron clamp to hold bodies of cans while soldering the seams, 1 triangular piece of cast iron about three-eighths of an inch in thickness and 6 inches in length, with a wooden handle attached to the apex, also used for holding can bodies in place while being seamed.

^a The first salmon cannery. By R. D. Hume. *Pacific Fisherman*, vol. II, no. 1, January, 1904, p. 19-21.

The process of canning was as follows: The bodies of the cans were first cut to proper size by the 'squaring shears, a line was then scribed with a gage about three-sixteenths of an inch from one edge, and they were next formed into cylindrical shape by the rolls. They were then taken to the soldering bench, and one edge lapped by the other until the edge met the line that had been scribed and fastened there by being soldered a small part of the length to hold them in place for the further purpose of seaming. They were then placed either in the iron clamp, which had a piece of wood attached to its under side, and held firmly, the clamp being closed by the operation of a treadle, or were slipped on a piece of wood, which was bolted to the bench, while being held in place by the triangular hand seamer, which was pressed down on the lap of the seam by the left hand of the operator. When this had been done a piece of solder, which had been prepared by shaking in a can together with rosin, was placed on the seam, and melted and rubbed lengthwise of the seam. After cooling the bodies were ready for the end or bottom, which operation was brought about by first cutting out circular blanks with the rotary shears, and then placing them in the cast-iron die, and bringing the handle of the screw press around with a swing with force enough to form up the end or bottom. In this operation there were many difficulties, as the ends or bottoms would many times stick to the upper part of the die and refuse to come off, and finger nails were pretty short in those days. To get the ends out of the lower part of the die was not so bad, as a wooden plunger operated by a treadle knocked them out, but sometimes they were in pretty bad shape. When the bottoms or ends were ready they were slipped on the bodies, and the edge of the bottom rolled about in a pan of powdered rosin until the seam was well dusted. A piece of solder similar in size and preparation as used for the side seam was placed in the can. They were then placed on the smooth side of the cast-iron slabs, and the operator, with a hot soldering copper shaped to fit the circle of the can, melted the solder and, by turning the can rapidly, soldered the full circumference. The output of this can factory was very imperfect, as at least one-half of the seams burst, owing to the lack of experience of the manager or want of good judgment.

When the can making was well underway Mr. Hapgood then turned his attention to getting the apparatus for canning on board the house boat. This in the cooking department consisted of a kettle made of boiler iron about 36 inches in diameter and 5 feet in depth, set in a brick furnace and fired from underneath. Alongside was a round bottom cast-iron pot holding about 60 gallons of water and heated in the same manner. These kettles, with a dozen coolers or circular sheet-iron pans with ropes attached and with holes cut in the bottoms for drainage, a set of 5-inch blocks and tackle, with a sheet-iron fire pot and a scratch awl, completed the bathroom outfit. The can filling and soldering room was furnished with a table through the center, where cutting the salmon in pieces to suit and the filling of the cans was done. On each side of the room there was a bench running the full length, on the end of one of which the cans were placed to receive the pickle, which was used at that time instead of the small quantity of salt that is placed in the cans during the operations of these later days. After the salmon had been cleaned by removing the entrails and washing them outside the covered portion of the scow, they were brought inside and placed on the table, and a man with a butcher knife in one hand and a stick in the other, which had a mark showing the length of the pieces desired, cut gashes in the side of the salmon as a guide, and then cut the fish into sections corresponding to the length of the mark on the stick. He

then proceeded to cut the sections in pieces to suit the cans. Then three or four operators placed the salmons in the cans and shoved them along the table to where a boy wiped the top edge and passed them along to two others who placed tops which fitted inside of the rim. The cans were then taken in wooden trays to the bench opposite the starting point, which was fitted with four sheet-iron pots, and at the one nearest the entrance to the house on the scow a man put a soldering flux on the top edge, which was made by adding zinc to muriatic acid, and then with a pointed soldering copper and a stick of solder melted the solder until a small portion could be drawn around the groove formed by the edge of the can and the bevel of the top. From there the cans were taken to the other parts of the bench, where two men finished soldering the head in, and then taken to the third man, who soldered, or, as it was called, buttoned the end of the seam lap. The cooking department or bathroom, as it was called, was separated from the filling and soldering room by a partition. The cans were shoved through a hole in the partition.

At this time the process was a secret. Mr. Hapgood did the cooking and all the work done inside, no one but a member of the firm being allowed to go in. This privacy was continued until the firm moved to the Columbia River and, the labor becoming too arduous for Mr. Hapgood to perform alone, a boy by the name of Charlie Taylor was taken in as an assistant. * * *

But to return to the original proposition: When the filled cans had been soldered and entered the bathroom they were put in the coolers and lowered into the cast-iron pot, one cooler of cans being cooked at a time. The cooler was lowered into the boiling fresh water until the cans were submerged to within 1 inch of the top ends and left to cook for one hour; then they were hoisted out and the vent holes in the center of the top soldered up, after which they were dumped into the boiler-iron kettle, which held a solution of salt and water of density sufficient to produce, when boiling, a heat of 228° to 230° F. They were cooked in this solution for one hour and then taken out of the kettle with an iron scoop shaped like a dip net, with a wooden handle about 6 feet in length. They were dumped into a tank of water on the other side of the partition which separated the bathroom from the packing room through an opening in the partition, receiving many a bump and bruise in the operation. Then they were washed with soap and rag to remove the dirt and grease, each can being handled separately. When this was done they were piled on the floor of the packing room and in a few days were painted with a mixture of red lead, turpentine, and linseed oil, for at that time buyers would have no canned salmon, no matter how good the quality, unless the cans were painted red.

When packs of 10,000 to 15,000 cases were made in a season only the absolutely essential machinery was used, the rest of the work, such as cutting and cleaning the fish and placing them in the cans, being done by hand. When larger canneries were constructed, especially in Alaska, where labor is expensive and difficult to obtain, the greater part of the workmen having to be brought up from the States, machinery to do as much as possible of the work became absolutely essential. The inventive genius of the country came to the rescue and one by one machines for cutting and cleaning the fish, filling the cans, putting the tops on, and washing them, were invented and put into use, while automatic weighing machines were produced and extensive improvements and alterations were made in the machines previously in use. There are to-day many large manufacturing es-

tablishments which devote all or the greater part of their facilities to furnishing machinery and supplies to this giant branch of the salmon industry.

When salmon canning was in its infancy a pack of from 150 to 200 cases was considered a good day's work. Now it is not an uncommon occurrence for a cannery to turn out from 1,500 to 2,000 cases in one day, and there are a few which have even greater capacity.

During the height of the salmon run a cannery is an exceedingly busy and interesting place, and a description of the methods used at the present time will show the giant strides the industry has made since the days of Hapgood, Hume & Company.

HANDLING THE SALMON.

At convenient spots near the fishing grounds large scows and lighters are anchored and the fishing crews deliver their catches aboard these, the tallyman on each scow keeping a record and giving the crew a receipt. Men fishing near the cannery deliver their catch alongside. Steamers and launches are used to tow out empty scows and bring in those filled. In the old days the fish were pitched by hand into bins on the wharves, but this laborious method has been superseded by the use of an elevator, which extends from a short distance above the top of the wharf to the water's edge, provision being made for raising or lowering the lower end according to the stage of the tide. This elevator is slanting, and is made of an endless chain operating in a shallow trough. About every 2 feet there is attached to the chain a crosspiece of wood. At the top of the elevator are chutes which deliver the fish at various convenient spots on the cutting-room floor.

At a few places tracks have been run down to the low-water stage and the steamers, launches, and scows come alongside these, small cars being run down to meet them, and be filled by men pitching the fish from the boats, the cars when filled being run up into the cutting room and dumped upon the floor. At other places men armed with pews (single-tined forks) pitch the fish up to the wharf, where other men pitch them to the cutters.

If the salmon have been in the scows for from 20 to 24 hours they are used as soon as possible after being delivered at the cannery; otherwise that length of time is usually allowed to elapse, the cannerymen claiming that if not allowed to shrink the fish will be in such condition that when packed much juice will be formed, so that in "blowing," after cooking, light-weight cans will be produced. The danger of canning fish that are too fresh, however, is of minor importance as compared with the tendency in the other direction.

Before dressing the fish a stream of water is kept playing over them in order to remove the dirt and slime, after which men with pews separate the different species into piles.

DRESSING.

The majority of the canneries still use the old hand method of dressing the fish, and in such places the selection of the butchering or dressing gangs is of prime importance. Two men constitute a "butcher's gang," and the number of these gangs is dependent upon the output of the plant. Boys place the fish, with the head out, upon the cutting tables. One man cuts off the heads, and is followed by another who removes the fins, tails, and viscera. The offal is thrown into a chute, whence it passes into the water under the cannery, while the dressed fish is transferred to a tank of water, to be scaled, washed, and scraped. It is then passed to another tank of water, where it receives a second washing, scraping, and final brushing with a whisklike broom, which removes any offal, blood, and scales that were overlooked in the first washing, after which it is removed to large bins on either side of the cutting machine.

The most useful cannery inventions in recent years have been of machines for doing the work of the dressing gangs. Several have been invented and work more or less satisfactorily. The one now in general use in canneries where such machines are employed was first used in 1903 at Fairhaven (now Bellingham), Wash. It removes the head, tail, and fins and opens and thoroughly cleans the fish ready to cut into pieces for the cans. By the use of these machines the dressing gang is almost entirely done away with, dispensing with 15 to 20 men.

CUTTING.

The usual method of cutting the salmon is by a machine. This is generally a large wooden cylindrical carrier, elliptical in shape, thus having a larger carrying capacity. Ledges or rests on the outside the length of the carrier are wide enough to hold the fish, and are slit in cross section through the ledges and outer casing to receive the gang knives. The latter are circular, fixed on an axle at the proper distances apart, and revolve at the highest point reached by the carrier and independently of the latter. The carrier and gang knives are set in motion, each revolving on its own shaft. As a rest on the carrier comes to a horizontal position, men stationed at the fish bins lay a fish on each ledge as it passes. Thence it is conveyed to the revolving gang knives and, after being divided, passes through on the downward course, sliding off the rest into the filling chute. The knives in these machines are so arranged as to cut the fish transversely in sections the exact length of the cans to be filled.

The rotary cutter shunts the tail pieces to one side, and these are carried by means of a chute to baskets. But few of the larger tail pieces are canned, the rest being thrown away, this forming a con-

siderable part of, the tremendous annual waste of the salmon canneries. As the tail portion is much smaller, with less meat, it can not be placed in the cans with the middle and head sections without detracting from their value, but if packed under a distinct and separate label, as is now done in a few canneries, there is no reason why the tails should not supply the demand for a cheap grade of fish.

In some of the smaller canneries, especially in those packing flat cans, the gang knives are worked by hand. In this case the knives are not circular, but elongated or semicircular in shape, tapering at the outer ends. They are mounted on an axle having a large iron lever at one end, and when this lever is raised the ends of the gang knives are thrown up and back. The fish is then placed in position under them and the lever pulled forward, the knives, with a scimitar-like movement, dividing the fish.

The original method of cutting was by means of a long knife wielded by a Chinaman who stood at a regular butcher's block. Although his strokes were incredibly quick, the rotary cutting machine is a vast improvement over the old way.

SALTING.

Every can of salmon is seasoned with one-fourth of an ounce of salt, which, to insure uniformity, is added by mechanical means. A table is used, in the top of which are holes equal distances apart. On the under side of the top is a sheet-iron plate, with an equal number of holes, which slides in a groove at the sides, and is worked either by a hand or foot lever. Just below is an open space large enough to accommodate a tray holding 36 or 48 cans. A workman stands in front of the table and slides a tray of cans into the open space. He then throws a quantity of salt upon the table and immediately scrapes this off with a thin piece of wood, each hole being filled in the operation, and the salt being prevented from falling through by the iron plate underneath. The lever is then pressed, the iron plate moves forward until the holes in it are directly under the table top, when the salt drops through into the cans. This operation can be repeated four or five times in a minute.

FILLING THE CANS.

Most canneries now use filling machines, although a few, more particularly those packing flat and odd-sized cans, still fill by hand.

The filling machine consists of a chute with a belt to which are attached wire racks about 4 inches apart, set at an angle to prevent the salt from spilling out, into which the salted cans are fed from the floor above and pass into the machine. At the same time the divided sections of salmon pass down another chute into the mouth of what

looks like a hand coffee mill. They pass through here down a smaller chute and are forced by two dogs into a receptacle through which the plunger, or filler, passes. Here the plunger comes opposite the open mouth of the empty can, which when it reaches this point is caught by a clasp or hook and held in front of the plunger, which is immediately thrust forward through a chamber filled with salmon, cutting the fish longitudinally and at the same time filling the can. The next movement forces the can out upon a table. When running at full speed one of these machines will fill about 80 cans a minute.

On being released by the clamp the cans roll upon a long table and are picked up by a man stationed here, who strikes each one upon a square piece of lead set in the table, in order to settle the contents down into the can and for the purpose of detecting any deficiency in weight. If not quite full the cans are pushed to the other side of the table, where a man adds the quantity of fish needed, a supply of small bits being kept at hand for this purpose. Generally the cans overrun in weight, frequently as much as an ounce. Occasionally a can is weighed in order to see that the machine is in perfect adjustment.

In the hand method the fillers stand on each side of a long table with a trough running down the middle from end to end. This is filled with the cut pieces of salmon, and the fillers, usually women and children, put into the cans large pieces at first and then smaller pieces to occupy the vacant spaces.

WASHING THE CANS.

The cans are put upon an endless belt by a workman and pass from the filling-machine table to the washing machine. This is a rotating apparatus, consisting of an iron framework holding 10 rests or stands on which the cans sit. Immediately overhead are small perpendicular shafts with an iron cap, the diameter of a can, fixed to the end of each. Each can as it reaches the machine is caught by one of the washers and the cap brought down over the top, a tight-fitting flange preventing water from getting inside. Revolving rapidly as it goes, with a stream of water against it of sufficient force to remove the dirt and grease, the can is carried until the machine has revolved 180 degrees, when it is released and passes out on a belt. A more modern method is to use jets of steam for washing, while one of the latest devices is to clean the cans by a cold-air blast which strikes directly on the top edge. A set of brushes against which the cans revolve is used in a few canneries.

After being washed the cans continue on an endless belt and pass two children whose duty is to put a small piece of scrap tin on the top of each. These pieces are called "chips," are from 1½

to 2 inches, and are scraps from the sheet tin used in making the tops of the cans. The shape is of no particular importance so long as the pieces are long enough to cover the hole in the top of the can, or the cap as it is called.

CAPPING.

The endless belt delivers the can to the capping or topping machine. On reaching this the can passes under a cap holding a top, the latter being fed in through a separate aperture, and the cap immediately falls with just sufficient force to put the top on the can without injuring either. The can is then forced out from under the capper by the rotation of the machine, and the next capper is brought around to receive another can. As the cans revolve they are carried under a crimper, situated directly opposite the capper, which presses the edge firmly around the body. While one can is being topped another is being crimped, after which it rolls out upon a belt on its side, and is taken through the acid trough. Before the tops are sealed the edges must be treated with a solution of muriatic acid, which is in a glass receptacle and is applied as the cans are rolled through the acid trough on the endless belt.

SOLDERING.

For many years the tops and also all other parts of a can were soldered by hand, a long, tedious, and expensive process, which eventually gave way to the soldering machine. This is composed of an endless chain about 6 feet long, revolving around two shafts at either end of an iron trough. In the bottom of the trough is the solder, which is kept at molten heat by a row of oil blast jets underneath. Between the lower part of the chain and trough is just enough room for a can to pass without jamming, and they are forced along the trough by a chain in contact with their sides. They enter the trough at an angle, their bottoms slightly inclined, which causes the top rim to be submerged in solder, thus distributing it evenly all around the edge.

In passing through the trough the cans make about half a dozen revolutions, which cause the tops to become very hot, and it is to prevent them from being blown off by the pressure of the steam which quickly generates that the center hole in the top is made. The "chip" previously mentioned prevents the hole from being choked with salmon.

A soldering machine having, instead of the endless chain to give motion to the cans, a metal spiral running the length of the machine and revolving on an axle through the center, is used in some canneries. Each loop grasps a can and follows it to the end, thus giving the cans the proper motion and preventing them from rolling

side by side and lapping the solder over the ends, as is frequently the case with the chain machines.

A few canneries use a revolving cooler, which has a disk upon which the cans rest. This disk is filled with running water, and after it makes two revolutions the cans are forced into an inclined trough under a stream of water. The usual method, however, is for the cans on leaving the soldering machine to pass under several jets of water to set the solder and at the end of the belt to be transferred by workmen to coolers or crates, which are made of flat strap iron, square shaped, and hold about 96 cans. The crate having been filled, it is placed upon a square truck and rolled aside, where the vent holes are stopped with a drop of solder.

TESTING.

The testing tank is a square wooden tank filled with water heated almost to the boiling point by steam pipes arranged in a coil at the bottom. The crates are hoisted into the test tank by a block and tackle attached to an overhead track, which permits the coolers to be swung to any place desired.

This test is for the purpose of detecting leaks due to imperfect soldering and is conducted by two workmen skilled in this operation. The slightest leak is detected by the appearance of small bubbles issuing from the cans. The spots where the bubbles appear are marked with a small iron tool held in the hand, and the cans are taken out and placed in small wooden trays, in which they are carried to the bench men, whose duty it is to mend them. Cans that have been mended are again tested as before. The bench men are located in front of a long bench on which are numerous fire pots, supplied with oil and air led through small tubes, in which the soldering irons are kept heated, the heat and air being regulated by connecting valves. Kerosene oil and gasoline are the fuels generally used now.

COOKING.

The salmon are invariably cooked in rectangular retorts which rest in a bed and have a track running the long way. In front of each is a turntable for the purpose of receiving trucks coming from any direction. Four trucks each holding 6 crates of cans, piled one upon another, are run into the retort, which is then closed and steam turned on, entering at the bottom. The amount of pressure is from 6 to 12 pounds, the heat 250° F. In most establishments the first cooking is continued about 60 minutes.

After the first cooking the crates are taken out and placed on a long table called a "venting table," where the cans are pricked with a wooden-headed hammer fitted with a small brad, to allow the steam

and superfluous water to escape. After the venting has been done the holes are soldered up, the crates again loaded on a truck and rolled into the second retort, where they are subjected to the same pressure of steam and heat as in the first cooking and for a period of about 60 minutes.

In some canneries the retorts for first cooking are made of heavy plank, well bolted to resist the steam pressure.

In the early days much secrecy and mystery was thrown about the cooking, and the work was carried on in a separate room, known as the "bathroom," under lock and key. The first cooking was done in common tubs. The early retorts were made of wood. Later, round iron kettles were substituted, nearly one-half consisting of cover, and round crates were used for holding the cans.

For many years cannery men believed that the double cooking of salmon was absolutely necessary, but in 1898 Mr. F. A. Seufert, at his cannery on the Columbia River, at Seuferts, Oreg., a short distance above The Dalles, discarded this idea, and has since used a one-cooking method. By the new process the cans are tested for leaks after the center hole in the top is soldered up, as before, and are left in the retort 70 minutes at 245° F. and 12 pounds steam pressure. According to its originator, this method saves more than one-half the labor in the bathroom, saves nearly one-half the labor in washing the cans after cooking, and also better retains the color of the fish.

SANITARY, OR SOLDERLESS, CANS.

A recent improvement in the canning business, and one which accomplishes the same purpose as the single cooking in retorts, is that of "sanitary cans," so called. In order to use these cans a quite radical, but economical, change in machinery is necessary. As the cans leave the filling machine they pass into a steam exhauster, consisting of a box about 30 feet in length, in which are three endless-chain belts running side by side. Under and over each belt are steam coils, and under each of the lower coils are single pipes, which through small holes throw jets of live steam upon the coils, creating an intense heat. The cans pass along the first belt, are then transferred to the second belt, on which they return to the entrance of the box, whence they pass to the third belt, and continuing along this to the end pass out to the topper and crimper, the whole operation occupying five minutes' time. One style of exhauster has 10 ovals formed by the pipe, and the cans pass along these from side to side of the exhauster until discharged at the far end. By this means the contents of the can are heated and the greater part of the air exhausted, which is the object of the first cooking in the retort under the method in general use.

The topper and crimper is a circular machine with six rests for the cans. The first work performed by the machine is to "true up" the upper edge of the can, which is done by a plunger that presses the upper flange of the can upon a shoulder. In the meantime the top, which is coated around the outer edge with cement, has been automatically fed into the machine, is now clamped on the can, and by another operation is crimped on tight. The cans then leave the machine on an endless conveyer and pass to the men who transfer them to the coolers, and these are immediately placed upon the trucks and run into the retort for the one cooking they are to receive. The time they are to remain here is somewhat variable, 70 to 125 minutes with a temperature of 242° F. being the common period.

By the use of these cans the soldering machine, and in fact all use of solder and acid, is done away with, a distinct sanitary improvement, for sometimes the substances would get into the can and cause a deleterious chemical change in the contents. It also does away with the first cooking and the subsequent venting and soldering, a saving both in labor and time consumed.

REPAIRING CANS.

Imperfect cans which are repaired before the first cooking are naturally in the same condition as if there had been no defects. If the leaks are discovered after cooking and are repaired at once and the contents recooked, they are still very good, the only difficulty being that by blowing or venting them a second time they lose weight. The above goods usually go in with the regular pack of their kind and are not classed as regular "do-overs."

When, however, a cannery is running at full capacity, defective cans can not always be repaired and recooked at once and are sometimes set aside for days. Decomposition follows, of course, as with any other meat that is exposed to the air, and the fish becomes unfit for food. When recooked the meat becomes mushy and the blowing or venting makes the cans very light, a defect which is frequently corrected by adding salt water. This, the "do-over," is the lowest class of goods. In the old days, and even yet to some extent, such cans are sold without labels to brokers, or else are given some indefinite label, perhaps with the name of some fictitious cannery, and sold in the lumber, mining, or negro districts, or shipped to foreign countries with less fastidious tastes in the matter of salmon. In 1910 one of the leading companies of Alaska adopted the policy of throwing overboard all "do-overs."

On coming from the second retort the crates are lowered into a bath of lye, or, as in some canneries, the cans are run through such a bath on an endless belt, which, with the aid of a slight rinsing and a

few rubs with a brush over the top, removes from the can all the grease and other material, and then passes them into another bath where the lye is washed off in hot fresh water. The cans then go to the cooling room, where a stream of water is played upon them, or, during rainy weather are placed out of doors upon the wharf, and there allowed to cool.

The top and bottom of the cans contract in cooling, and for several hours a sharp popping noise is heard. Here, as in nearly every process through which they pass, the cans are again tested, this time by tapping the tops with a small piece of iron about 6 inches long, or, sometimes, a 12-penny nail. The sound conveys to the ear of the tester an unmistakable meaning as to the condition of the can, and the faulty cans that escape notice during the other tests are invariably found in this one.

LACQUERING.

An almost universal custom in the salmon-canning industry, but one that is not common in the canning of vegetables, fruits, etc., is that of lacquering the cans. This idea of protecting the can on the outside has been followed from the very beginning, for two reasons: (1) That the English market which, at that time especially, absorbed the greater part of these goods, insisted on their shipments being finished in this way, and (2) from the fact, as these canners speedily found out, that if they did not protect their cans in some way enormous losses through rust would ensue.

The first experiment of this nature was to paint the cans by hand with red paint, treating each singly. Next a composition of logwood extract and alcohol was tried, which, however, did not produce satisfactory results for a very plain reason—the can was dyed instead of being lacquered. The next attempt was to varnish the cans with a japan varnish reduced with alcohol, but this was found to dry too slowly for speedy handling. After extended experimentation the quick-drying brown lacquer of the present time was evolved, which carries asphaltum in the form of an asphalt varnish as its base, this being supplanted in some cases by gilsonite. This lacquer can be procured in either a heavy or light body, is generally reduced with benzine or gasoline, and is applied according to the requirements of the market, which in some localities demands a heavy coating and in others a much lighter finish, the latter giving a rich golden brown color. Some experiments have also been made in using brighter colored lacquers for this work. Several of these, made to give a bright golden, copper, or other color, are extremely attractive in appearance, while at the same time protecting the tin against rust quite as well as the brown.

The industry soon outgrew the hand method of lacquering, and the process which for a number of years was universal in the trade, and is still used by some canneries, succeeded it. For this there are a number of rectangular box vats about 40 by 80 inches and 18 inches in depth, the number varying with the capacity of the cannery. These are usually lined with galvanized metal and provided with a grid-iron-shaped iron frame, hung from a windlass or other tackle for lifting or lowering from top to bottom of the vat. The cans are loaded on this gridiron, being placed in an inclined position to allow the draining of the lacquer, and are lowered in the vat sufficiently to submerge them in the lacquer with which the vat is charged to a depth of 7 to 10 inches. The loaded gridiron is then raised to the top of the vat and the cans allowed to drain and dry before piling. This method, while being more effective in regard to the volume of work, was still of necessity a very slow and tedious operation. In damp or rainy weather, especially when it is not possible to open warehouse doors and windows, the gas arising from a number of these vats makes effective drying almost impossible.

Another principal objection to this method of lacquering, which applied also to all earlier attempts, was the impossibility of obtaining an even coat of lacquer when the can was allowed to dry in any stationary position. There was also a large waste by evaporation.

Notwithstanding repeated efforts at invention, however, it was not until 1901 that an effective machine for handling this difficult work was put on the market. The apparatus now in use by a number of canneries receives the cans on a revolving wheel fitted with rests for holding them while passing through the lacquer bath. From here they roll upon an endless chain which revolves the cans as they pass through a long box in which a hot blast dries them before they reach the end of the machine. The rotating or rolling motion given to the can after the lacquer bath, preventing the lacquer from draining to and consequently accumulating on any part of its surface, also has the effect of distributing the lacquer evenly and results in a clean and neatly finished can. The air blast facilitates the work of drying to such an extent that it requires only about two minutes after being deposited on the drying bed of the machine for the cans to be ready for handling, while the quantity of cans which can be handled in a day is vastly greater than by the old method.

A few flat and oval cans are not lacquered, but are protected from rust by wrapping in tissue paper, over which the label is placed.

LABELING.

While machines have been made for this purpose, and some of them are in use, the work is usually done by hand. A number of men

seat themselves about 4 feet apart in front of the pile of cans. Each man has in front of him a package of several hundred labels, and by bunching them on a slant so that successive margins protrude beyond each preceding, he can apply paste to the entire number with one stroke of the brush. A can is placed on the label, is quickly rolled, and the label is on much quicker than one can tell it. Each man places to his right the cans he labels, forming a pile of length and width equal to his unlabeled pile, and when the entire lot has been labeled it has been shifted only about 4 feet. Cans of fancy brands of salmon put up on the Columbia River and in the Puget Sound region are wrapped in colored tissue paper before the label is put on. Cartons similar to those used by the sardine packers would make good containers for fancy brands and would be much cheaper than the present method.

Several attempts have been made to popularize salmon packed in glass and porcelain jars, and while these have met with some favor, it was not sufficient to warrant a continuance of the practice for any length of time. None are being so packed at the present time.

BRANDS.

A very important feature of the canning industry is the selection of appropriate brands or labels for the various grades of salmon. Each company has a number of these, which it has acquired either by designing them or by absorbing another company which owned them. A well-known brand has a value in itself and sometimes is a very important asset. A company will sometimes market a considerable part of its product in one section, and here, where the consumer has become familiar with the brand and pleased with the contents of the can, he will ask for and accept no other, despite the fact that the latter might be, and probably is, the equal of the product he has been using.

Up to a few years ago one of the most serious evils in the trade was the use of misleading and lying brands. The high-grade product would almost invariably be correctly and fully branded, but "chums" and "pinks" were usually branded as "Fresh salmon," "Choice salmon," etc., which would deceive all persons but those well acquainted with the industry. "Do-overs" and very poor fish were usually marketed under a brand which bore the name of a fictitious company or of no company at all.

The passage of State laws of varying degrees of efficiency governing the branding of salmon helped slightly to remedy this condition of affairs, but it was not until the Pure Food and Drugs Act, approved June 30, 1906, was put into force by the Government that any radical improvement was noticeable. At the present time but few misleading brands are in use.

BOXING OR CASING.

A case of salmon generally contains 48 one-pound cans or their equivalent, i. e., 24 two-pound cans or 96 half-pound cans. Some canneries pack their half-pound cans in cases of 48. These cases are usually made of wood and cost from 9 to 11 cents each knocked down.

CAN MAKING.

Some of the canneries in the coast States purchase their cans ready made, but the usual method is to purchase the sheet tin and make up the cans in the canneries. This is especially necessary in Alaska, as it would be impossible to find room on the cannery ships for such a bulk as they would make in addition to the other supplies necessary. Furthermore, the making of cans provides work for a large part of the crew, otherwise unemployed while the rest are getting ready the other necessary paraphernalia. The work is done by machinery and occupies several weeks' time.

MILD CURING.

The beginning of the business of mild curing salmon, or "sweet pickling," as it is sometimes called, is of comparatively recent date.

In 1889 a German dealer came to the Columbia River and tried to interest some of the cannery men in the business. Messrs. J. O. Hanthorn, M. J. Kinney, and J. W. Cook were persuaded to prepare some, and the plant of the Northwest Cold Storage Company, at Portland, was used to keep the fish at a low temperature during repacking and preparation for shipment. These fish were shipped to Germany, but the shippers received no financial returns, word coming back that the fish were not satisfactory.

Owing to this lack of success from the first effort no further attempt was made until 1894, when Mueller & Loring, of Chicago, put up a carload of mild-cured salmon at Kalama, Wash., and shipped it to Germany. In 1896 Charles Ruckles and Wallace Brothers, of Kalama, packed several carloads for the German market. It was not until 1898 that the business was permanently established on the Columbia, the Trescott Packing Company and S. Schmidt & Sons putting up plants at Warrenton and Astoria, respectively.

In 1900 the Trescott Packing Company began packing the spring and fall runs, and the Sacramento River Packers' Association packed the fall run, on the Sacramento River, the business being carried on here every year since.

In 1901 the Sacramento River Packers' Association began at Monterey the mild curing of the spring salmon that were taken with hook and line in the open ocean.

S. Ellmore & Company started the industry in 1902 at Tillamook, and the business began on Puget Sound in 1901, when the San Juan Fishing & Packing Company and the Seattle Fish Company took it up.

Prior to 1906 several of the Alaska cannery men put up each season a few tierces of mild-cured salmon, but it was not until this time that the industry really began as such. In that year J. Lindenberger (Inc.) started packing at Ketchikan, Alaska. The following year several other plants were started, and in 1910 almost all of the king salmon taken in southeast Alaska were mild cured.

In mild curing the fish are split down the middle, the head, tail, and all fins except the pectorals removed, and the backbone cut out. The fish is then in two halves. Each of these halves, or sections, is then scored on the outside eight or nine times with the knife. They are then thrown into a cleaning vat, and here the inner side of each section is carefully scraped clear of blood and membrane with a knife, while the outside is thoroughly cleaned with a scrubbing brush. The sections are then laid carefully inner side up in another vat partly filled with clear, cold, running water, or into a tierce partly filled with fresh water and cracked ice, in which they remain for an hour. Formerly the fish were put into brine, but it has been found that ice water answers the purpose much better. After being thoroughly cooled, the sections are salted down in the tierces, each one being laid with its tail toward the center. Usually about 50 whole fish are required to fill a tierce. The fish are but lightly salted, and owing to this fact must be kept in cold storage until used.

In the early days of the industry different preparations, which included salicylic and boracic acids, were used to help preserve the fish. This caused much complaint from the Germans, and finally their Government subjected our product to a rigid inspection, with most salutary results, as now it is one of the purest and best products put up on this coast, the use of acids being done away with entirely.

The king salmon is almost invariably the species mild cured, being the only one large enough to answer the requirements of the trade. In 1907 a Ketchikan, Alaska, packer put up a quantity of coho, dog, and humpback salmon, but he found so much difficulty in disposing of the product that he abandoned further efforts in this line.

The principal consumers of the mild-cured salmon are the smokers, who take them from the tierce, wash them for a few minutes, and then have a practically fresh fish to smoke, and not, as in the days when hard-pickled salmon were used, one that had lost most of its oil and flavor through the excessive amount of salt needed to preserve it.

The greater part of the product put up on this coast goes to Europe, Germany being the principal consumer, but considerable quantities are sold in Norway, Sweden, and other countries, while the smokers of the cities east of the Rocky Mountains use large shipments every year.

PICKLING.

The earliest method of preserving salmon on the coast was by pickling. At times this industry attained to large proportions, but during the last 10 years it has been declining, largely because of the increasing popularity of mild-cured salmon. All species of salmon are pickled, but the most popular is the red salmon.

In dressing salmon for pickling the heads are removed, the fish split along the belly, the cut ending with a downward curve on the tail. The viscera and two-thirds of the backbone are removed, and the blood, gurry, and black stomach membrane scraped away. The fish are then thoroughly scrubbed and washed in cold water. They are next placed in pickling butts with about 15 pounds of salt to every 100 pounds of fish. The fish remain here about one week, when they are removed, rubbed clean with a scrub brush, and repacked in market barrels, one sack of salt being used to every three barrels of 200 pounds each. About 40 to 52 red salmon, 25 to 35 coho salmon, 70 to 80 humpback salmon, 10 to 14 king salmon, and 25 to 30 dog salmon are required in packing a barrel of pickled salmon.

A few salteries also pack "bellies." This product is merely the belly of the fish, which is the fattest portion, and as most of the packers threw away the rest of the fish, thus causing a very large waste of choice food, this method has come under the ban of the law in some of the coast States and in Alaska. As a result but few "bellies" are packed now, and most of these only when some economic use is made of the remainder. Humpback salmon furnish the major part of the "belly" pack.

DRY SALTING.

During the progress of the Russian-Japanese War the preparation of dry-salted dog salmon became an important industry, but as soon as the Japanese fishermen resumed their former occupations the demand fell off so much that the industry was virtually abandoned in the United States, although a number of Japanese continue it in British Columbia. The fish, after being dressed, were packed in boxes, in salt, these boxes holding about 560 pounds of fish, and were shipped in this condition to Japan.

At a number of places in Alaska the bellies of red and coho salmon are cut out and salted, after which the backs are dried in the sun and,

thus cured, are used for fox food at the numerous fox ranches. This product is called "ukalu."

SMOKING.

The smoking of salmon is virtually a continuation of the pickling, as the fish must be pickled before being smoked, the main purpose of the pickling being to preserve them until the time arrives for smoking, which may be weeks or months after the fish are caught. For smoking them the salmon are taken out of the barrel and soaked until as much as possible of the salt is removed. They are then put into the smokehouses and subjected to the heat and smoke of a fairly hot fire for about two days in order that they may be thoroughly dried and hardened. Exposure to a smoldering fire (alderwood is a favorite fuel) for about three days completes the process.

For shipment smoked salmon are packed in wooden boxes, oil-paper being placed between the fish.

A variation of the smoking process is known as "kippering." With this method the salmon are dried in a hot fire for about 20 hours and then smoked over another hot fire for about 24 hours. The "buckling" process is also similar to this.

Dog and king salmon are often cut into steaks and kippered. As the sale of white-meated king salmon is somewhat hampered by the whiteness, the smokers use a coloring preparation, known in the trade as Zanzibar carmine. This gives the outside of the fish a deep-colored red gloss, but leaves the inside its natural white color. The steaks are wrapped in paper and packed in baskets holding 10 pounds each.

A smoked product known locally as "beleke," is put up at Kodiak, Alaska, from red and coho salmon. Steelhead trout are the best for this purpose, but are not often utilized owing to their scarcity in this region. In preparing "beleke" only the backs of the fish are used, the belly part being cut out and pickled separately. The backs are divided into three grades, according to size, viz, "small," "medium," and "large." They are first put into a brine, the "large" being put in first, followed by the "medium" and "small" at intervals of 1 hour each, so that all will be cured at about the same time. The coho backs, being the largest, are kept in the brine from 19 to 20 hours, while the red salmon backs, which are smaller, remain in the brine only about 16 hours. After being thoroughly salted the backs are removed from the brine and rinsed in fresh water, then hung in the air for about 24 hours to dry and to allow a thin skin to form on the outside. They are then hung in the smokehouse, in the presence of a little fire of cottonwood or alder. On dry days the gable windows are thrown open and the wind allowed to

pass through while the smoking is going on. The smoking must be done slowly, two weeks being devoted to it.

There is a good demand for this product locally, the fish selling for from 15 to 20 cents a pair, but little effort has been made to extend its sale outside of central Alaska.

FREEZING.

The process of preserving fish by freezing was first introduced in 1888. Previous to this the comparatively ancient method of packing with ice, or in rare instances letting the fish freeze naturally during the winter months, was followed. Packing with ice is in quite general use to-day for shipments of fish which are to be preserved for short periods of time. Cooling with ice never results in a temperature lower than 32° F., which, of course, does not freeze the fish.

The freezing of salmon and steelhead trout began on the Sacramento and Columbia Rivers in the late eighties. It was taken up in a small way on Puget Sound in 1892. That year Wallace Bros. and Ainsworth & Dunn froze a small lot, and the venture was so successful that the next year nearly all of the wholesale dealers on the Sound took up the business. In Alaska the preparing of frozen salmon began in 1902. The San Juan Fishing & Packing Company, soon to be succeeded by the Pacific Cold Storage Company, put up a cannery and cold-storage plant at Taku Harbor, in southeast Alaska, in 1901, though it did not operate the cold-storage portion until 1902. This is the only plant which has operated in Alaska, although the New England Fish Company erected in 1909 a large plant at Ketchikan for the freezing of halibut primarily, but will probably freeze salmon also.

The freezing of salmon is almost invariably carried on in connection with other methods of handling and preserving, and the purpose is usually to secure the fish when numerous and cheap, freeze them, and then hold them until the runs are over and the fish are once more in good demand at high prices. The business proved so profitable, however, that the dealers began to look for wider markets for their product. Europe, more especially Germany, was prospected and a profitable market soon developed, with the result that to-day frozen Pacific salmon can be secured in nearly every town of any size in western Europe, while large quantities are marketed all over our own country.

There are four important features in packing and using frozen salmon: (1) To get fresh fish; (2) to keep them cold (about 15° above zero) after they are frozen; (3) to keep a coat of ice on them, and (4) to allow them to thaw slowly in cold water before cooking.

In selecting salmon for freezing only the finest and freshest of each species are used. The current belief that freezing destroys the flavor of the fish is erroneous, the flavor depending entirely upon the condition before freezing, and the quicker they are frozen after being caught the better will the natural flavor of the fish be preserved. Frozen salmon are just as wholesome as fresh, and their chemical constituents are almost identical. The danger lies in the temptation to freeze the fish after decomposition has set in, but, fortunately, this is now very rarely practiced in the salmon industry.

The coho, or silver, and the chum, or dog, salmon are the choicest of the salmons for freezing. The other species except the red, or sockeye, which is too oily and rarely frozen, are also frozen in varying quantities. The steelhead trout, which is ranked by the Pacific coast dealers among the salmon, is considered the choicest fish of all for freezing.

One of the most modern plants on the coast—that of the New England Fish Company, at Ketchikan, Alaska—has four freezers, each 25 feet by 10 feet 6 inches, in which a temperature of from 25° to 30° F. below zero can be maintained if desired, although a temperature of more than 10° below zero is rarely ever required. All freezing is by direct expansion and each freezer is piped with about 2 feet of 1¼-inch pipe per cubic foot of freezing space. The bunkers in the freezers are in pairs, each nine pipes wide, spaced 10 inches apart. This leaves a 3½-foot passage through the center of each freezer opposite the 3½ by 6½ foot swing doors. The salmon are laid on pans, which are placed on the tiers of pipes.

After freezing, the salmon are passed through openings in the rear of the freezers into the glazing room, which has a temperature of about 20°F., where they are dipped into water, and when removed are covered with a thin glaze of ice, which may be thickened by repeated dippings. This is an extra precaution to exclude the air from the fish.

After being thoroughly frozen and glazed, each fish is covered first with a parchment, like rolls of butter, and then with a piece of heavy brown paper. They are then packed in boxes holding about 250 pounds each, placed in the cold-storage cars and shipped.

MISCELLANEOUS PRODUCTS.

A few years ago a company on the Columbia River put up what was known as "fish pudding." In preparing this the salmon was ground fine, mixed with milk and eggs, and then packed in tin cans. The preparation was soon abandoned.

In 1903 one of the Point Roberts canneries packed a new product which was called "salmon paste." For this the fish was ground up,

cooked, seasoned with spices, etc., and made into fish balls, a very palatable dish when warmed over.

In 1905 a Seattle concern began the manufacture of wienerwurst sausages from halibut and salmon.

The Indians in the Bristol Bay region of Alaska occasionally dress the skins of salmon and make of them leather for the tops of boots, also bags and other small articles.

Every year immense quantities of salmon roe are thrown away in the fisheries of the west coast, though there is but little doubt that, if properly prepared, a market could be found for this now waste part of the fish. In France there is a good market for a product known as "rogue," which is the spawn of cod, haddock, hake, and pollock salted in casks, and which is used as bait in the sardine fisheries. Salmon spawn is the choicest and most successful bait used on this coast, and if properly prepared would undoubtedly answer the purpose as well, if not better, owing to its oiliness and attractive color, than the regular "rogue." The roes should be soaked for some days in old brine and then packed in strong casks holding about 25 gallons each. It might also prove to be a good bait for tolling mackerel on the Atlantic coast.

In 1910 a considerable quantity of salmon roe was prepared in Siberia and sold in competition with caviar, which is prepared from sturgeon eggs. The product met with favor in Europe and several Alaska firms are preparing to put it up in 1912. It should be prepared in the same manner as caviar.

Several establishments are putting up these eggs in jars and hermetically sealed cans for use as bait in sport fishing.

A product which was first made in Norway is prepared by means of an invention which quickly dries and pulverizes the flesh of fresh fish. The resulting powder, called "fish flour," is easy to transport from one place to another and has great nutritive value. It is probable that the tailpieces of the fish, which are at present thrown away, and the cheaper grades of salmon might be prepared in this way and thus furnish another market for salmon.

OIL AND FERTILIZER.

As early as 1888 there was a small plant at Astoria, Oreg., where the refuse of the canneries was utilized for the manufacture of oil and fertilizer. In that year 8,000 gallons of oil (chiefly from salmon heads), and 90 tons of fertilizer were prepared. The oil was worth 22½ cents per gallon and the fertilizer had a market value of \$20 per ton. Most of the refuse was dumped into the river, however. In 1898 a similar plant was established in the Puget Sound district

of Washington, but for some reason the industry has languished almost from the start.

In 1882 the Alaska Oil & Guano Company established a fertilizer plant at Killisnoo, Alaska, for the extraction of oil and fertilizer from herring, and has operated the plant continuously ever since. In some years large quantities of whole salmon have been handled at this plant, and the resulting product was found to sell as well as that from herring.

Probably the most serious evil in the salmon industry to-day is the enormous wastage which annually occurs. About one-fourth of the total weight of each fish handled at the various packing plants is thrown away. With the exception of the tailpiece, which is discarded at most canneries owing to the excessive amount of bone which would be in the product if canned, this waste material could not be utilized as food, comprising as it does the head, viscera, fins, and tail. When not conveniently near the very few fertilizer plants at present in operation this product is either allowed to pass through chutes into the water under the cannery, or is dumped into scows and towed to the ocean or the deeper waters of the sounds, and here thrown overboard. This procedure, not only exceedingly wasteful, is also far from beneficial to the waters where deposited.

The great desideratum in the salmon fisheries of the Pacific coast at the present time is the invention of a small odorless-fertilizer plant, costing not more than \$2,500 or \$3,000, which can be installed at the various salmon canneries and salteries. The offal from the cannery could there be utilized and the product obtained would doubtless net a fair return on such an investment, while at the same time the present (in the aggregate) enormous waste would be stopped, and the waters adjacent to the canneries rendered far more agreeable to the fishes as well as to the people on shore. It is absolutely essential that the plant shall be odorless, as the smell of the ordinary fertilizer establishment would be very offensive to persons visiting the cannery and would not enhance the demand for canned salmon. At the present time the cheapest plant available costs about \$10,000, and very few canneries can afford to invest this sum of money in the disposal of their own offal alone.

VII. STATISTICS OF THE PACIFIC SALMON INDUSTRY IN 1909.

This is the first report in which detailed statistics of the salmon fisheries of Washington, Oregon, California, and Alaska have been shown for the same year. Partial statistics of British Columbia and Yukon Territory of the Dominion of Canada are also included.

PERSONS EMPLOYED.

The large army of 28,945 men, women, and children were employed in the salmon fisheries of Alaska and the three coast States. Alaska leads with 11,433, followed by Washington, Oregon, and California in the order named. Over two-thirds of the grand total is made up of whites. The Chinese and Japanese have almost the same number, while 2,803 Indians were employed.

PERSONS ENGAGED IN THE SALMON FISHERIES OF THE PACIFIC COAST STATES AND ALASKA IN 1909.

Occupation and race.	Alaska.	Washing- ton.	Oregon.	Califor- nia.	Total.
Fishermen:					
Whites.....	2,486	4,426	4,179	2,114	13,205
Indians.....	1,176	221			1,397
Chinese.....				15	15
Japanese.....	13			168	181
Total.....	3,675	4,647	4,179	2,297	14,798
Shoresmen:					
Whites.....	1,911	2,091	404	276	4,682
Indians.....	1,246	115		15	1,376
Chinese.....	1,992	1,270	411		3,673
Japanese.....	2,136	1,102	256	5	3,499
Total.....	7,285	4,578	1,071	296	13,230
Transporters:					
Whites.....	443	292	70	82	887
Indians.....	30				30
Total.....	473	292	70	82	917
Total:					
Whites.....	4,840	6,809	4,653	2,472	18,774
Indians.....	2,452	336		15	2,803
Chinese.....	1,992	1,270	411	15	3,688
Japanese.....	2,149	1,102	256	173	3,680
Grand total.....	11,433	9,517	5,320	2,675	28,945

INVESTMENT.

The total investment in the salmon fisheries was \$25,157,813, of which Alaska furnishes more than one-half. Gill nets are the principal form of apparatus in use, followed by stationary traps, or pounds, diver nets, haul seines, purse seines, etc.

INVESTMENT IN THE SALMON FISHERIES OF THE PACIFIC COAST STATES AND ALASKA
IN 1909.

Items.	Alaska.		Washington.		Oregon.	
	Number.	Value.	Number.	Value.	Number.	Value.
Transporting vessels:						
Power vessels.....	133	\$1,067,944	93	\$40,500	30	\$119,900
Tonnage.....	5,891		1,158		288	
Outfit.....		266,986		135,625		25,350
Sailing vessels.....	43	1,085,400				
Tonnage.....	59,761					
Outfit.....		108,540				
Power boats.....	17	24,840	5	3,550	15	28,900
Fishing boats, power.....	60	30,000	464	472,650	287	139,600
Fishing boats, sail and row.....	1,821	211,671	2,244	128,945	1,890	224,545
Scows and house boats.....	310	171,005	398	168,673	114	45,050
Pile drivers.....	43	90,555	62	124,350	2	1,800
Apparatus, shore fisheries:						
Purse seines.....	98	27,188	101	44,150		
Haul seines.....	94	27,731	246	28,955	48	16,280
Gill nets, drift.....	1,209	111,756	1,620	168,831	2,818	523,331
Gill nets, set.....			1,624	37,259	1,122	27,614
Diver nets.....			48	10,160	418	22,375
Traps, stationary.....	73	130,794	525	1,324,968	21	25,750
Traps, floating.....	15	21,250	1	2,000		
Reef nets.....			9	4,500		
Wheels, stationary.....			13	76,000	26	313,000
Wheels, scow.....			3	8,500	9	22,000
Spears.....	20	30				
Lines, trolling.....		523		261		
Lines, hand.....						
Shore and accessory property.....		5,601,259		1,730,030		1,554,780
Cash capital.....		4,970,799		1,424,500		551,500
Total.....		13,948,271		6,334,807		3,641,775

Items.	California.		Total.	
	Number.	Value.	Number.	Value.
Transporting vessels:				
Power vessels.....	4	\$37,748	260	\$1,666,092
Tonnage.....	56		7,393	
Outfit.....		3,920		431,881
Sailing vessels.....			43	1,085,400
Tonnage.....			59,761	
Outfit.....				108,540
Power boats.....	41	63,300	78	120,990
Fishing boats, power.....	171	91,050	982	733,300
Fishing boats, sail and row.....	1,158	128,245	7,113	693,406
Scows and house boats.....	50	13,925	872	398,653
Pile drivers.....			107	216,705
Apparatus, shore fisheries:				
Purse seines.....			a 199	71,338
Haul seines.....	47	5,650	b 435	78,616
Gill nets, drift.....	1,086	167,570	c 6,733	971,488
Gill nets, set.....			d 2,746	64,873
Diver nets.....			e 406	32,535
Traps, stationary.....			619	1,481,512
Traps, floating.....			16	23,250
Reef nets.....			9	4,500
Wheels, stationary.....			39	389,000
Wheels, scow.....			12	30,500
Spears.....			20	30
Lines, trolling.....		1,149		1,933
Lines, hand.....		10		10
Shore and accessory property.....		497,393		9,383,462
Cash capital.....		223,000		7,169,799
Total.....		1,232,960		25,157,813

a Aggregate length of 104,570 yards.
b Aggregate length of 111,558 yards.
c Aggregate length of 2,356,847 yards.

d Aggregate length of 151,655 yards.
e Aggregate length of 65,800 yards.

PRODUCTS.

The total products amount to 365,336,482 pounds, which returned the fishermen \$7,224,024. Bluebacks, sockeyes, or red salmon were most numerous in Alaska and Washington, chinooks in California, coho or silver, dog or chum, and steelhead trout in Washington, while humpbacks were taken commercially in Alaska and Washington alone, being especially numerous in Alaska.

PRODUCTS OF THE SALMON FISHERIES OF ALASKA AND THE PACIFIC COAST STATES
IN 1909.

Species.	Alaska.		Washington.		Oregon.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Blueback, sockeye or red.....	116,014,486	\$1,029,079	77,280,989	\$2,835,666	844,324	\$34,703
Chinook, king or spring.....	8,959,544	151,984	11,016,476	604,906	13,952,814	736,456
Coho, silver or white.....	3,526,404	41,233	21,328,466	554,157	5,184,520	127,204
Dog or chum.....	9,456,048	15,583	25,520,426	164,300	699,348	3,818
Humpback or pink.....	37,965,928	95,065	17,495,586	46,187
Steelhead trout.....	11,650	400	2,427,251	130,486	1,510,285	66,802
Total.....	175,934,060	1,333,344	155,069,194	4,335,702	22,191,291	968,983

Species.	California.		Total.	
	Pounds.	Value.	Pounds.	Value.
Blueback, sockeye or red.....	21,000	\$689	194,160,799	\$3,900,137
Chinook, king or spring.....	11,962,248	580,094	45,891,082	2,073,440
Coho, silver or white.....	145,500	4,575	30,184,890	727,169
Dog or chum.....	4,200	84	35,680,022	183,785
Humpback or pink.....	55,461,514	141,252
Steelhead trout.....	8,989	553	3,958,175	198,241
Total.....	12,141,937	585,995	365,336,482	7,224,024

NOTE.—In addition to the above, British Columbia produced 89,852,089 pounds, which returned the fishermen \$1,832,573, and the Yukon Territory (Yukon River), 80,565 pounds, which returned the white fishermen \$10,209.

PRODUCTS CANNED.

In order to show the total pack of the Pacific coast of the North American Continent, the pack of British Columbia has been included. The total pack reduced to a common basis of forty-eight 1-pound cans amounted to 5,392,306½ cases, valued at \$25,518,669. Alaska leads in the total pack, with Washington second. Alaska also leads in the pack of sockeyes, humpbacks, and chums. Washington leads in the pack of cohoes and Oregon in the pack of chinooks and steelhead trout.

SALMON CANNED IN ALASKA, BRITISH COLUMBIA, WASHINGTON, OREGON, AND CALIFORNIA IN 1909.

Products.	Alaska.		British Columbia.		Washington.	
	Cases.	Value.	Cases.	Value.	Cases.	Value.
Chinook, king, or spring:						
½-pound flat.....			360	\$1,440	23,550	\$98,780
1-pound flat.....			1,214	7,314	40,730	268,849
1-pound flat exports.....					606	4,242
½-pound tall.....			176	516		
1-pound tall.....	48,034	\$207,624	17,613	94,110	21,426	116,593
½-pound oval.....						
1-pound oval.....			444	2,886	1,110	10,212
2-pound nominal.....						
Total.....	48,034	207,624	19,807	106,266	87,422	498,676
Coho, silver, or silverside:						
½-pound flat.....			2,132	5,969	34,292	94,417
1-pound flat.....	1,206	5,543	5,911	28,373	28,885	134,755
1-pound tall.....	55,350	225,486	61,520	258,400	137,008	570,030
2-pound nominal.....					427	2,562
Total.....	56,556	231,029	69,563	292,742	200,612	801,764
Chum, or dog:						
½-pound flat.....					1,300	1,950
1-pound flat.....					219	591
1-pound tall.....	120,712	274,110	16,573	39,775	83,664	197,932
Total.....	120,712	274,110	16,573	39,775	85,183	200,473
Humpback, or pink:						
1-pound flat.....			2,267	6,234	2,030	5,585
1-pound tall.....	464,873	1,114,839	27,722	66,581	368,963	896,757
Total.....	464,873	1,114,839	29,989	72,815	370,993	902,342
Sockeye, blueback, or red:						
½-pound flat.....	16,385	63,888	483,760	1,935,040	229,502	927,967
1-pound flat.....	85,193	236,609	314,706	1,888,236	456,712	2,746,667
½-pound tall.....			12,880	42,504		
1-pound tall.....	1,611,916	7,310,053	277,893	1,500,623	487,479	2,558,993
½-pound oval.....			17,650	75,013		
1-pound oval.....			406	2,639		
1-pound squats.....			8,312	49,872		
Total.....	1,713,494	7,610,550	1,115,607	5,493,927	1,173,693	6,233,627
Steelhead trout:						
½-pound flat.....					945	2,937
1-pound flat.....					3,794	19,422
1-pound tall.....					3,897	22,602
Total.....					8,636	44,961
Grand total.....	2,403,669	9,438,152	1,251,539	6,005,525	1,926,539	8,681,843

SALMON CANNED IN ALASKA, BRITISH COLUMBIA, WASHINGTON, OREGON, AND CALIFORNIA IN 1909—Continued.

Products.	Oregon.		California.		Total.	
	Cases.	Value.	Cases.	Value.	Cases.	Value.
Chinook, king, or spring:						
½-pound flat.....	69,557	\$289,534			93,467	\$389,754
1-pound flat.....	54,591	396,809	5,663	\$28,315	102,198	701,287
1-pound flat exports.....					606	4,242
½-pound tall.....					176	516
1-pound tall.....	23,057	148,815			110,130	567,142
½-pound oval.....	534	2,670			534	2,670
1-pound oval.....	848	8,242			2,402	21,340
2-pound nominal.....	458	1,833			458	1,833
Total.....	149,045	847,903	5,663	28,315	309,971	1,688,784
Coho, silver, or silverside:						
½-pound flat.....	20,331	56,928			56,755	157,314
1-pound flat.....	11,755	51,702			47,757	220,373
1-pound tall.....	39,326	157,856			293,204	1,211,802
2-pound nominal.....	315	945			742	3,507
Total.....	71,727	267,461			398,458	1,592,996
Chum, or dog:						
½-pound flat.....					1,300	1,950
1-pound flat.....					219	591
1-pound tall.....	9,225	21,218			230,174	533,035
Total.....	9,225	21,218			231,693	535,576
Humpback, or pink:						
1-pound flat.....					4,297	11,819
1-pound tall.....	55	132			861,613	2,078,309
Total.....	55	132			865,910	2,090,128
Sockeye, blueback, or red:						
½-pound flat.....	32,071	133,095			761,718	3,059,990
1-pound flat.....	6,645	39,870			863,256	4,911,382
½-pound tall.....					12,880	42,504
1-pound tall.....	50	320			2,377,338	11,369,989
½-pound oval.....					17,650	75,013
1-pound oval.....					406	2,639
1-pound squats.....					8,312	49,872
Total.....	38,766	173,285			4,041,560	19,511,389
Steelhead trout:						
½-pound flat.....	7,064	22,084			8,009	25,021
1-pound flat.....	1,365	7,695			5,159	27,117
1-pound tall.....	4,320	25,056			8,217	47,658
Total.....	12,749	54,835			21,385	99,796
Grand total.....	281,567	1,364,834	5,663	28,315	5,868,977	25,518,669

^a All 1-pound cases contain forty-eight 1-pound cans; the ½-pound cases contain forty-eight ½-pound cans. Reduced to a common basis of cases containing forty-eight 1-pound cans, the aggregate pack amounts to 5,392,306½ cases.

MISCELLANEOUS PRODUCTS.

The total miscellaneous secondary products prepared amounted to 29,808,129 pounds, valued at \$2,096,030. Of these the largest quantity and value is represented in the mild-cured pack. The pickled pack is second in quantity but is exceeded in value by the frozen pack. Alaska leads Washington very slightly in the quantity of products prepared, but both are exceeded in value of products by Oregon.

MISCELLANEOUS SECONDARY PRODUCTS PREPARED IN ALASKA AND THE PACIFIC COAST STATES IN 1909.

Products.	Alaska.		Washington.		Oregon.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Frozen:						
Chinook, king, or spring.....			74,183	\$7,418	14,000	\$1,400
Coho, silver, or silverside.....	35,721	\$1,072	528,477	30,149	216,175	13,868
Dog, or chum.....	77,882	1,558	1,364,672	67,161		
Humpback, or pink.....			62,945	1,888		
Steelhead trout.....	9,450	473	504,165	46,615	1,446,685	144,658
Total.....	123,053	3,103	2,534,442	153,231	1,676,860	159,926
Mild-cured:						
Chinook, king, or spring.....	1,833,600	149,300	2,292,800	273,826	4,365,442	434,825
Pickled:						
Chinook, king, or spring.....	88,200	3,798	1,000	540	400	24
Chinook bellies.....	7,000	175	6,750	671		
Coho, silver, or silverside.....	63,600	2,485			2,600	130
Coho bellies.....	227,750	3,843				
Dog, or chum.....	7,000	190	50,000	175		
Humpback, or pink.....	311,400	9,405	1,615,000	48,450		
Humpback backs.....	11,200	224				
Humpback bellies.....	169,480	7,396	172,400	8,620		
Sockeye, blueback, or red.....	5,301,500	167,298				
Sockeye bellies.....	783,600	13,902				
Total.....	6,970,730	208,716	1,845,150	58,456	3,000	154
Dry-salted and dried:						
Chinook, king, or spring.....	800	45				
Coho, silver, or silverside, backs..	14,500	549				
Dog, or chum.....	71,600	1,038				
Humpback backs.....	51,500	545				
Sockeye, blueback, or red, backs.	83,000	2,302				
Total.....	221,400	4,479				
Smoked:						
Chinook, king, or spring.....			30,165	2,413	127,700	19,155
Chinook, white-meated, kippered.....			190,500	16,050		
Coho, silver, or silverside.....			30,000	1,800	20,000	2,000
Coho backs.....	4,000	400				
Dog, or chum.....	585	43	517,245	25,862		
Dog, kippered.....			5,000	500		
Humpback backs, kippered.....			100,000	5,000		
Sockeye, blueback, or red, backs.	40,300	2,780				
Total.....	44,885	3,223	872,910	51,625	147,700	21,155
Fertilizer.....	159,224	2,287	1,210,000	18,610		
Oil.....	120,113	3,216	380,648	14,161		
Grand total.....	9,473,005	374,324	9,135,950	569,909	6,193,002	616,060

MISCELLANEOUS SECONDARY PRODUCTS PREPARED IN ALASKA AND THE PACIFIC COAST STATES IN 1909—Continued.

Products.	California.		Total.	
	Pounds.	Value.	Pounds.	Value.
Frozen:				
Chinook, king, or spring.....			88,183	\$8,818
Coho, silver, or silverside.....			780,373	45,089
Dog, or chum.....			1,442,551	68,719
Humpback, or pink.....			62,945	1,888
Steelhead trout.....			1,960,300	191,746
Total.....			4,334,355	316,260
Mild-cured:				
Chinook, king, or spring.....	4,887,962	\$520,468	13,379,804	1,378,419
Pickled:				
Chinook, king, or spring.....			89,600	4,362
Chinook bellies.....			13,750	846
Coho, silver, or silverside.....			66,200	2,615
Coho bellies.....			227,750	3,843
Dog, or chum.....			57,000	365
Humpback, or pink.....			1,926,400	57,855
Humpback backs.....			11,200	224
Humpback bellies.....			341,880	16,016
Sockeye, blueback, or red.....			5,301,500	167,298
Sockeye bellies.....			783,600	13,902
Total.....			8,818,880	267,326
Dry-salted and dried:				
Chinook, king, or spring.....			800	45
Coho, silver, or silverside backs.....			14,500	549
Dog, or chum.....			71,600	1,038
Humpback backs.....			51,500	545
Sockeye, blueback, or red, backs.....			83,000	2,302
Total.....			221,400	4,479
Smoked:				
Chinook, king, or spring.....	110,550	14,643	268,415	36,211
Chinook, white-meated, kippered.....			190,500	16,050
Coho, silver or silverside.....	7,660	626	57,660	4,426
Coho backs.....			4,000	400
Dog, or chum.....			517,830	25,905
Dog, kippered.....			5,000	500
Humpback backs, kippered.....			100,000	5,000
Sockeye, blueback, or red, backs.....			40,300	2,780
Total.....	118,210	15,269	1,183,705	91,272
Fertilizer.....			1,369,224	20,897
Oil.....			a 500,761	17,377
Grand total.....	5,006,172	535,737	29,808,129	2,096,030

a Represents 66,728 gallons.

WASHINGTON.

Owing to the quadrennially heavy run of sockeye salmon and the biennial run of humpback salmon into Puget Sound occurring in 1909, the catch of both species of salmon was very heavy. The purse seiners made exceptionally heavy catches of sockeye salmon, while the traps had so many humpbacks in them that the greater part were turned out, it being impossible to find a market for them. In many places people were allowed to take away with them, free of charge, as many humpbacks as they wished.

In Grays Harbor the run of salmon was fairly good. On the Quinault River the Indians made very successful catches. Early in the season a meeting of the tribe was held, and it was decided that a

50-foot runway in the center of the stream should be kept clear of nets so as to allow the fish an opportunity to reach the spawning beds in the lake.

In Willapa Harbor the run was fair.

On the Columbia River the catch was not as large as in 1908, which was due partly to the shortening of the open fishing season.

STATISTICS BY COUNTIES.

Persons employed.—The total number of persons employed was 9,517, of which the large majority were whites.

PERSONS EMPLOYED IN THE SALMON FISHERIES OF WASHINGTON, BY COUNTIES AND NATIONALITIES, IN 1909.

Counties.	Fishermen.			Shoresmen.				
	Whites.	Indians.	Total.	Whites.	Chinese.	Japanese.	Indians.	Total.
Whatcom.....	643		643	1,056	631	488	55	2,230
San Juan.....	193	12	205	42	40	40		122
Skagit.....	303		303	569	290	414	40	1,313
Island.....	273		273	2				2
Snohomish.....	284		284	6				6
King.....	527		527	55				55
Pierce.....	276		276	12				12
Thurston.....	50		50					
Mason.....	67		67					
Kitsap.....	241		241					
Clallam.....	56	176	232	63	20	12		95
Jefferson.....	68		68	163	70	50	20	303
Chehalis.....	112	33	145	16	45	15		76
Pacific.....	616		616	18	40	20		78
Wahkiakum.....	533		533	80	134	63		277
Cowlitz.....	61		61	6				6
Clarke.....	13		13					
Skamania.....	82		82	2				2
Klickitat.....	28		28	1				1
Total.....	4,426	221	4,647	2,091	1,270	1,102	115	4,578

Counties.	Trans- porters.	Total employed.				Grand total.
	Whites.	Whites.	Chinese.	Japanese.	Indians.	
Whatcom.....	129	1,828	631	488	55	3,002
San Juan.....	9	244	40	40	12	336
Skagit.....	75	947	290	414	40	1,691
Island.....		275				275
Snohomish.....		290				290
King.....	19	601				601
Pierce.....	2	290				290
Thurston.....		50				50
Mason.....		67				67
Kitsap.....		241				241
Clallam.....	6	125	20	12	176	333
Jefferson.....	12	243	70	50	20	383
Chehalis.....	3	131	45	15	33	224
Pacific.....	11	645	40	20		705
Wahkiakum.....	25	638	134	63		835
Cowlitz.....	1	68				68
Clarke.....		13				13
Skamania.....		84				84
Klickitat.....		29				29
Total.....	292	6,809	1,270	1,102	336	9,517

Investment, apparatus, etc.—The total investment in the fisheries amounted to \$6,334,807. Whatcom County has the largest investment, nearly one-third of the total.

INVESTMENT IN THE SALMON FISHERIES OF WASHINGTON, BY COUNTIES, IN 1909.

Items.	Whatcom.		San Juan.		Skagit.		Island.		Snohomish.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Transporting vessels:										
Power vessels.....	36	\$192,500	3	\$15,500	18	\$108,900				
Tonnage.....	517		48		293					
Outfit.....		60,500		4,600		32,400				
Fishing boats, power.....	40	59,850	8	21,250	43	37,250	22	\$13,900	26	\$16,400
Fishing boats, sail and row.....	247	8,210	73	3,190	207	7,410	85	3,210	203	6,380
Scows and house boats..	188	101,350	47	15,833	31	9,150	63	18,200	17	3,800
Pile drivers.....	13	61,000	5	23,600			2	9,000		
Apparatus, shore fisheries:										
Purse seines.....	9	3,900	7	2,550	4	1,500	1	500	4	1,000
Haul seines.....	9	1,100			17	1,285	27	2,010	20	3,005
Gill nets, drift.....	71	12,250	3	225	338	26,270	1	300	130	1,036
Gill nets, set.....	96	6,200	18	310	336	5,700	1	10	537	6,317
Trap nets, stationary	72	372,540	23	116,178	12	46,500	29	176,500	8	35,000
Reef nets.....	2	1,000	7	3,500						
Lines, trolling.....										15
Shore and accessory property.....		600,003		37,350		382,044		5,250		6,245
Cash capital.....		679,000		45,000		309,000				
Total.....		2,159,403		289,086		967,409		228,880		79,198

Items.	King.		Pierce.		Thurston.		Mason.		Kitsap.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Transporting vessels:										
Power vessels.....	7	\$23,300	1	\$2,500						
Tonnage.....	56		5							
Outfit.....		11,400		800						
Power boats.....			1	200						
Fishing boats, power.....	60	107,900	23	60,200	1	\$2,500	4	\$3,800	26	\$36,900
Fishing boats, sail and row.....	234	7,350	88	3,300	27	880	29	1,310	85	3,055
Scows and house boats..									6	850
Pile drivers.....									1	2,000
Apparatus, shore fisheries:										
Purse seines.....	37	18,500	22	8,500	2	1,000	1	500	12	5,700
Haul seines.....	52	4,650	25	1,950	8	600	13	1,025	36	2,930
Gill nets, drift.....	193	8,760	73	1,900	4	100	1	300	7	1,950
Gill nets, set.....	82	820	143	3,600	16	400	22	600	8	88
Trap nets, stationary									4	13,500
Shore and accessory property.....		166,800		15,375		200		300		850
Cash capital.....		60,000		5,000						
Total.....		409,480		103,325		5,680		7,835		67,823

INVESTMENT IN THE SALMON FISHERIES OF WASHINGTON, BY COUNTIES, IN 1909—Continued.

Items.	Clallam.		Jefferson.		Chehalis.		Pacific.		Wahkiakum.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Transporting vessels:										
Power vessels.....	3	\$12,000	4	\$29,000	1	\$3,000	6	\$16,700	13	\$36,100
Tonnage.....	27		50		8		48		101	
Outfit.....		4,000		6,100		500		4,315		10,660
Power boats.....	1	1,500					2	1,800	1	450
Fishing boats, power....	2	1,600	5	2,200	5	2,500	88	46,800	72	43,500
Fishing boats, sail and row.....	212	9,580	29	940	115	8,350	317	22,820	191	38,735
Scows and house boats..	2	1,000	16	5,050	1	400	9	3,300	16	8,990
Pile drivers.....			1	5,000	3	450	37	23,300		
Apparatus, shore fisheries:										
Purse seines.....							2	500		
Haul seines.....	8	900	11	800			2	350	11	5,500
Gill nets, drift.....	70	700	5	340	100	8,000	207	36,000	417	70,700
Gill nets, set.....	8	100	25	430	189	9,724	46	1,340	33	615
Trap nets, stationary.....			2	8,000	15	3,400	280	506,400	52	36,800
Trap nets, floating.....			1	2,000						
Lines, trolling.....		246								
Shore and accessory property.....		20,325		60,345		36,753		59,625		310,455
Cash capital.....		20,000		50,000		20,000		38,000		190,500
Total.....		71,951		170,205		93,077		761,250		753,005

Items.	Cowlitz.		Clarke.		Skamania.		Klickitat.		Total.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Transporting vessels:										
Power vessels.....	1	\$1,000							93	\$440,500
Tonnage.....	5								1,158	
Outfit.....		350								135,625
Power boats.....									5	3,950
Fishing boats, power....	28	11,700	1	\$400	10	\$4,000			464	472,650
Fishing boats, sail and row.....	29	1,200	12	360	46	1,945	15	\$720	2,244	128,945
Scows and house boats..					2	750			398	168,673
Pile drivers.....									62	124,350
Apparatus, shore fisheries:										
Purse seines.....									a 101	44,150
Haul seines.....	1	150			4	1,500	2	1,200	b 246	28,955
Gill nets, drift.....									c 1,620	168,831
Gill nets, set.....	18	350	12	180	32	455	2	20	d 1,624	37,259
Diver nets.....	29	5,300	6	1,210	13	3,650			e 48	10,160
Trap nets, stationary.....	21	8,400			3	750	4	1,000	525	1,324,968
Trap nets, floating.....									1	2,000
Reef nets.....									9	4,500
Wheels, stationary..					10	44,000	3	32,000	13	76,000
Wheels, scow.....					2	7,000	1	1,500	3	8,500
Lines, trolling.....										261
Shore and accessory property.....		21,800		10		5,075		1,225		1,730,030
Cash capital.....		8,000								1,424,500
Total.....		58,250		2,160		69,125		37,665		6,334,807

a Aggregate length of 68,900 yards.

b Aggregate length of 44,824 yards.

c Aggregate length of 429,115 yards.

d Aggregate length of 92,030 yards.

e Aggregate length of 19,200 yards.

Products.—The total catch amounted to 155,069,194 pounds, valued at \$4,335,702. Whatcom County leads in the catch. Sockeye salmon constitute about one-half of the total catch.

PRODUCTS OF THE SALMON FISHERIES OF WASHINGTON, BY APPARATUS, SPECIES, AND COUNTIES, IN 1909.

Apparatus and species.	Whatcom.		San Juan.		Skagit.		Island.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
PURSE SEINES.								
Chinook, or king.....	37,568	\$1,514	24,094	\$840	12,000	\$540	2,000	\$100
Coho, or silver.....	346,000	8,880	280,008	7,000	200,000	5,000	42,000	1,050
Dog, or chum.....	496,000	2,480	280,000	1,400	160,000	800	112,000	560
Humpback, or pink.....			175,000	350	30,000	300		
Sockeye, or blueback.....	1,146,000	43,600	973,000	35,000	650,000	26,000	140,000	5,666
Steelhead trout.....					2,000	100	800	40
Total.....	2,025,568	56,474	1,732,102	44,590	1,054,000	32,740	296,800	7,416
HAUL FINES.								
Chinook, or king.....					154,400	7,060		
Coho, or silver.....	21,000	630			110,000	2,750	560,000	16,800
Dog, or chum.....	39,000	195			590,000	2,950	1,280,000	7,710
Humpback, or pink.....	14,000	35			20,000	50		
Steelhead trout.....	7,000	350			2,428	121		
Total.....	81,000	1,210			876,828	12,931	1,840,000	24,510
GILL NETS.								
Chinook, or king.....	22,332	967	47,300	1,880	617,362	25,753		
Coho, or silver.....	1,122,000	29,200	79,200	1,980	662,376	20,873	1,500	45
Dog, or chum.....	70,000	350	4,800	24	673,838	3,573	3,000	30
Humpback, or pink.....					17,800	221		
Sockeye, or blueback.....	1,328,450	51,158	44,500	1,780	384,750	12,510	30,000	1,200
Steelhead trout.....					124,200	8,004		
Total.....	2,542,782	81,675	175,800	5,664	2,480,326	70,934	34,500	1,275
REEF NETS.								
Chinook, or king.....	5,000	250	40,000	2,000				
Coho, or silver.....	27,000	810	109,000	3,270				
Dog, or chum.....	6,000	50	90,000	450				
Sockeye, or blueback.....	75,000	3,000	290,000	11,600				
Total.....	113,000	4,110	529,000	17,320				
TRAP NETS.								
Chinook, or king.....	1,378,391	66,229	574,072	25,697	354,929	18,270	1,272,680	111,735
Coho, or silver.....	3,387,624	73,940	718,124	17,967	482,116	12,271	1,615,314	42,876
Dog, or chum.....	570,412	2,852	229,408	1,148	1,227,536	6,457	857,760	4,789
Humpback, or pink.....	8,440,850	21,102	4,205,320	11,585	1,613,188	4,179	2,381,428	5,954
Sockeye, or blueback.....	41,032,910	1,558,804	7,665,005	187,312	2,881,185	108,398	4,574,145	168,468
Steelhead trout.....			272	13	4,000	260	45,310	2,266
Total.....	54,810,187	1,722,927	13,392,201	243,722	6,562,954	149,835	10,746,637	336,088
TOTAL.								
Chinook, or king.....	1,443,291	68,960	685,466	30,417	1,138,691	51,623	1,274,680	111,835
Coho, or silver.....	4,903,624	113,460	1,186,332	30,217	1,454,492	40,894	2,218,814	60,771
Dog, or chum.....	1,181,412	5,927	604,208	3,022	2,651,374	13,780	2,252,760	13,089
Humpback, or pink.....	8,454,850	21,137	4,380,320	11,935	1,680,988	4,750	2,381,428	5,954
Blueback, or sockeye.....	43,582,360	1,656,562	8,972,505	235,692	3,915,935	146,908	4,744,145	175,334
Steelhead trout.....	7,000	350	272	13	132,628	8,485	46,110	2,306
Grand total.....	59,573,537	1,866,396	15,829,103	311,296	10,974,108	266,440	12,917,937	369,289

PRODUCTS OF THE SALMON FISHERIES OF WASHINGTON, BY APPARATUS, SPECIES, AND COUNTIES, IN 1909—Continued.

Apparatus and species.	Snohomish.		King.		Pierce.		Thurston.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
PURSE SEINES.								
Chinook, or king.....	8,000	\$400			82,285	\$4,400	1,250	\$50
Coho, or silver.....	159,998	4,400	766,000	\$21,175	513,340	13,833	54,396	1,510
Dog, or chum.....	350,000	1,700	1,640,000	14,500	2,482,000	12,410	570,000	3,600
Sockeye, or blueback.....	800,000	28,800	7,050,000	282,000	4,394,995	158,220	125,000	5,000
Steelhead trout.....			14,100	987	10,400	520	400	20
Total.....	1,317,998	35,300	9,470,100	318,662	7,483,020	189,383	751,046	10,180
HAUL SEINES.								
Chinook, or king.....			65,500	4,585	18,743	1,312		
Coho, or silver.....	155,250	3,125	364,000	11,000	462,000	13,000	60,000	2,000
Dog, or chum.....	399,000	1,995	808,000	10,100	1,293,000	8,750	340,000	6,800
Humpback, or pink.....	202,000	503						
Total.....	756,250	5,623	1,237,500	25,685	1,773,743	23,062	400,000	8,800
GILL NETS.								
Chinook, or king.....	337,900	12,164	49,500	3,960	30,000	2,400		
Coho, or silver.....	438,256	16,480	555,000	18,500	246,000	10,250	90,000	3,000
Dog, or chum.....	101,380	731	70,400	440	32,000	200	48,000	240
Humpback, or pink.....			42,000	525				
Sockeye, or blueback.....			335,500	13,420				
Steelhead trout.....	142,243	9,293	205,000	12,300	100,000	10,000	30,000	1,500
Total.....	1,019,779	38,668	1,257,400	49,145	408,000	22,850	168,000	4,740
TRAP NETS.								
Chinook, or king.....	385,150	16,716						
Coho, or silver.....	908,764	23,167						
Dog, or chum.....	813,200	4,066						
Humpback, or pink.....	354,000	1,383						
Steelhead trout.....	27,000	1,350						
Total.....	2,488,114	46,682						
LINES.								
Coho, or silver.....	281,250	7,500						
TOTAL.								
Chinook, or king.....	731,050	29,280	115,000	8,545	131,028	8,112	1,250	50
Coho, or silver.....	1,943,518	54,672	1,685,000	50,675	1,221,340	37,083	204,396	6,510
Dog, or chum.....	1,663,580	8,492	2,518,400	25,040	3,807,000	21,360	958,000	10,640
Humpback, or pink.....	556,000	1,886	42,000	525				
Blueback, or sockeye.....	800,000	28,800	7,385,500	295,420	4,394,995	158,220	125,000	5,000
Steelhead trout.....	169,243	10,643	219,100	13,287	110,400	10,520	30,400	1,520
Grand total.....	5,863,391	133,773	11,965,000	393,492	9,664,763	235,295	1,319,046	23,720

PRODUCTS OF THE SALMON FISHERIES OF WASHINGTON, BY APPARATUS, SPECIES,
AND COUNTIES, IN 1909—Continued.

Apparatus and species.	Mason.		Kitsap.		Clallam.		Jefferson.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
PURSE SEINES.								
Chinook, or king.....			40,000	\$2,000				
Coho, or silver.....	108,000	\$2,700	613,990	15,350				
Dog, or chum.....	400,000	2,000	2,540,000	13,700				
Sockeye, or blueback.....	100,000	4,000	2,045,000	\$1,800				
Steelhead trout.....	600	42	4,900	245				
Total.....	608,600	8,742	5,243,890	113,095				
HAUL SEINES.								
Chinook, or king.....			12,000	600	31,000	\$1,550	21,000	\$1,050
Coho, or silver.....	437,998	11,480	378,000	9,990	110,000	3,300	122,000	3,760
Dog, or chum.....	756,000	4,370	1,129,000	8,970	39,000	330	227,000	2,488
Sockeye, or blueback.....							8,000	400
Steelhead trout.....	3,000	210	17,080	854	14,200	710	5,200	260
Total.....	1,196,998	16,060	1,536,080	20,414	194,200	5,890	383,500	7,958
GILL NETS.								
Chinook, or king.....					75,000	3,750	17,000	970
Coho, or silver.....	40,000	1,200	18,000	490	60,515	1,578	74,000	2,220
Dog, or chum.....	81,000	640	33,000	395	30,000	150	48,000	240
Sockeye, or blueback.....	25,000	1,000	154,000	6,140			24,500	980
Steelhead trout.....	4,000	240	2,300	115	33,055	1,653	7,000	350
Total.....	150,000	3,080	207,300	7,140	198,570	7,131	170,500	4,760
TRAP NETS.								
Chinook, or king.....			106,225	5,305			4,282	199
Coho, or silver.....			504,074	13,020			265,662	6,642
Dog, or chum.....			1,333,704	6,669			1,036,472	5,182
Steelhead trout.....							1,735	87
Total.....			1,944,003	24,994			1,308,151	12,110
LINES.								
Chinook, or king.....					110,880	4,800		
Coho, or silver.....					571,284	17,649		
Dog, or chum.....					4,000	20		
Total.....					686,164	22,469		
TOTAL.								
Chinook, or king.....			158,225	7,905	216,880	10,100	42,282	2,219
Coho, or silver.....	585,998	15,380	1,514,064	38,850	741,799	22,527	461,662	12,622
Dog, or chum.....	1,237,000	7,010	5,035,704	29,734	73,000	500	1,312,072	7,910
Blueback, or sockeye.....	125,000	5,000	2,199,000	87,940			32,500	1,380
Steelhead trout.....	7,600	492	24,280	1,214	47,255	2,363	13,935	697
Grand total.....	1,955,598	27,882	8,931,273	165,643	1,078,934	35,490	1,862,451	24,828

PRODUCTS OF THE SALMON FISHERIES OF WASHINGTON, BY APPARATUS, SPECIES, AND COUNTIES, IN 1909—Continued.

Apparatus and species.	Chehalis.		Pacific.		Wahkiakum.		Cowlitz.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
PURSE SEINES.								
Chinook, or king			8,919	\$535				
Coho, or silver			2,184	44				
Sockeye, or blueback			1,090	49				
Steelhead trout			4,742	190				
Total			16,935	818				
HAUL SEINES.								
Chinook, or king			11,500	345	312,616	\$18,957	50,000	\$3,000
Coho, or silver					42,417	848		
Sockeye, or blueback					19,722	888	12,000	600
Steelhead trout					112,221	5,411	28,000	1,400
Total			11,500	345	486,976	26,104	90,000	5,000
GILL NETS.								
Chinook, or king	571,586	\$15,840	813,978	47,253	1,100,511	66,031		
Coho, or silver	641,858	16,571	187,000	5,500	316,274	6,325		
Dog, or chum	306,256	1,889	57,800	432	400,224	2,354		
Sockeye, or blueback	638,000	23,200	4,500	203				
Steelhead trout	118,000	4,066	45,142	2,328	139,877	6,994	13,000	620
Total	2,275,700	61,566	1,108,420	55,716	1,956,886	81,704	13,000	620
DIVER NETS.								
Chinook, or king							172,667	10,820
Steelhead trout							76,533	3,827
Total							249,200	14,647
TRAP NETS.								
Chinook, or king	49,000	1,113	1,208,963	67,996	31,669	492	69,690	303
Coho, or silver	165,000	3,875	620,461	9,649	458,571	9,172	203,000	4,290
Dog, or chum	36,000	225	725,652	8,996	634,384	3,490	65,600	410
Sockeye, or blueback			113,195	5,093				
Steelhead trout			431,615	21,779	32,416	1,621	6,800	290
Total	250,000	5,213	3,099,886	113,513	1,157,040	14,775	345,090	5,293
TOTAL.								
Chinook, or king	620,586	16,953	2,043,360	116,129	1,444,796	85,480	292,357	14,123
Coho, or silver	806,858	20,446	809,645	15,193	817,262	16,345	203,000	4,290
Dog, or chum	342,256	2,114	783,452	9,428	1,034,608	5,844	65,600	410
Blueback, or sockeye	638,000	23,200	118,785	5,345	19,722	888	12,000	600
Steelhead trout	118,000	4,066	481,499	24,297	284,514	14,026	124,333	6,137
Grand total	2,525,700	66,779	4,236,741	170,392	3,600,902	122,583	697,290	25,560

PRODUCTS OF THE SALMON FISHERIES OF WASHINGTON, BY APPARATUS, SPECIES,
AND COUNTIES, IN 1909—Continued.

Apparatus and species.	Clarke.		Skamania.		Klickitat.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
PURSE SEINES.								
Chinook, or king.....							216,116	\$10,379
Coho, or silver.....							3,085,916	80,942
Dog, or chum.....							9,030,000	53,150
Humpback, or pink.....							295,000	650
Sockeye, or blueback.....							17,425,085	670,135
Steelhead trout.....							37,942	2,144
Total.....							30,000,059	\$17,400
HAUL SEINES.								
Chinook, or king.....			180,000	\$12,600			856,759	51,059
Coho, or silver.....					200,000	\$6,000	3,022,665	84,683
Dog, or chum.....							6,900,600	54,658
Humpback, or pink.....							236,000	588
Sockeye, or blueback.....			24,000	1,200			63,722	3,088
Steelhead trout.....			18,000	900	300,480	15,024	507,609	25,240
Total.....			222,000	14,700	500,480	21,024	11,587,355	219,316
GILL NETS.								
Chinook, or king.....	3,000	\$210	15,944	1,115	500	50	3,702,213	182,343
Coho, or silver.....	8,015	244	6,216	186	1,000	30	4,547,210	134,672
Dog, or chum.....							1,959,698	11,688
Humpback, or pink.....							59,500	746
Sockeye, or blueback.....			2,850	143			2,972,050	111,734
Steelhead trout.....	9,700	485	9,150	458	600	36	983,267	58,442
Total.....	20,715	939	34,160	1,902	2,400	116	14,224,238	499,625
DIVER NETS.								
Chinook, or king.....	14,000	980	77,614	5,433			264,281	17,233
Coho, or silver.....			2,000	60			2,000	60
Steelhead trout.....	3,000	150	3,000	150			82,533	4,127
Total.....	17,000	1,130	82,614	5,643			348,814	21,420
REEF NETS.								
Chinook, or king.....							45,000	2,250
Coho, or silver.....							136,000	4,080
Dog, or chum.....							96,000	500
Sockeye, or blueback.....							365,000	14,600
Total.....							642,000	21,430
TRAP NETS.								
Chinook, or king.....			4,200	294	14,600	1,022	5,453,851	315,371
Coho, or silver.....			3,000	90	17,600	528	9,349,310	217,487
Dog, or chum.....							7,530,128	44,284
Humpback, or pink.....							16,994,786	44,203
Sockeye, or blueback.....			800	40	2,250	128	56,269,490	2,023,243
Steelhead trout.....			3,600	180	6,600	366	559,348	28,212
Total.....			11,600	604	41,050	2,044	96,156,913	2,677,800
WHEELS.								
Chinook, or king.....			261,736	16,039	105,640	5,432	367,376	21,471
Coho, or silver.....			18,751	666	314,080	6,418	332,831	7,084
Sockeye, or blueback.....			173,842	7,358	11,800	508	185,642	7,866
Steelhead trout.....			52,552	2,081	204,000	10,240	256,552	12,321
Total.....			506,881	26,144	635,520	22,598	1,142,401	48,742
LINES.								
Chinook, or king.....							110,880	4,800
Coho, or silver.....							852,534	25,149
Dog, or chum.....							4,000	20
Total.....							967,414	29,969

PRODUCTS OF THE SALMON FISHERIES OF WASHINGTON, BY APPARATUS, SPECIES, AND COUNTIES, IN 1909—Continued.

Apparatus and species.	Clarke.		Skamania.		Klickitat.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
TOTAL.								
Chinook, or king.....	17,000	\$1,190	539,494	\$35,481	121,040	\$6,504	11,016,476	\$604,906
Coho, or silver.....	8,015	244	29,967	1,002	532,680	12,976	21,328,466	554,157
Dog, or chum.....							25,520,426	164,300
Humpback, or pink.....							17,495,586	46,187
Blueback, or sockeye.....			201,492	8,741	14,050	636	77,280,989	2,835,666
Steelhead trout.....	12,700	635	86,302	3,769	511,680	25,666	2,427,251	130,486
Grand total.....	37,715	2,069	857,255	48,993	1,179,450	45,782	155,069,194	4,335,702

STATISTICS BY WATERS.

Persons employed.—Puget Sound leads in the number of persons employed in all branches of the industry, followed by Columbia River, Grays Harbor, and Willapa Harbor in the order named.

PERSONS EMPLOYED IN THE SALMON FISHERIES OF WASHINGTON, BY WATERS AND NATIONALITIES, IN 1909.

Occupation and race.	Puget Sound.	Grays Harbor.	Willapa Harbor.	Columbia River.	Total.
Fishermen:					
Whites.....	2,981	112	130	1,203	4,426
Indians.....	188	33			221
Total.....	3,169	145	130	1,203	4,647
Shoresmen:					
Whites.....	1,968	16	10	97	2,091
Indians.....	115				115
Chinese.....	1,051	45	10	164	1,270
Japanese.....	1,004	15	10	73	1,102
Total.....	4,138	76	30	334	4,578
Transporters:					
Whites.....	252	3	4	33	292
Total:					
Whites.....	5,201	131	144	1,333	6,809
Indians.....	303	33			336
Chinese.....	1,051	45	10	164	1,270
Japanese.....	1,004	15	10	73	1,102
Grand total.....	7,559	224	164	1,570	9,517

Investment, apparatus, etc.—Puget Sound leads in the total investment. The principal forms of apparatus used in the waters of Washington are gill nets, haul and purse seines, traps, and wheels.

INVESTMENT IN THE SALMON FISHERIES OF WASHINGTON, BY WATERS, IN 1909.

Items.	Puget Sound.		Grays Harbor.		Willapa Harbor.		Columbia River.		Total.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Transporting vessels:										
Power vessels.....	72	\$383,700	1	\$3,000	2	\$8,500	18	\$45,300	93	\$440,500
Tonnage.....	996		8		19		135		1,158	
Outfit.....		119,860		500		2,190		13,075		135,625
Power boats.....	2	1,700					3	2,250	5	3,950
Fishing boats, power..	260	363,750	5	2,500	24	7,800	175	98,600	464	472,650
Fishing boats, sail and row.....	1,519	54,815	115	8,350	48	6,340	562	59,440	2,244	128,945
Scows and house boats.	370	155,233	1	400	8	2,800	19	10,240	398	168,673
Pile drivers.....	22	100,600	3	450	2	1,800	35	21,500	62	124,350
Apparatus, shore fisheries:										
Purse seines.....	a 99	43,650					b 2	500	101	44,150
Haul seines.....	c 226	20,255			d 2	350	e 18	8,350	246	28,955
Gill nets, drift.....	f 896	54,131	g 100	8,000	h 80	5,600	i 544	101,100	1,620	168,831
Gill nets, set.....	j 1,292	24,575	k 189	9,724	l 12	360	m 131	2,600	1,624	37,259
Diver nets.....							n 48	10,160	48	10,160
Trap nets, stationary.....	150	768,218	15	3,400	35	16,400	325	536,950	525	1,324,968
Trap nets, floating.....	1	2,000							1	2,000
Reef nets.....	9	4,500							9	4,500
Wheels, stationary.....							13	76,000	13	76,000
Wheels, scow.....							3	8,500	3	8,500
Lines, trolling.....		261								261
Shore and accessory property.....		1,295,087		36,753		50,000		348,190		1,730,030
Cash capital.....		1,168,000		20,000		18,000		218,500		1,424,500
Total.....		4,560,335		93,077		120,140		1,561,255		6,334,807

a Aggregate length of 68,100 yards.

b Aggregate length of 800 yards.

c Aggregate length of 35,841 yards.

d Aggregate length of 300 yards.

e Aggregate length of 8,683 yards.

f Aggregate length of 112,915 yards.

g Aggregate length of 20,000 yards.

h Aggregate length of 28,000 yards.

i Aggregate length of 268,200 yards.

j Aggregate length of 57,980 yards.

k Aggregate length of 27,960 yards.

l Aggregate length of 720 yards.

m Aggregate length of 5,370 yards.

n Aggregate length of 19,200 yards.

Products.—The total catch amounted to 155,069,194 pounds, valued at \$4,335,702, of which Puget Sound produced 141,934,141 pounds, valued at \$3,853,544. Trap nets were the most effective. No humpbacks were taken commercially elsewhere than in Puget Sound, while no sockeyes or bluebacks were taken commercially in Willapa Harbor.

PRODUCTS OF THE SALMON FISHERIES OF WASHINGTON, BY APPARATUS, SPECIES, AND WATERS, IN 1909.

Apparatus and species.	Puget Sound.		Grays Harbor.		Willapa Harbor.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
PURSE SEINES.						
Chinook, or king.....	207,197	\$9,844				
Coho, or silver.....	3,083,732	80,898				
Dog, or chum.....	9,030,000	53,150				
Humpback, or pink.....	205,000	650				
Sockeye, or blueback.....	17,423,995	670,086				
Steelhead trout.....	33,200	1,954				
Total.....	29,983,124	816,582				
HAUL SEINES.						
Chinook, or king.....	302,643	16,157			11,500	\$345
Coho, or silver.....	2,780,248	77,835				
Dog, or chum.....	6,900,600	54,658				
Humpback, or pink.....	236,000	588				
Sockeye, or blueback.....	8,000	400				
Steelhead trout.....	48,908	2,505				
Total.....	10,276,399	152,143			11,500	345
GILL NETS.						
Chinook, or king.....	1,196,394	51,844	571,586	\$15,840	40,000	1,200
Coho, or silver.....	3,386,847	105,816	641,858	16,571	22,000	2,200
Dog, or chum.....	1,195,418	7,013	306,256	1,889	9,800	162
Humpback, or pink.....	59,800	746				
Sockeye, or blueback.....	2,326,700	88,188	638,000	23,200		
Steelhead trout.....	647,798	43,455	118,000	4,066	16,000	800
Total.....	8,812,957	297,062	2,275,700	61,566	87,800	4,362
REEF NETS.						
Chinook, or king.....	45,000	2,250				
Coho, or silver.....	136,000	4,080				
Dog, or chum.....	96,000	500				
Sockeye, or blueback.....	365,000	14,600				
Total.....	642,000	21,430				
TRAP NETS.						
Chinook, or king.....	4,075,729	244,151	49,000	1,113	187,799	6,890
Coho, or silver.....	7,881,678	189,883	165,000	3,875	262,271	2,485
Dog, or chum.....	6,068,492	31,163	36,000	225	643,332	8,482
Humpback, or pink.....	16,994,786	44,203				
Sockeye, or blueback.....	56,153,245	2,022,982				
Steelhead trout.....	78,317	3,976			170	7
Total.....	91,252,247	2,536,358	250,000	5,213	1,093,572	17,864
LINES.						
Chinook, or king.....	110,880	4,800				
Coho, or silver.....	852,534	25,149				
Dog, or chum.....	4,000	20				
Total.....	967,414	29,969				
TOTAL.						
Chinook, or king.....	5,937,843	329,046	620,586	16,953	239,299	8,435
Coho, or silver.....	18,121,039	483,661	806,858	20,446	284,271	4,685
Dog, or chum.....	23,294,510	146,504	342,256	2,114	653,132	8,644
Humpback, or pink.....	17,495,586	46,187				
Sockeye, or blueback.....	76,276,940	2,796,256	638,000	23,200		
Steelhead trout.....	808,223	51,890	118,000	4,066	16,170	807
Grand total.....	141,934,141	3,853,544	2,525,700	66,779	1,192,872	22,571

PRODUCTS OF THE SALMON FISHERIES OF WASHINGTON, BY APPARATUS, SPECIES,
 AND WATERS, IN 1909—Continued.

Apparatus and species.	Columbia River.		Total.	
	Pounds.	Value.	Pounds.	Value.
PURSE SEINES.				
Chinook, or king.....	8,919	\$535	216,116	\$10,379
Coho, or silver.....	2,184	44	3,085,916	80,942
Dog, or chum.....			9,030,000	53,150
Humpback, or pink.....			205,000	650
Sockeye, or blueback.....	1,090	49	17,425,085	670,135
Steelhead trout.....	4,742	190	37,942	2,144
Total.....	16,935	818	30,000,059	817,400
HAUL SEINES.				
Chinook, or king.....	542,616	34,557	856,759	51,059
Coho, or silver.....	242,417	6,848	3,022,665	84,683
Dog, or chum.....			6,900,000	54,658
Humpback, or pink.....			236,000	588
Sockeye, or blueback.....	55,722	2,688	63,722	3,088
Steelhead trout.....	458,701	22,735	507,609	25,240
Total.....	1,299,456	66,828	11,587,355	219,316
GILL NETS.				
Chinook, or king.....	1,894,233	113,459	3,702,213	182,343
Coho, or silver.....	496,505	10,085	4,547,210	134,672
Dog, or chum.....	448,224	2,624	1,959,698	11,688
Humpback, or pink.....			59,800	746
Sockeye, or blueback.....	7,350	346	2,972,050	111,734
Steelhead trout.....	201,469	10,121	983,267	58,442
Total.....	3,047,781	136,635	14,224,238	499,625
DIVER NETS.				
Chinook, or king.....	264,281	17,233	264,281	17,233
Coho, or silver.....	2,000	60	2,000	60
Steelhead trout.....	82,533	4,127	82,533	4,127
Total.....	348,814	21,420	348,814	21,420
REEF NETS.				
Chinook, or king.....			45,000	2,250
Coho, or silver.....			136,000	4,080
Dog, or chum.....			96,000	500
Sockeye, or blueback.....			365,000	14,600
Total.....			642,000	21,430
TRAP NETS.				
Chinook, or king.....	1,141,323	63,217	5,453,851	315,371
Coho, or silver.....	1,040,361	21,244	9,349,310	217,487
Dog, or chum.....	782,304	4,414	7,530,128	44,284
Humpback, or pink.....			16,994,786	44,203
Sockeye, or blueback.....	116,245	5,261	56,269,490	2,028,243
Steelhead trout.....	480,861	24,229	559,348	28,212
Total.....	3,561,094	118,365	96,156,913	2,677,800
WHEELS.				
Chinook, or king.....	367,376	21,471	367,376	21,471
Coho, or silver.....	332,831	7,084	332,831	7,084
Sockeye, or blueback.....	185,642	7,866	185,642	7,866
Steelhead trout.....	256,552	12,321	256,552	12,321
Total.....	1,142,401	48,742	1,142,401	48,742
LINES.				
Chinook, or king.....			110,880	4,800
Coho, or silver.....			852,534	25,149
Dog, or chum.....			4,000	20
Total.....			967,414	29,969
TOTAL.				
Chinook, or king.....	4,218,748	250,472	11,016,476	604,906
Coho, or silver.....	2,116,298	45,365	21,328,466	554,157
Dog, or chum.....	1,230,528	7,038	25,520,426	164,300
Humpback, or pink.....			17,495,586	46,187
Sockeye, or blueback.....	366,049	16,210	77,280,989	2,835,666
Steelhead trout.....	1,484,858	73,723	2,427,251	130,486
Grand total.....	9,416,481	392,808	155,069,194	4,335,702

Products canned.—Of the total pack of 1,926,539 cases, valued at \$8,681,843, 1,757,539 cases, valued at \$7,917,608, were packed on Puget Sound. One of the canneries operating on the Columbia River brought some sockeyes from Puget Sound, and the Puget Sound packers could have packed many more humpbacks than they did; but refrained from doing so because of the low prices prevailing at the time for canned humpbacks.

PACK OF CANNED SALMON IN WASHINGTON IN 1909.

Products.	Puget Sound.		Grays Harbor.		Willapa Harbor.	
	Cases.	Value.	Cases.	Value.	Cases.	Value.
Chinook, or king, red:						
½-pound flat.....	655	\$2,620				
1-pound flat.....	8,278	49,668			197	\$837
1-pound flat exports.....						
1-pound tall.....	2,003	10,817	3,544	\$15,594	1,258	5,032
Total.....	10,936	63,105	3,544	15,594	1,455	5,869
Chinook, or king, white:						
1-pound flat.....	2,033	8,210				
1-pound tall.....	378	1,289	2,177	5,225		
Total.....	2,411	9,499	2,177	5,225		
Coho, or silver:						
½-pound flat.....	24,061	65,771	1,088	3,046		
1-pound flat.....	21,431	103,268	1,176	5,174		
1-pound tall.....	109,249	458,845	7,299	29,926	4,822	17,359
2-pound nominal.....	427	2,562				
Total.....	155,168	630,446	9,563	38,146	4,822	17,359
Chum, or dog:						
½-pound flat.....					1,300	1,950
1-pound flat.....	219	591				
1-pound tall.....	53,469	128,325	5,047	11,608	5,097	11,213
Total.....	53,688	128,916	5,047	11,608	6,397	13,163
Humpback, or pink:						
1-pound flat.....	2,030	5,585				
1-pound tall.....	368,963	896,757				
Total.....	370,993	902,342				
Sockeye, or blueback:						
½-pound flat.....	224,455	906,770				
1-pound flat.....	454,381	2,728,186	244	1,464		
1-pound tall.....	485,507	2,548,344	1,405	7,587		
Total.....	1,164,343	6,183,300	1,649	9,051		
Grand total.....	1,757,539	7,917,608	21,980	79,624	12,674	36,391

PACK OF CANNED SALMON IN WASHINGTON IN 1909—Continued.

Products.	Columbia River.		Total.	
	Cases.	Value.	Cases.	Value.
Chinook, or king, red:				
½-pound flat.....	22,895	\$96,160	23,550	\$98,780
1-pound flat.....	30,222	210,134	38,697	260,639
1-pound flat exports.....	606	4,242	606	4,242
1-pound tall.....	12,066	78,636	18,871	110,079
1-pound oval.....	1,110	10,212	1,110	10,212
Total.....	66,899	399,384	82,834	483,952
Chinook, or king, white:				
1-pound flat.....			2,033	8,210
1-pound tall.....			2,555	6,514
Total.....			4,588	14,724
Coho, or silver:				
½-pound flat.....	9,143	25,600	34,292	94,417
1-pound flat.....	6,278	26,313	28,885	134,755
1-pound tall.....	15,638	63,900	137,008	570,030
2-pound nominal.....			427	2,562
Total.....	31,059	115,813	200,612	801,764
Chum, or dog:				
½-pound flat.....			1,300	1,950
1-pound flat.....			219	591
1-pound tall.....	20,051	46,786	83,664	197,932
Total.....	20,051	46,786	85,183	200,473
Humpback, or pink:				
1-pound flat.....			2,030	5,585
1-pound tall.....			368,963	896,757
Total.....			370,993	902,342
Sockeye, or blueback:				
½-pound flat.....	^a 5,047	21,197	229,502	927,967
1-pound flat.....	2,087	17,017	456,712	2,746,667
1-pound tall.....	567	3,062	487,479	2,558,993
Total.....	7,701	41,276	1,173,693	6,233,627
Steelhead trout:				
1-pound flat.....	945	2,937	945	2,937
1-pound flat.....	3,794	19,422	3,794	19,422
1-pound tall.....	3,897	22,602	3,897	22,602
Total.....	8,636	44,961	8,636	44,961
Grand total.....	134,346	648,220	^b 1,926,539	8,681,843

^a Includes 997 cases, valued at \$4,187, packed with sockeyes from Puget Sound.

^b All 1-pound cases contain 48 1-pound cans; the ½-pound cases contain 48 ½-pound cans. Reduced to a common basis of cases containing 48 1-pound cans, the pack is 1,781,317½ cases.

Miscellaneous products.—By far the greater part of the miscellaneous secondary products were prepared on Puget Sound. Pickled salmon predominate in quantity, but mild-cured salmon represent the greatest value.

MISCELLANEOUS SECONDARY PRODUCTS PACKED IN WASHINGTON IN 1909.

NOTE.—Mild-cured salmon have been figured on a basis of 800 pounds to the tierce and pickled fish on a basis of 200 pounds to the barrel.

Products.	Puget Sound.		Grays Harbor.		Willapa Harbor.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Frozen:						
Coho, or silver, round.....	396,477	\$21,989				
Coho, or silver, dressed.....	60,000	4,200				
Dog, or chum, round.....	1,099,985	55,250				
Dog, or chum, dressed.....	264,687	11,911				
Humpback, round.....	62,945	1,888				
King, or spring, round.....	70,183	7,018				
King, or spring, dressed.....	4,000	400				
Steelhead trout, round.....	202,165	18,195	70,000	\$6,300		
Total.....	2,160,442	120,851	70,000	6,300		
Mild cured:						
King, or spring.....	1,687,200	210,770	60,000	9,000	23,200	\$1,856
Pickled:						
King, or spring.....			1,000	540		
King, or spring, bellies.....						
Dog, or chum.....	50,000	175				
Humpback.....	1,615,000	48,450				
Humpback bellies.....	172,400	8,620				
Total.....	1,837,400	57,245	1,000	540		
Smoked:						
Coho, or silver.....	30,000	1,800				
Dog, or chum.....	517,245	25,862				
Dog, or chum, kippered.....	5,000	500				
Humpback backs, kippered.....	100,000	5,000				
King, or spring.....	30,165	2,413				
King, or spring, white, kippered.....	190,500	16,050				
Total.....	872,910	51,625				
Fertilizer.....	1,210,000	18,610				
Oil.....	380,648	14,161				
Grand total.....	8,148,600	473,262	131,000	15,840	23,200	1,856

MISCELLANEOUS SECONDARY PRODUCTS PACKED IN WASHINGTON IN 1909—Continued.

Products.	Columbia River.		Total.	
	Pounds.	Value.	Pounds.	Value.
Frozen:				
Coho, or silver, round.....	72,000	\$3,960	468,477	\$25,949
Coho, or silver, dressed.....			60,000	4,200
Dog, or chum, round.....			1,099,985	55,250
Dog, or chum, dressed.....			264,687	11,911
Humpback, round.....			62,945	1,888
King, or spring, round.....			70,183	7,018
King, or spring, dressed.....			4,000	400
Steelhead trout, round.....	232,000	22,120	504,165	46,615
Total.....	304,000	26,080	2,534,442	153,231
Mild cured:				
King, or spring.....	522,400	52,200	2,292,800	273,826
Pickled:				
King, or spring.....			1,000	540
King, or spring, bellies.....	6,750	671	6,750	671
Dog, or chum.....			50,000	175
Humpback.....			1,615,000	48,450
Humpback bellies.....			172,400	8,620
Total.....	6,750	671	1,845,150	58,456
Smoked:				
Coho, or silver.....			30,000	1,800
Dog, or chum.....			517,245	25,862
Dog, or chum, kippered.....			5,000	500
Humpback backs, kippered.....			100,000	5,000
King, or spring.....			30,165	2,413
King, or spring, white, kippered.....			190,500	16,050
Total.....			872,910	51,625
Fertilizer.....			1,210,000	18,610
Oil.....			^a 380,648	14,161
Grand total.....	833,150	78,951	9,135,950	569,909

^a Represents 50,713 gallons.

COLUMBIA RIVER.

As the Columbia River forms the boundary between Oregon and Washington and the citizens of both States operate in the river, for convenience tables showing persons employed, investment, catch, and the packs of canned salmon and miscellaneous secondary products on both sides of the river are combined in the tables given below, in addition to showing most of these data in the regular state tables.

PERSONS EMPLOYED IN THE SALMON FISHERIES OF THE COLUMBIA RIVER IN 1909.

Occupation and race.	Number.	Occupation and race.	Number.
Fishermen: Whites.....	4,443	Transporters: Whites.....	80
Shoresmen:		Total:	
Whites.....	426	Whites.....	4,949
Chinese.....	417	Chinese.....	417
Japanese.....	268	Japanese.....	268
Total.....	1,111	Grand total.....	5,634

INVESTMENT IN THE SALMON FISHERIES OF THE COLUMBIA RIVER IN 1909.

Items.	Number.	Value.	Items.	Number.	Value.
Transporting vessels:			Apparatus, shore fisheries—Con.		
Power vessels.....	39	\$118,400	Gill nets, drift.....	2,755	\$571,305
Tonnage.....	335		Gill nets, set.....	443	8,163
Outfit.....		29,875	Diver nets.....	166	32,535
Power boats.....	14	26,550	Trap nets.....	346	562,700
Fishing boats, power.....	425	222,700	Wheels, stationary.....	39	389,000
Fishing boats, sail and row.....	1,923	254,395	Wheels, scow.....	12	30,500
Scows and house boats.....	110	51,950	Shore and accessory property.....		1,577,300
Pile drivers.....	37	23,300	Cash capital.....		647,000
Apparatus, shore fisheries:			Total.....		4,567,423
Haul seines.....	52	21,250			
Purse seines.....	2	500			

CATCH, BY APPARATUS AND SPECIES, IN THE SALMON FISHERIES OF THE COLUMBIA RIVER IN 1909.

Apparatus and species.	Pounds.	Value.	Apparatus and species.	Pounds.	Value.
PURSE SEINES.			TRAP NETS.		
Chinook, or king.....	8,919	\$535	Blueback, or sockeye.....	141,265	\$6,387
Coho, or silver.....	2,184	44	Chinook, or king.....	1,198,383	65,823
Blueback, or sockeye.....	1,090	49	Dog, or chum.....	931,564	5,188
Steelhead trout.....	4,742	190	Silver, or coho.....	1,602,581	32,888
Total.....	16,935	818	Steelhead trout.....	527,071	26,540
HAUL SEINES.			Total.....	4,400,864	136,826
Blueback, or sockeye.....	110,503	5,183	WHEELS.		
Chinook, or king.....	1,392,377	85,261	Blueback, or sockeye.....	949,165	38,898
Dog, or chum.....	24,000	150	Chinook, or king.....	1,091,751	64,082
Silver, or coho.....	506,439	12,135	Silver, or coho.....	603,453	12,683
Steelhead trout.....	1,078,118	52,562	Steelhead trout.....	592,819	27,835
Total.....	3,111,437	155,291	Total.....	3,237,188	143,498
GILL NETS.			TOTAL.		
Blueback, or sockeye.....	8,350	396	Blueback, or sockeye.....	1,210,373	50,913
Chinook, or king.....	11,958,512	667,221	Chinook, or king.....	16,534,480	938,808
Dog, or chum.....	542,472	3,223	Dog, or chum.....	1,498,036	8,561
Silver, or coho.....	792,774	16,504	Silver, or coho.....	3,509,431	74,314
Steelhead trout.....	515,940	25,292	Steelhead trout.....	2,803,023	136,636
Total.....	13,818,048	712,636	Grand total.....	25,555,343	1,209,232
DIVER NETS.					
Chinook, or king.....	884,538	55,886			
Silver, or coho.....	2,000	60			
Steelhead trout.....	84,333	4,217			
Total.....	970,871	60,163			

CANNED PACK ON BOTH SIDES OF THE COLUMBIA RIVER IN 1909.

Products.	Cases. ^a	Value.	Products.	Cases. ^a	Value.
Blueback, or sockeye:			Humpback, or pink:		
½-pound flat.....	b 37,118	\$154,292	1-pound tall.....	d 55	\$132
1-pound flat.....	8,732	56,887	Silverside, coho, or white:		
1-pound tall.....	c 617	3,382	½-pound flat.....	12,447	34,852
Total.....	46,467	214,561	1-pound flat.....	14,498	62,468
Chinook, or king:			1-pound tall.....	21,455	87,750
½-pound flat.....	90,281	379,181	Total.....	48,400	185,070
1-pound flat.....	84,212	603,651	Steelhead trout:		
1-pound flat exports.....	606	4,242	½-pound flat.....	8,009	25,021
1-pound tall.....	29,519	193,827	1-pound flat.....	5,159	27,117
½-pound oval.....	534	2,670	1-pound tall.....	8,217	47,658
1-pound oval.....	1,919	18,142	Total.....	21,385	99,796
2-pound nominal.....	458	1,833	Grand total.....	348,378	1,760,220
Total.....	207,529	1,203,546			
Chum, or dog:					
1-pound tall.....	24,542	57,115			

^a All 1-pound cases contain 48 1-pound cans; the ½-pound cases contain 48 ½-pound cans.

^b Of these, 5,592 cases, valued at \$22,883, were filled with sockeyes brought from Puget Sound, Wash.

^c Of these, 50 cases, valued at \$320, were filled with sockeyes brought from Puget Sound, Wash.

^d Filled with fish brought from Puget Sound, Wash.

PACK OF MISCELLANEOUS PRODUCTS ON BOTH SIDES OF THE COLUMBIA RIVER IN 1909.

Products.	Pounds.	Value.	Products.	Pounds.	Value.
Frozen:			Smoked:		
Chinook.....	14,000	\$1,400	Chinook.....	127,700	\$19,155
Silverside.....	288,175	17,828	Silverside.....	20,000	2,000
Steelhead trout.....	1,646,662	163,887	Total.....	147,700	21,155
Total.....	1,948,837	183,115	Grand total.....	6,535,533	648,125
Mild-cured:					
Chinook.....	4,432,246	443,184			
Pickled:					
Chinook bellies.....	6,750	671			

OREGON.

The catch of salmon in the Columbia River in 1909 was only fair, owing partly to the shortening of the open fishing season. On the coast streams conditions were far from favorable. Low water at one time kept the salmon from entering the streams; afterwards freshets and storms made fishing impossible at times. A few places, however, show increases over the previous year.

STATISTICS BY COUNTIES.

Persons employed.—The total number of persons employed was 5,320. All of the fishermen and transporters were whites. Clatsop County, in which Astoria is located, has more than half of the persons employed.

PERSONS EMPLOYED IN THE SALMON FISHERIES OF OREGON, BY COUNTIES AND NATIONALITIES, IN 1909.

Counties.	Fisher- men.	Shoresmen				Trans- porters.	Total.				
	Whites.	Whites.	Chi- nese.	Japa- nese.	Total.	Whites.	Whites.	Chi- nese.	Japa- nese.	Total.	
Wasco.....	48	21	33	8	62	69	33	8	110	
Hood River.....	6	6	6	
Multnomah.....	88	29	68	42	139	2	119	68	42	229	
Clackamas.....	86	86	86	
Columbia.....	149	21	21	8	178	178	
Clatsop.....	2,863	258	152	145	555	37	3,158	152	145	3,455	
Tillamook.....	154	11	50	9	70	4	169	50	9	228	
Lincoln.....	144	9	19	14	42	153	19	14	186	
Lane.....	121	7	30	14	51	2	130	30	14	174	
Douglas.....	100	5	19	10	34	2	107	19	10	136	
Coos.....	276	26	36	14	76	10	312	36	14	362	
Curry.....	33	15	4	19	5	53	4	57	
Josephine.....	111	2	2	113	113	
Total.....	4,179	404	411	256	1,071	70	4,653	411	256	5,320	

Investment, apparatus, etc.—The total investment amounted to \$3,641,775, of which more than one-half is contributed by Clatsop County. The gill net is the principal form of apparatus used in most counties.

INVESTMENT IN THE SALMON FISHERIES OF OREGON, BY COUNTIES, IN 1909.

Items.	Wasco.		Hood River.		Multnomah.		Clackamas.		Columbia.	
	Num- ber.	Value.	Num- ber.	Value.	Num- ber.	Value.	Num- ber.	Value.	Num- ber.	Value.
Transporting vessels:										
Power vessels.....	1	\$4,000	4	\$10,900
Tonnage.....	11	26
Outfit.....	600	1,570
Power boats.....	1	1,000	2	1,800
Fishing boats, power.....	1	\$2,000	16	7,900	76	17,100
Fishing boats, sail and row.....	16	800	6	\$240	53	2,300	43	\$1,290	33	1,810
Scows and house boats.....	5	1,350	4	1,500
Apparatus, shore fisheries:										
Haul seines.....	1	500	1	400	4	1,400
Gill nets, drift.....	8	560	72	3,470
Gill nets, set.....	4	70	20	360	52	871	71	792	50	920
Diver nets.....	26	6,250	89	15,825
Pound nets.....	10	6,750
Wheels, stationary.....	14	260,000	12	53,000
Wheels, scow.....	4	6,000	5	16,000
Shore and accessory property.....	261,600	123,015	115	69,565
Cash capital.....	45,000	103,500	15,000
Total.....	575,970	600	320,746	5,667	144,140

INVESTMENT IN THE SALMON FISHERIES OF OREGON, BY COUNTIES, IN 1909—
Continued.

Items.	Clatsop.		Tillamook.		Lincoln.		Lane.		Douglas.	
	Num-ber.	Value.	Num-ber.	Value.	Num-ber.	Value.	Num-ber.	Value.	Num-ber.	Value.
Transporting vessels:										
Power vessels.....	16	\$58,200	2	\$7,300			1	\$3,000	1	\$2,000
Tonnage.....	163		16				7		5	
Outfit.....		14,630		1,750				950		400
Power boats.....	8	21,500	1	2,000	2	\$600				
Fishing boats, power.....	157	97,100	3	600	3	1,500	6	1,200		
Fishing boats, sail and row.....	1,210	188,515	74	5,550	73	5,925	90	2,670	50	2,100
Scows and house boats.....	82	38,860					7	1,020		
Pile drivers.....	2	1,800								
Apparatus, shore fisheries:										
Haul seines.....	28	10,600					1	130		
Gill nets, drift.....	2,131	466,175	63	8,230	112	10,400	51	6,195	30	2,125
Gill nets, set.....	115	2,550	151	4,530	153	4,490	108	1,502	116	4,420
Diver nets.....	3	300								
Pound nets.....	11	19,000								
Shore and accessory property.....		774,815		69,883		41,848		17,100		21,589
Cash capital.....		265,000		28,000		12,500		13,500		12,000
Total.....		1,959,045		127,843		77,263		47,267		44,634

Items.	Coos.		Curry.		Josephine.		Total.	
	Num-ber.	Value.	Num-ber.	Value.	Num-ber.	Value.	Num-ber.	Value.
Transporting vessels:								
Power vessels.....	4	\$24,500	1	\$10,000			30	\$119,900
Tonnage.....	34		26				288	
Outfit.....		4,100		1,350				25,350
Power boats.....			1	2,000			15	28,900
Fishing boats, power.....	25	12,200					287	139,600
Fishing boats, sail and row.....	164	8,125	22	3,300	56	\$1,920	1,890	224,545
Scows and house boats.....	16	2,320					114	45,050
Pile drivers.....							2	1,800
Apparatus, shore fisheries:								
Haul seines.....	8	2,350	1	300	4	600	^a 48	16,280
Gill nets, drift.....	279	23,176	6	800	66	2,200	^b 2,818	523,331
Gill nets, set.....	166	4,720	102	2,305	14	84	^c 1,122	27,614
Diver nets.....							^d 418	22,375
Pound nets.....							21	25,750
Wheels, stationary.....							26	313,000
Wheels, scow.....							9	22,000
Shore and accessory property.....		67,400		100,400		7,450		1,554,780
Cash capital.....		42,000		15,000				551,500
Total.....		190,891		135,455		12,254		3,641,775

^a Aggregate length of 22,855 yards.^b Aggregate length of 1,187,832 yards.^c Aggregate length of 59,625 yards.^d Aggregate length of 46,600 yards.

Products.—The total catch amounted to 22,191,291 pounds, valued at \$968,983, of which Clatsop County contributed more than one-half. Gill nets catch more than two-thirds of the total. Chinook salmon constitute more than one-half of the total catch.

PRODUCTS OF THE SALMON FISHERIES OF OREGON, BY SPECIES AND APPARATUS, IN 1909.

Apparatus and species.	Wasco.		Hood River.		Multnomah.		Clackamas.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
SEINES.								
Blueback.....					6,000	\$300		
Chinook, fresh.....					41,000	2,870		
Silver.....	206,000	\$4,120						
Steelhead trout.....	105,280	4,120			4,000	200		
Total.....	311,280	8,240			51,000	3,370		
GILL NETS.								
Blueback.....					1,000	50		
Chinook, fresh.....	1,800	144	9,700	\$679	18,000	770	208,000	\$8,320
Silver.....	2,600	78	14,700	521	17,100	513	7,000	210
Steelhead trout.....	800	48	5,500	306	20,900	975	24,000	720
Total.....	5,200	270	29,900	1,506	57,000	2,308	239,000	9,250
DIVER NETS.								
Chinook.....					131,757	9,223		
Steelhead trout.....					1,800	90		
Total.....					133,557	9,313		
WHEELS.								
Blueback.....	534,555	21,382			228,968	9,650		
Chinook, fresh.....	497,805	28,998			226,570	13,613		
Silver.....	243,000	4,860			27,622	739		
Steelhead trout.....	272,835	13,232			63,432	2,282		
Total.....	1,548,195	68,472			546,592	26,284		
TOTAL.								
Blueback.....	534,555	21,382			235,968	10,000		
Chinook, fresh.....	499,605	29,142	9,700	679	417,327	26,476	208,000	8,320
Silver.....	451,600	9,058	14,700	521	44,722	1,252	7,000	210
Steelhead trout.....	378,915	17,400	5,500	306	90,132	3,547	24,000	720
Grand total.....	1,864,675	76,982	29,900	1,506	788,149	41,275	239,000	9,250

PRODUCTS OF THE SALMON FISHERIES OF OREGON, BY SPECIES AND APPARATUS, IN 1909—Continued.

Apparatus and species.	Columbia.		Clatsop.		Tillamook.		Lincoln.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
SEINES.								
Blueback			48,781	\$2,195				
Chinook, fresh	64,115	\$3,506	744,646	44,328				
Dog			24,000	150				
Silver	5,419	108	52,603	1,059				
Steelhead	83,073	4,154	427,054	21,353				
Total	152,607	7,768	1,297,094	69,085				
GILL NETS.								
Chinook, fresh			9,826,779	543,849	417,827	\$11,916	255,268	\$12,073
Dog			94,248	599	323,480	1,617	72,360	453
Silver			254,869	5,097	421,587	12,244	580,182	16,755
Steelhead	129,200	6,460	134,071	6,662	5,000	100	6,200	248
Total	129,200	6,460	10,309,967	556,207	1,167,894	25,877	914,010	29,529
DIVER NETS.								
Chinook	476,500	28,710	12,000	720				
POUND NETS.								
Blueback			25,020	1,126				
Chinook, fresh	13,450	59	43,610	2,547				
Dog	145,100	748	4,160	26				
Silver	544,000	11,280	18,220	364				
Steelhead	13,600	680	32,610	1,631				
Total	716,150	12,767	123,620	5,694				
TOTAL.								
Blueback			73,801	3,321				
Chinook, fresh	554,065	32,275	10,627,035	591,444	417,827	11,916	255,268	12,073
Dog	145,100	748	122,408	775	323,480	1,617	72,360	453
Silver	549,419	11,388	325,692	6,520	421,587	12,244	580,182	16,755
Steelhead trout	225,873	11,294	593,745	29,646	5,000	100	6,200	248
Grand total	1,474,457	55,705	11,742,681	631,706	1,167,894	25,877	914,010	29,529

Apparatus and species.	Lane.		Douglas.		Coos.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
SEINES.						
Chinook, fresh	5,000	\$125			16,200	\$466
Silver	8,000	200			176,452	4,411
Steelhead					3,900	78
Total	13,000	325			196,552	4,955
GILL NETS.						
Chinook, fresh	\$2,304	2,057	62,912	\$1,573	127,581	3,497
Chinook, salted	12,000	480				
Dog			36,000	225		
Silver	970,348	24,256	351,072	8,728	1,210,048	30,251
Steelhead			13,000	260	55,000	1,100
Total	1,064,652	26,793	462,984	10,786	1,392,629	34,848
TOTAL.						
Chinook, fresh	87,304	2,182	62,912	1,573	143,781	3,963
Chinook, salted	12,000	480				
Dog			36,000	225		
Silver	978,348	24,456	351,072	8,728	1,386,500	34,662
Steelhead trout			13,000	260	58,900	1,178
Grand total	1,077,652	27,118	462,984	10,786	1,589,181	39,803

PRODUCTS OF THE SALMON FISHERIES OF OREGON, BY SPECIES AND APPARATUS, IN 1909—Continued.

Apparatus and species.	Curry.		Josephine.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
SEINES.						
Blueback.....					54,781	\$2,495
Chinook, fresh.....	25,652	\$292	5,248	\$330	901,861	51,917
Dog.....					24,000	150
Silver.....					448,474	9,898
Steelhead.....					623,317	29,905
Total.....	25,652	292	5,248	330	2,052,433	94,365
GILL NETS.						
Blueback.....					1,000	50
Chinook, fresh.....	462,000	4,620	165,090	10,691	11,637,261	600,189
Chinook, salted.....					12,000	480
Dog.....					526,088	2,894
Silver.....	72,000	1,200	1,698	210	3,903,204	100,063
Steelhead.....	107,100	2,018	3,920	85	502,691	18,982
Total.....	641,100	7,838	168,708	10,986	16,582,244	722,658
DIVER NETS.						
Chinook, fresh.....					620,257	38,653
Steelhead.....					1,800	90
Total.....					622,057	38,743
POUND NETS.						
Blueback.....					25,020	1,126
Chinook, fresh.....					57,060	2,606
Dog.....					149,260	774
Silver.....					562,220	11,644
Steelhead.....					46,210	2,311
Total.....					839,770	18,461
WHEELS.						
Blueback.....					763,523	31,032
Chinook, fresh.....					724,375	42,611
Silver.....					270,622	5,599
Steelhead.....					336,267	15,514
Total.....					2,094,787	94,756
TOTAL.						
Blueback.....					844,324	34,703
Chinook, fresh.....	487,652	4,912	170,338	11,021	13,940,814	735,976
Chinook, salted.....					12,000	480
Dog.....					699,348	3,818
Silver.....	72,000	1,200	1,698	210	5,184,520	127,204
Steelhead trout.....	107,100	2,018	1,920	85	1,510,285	66,802
Grand total.....	666,752	8,130	173,956	11,316	22,191,291	968,983

STATISTICS BY WATERS.

Persons employed.—The Columbia River furnishes about four-fifths of the total number of persons employed. The Coquille River is second and the Siuslaw River third in this respect.

PERSONS EMPLOYED IN THE SALMON FISHERIES OF OREGON, BY WATERS AND NATIONALITIES, IN 1909.

Occupation and nationality.	Columbia River.	Nehalem River.	Tillamook Bay.	Nestueca River.	Siletz River.	Yaquina Bay and River.	Alesea Bay and River.
Fishermen: Whites.....	3,240	48	46	60	16	63	65
Shoresmen:							
Whites.....	329	5	6		2	2	5
Chinese.....	253	23	27			5	14
Japanese.....	195	6	3			5	9
Total.....	777	34	36		2	12	28
Transporters: Whites.....	47		4				
Total:							
Whites.....	3,616	53	56	60	18	65	70
Chinese.....	253	23	27			5	14
Japanese.....	195	6	3			5	9
Grand total.....	4,064	82	86	60	18	75	93

Occupation and nationality.	Siuslaw River.	Umpqua River.	Coos Bay.	Coquille River.	Rogue River.	Total.
Fishermen: Whites.....	121	100	114	162	144	4,179
Shoresmen:						
Whites.....	7	5	14	12	17	404
Chinese.....	30	19	14	22	4	411
Japanese.....	14	10	4	10		256
Total.....	51	34	32	44	21	1,071
Transporters: Whites.....	2	2	10		5	70
Total:						
Whites.....	130	107	138	174	166	4,653
Chinese.....	30	19	14	22	4	411
Japanese.....	14	10	4	10		256
Grand total.....	174	136	156	206	170	5,320

Investment, apparatus, etc.—More than two-thirds of the investment is found on the Columbia River, and this is the only river on which diver nets, pound or trap nets, and wheels are employed.

INVESTMENT IN THE SALMON FISHERIES OF OREGON, BY WATERS, IN 1909.

Items.	Columbia River.		Nehalem River.		Tillamook Bay.		Nestucca River.		Siletz River.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Transporting vessels:										
Power vessels.....	21	\$73,100			2	\$7,300				
Tonnage.....	200				16					
Outfit.....		16,800				1,750				
Power boats.....	11	24,300			1	2,000			1	\$200
Fishing boats, power.....	250	124,100			3	600				
Fishing boats, sail and row.....	1,361	194,955	24	\$1,800	20	1,500	30	\$2,250	9	1,425
Scows and house boats.....	91	41,710								
Pile drivers.....	2	1,800								
Apparatus, shore fisheries:										
Haul seines.....	34	12,900								
Gill nets, drift.....	2,211	470,205	17	1,980	26	3,250	20	3,000	3	300
Gill nets, set.....	312	5,563	70	2,100	31	930	50	1,500	8	240
Diver nets.....	118	22,375								
Pound, or trap, nets.....	21	25,750								
Wheels, stationary.....	26	313,000								
Wheels, scow.....	9	22,000								
Shore and accessory property.....		1,229,110		53,078		16,605		200		17,174
Cash capital.....		428,500		10,000		18,000				1,000
Total.....		3,006,168		68,958		51,935		6,950		20,339

Items.	Yaquina Bay and River.		Alsea Bay and River.		Siuslaw River.		Umpqua River.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Transporting vessels:								
Power vessels.....					1	\$3,000	1	\$2,000
Tonnage.....					7		5	
Outfit.....						950		400
Power boats.....			1	\$400				
Fishing boats, power.....	3	\$1,500			6	1,200		
Fishing boats, sail and row.....	30	2,600	34	1,900	90	2,670	50	2,100
Scows and house boats.....					7	1,020		
Apparatus, shore fisheries:								
Haul seines.....					1	130		
Gill nets, drift.....	60	5,200	49	4,900	51	6,195	30	2,125
Gill nets, set.....	80	2,300	65	1,950	108	1,502	116	4,420
Shore and accessory property.....		5,500		19,174		17,100		21,589
Cash capital.....		1,000		10,500		13,500		12,000
Total.....		18,100		38,824		47,267		44,634

Items.	Coos Bay.		Coquille River.		Rogue River.		Total.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Transporting vessels:								
Power vessels.....	4	\$24,500			1	\$10,000	30	\$119,900
Tonnage.....	34				26		288	
Outfit.....		4,100				1,350		25,350
Power boats.....					1	2,000	15	28,900
Fishing boats, power.....	22	11,600	3	\$600			287	139,600
Fishing boats, sail and row.....	26	3,325	138	4,800	78	5,220	1,892	224,545
Scows and house boats.....	5	890	11	1,430			114	45,050
Pile drivers.....							2	1,800
Apparatus, shore fisheries:								
Haul seines.....	2	550	6	1,800	5	900	48	16,280
Gill nets, drift.....	165	14,176	114	9,000	72	3,000	2,818	523,331
Gill nets, set.....	46	1,120	120	3,600	116	2,389	1,122	27,614
Diver nets.....							418	22,375
Pound, or trap, nets.....							21	25,750
Wheels, stationary.....							26	313,000
Wheels, scow.....							9	22,000
Shore and accessory property.....		46,000		21,400		107,850		1,554,780
Cash capital.....		17,000		25,000		15,000		551,500
Total.....		123,261		67,630		147,709		3,641,775

Catch.—The Columbia River produces more than two-thirds of the total catch, the Siuslaw River is second, and Coos Bay third. Bluebacks are taken on the Columbia River alone. The gill net is the only form of apparatus employed in most of the rivers.

PRODUCTS OF THE SALMON FISHERIES OF OREGON, BY APPARATUS, SPECIES, AND WATERS, IN 1909.

Apparatus and species.	Columbia River.		Nehalem River.		Tillamook Bay.		Nestucca River.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
HAUL SEINES.								
Blueback, or sockeye.....	54,781	\$2,495						
Chinook, or king, fresh.....	849,761	50,704						
Dog, or chum.....	24,000	150						
Silver, or coho.....	264,022	5,287						
Steelhead trout.....	619,417	29,827						
Total.....	1,811,981	88,463						
GILL NETS.								
Blueback, or sockeye.....	1,000	50						
Chinook, or king, fresh.....	10,064,279	553,762	50,284	\$1,509	314,810	\$7,870	52,733	\$2,537
Dog, or chum.....	94,248	599			259,856	1,299		
Silver, or coho.....	296,269	6,419	206,826	5,171	146,592	3,665	68,169	3,408
Steelhead trout.....	314,471	15,171	63,624	318	5,000	100		
Total.....	10,770,267	576,001	320,734	6,998	726,258	12,934	120,902	5,945
DIVER NETS.								
Chinook, or king, fresh.....	620,257	38,653						
Steelhead trout.....	1,800	90						
Total.....	622,057	38,743						
POUND NETS.								
Blueback, or sockeye.....	25,020	1,126						
Chinook, or king, fresh.....	57,060	2,606						
Dog, or chum.....	149,260	774						
Silver, or coho.....	562,220	11,644						
Steelhead trout.....	46,210	2,311						
Total.....	839,770	18,461						
WHEELS.								
Blueback, or sockeye.....	763,523	31,032						
Chinook, or king, fresh.....	724,375	42,611						
Silver, or coho.....	270,622	5,599						
Steelhead trout.....	336,267	15,514						
Total.....	2,094,787	94,756						
TOTAL.								
Blueback, or sockeye.....	844,324	34,703						
Chinook, or king, fresh.....	12,315,732	688,336	50,284	1,509	314,810	7,870	52,733	2,537
Dog, or chum.....	267,508	1,523			259,856	1,299		
Silver, or coho.....	1,393,133	28,949	206,826	5,171	146,592	3,665	68,169	3,408
Steelhead trout.....	1,318,165	62,913	63,624	318	5,000	100		
Grand total.....	16,138,862	816,424	320,734	6,998	726,258	12,934	120,902	5,945

PRODUCTS OF THE SALMON FISHERIES OF OREGON, BY APPARATUS, SPECIES, AND WATERS, IN 1909—Continued.

Apparatus and species.	Siletz River.		Yaquina Bay and River.		Alsea Bay and River.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
GILL NETS.						
Chinook, or king, fresh.....	53,690	\$2,148	33,722	\$1,532	167,856	\$8,393
Dog, or chum.....			42,640	267	29,720	186
Silver, or coho.....			246,738	6,752	333,444	10,003
Steelhead trout.....					6,200	248
Total.....	53,690	2,148	323,100	8,551	537,220	18,830
TOTAL.						
Chinook, or king, fresh.....	53,690	2,148	33,722	1,532	167,856	8,393
Dog, or chum.....			42,640	267	29,720	186
Silver, or coho.....			246,738	6,752	333,444	10,003
Steelhead trout.....					6,200	248
Grand total.....	53,690	2,148	323,100	8,551	537,220	18,830
Apparatus and species.	Siuslaw River.		Umpqua River.		Coos Bay.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
HAUL SEINES.						
Chinook, or king, fresh.....	5,000	\$125			12,100	\$363
Silver, or coho.....	8,000	200			39,000	975
Steelhead trout.....					3,900	78
Total.....	13,000	325			55,000	1,416
GILL NETS.						
Chinook, or king, fresh.....	82,304	2,057	62,912	\$1,573	100,181	2,812
Chinook, or king, salted.....	12,000	480				
Dog, or chum.....			36,000	225		
Silver, or coho.....	970,348	24,256	351,072	8,728	660,240	16,506
Steelhead trout.....			13,000	260	49,000	980
Total.....	1,064,652	26,793	462,984	10,786	809,421	20,298
TOTAL.						
Chinook, or king, fresh.....	87,304	2,182	62,912	1,573	112,281	3,175
Chinook, or king, salted.....	12,000	480				
Dog, or chum.....			36,000	225		
Silver, or coho.....	978,348	24,456	351,072	8,728	699,240	17,481
Steelhead trout.....			13,000	260	52,900	1,058
Grand total.....	1,077,652	27,118	462,984	10,786	864,421	21,714

PRODUCTS OF THE SALMON FISHERIES OF OREGON, BY APPARATUS, SPECIES, AND WATERS, IN 1909—Continued.

Apparatus and species.	Coquille River.		Rogue River.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
HAUL SEINES.						
Blueback, or sockeye.....					54,781	\$2,495
Chinook, or king, fresh.....	4,100	\$103	30,900	\$622	901,861	51,917
Dog, or chum.....					24,000	150
Silver, or coho.....	137,452	3,436			448,474	9,898
Steelhead trout.....					623,317	29,905
Total.....	141,552	3,539	30,900	622	2,052,433	94,365
GILL NETS.						
Blueback, or sockeye.....					1,000	50
Chinook, or king, fresh.....	27,400	685	627,090	15,311	11,637,261	600,189
Chinook, or king, salted.....					12,000	480
Dog, or chum.....					526,088	2,894
Silver, or coho.....	549,808	13,745	73,698	1,410	3,903,204	100,063
Steelhead trout.....	6,000	120	109,020	2,103	502,691	18,982
Total.....	583,208	14,550	809,808	18,824	16,582,244	722,658
DIVER NETS.						
Chinook, or king, fresh.....					620,257	38,653
Steelhead trout.....					1,800	90
Total.....					622,057	38,743
POUND NETS.						
Blueback, or sockeye.....					25,020	1,126
Chinook, or king, fresh.....					57,060	2,606
Dog, or chum.....					149,260	774
Silver, or coho.....					562,220	11,644
Steelhead trout.....					46,210	2,311
Total.....					839,770	18,461
WHEELS.						
Blueback, or sockeye.....					763,523	31,032
Chinook, or king, fresh.....					724,375	42,611
Silver, or coho.....					270,622	5,599
Steelhead trout.....					336,267	15,514
Total.....					2,094,787	94,756
TOTAL.						
Blueback, or sockeye.....					844,324	34,703
Chinook, or king, fresh.....	31,500	788	657,990	15,933	13,940,814	735,976
Chinook, or king, salted.....					12,000	480
Dog, or chum.....					635,724	3,500
Silver, or coho.....	687,260	17,181	73,698	1,410	5,184,520	127,204
Steelhead trout.....	6,000	120	109,020	2,103	1,573,909	67,120
Grand total.....	724,760	18,089	840,708	19,446	22,191,291	968,983

Products canned.—As in other branches of the industry the Columbia River leads, producing more than two-thirds of the pack of canned salmon. But little was done on the Rogue River, owing to the recent death of Mr. R. D. Hume, owner of the principal cannery. Bluebacks and steelheads were packed on the Columbia River alone. All of the humpbacks and part of the sockeyes packed on the Columbia River were brought from Puget Sound, Wash.

PACK OF CANNED SALMON IN OREGON, BY WATERS, IN 1909.

Products.	Columbia River.		Nehalem River.		Tillamook Bay.		Yaquina River and Bay.	
	Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
Blueback, or sockeye:								
½-pound flat.....	a 32,071	\$133,095						
1-pound flat.....	6,645	39,870						
1-pound tall.....	b 50	329						
Total.....	38,766	173,285						
Chinook, or king:								
½-pound flat.....	67,386	283,021	228	\$684	965	\$2,895		
1-pound flat.....	53,990	393,517						
1-pound tall.....	17,453	115,191	1,643	9,858	2,128	12,768		
½-pound oval.....	534	2,670						
1-pound oval.....	809	7,930						
2-pound nominal.....	458	1,833						
Total.....	140,630	804,162	1,871	10,542	3,093	15,663		
Chum, or dog:								
1-pound tall.....	4,491	10,329	909	2,091	3,712	8,538	33	\$76
Humpback, or pink:								
1-pound tall.....	c 55	132						
Silverside, coho, or white:								
1-pound flat.....	3,304	9,252	2,546	7,129	2,119	5,933		
1-pound flat.....	8,220	36,155						
1-pound tall.....	5,817	23,850	3,281	13,124	3,909	15,876	1,139	4,556
Total.....	17,341	69,257	5,827	20,253	6,088	21,809	1,139	4,556
Steelhead trout:								
½-pound flat.....	7,064	22,084						
1-pound flat.....	1,365	7,695						
1-pound tall.....	4,320	25,056						
Total.....	12,749	54,835						
Grand total.....	214,032	1,112,090	8,607	32,886	12,893	46,010	1,172	4,632

Products.	Alesea River and Bay.		Siuslaw River.		Umpqua River.		Coos Bay.	
	Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
Chinook, or king:								
½-pound flat.....	928	\$2,784					50	\$150
1-pound flat.....							211	1,013
1-pound tall.....	655	3,930	632	\$3,792	500	\$3,000		
1-pound oval.....							39	312
Total.....	1,583	6,714	632	3,792	500	3,000	300	1,475
Chum, or dog:								
1-pound tall.....	80	184						
Silverside, coho, or white:								
½-pound flat.....	2,601	7,283	4,017	11,248			2,088	5,846
1-pound flat.....							1,841	8,100
1-pound tall.....	4,186	16,734	5,427	21,708	7,753	31,012	759	3,036
2-pound nominal.....							315	945
Total.....	6,787	24,027	9,444	32,956	7,753	31,012	5,003	17,927
Grand total.....	8,450	30,925	10,076	36,748	8,253	34,012	5,303	19,402

a Of these, 4,595 cases, valued at \$18,696, were filled with sockeyes brought from Puget Sound, Wash.

b Packed with sockeye salmon from Puget Sound, Wash.

c Packed with humpback salmon from Puget Sound, Wash.

PACK OF CANNED SALMON IN OREGON, BY WATERS, IN 1909—Continued.

Products.	Coquille River.		Rogue River.		Total.	
	Cases.	Value.	Cases.	Value.	Cases.	Value.
Blueback, or sockeye:						
½-pound flat					32,071	\$133,095
1-pound flat					6,645	39,870
1-pound tall					50	320
Total					38,766	173,285
Chinook, or king:						
½-pound flat					69,557	289,534
1-pound flat	204	\$979	186	\$1,300	54,591	396,809
1-pound tall	46	276			23,057	148,815
½-pound oval					534	2,670
1-pound oval					848	8,242
2-pound nominal					458	1,833
Total	250	1,255	186	1,300	149,045	847,903
Chum, or dog:						
1-pound tall					9,225	21,218
Humpback, or pink:						
1-pound tall					55	132
Silverside, coho, or white:						
½-pound flat	3,656	10,237			20,331	56,928
1-pound flat	1,226	5,394	468	2,053	11,755	51,702
1-pound tall	6,764	27,056	231	924	39,326	157,886
2-pound nominal					315	945
Total	11,646	42,687	699	2,977	71,727	267,461
Steelhead trout:						
½-pound flat					7,064	22,084
1-pound flat					1,365	7,695
1-pound tall					4,320	25,056
Total					12,749	54,835
Grand total.....	11,896	43,942	885	4,277	a 281,567	1,364,834

a All 1-pound cases contain 48 1-pound cans; the ½-pound cases contain 48 ½-pound cans. Reduced to a common basis of 48 1-pound cans the pack is 216,788½ cases.

Miscellaneous secondary products.—The Columbia River produces a large part of the miscellaneous secondary products. Mild-cured salmon form the greater part of the pack, followed by frozen, smoked, and pickled salmon in the order named.

PACK OF MISCELLANEOUS SECONDARY PRODUCTS IN OREGON, BY WATERS, IN 1909.

Products.	Columbia River.		Nehalem River.		Tillamook Bay.		Siletz River.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Frozen:								
Chinook	14,000	\$1,400						
Silverside	216,175	13,868						
Steelhead trout.....	1,414,662	141,767						
Total	1,644,837	157,035						
Mild-cured:								
Chinook	3,909,846	390,984	15,485	\$1,239	59,595	\$4,768	41,575	\$4,003
Smoked:								
Chinook	127,700	19,155						
Silverside	20,000	2,000						
Total	147,700	21,155						
Grand total.....	5,702,383	569,174	15,485	1,239	59,595	4,768	41,575	4,003

PACK OF MISCELLANEOUS SECONDARY PRODUCTS IN OREGON, BY WATERS, IN 1909—Continued.

Products.	Alsea River and Bay.		Siuslaw River.		Umpqua River.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Mild-cured:						
Chinook.....	32,386	\$3,158	12,000	\$960	4,002	\$240
Pickled:						
Chinook.....			400	24		
Silverside.....			2,600	130		
Total.....			3,000	154		
Grand total.....	32,386	3,158	15,000	1,114	4,002	240

Products.	Coos Bay.		Rogue River.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Frozen:						
Chinook.....					14,000	\$1,400
Silverside.....					216,175	13,868
Steelhead trout.....			32,023	\$2,891	1,446,685	144,658
Total.....			32,023	2,891	1,676,860	159,926
Mild-cured:						
Chinook.....	48,000	\$4,800	242,553	24,673	4,365,442	434,825
Pickled:						
Chinook.....					400	24
Silverside.....					2,600	130
Total.....					3,000	154
Smoked:						
Chinook.....					127,700	19,155
Silverside.....					20,000	2,000
Total.....					147,700	21,155
Grand total.....	48,000	4,800	274,576	27,564	6,193,002	616,060

CALIFORNIA.

In Eel River the runs of all species of salmon were very poor. For the first few days of the season the catch was very heavy, after which the run dwindled down to almost nothing. Nearly all of these were shipped fresh to San Francisco, where the dealers claimed that most of them arrived in bad condition.

In the Sacramento River the run was a very fair one, and all of the product was marketed in either a fresh, mild-cured, or smoked condition, none being canned. The interesting table following shows the daily deliveries of chinook salmon to one of the mild-curing plants on the river, and the total and average weights of same.

DAILY DELIVERIES OF CHINOOK SALMON TO A MILD-CURING PLANT ON THE SACRAMENTO RIVER, SEASON OF 1909.

Date.	Number.	Total weight.	Average.	Date.	Number.	Total weight.	Average.
SPRING, 1909.				FALL, 1909.			
Apr. 16.....	21	421	20.0	Aug. 17.....	279	6,658	23.8
17.....	13	297	22.0	18.....	325	8,021	24.6
19.....	109	2,411	22.0	19.....	147	4,018	27.3
20.....	305	7,512	24.6	20.....	185	4,954	26.7
21.....	111	2,826	25.4	21.....	39	1,011	25.9
22.....	183	4,510	24.6	23.....	1,731	42,829	24.7
23.....	331	7,708	23.2	24.....	458	11,885	26.0
24.....	163	3,919	24.0	25.....	279	7,444	26.7
26.....	284	5,918	23.8	26.....	315	8,250	26.0
27.....	75	1,788	23.8	27.....	145	3,747	25.8
28.....	104	2,391	23.0	28.....	86	2,309	28.0
29.....	116	2,716	23.2	30.....	1,300	32,926	25.3
30.....	358	8,059	23.0	31.....	812	21,018	25.8
May 1.....	251	5,739	22.8	Sept. 1.....	628	16,331	26.0
3.....	171	4,016	23.4	2.....	356	9,654	27.1
4.....	175	4,128	23.5	3.....	242	6,582	27.1
5.....	107	2,490	22.6	4.....	105	2,885	27.4
6.....	66	1,680	25.4	6.....	1,176	31,640	26.9
7.....	132	2,957	22.4	7.....	915	24,277	26.5
8.....	96	2,287	23.8	8.....	758	19,874	26.2
10.....	308	7,302	23.3	9.....	704	18,851	26.7
12.....	152	3,717	24.4	10.....	677	18,204	26.8
13.....	89	2,056	23.1	11.....	369	9,592	26.0
14.....	274	6,635	24.2	13.....	1,917	49,781	25.9
15.....	254	6,201	24.4	14.....	1,343	35,555	26.4
17.....	310	7,378	23.8	15.....	751	20,097	26.7
18.....	323	7,844	24.2	16.....	647	17,328	26.7
19.....	210	5,037	23.9	17.....	1,493	35,883	24.0
20.....	226	5,246	23.2	Total.....	18,182	471,607	25.9
21.....	154	3,778	24.5	Grand total.....	26,201	661,699	25.45
22.....	166	4,150	25.0				
24.....	315	7,290	23.1				
25.....	422	9,917	23.5				
26.....	342	7,767	22.7				
27.....	245	5,900	24.0				
28.....	268	6,496	24.2				
29.....	197	4,826	24.5				
30.....	330	7,529	22.8				
June 1.....	299	7,250	24.2				
Total.....	8,019	190,092	23.7				

The southernmost point on our coast where salmon are taken commercially is in Monterey Bay, and it is here that trolling was first engaged in to any extent. Yearly the chinooks come into Monterey and Santa Cruz Bays, where they sometimes remain feeding for months. When they strike in, which in numbers they usually do the latter part of April, they are in the pursuit of squid, sardines, anchovies, and other small fish, and their presence is first indicated to the fishermen by the occasional disturbances of the surface by the small fish. It is a signal for the fishermen and sportsmen, who go out in both sail and row boats.

During 1909 most of the catch was made in the vicinity of Monterey, the salmon appearing in but small numbers in Santa Cruz Bay.

While evidently coming in schools at first, salmon soon scatter about in pursuit of their prey, thus making the use of nets unprofitable. In a dead calm troll fishing practically ceases, but with the return of the breeze the fish resume biting.

The silver salmon come into Monterey Bay in July and are usually taken in that one month alone. Some of them run as large as 12 to 13 pounds each and all are feeding.

During 1909 the dealers had an agreement with their fishermen, who are mostly Japanese, under which they kept back a certain percentage of the price until the end of the season. This was done in order to make certain that the fishermen would not go off and sell to some one else the better fish and bring them the poorer quality.

The following table shows the daily receipts of chinook salmon at the mild-curing plant of one of the companies operating at Monterey during 1909. The table also shows the number of boats fishing, the number of fish caught, and the total weight of same, and the average weight per fish:

DAILY DELIVERIES OF CHINOOK SALMON AT A MILD-CURING PLANT ON MONTEREY BAY, SEASON OF 1909.

Date.	Number of boats.	Number of fish.	Total weight.	Average weight.	Date.	Number of boats.	Number of fish.	Total weight.	Average weight.
1909.					1909.				
Apr. 30.....	70	966	10,002	18.3	June 21.....	106	1,808	30,090	16.6
May 1.....	69	319	4,096	12.8	22.....	110	1,678	20,576	12.2
3.....	12	20	369	18.4	23.....	104	1,135	15,964	14.0
4-5.....	30	152	2,512	16.5	24.....	111	1,811	26,826	14.5
6.....	41	126	1,758	14.0	25.....	100	595	9,549	16.0
7.....	35	93	1,084	11.6	26.....	108	615	9,645	15.0
8.....	23	47	1,602	13.0	27.....	46	142	1,831	12.7
10.....	15	47	633	13.0	28.....	44	212	2,719	12.8
11.....	28	56	770	13.4	29.....	88	566	7,030	12.5
12.....	82	642	8,210	12.5	30.....	101	1,175	14,499	13.0
13.....	83	613	6,250	10.2	July 1.....	111	1,416	18,363	13.0
14.....	93	847	9,993	11.8	2.....	100	634	8,576	13.5
15.....	103	615	7,835	12.7	3.....	108	1,313	16,060	12.2
16.....	16	26	429	16.0	6.....	113	1,687	24,508	15.0
17.....	107	1,152	14,612	12.7	7.....	114	1,568	20,054	13.0
18.....	87	318	4,607	15.0	8.....	116	1,428	20,401	14.25
19.....	63	135	1,673	12.5	9.....	80	971	13,350	14.0
22.....	31	46	667	15.0	10.....	114	973	13,236	13.5
23.....	82	476	6,043	12.7	11.....	88	581	8,184	14.0
24.....	107	1,652	23,600	14.3	12.....	79	400	5,196	13.0
25.....	114	3,390	50,621	15.0	13.....	62	407	4,847	12.0
26.....	118	1,190	17,590	12.0	14.....	91	466	5,469	11.7
27.....	54	94	1,619	17.0	15.....	98	513	6,166	12.0
28.....	68	222	3,458	15.5	16.....	85	495	5,713	11.6
29.....	93	650	9,874	15.5	17.....	85	506	5,697	11.25
30.....	118	2,852	38,567	13.5	19.....	55	257	3,187	12.4
June 1.....	119	1,005	14,625	14.0	20.....	91	422	5,565	13.1
2.....	95	493	8,273	17.0	21.....	62	205	3,252	15.75
3.....	115	1,245	20,256	17.0	22.....	68	356	5,178	15.0
4.....	109	1,000	14,304	14.0	23.....	79	460	6,237	13.5
5.....	112	724	10,437	14.0	24.....	95	1,284	15,391	12.0
6.....	96	1,615	22,571	14.0	26.....	108	1,176	16,437	14.0
7.....	114	988	12,901	13.0	27.....	104	1,487	22,766	15.30
8.....	95	485	7,042	14.5	28.....	105	961	18,576	19.5
9.....	80	307	4,804	16.0	29.....	88	267	5,521	20.7
10.....	68	200	3,437	17.0	30.....	59	114	2,548	22.7
11.....	66	243	4,786	22.0	Aug. 1.....	47	144	2,832	19.9
12.....	83	348	6,187	19.0	2.....	79	287	4,908	17.0
13.....	95	623	10,218	16.0	3.....	43	78	1,574	20.0
14.....	106	499	7,965	16.0	4.....	21	71	1,366	19.0
15.....	89	390	6,655	18.0	5.....	43	170	3,546	20.9
16.....	112	1,729	27,524	16.0	6.....	70	274	4,845	18.0
17.....	115	3,092	48,138	15.4	7.....	52	114	2,156	19.0
18.....	105	1,395	24,436	17.6	9-12.....	12	20	502	25.0
19.....	117	3,725	61,789	16.7					
20.....	112	2,083	35,265	17.0	Total.....		71,619	1,043,358	14.6
	111	1,442	23,335	16.2					

STATISTICS BY COUNTIES.

Persons employed.—The total number of persons employed was 2,675, Contra Costa County leading with 774 persons.

PERSONS ENGAGED IN THE CALIFORNIA SALMON FISHERIES, BY COUNTIES, IN 1909.

Counties.	Fishermen.				Shoresmen. ^a				Trans- porters (whites).	Grand total.
	Whites.	Japa- nese.	Chi- nese.	Total.	Whites.	Indians.	Japa- nese.	Total.		
Del Norte.....	84			84	17	15		32	3	119
Humboldt.....	339			339	19			19		358
Alameda.....					25			25		25
Marin.....	8			8						8
San Francisco.....	60			60	60			60	8	128
Solano.....	420			420	50			50	24	494
Contra Costa.....	654			654	78			78	42	774
San Joaquin.....	64	24		88						88
Yolo.....	42			42						42
Sacramento.....	178			178						178
Sutter.....	12			12						12
Butte.....	45			45					5	50
Glenn.....	20			20	1			1		21
Tehama.....	45			45			5	5		50
Shasta.....	10			10						10
Monterey.....	65	144	15	224	26			26		250
Santa Cruz.....	68			68						68
Total.....	2,114	168	15	2,297	276	15	5	296	82	2,675

^a All the shoresmen reported for Alameda County and part of those reported for San Francisco County are employed by one of the Alaskan canning companies and have been reported here, as they are employed here the whole year.

Investment, apparatus, etc.—The total investment amounts to \$1,232,960. The shore property reported for Alameda County belongs to one of the companies operating in Alaska. Contra Costa leads in the total investment. Gill nets, haul seines, and trolling lines are the principal forms of apparatus in use.

INVESTMENT IN THE SALMON FISHERIES OF CALIFORNIA, BY COUNTIES, IN 1909.

Items.	Del Norte.		Humboldt.		Alameda.		Marin.		San Francisco.	
	Num- ber.	Value.	Num- ber.	Value.	Num- ber.	Value.	Num- ber.	Value.	Num- ber.	Value.
Transporting vessels:										
Power vessels.....	1	\$3,248							1	\$25,000
Tonnage.....	9								32	
Outfit.....		750								1,240
Power boats.....									4	7,000
Fishing boats, power.....									15	18,000
Fishing boats, sail and row.....	54	2,640	253	\$6,625			4	\$400	15	1,500
House boats and scows.....			2	100						
Apparatus, shore fisheries:										
Haul seines.....	4	550	17	2,450						
Gill nets, drift.....	50	11,300	286	19,375			4	1,050	30	7,875
Shore and accessory property.....		17,020		7,750		\$159,550		50		155,320
Cash capital.....		10,000		4,500						43,500
Total.....		45,508		40,800		159,550		1,500		259,435

INVESTMENT IN THE SALMON FISHERIES OF CALIFORNIA, BY COUNTIES, IN 1909—Continued.

Items.	Solano.		Contra Costa.		San Joaquin.		Yolo.		Sacramento.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Transporting vessels:										
Power vessels.....	1	\$4,000	1	\$5,500						
Tonnage.....	10		5							
Outfit.....		1,000		930						
Power boats.....	14	19,500	23	36,800						
Fishing boats, power.....	30	10,400	32	21,000	28	\$8,400	4	\$1,600	17	\$6,800
Fishing boats, sail and row.....	183	36,400	300	58,500	16	2,900	17	990	77	5,170
House boats and scows.....	10	4,000	11	4,800			5	1,000	19	3,650
Apparatus, shore fisheries:										
Gill nets, drift.....	210	39,500	322	64,400	44	6,600	21	2,550	113	14,320
Hand lines.....				10						
Shore and accessory property.....		29,900		117,113		580		145		815
Cash capital.....		50,000		85,000						
Total.....		194,700		394,053		18,480		6,285		30,755

Items.	Sutter.		Butte.		Glenn.		Tehama.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Apparatus, shore fisheries:								
Fishing boats, sail and row.....	6	\$375	20	\$840	6	\$300	20	\$1,000
House boats and scows.....	3	375						
Haul seines.....			10	1,000	4	400	10	1,020
Gill nets, drift.....	6	600						
Shore and accessory property.....		50		2,075		600		2,150
Total.....		1,400		3,915		1,300		4,170

Items.	Shasta.		Monterey.		Santa Cruz.		Total.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Transporting vessels:								
Power vessels.....							4	\$37,748
Tonnage.....							56	
Outfit.....								3,920
Power boats.....							41	63,300
Fishing boats, power.....			24	\$13,850	21	\$11,000	171	91,050
Fishing boats, sail and row.....	4	\$200	170	7,805	13	2,600	1,158	128,245
House boats and scows.....							50	13,925
Apparatus, shore fisheries:								
Haul seines.....	2	230					^a 47	5,650
Gill nets, drift.....							^b 1,086	167,570
Trolling lines.....				886		263		1,149
Hand lines.....								10
Shore and accessory property.....		275		3,900		100		497,393
Cash capital.....				30,000				223,000
Total.....		705		56,441		13,963		1,232,960

^a Aggregate length of 13,449 yards.^b Aggregate length of 438,420 yards.

Catch.—The total catch amounts to 12,141,937 pounds, valued at \$585,995. Contra Costa County leads in catch, followed closely by Solano County. Nearly four-fifths of the catch was made with gill nets, while chinook salmon comprise almost all of the catch.

PRODUCTS OF THE SALMON FISHERIES OF CALIFORNIA, BY APPARATUS AND SPECIES, IN 1909.

Apparatus and species.	Del Norte.		Humboldt.		Marin.		San Francisco.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
GILL NETS.								
Blueback.....			9,300	\$317				
Chinook, fresh.....	524,225	\$8,532	463,649	16,970	5,380	\$310	91,063	\$4,055
Chinook, salted.....	27,000	1,220						
Silver, fresh.....	50,000	900	23,000	690				
Silver, salted.....	20,000	1,000						
Steelhead trout.....			4,700	235				
Total.....	621,225	11,652	500,649	18,212	5,380	310	91,063	4,055
SEINES.								
Blueback.....			11,700	372				
Chinook, fresh.....			301,600	12,064				
Chinook, salted.....	10,000	400	32,049	2,932				
Silver, fresh.....			12,000	360				
Silver, salted.....	24,000	800	2,000	100				
Dog.....			4,200	84				
Total.....	34,000	1,200	363,549	15,912				
TOTAL.								
Blueback.....			21,000	689				
Chinook, fresh.....	524,225	8,532	765,249	29,034	5,380	310	91,063	4,055
Chinook, salted.....	37,000	1,620	32,049	2,932				
Silver, fresh.....	50,000	900	35,000	1,050				
Silver, salted.....	44,000	1,800	2,000	100				
Dog.....			4,200	84				
Steelhead trout.....			4,700	235				
Grand total.....	655,225	12,852	864,198	34,124	5,380	310	91,063	4,055

Apparatus and species.	Solano.		Contra Costa.		San Joaquin.		Yolo.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
GILL NETS.								
Chinook, fresh.....	3,238,788	\$168,713	3,944,902	\$210,855	61,187	\$2,585	197,520	\$10,852
Steelhead trout.....			678	41				
Total.....	3,238,788	168,713	3,945,580	210,896	61,187	2,585	197,520	10,852
LINES.								
Steelhead trout.....			3,500	270				
Total.....			3,500	270				
TOTAL.								
Chinook, fresh.....	3,238,788	168,713	3,944,902	210,855	61,187	2,585	197,520	10,852
Steelhead trout.....			4,178	311				
Grand total.....	3,238,788	168,713	3,949,080	211,166	61,187	2,585	197,520	10,852

PRODUCTS OF THE SALMON FISHERIES OF CALIFORNIA, BY APPARATUS AND SPECIES,
IN 1909—Continued.

Apparatus and species.	Sacramento.		Sutter.		Butte.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
GILL NETS.						
Chinook, fresh.....	599,723	\$32,690	62,119	\$1,917		
Total.....	599,723	32,690	62,119	1,917		
SEINES.						
Chinook, fresh.....					163,022	\$8,285
Total.....					163,022	8,285
TOTAL.						
Chinook, fresh.....	599,723	32,690	62,119	1,917	163,022	8,285
Grand total.....	599,723	32,690	62,119	1,917	163,022	8,285
Apparatus and species.	Glenn.		Tehama.		Shasta.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
SEINES.						
Chinook, fresh.....	72,547	\$3,627	314,102	\$16,905	46,475	\$2,789
Total.....	72,547	3,627	314,102	16,905	46,475	2,789
TOTAL.						
Chinook, fresh.....	72,547	3,627	314,102	16,905	46,475	2,789
Grand total.....	72,547	3,627	314,102	16,905	46,475	2,789
Apparatus and species.	Monterey.		Santa Cruz.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
GILL NETS.						
Blueback.....					9,300	\$317
Chinook, fresh.....					9,188,556	457,479
Chinook, salted.....					27,000	1,220
Silver, fresh.....					73,090	1,590
Silver, salted.....					20,000	1,000
Steelhead trout.....					5,378	276
Total.....					9,323,234	461,882
SEINES.						
Blueback.....					11,700	372
Chinook, fresh.....					897,746	43,670
Chinook, salted.....					42,049	3,332
Silver, fresh.....					12,000	360
Silver, salted.....					26,000	900
Dog.....					4,200	84
Total.....					993,695	48,718
LINES.						
Chinook.....	1,769,524	\$72,634	37,373	\$1,759	1,806,897	74,393
Silver.....	10,000	500	4,500	225	14,500	725
Steelhead trout.....			111	7	3,611	277
Total.....	1,779,524	73,134	41,984	1,991	1,825,008	75,395
TOTAL.						
Blueback.....					21,000	689
Chinook, fresh.....	1,769,524	72,634	37,373	1,759	11,893,199	575,542
Chinook, salted.....					69,049	4,552
Silver, fresh.....	10,000	500	4,500	226	99,500	2,675
Silver, salted.....					46,000	1,900
Dog.....					4,200	84
Steelhead trout.....			111	7	8,989	553
Grand total.....	1,779,524	73,134	41,984	1,991	12,141,937	585,995

STATISTICS BY WATERS.

Persons employed.—Of the 2,675 persons employed in the industry, 1,880 were on the Sacramento River. The next largest number was employed on Monterey Bay.

PERSONS ENGAGED IN THE SALMON FISHERIES OF CALIFORNIA, BY WATERS AND NATIONALITIES, IN 1909.

Occupation and race.	Smith River.	Klamath River.	Mad River.	Eureka Bay.	Eel River.	Sacramento River.	Monterey Bay.	Total.
Fishermen:								
Whites.....	47	37	41	7	291	1,558	133	2,114
Chinese.....							15	15
Japanese.....						24	144	168
Total.....	47	37	41	7	291	1,582	292	2,297
Shoresmen:								
Whites.....	17			6	13	214	26	276
Indians.....	15							15
Japanese.....						5		5
Total.....	32			6	13	219	26	296
Transporters:								
Whites.....		3				79		112
Total:								
Whites.....	64	40	41	13	304	1,851	159	2,472
Indians.....	15							15
Chinese.....							15	15
Japanese.....						29	144	173
Grand total....	79	40	41	13	304	1,880	318	2,675

Investment, apparatus, etc.—More than nine-tenths of the total investment is represented in the Sacramento River. Trolling lines are used in Monterey Bay.

INVESTMENT IN THE SALMON FISHERIES OF CALIFORNIA, BY WATERS, IN 1909.

Items.	Smith River.		Klamath River.		Mad River.		Eureka Bay.	
	Num-ber.	Value.	Num-ber.	Value.	Num-ber.	Value.	Num-ber.	Value.
Transporting vessels:								
Power vessels.....			1	\$3,248				
Tonnage.....			9					
Outfit.....				750				
Fishing boats, sail and row.....	23	\$770	31	1,870	33	\$865	7	\$175
Apparatus, shore fisheries:								
Haul seines.....	4	550			4	500		
Gill nets, drift.....	15	800	35	10,500	37	1,800	7	525
Shore and accessory property.....		420		16,600		100		900
Cash capital.....				10,000				1,500
Total.....		2,540		42,968		3,265		3,100

INVESTMENT IN THE SALMON FISHERIES OF CALIFORNIA, BY WATERS, IN 1909—Continued.

Items.	Eel River.		Sacramento River.		Monterey Bay.		Total.	
	Number.	Value.	Number.	Value.	Number.	Value.	Number.	Value.
Transporting vessels:								
Power vessels.....			3	\$34,500			4	\$37,748
Tonnage.....			47				56	
Outfit.....				3,170				3,920
Power boats.....			41	63,300			41	63,300
Fishing boats, power.....			126	66,200	45	\$24,850	171	91,050
Fishing boats, sail and row.....	213	\$5,585	668	108,575	183	10,405	1,158	128,245
Scows and house boats.....	2	100	48	13,825			50	13,925
Apparatus, shore fisheries:								
Haul seines.....	13	1,950	26	2,650			47	5,650
Gill nets, drift.....	242	17,050	750	136,895			1,086	167,570
Trolling lines.....						1,149		1,149
Hand lines.....				10				10
Shore and accessory property.....		6,750		468,623		4,000		497,393
Cash capital.....		3,000		178,500		30,000		223,000
Total.....		34,435		1,076,248		70,404		1,232,960

Catch.—About four-fifths of the total catch was made on the Sacramento River; Monterey Bay was second and Eel River third. With the exception of Monterey Bay, gill nets take the largest part of the catch on all the waters. The catch of species other than chinook is very small.

PRODUCTS OF THE SALMON FISHERIES OF CALIFORNIA, BY APPARATUS, SPECIES, AND WATERS, IN 1909.

Apparatus and species.	Smith River.		Klamath River.		Mad River.		Eureka Bay.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
GILL NETS.								
Blueback.....					3,800	\$152		
Chinook, fresh.....	40,000	\$1,200	484,225	\$7,332	50,000	2,000	28,000	\$840
Chinook, salted.....	20,000	800	7,000	420				
Silver, fresh.....			50,000	900	12,000	360		
Silver, salted.....			20,000	1,000				
Total.....	60,000	2,000	561,225	9,652	65,800	2,512	28,000	840
HAUL SEINES.								
Blueback.....					2,100	84		
Chinook, fresh.....					28,000	1,120		
Chinook, salted.....	10,000	400			6,000	360		
Silver, fresh.....					7,000	210		
Silver, salted.....	24,000	800						
Total.....	34,000	1,200			43,100	1,774		
TOTAL.								
Blueback.....					5,900	236		
Chinook, fresh.....	40,000	1,200	484,225	7,332	78,000	3,120	28,000	840
Chinook, salted.....	30,000	1,200	7,000	420	6,000	360		
Silver, fresh.....			50,000	900	19,000	570		
Silver, salted.....	24,000	800	20,000	1,000				
Grand total.....	94,000	3,200	561,225	9,652	108,900	4,286	28,000	840

PRODUCTS OF THE SALMON FISHERIES OF CALIFORNIA, BY APPARATUS, SPECIES, AND WATERS, IN 1909—Continued.

Apparatus and species.	Eel River.		Sacramento River.		Monterey Bay.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
GILL NETS.								
Blueback.....	5,500	\$165					9,300	\$327
Chinook, fresh.....	385,649	14,130	8,200,682	\$431,977			9,185,556	457,479
Chinook, salted.....							27,000	1,220
Silver, fresh.....	11,000	330					73,000	1,590
Silver, salted.....							20,000	1,000
Steelhead trout.....	4,700	235	678	41			5,378	276
Total.....	406,849	14,860	8,201,360	432,018			9,323,234	461,892
HAUL SEINES.								
Blueback.....	9,600	288					11,700	372
Chinook, fresh.....	273,600	10,944	596,146	31,606			897,746	43,670
Chinook, salted.....	26,049	2,572					42,049	3,332
Dog, or chum.....	4,200	84					4,200	84
Silver, fresh.....	5,000	150					12,000	360
Silver, salted.....	2,000	100					26,000	900
Total.....	320,449	14,138	596,146	31,606			993,695	48,718
LINES.								
Chinook.....					1,806,897	\$74,393	1,806,897	74,393
Silver.....					14,500	725	14,500	725
Steelhead trout.....			3,500	270	111	7	3,611	277
Total.....			3,500	270	1,821,508	75,125	1,823,008	75,395
TOTAL.								
Blueback.....	15,100	453					21,000	689
Chinook, fresh.....	659,249	25,074	8,796,828	463,583	1,806,897	74,393	11,893,199	575,542
Chinook, salted.....	26,049	2,572					69,049	4,552
Dog, or chum.....	4,200	84					4,200	84
Silver, fresh.....	16,000	480			14,500	725	99,500	2,675
Silver, salted.....	2,000	100					46,000	1,900
Steelhead trout.....	4,700	235	4,178	311	111	7	8,989	553
Grand total.....	727,298	28,998	8,801,006	463,894	1,821,508	75,125	12,141,937	585,995

Products canned.—But one cannery was operated in 1909, and that at Requa, on the Klamath River. The pack of this cannery was 5,663 cases of 1-pound flat chinooks, which sold for \$28,315.

Miscellaneous secondary products.—Mild-cured and smoked salmon comprise the secondary products prepared.

PACK OF MISCELLANEOUS SECONDARY PRODUCTS IN CALIFORNIA, BY WATERS, IN 1909.

Products.	Eel River.		Sacramento River.		Monterey Bay.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Mild-cured:								
Chinook.....	64,000	\$6,400	4,095,162	\$450,019	728,800	\$64,049	4,887,962	\$520,468
Smoked:								
Chinook.....	50,000	5,000	56,550	8,943	4,000	700	110,550	14,643
Silver.....	3,000	300	4,660	326			7,660	626
Total.....	53,000	5,300	61,210	9,269	4,000	700	118,210	15,269
Grand total.....	117,000	11,700	4,156,372	459,288	732,800	64,749	5,006,172	535,737

ALASKA.

The season of 1909 was a very quiet one in Alaska. Owing to the expected quadrennial heavy run of sockeye salmon on Puget Sound, several cannery men who operate there and in Alaska shut down their Alaska plants and devoted all their energies to the Sound, which materially reduced the amount of fishing gear used in Alaska, and as a consequence the total quantity of products produced. In western Alaska the ice hampered operations in the early part of the season, but, with the exception of the Ugashik and Ugaguk Rivers, the runs were fairly good. The weather was very severe on Nusagak Bay and as a result eight fishermen lost their lives there by drowning. In Central Alaska the run of salmon in the neighborhood of Karluk fell off very materially as compared with 1908, but in Chignik the usual good run appeared. In southeast Alaska, except in the lower portion, the run was very good, but the cannery men packed no more of the cheaper grades than they felt could be disposed of at the then unremunerative prices prevailing.

Persons engaged.—The total number of persons engaged in the Alaska salmon fisheries was 11,433. Western Alaska leads in the total number, followed by southeast and central Alaska in the order named. A large number of Indians are employed in this industry.

PERSONS ENGAGED IN THE ALASKA SALMON FISHERIES IN 1909.

Occupation and race.	Southeast Alaska.	Central Alaska.	Western Alaska.	Total.
Fishermen:				
Whites.....	662	400	1,424	2,486
Indians.....	982	184	10	1,176
Japanese.....	13			13
Total.....	1,657	584	1,434	3,675
Shoresmen:				
Whites.....	442	277	1,192	1,911
Indians.....	815	124	307	1,246
Chinese.....	546	377	1,069	1,992
Japanese.....	348	356	1,432	2,136
Total.....	2,151	1,134	4,000	7,285
Transporters:				
Whites.....	148	108	187	443
Indians.....	13	17		30
Total.....	161	125	187	473
Total:				
Whites.....	1,252	785	2,803	4,840
Indians.....	1,810	325	317	2,452
Chinese.....	546	377	1,069	1,992
Japanese.....	361	356	1,432	2,149
Grand total.....	3,969	1,843	5,621	11,433

Investments, apparatus, etc.—The total investment amounted to \$13,948,271. Gill nets predominate, while purse and haul seines and stationary traps are important.

INVESTMENT IN THE ALASKA SALMON FISHERIES IN 1909.

Items.	Southeast Alaska.		Central Alaska.		Western Alaska.		Total.	
	Num-ber.	Value.	Num-ber.	Value.	Num-ber.	Value.	Num-ber.	Value.
Transporting vessels:								
Power vessels.....	69	\$263,256	25	\$213,019	39	\$591,669	133	\$1,067,944
Tonnage.....	1,173		1,482		3,236		5,891	
Outfit.....		65,814		53,255		147,917		266,986
Sailing vessels.....	5	158,000	9	289,000	29	638,400	43	1,085,400
Tonnage.....	7,434		14,270		38,057		59,761	
Outfit.....		15,800		28,900		63,840		108,540
Power boats.....	11	11,760	4	8,400	2	4,680	17	24,840
Fishing boats, power.....	60	30,000					60	30,000
Fishing boats, sail and row.....	766	25,981	300	21,215	755	164,475	1,821	211,671
Scows and house boats.....	98	38,175	79	30,930	133	101,900	310	171,005
Pile drivers.....	13	34,405	15	29,850	15	26,300	43	90,555
Apparatus, shore fisheries:								
Haul seines.....	45	12,451	49	15,280			a 94	27,731
Purse seines.....	98	27,188					b 98	27,188
Gill nets, drift.....	256	34,030	57	11,020	896	66,706	c 1,209	111,756
Traps, stationary.....	36	79,700	20	29,450	17	21,644	73	130,794
Traps, floating.....	14	19,750	1	1,500			15	21,250
Lines.....		523						523
Spears.....	20	30					20	30
Shore and accessory property.....		1,788,902		1,200,716		2,611,641		5,601,259
Cash capital.....		2,223,493		890,531		1,856,775		4,970,799
Total.....		4,829,258		2,823,066		6,295,947		13,948,271

^a Aggregate length of 30,430 yards.

^c Aggregate length of 301,480 yards.

^b Aggregate length of 35,670 yards.

Catch.—The total catch amounted to 175,934,060 pounds, valued at \$1,333,344. Red or sockeye salmon comprise almost two-thirds of the total catch. As compared with 1908, the catch of all species, except king salmon, decreased very materially, due to causes described elsewhere.

CATCH, BY SPECIES AND APPARATUS, IN THE SALMON FISHERIES OF ALASKA IN 1909.

Apparatus and species.	Southeast Alaska.		Central Alaska.		Western Alaska.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
SEINES.								
Coho, or silver.....	991,062	\$13,214	313,548	\$2,090			1,304,610	\$15,304
Dog, or chum.....	3,102,192	5,817					3,102,192	5,817
Humpback, or pink.....	22,288,020	55,720	510,196	957			22,798,216	56,677
King, or spring.....	6,446	193	85,954	195			92,400	388
Red, or sockeye.....	6,426,325	102,821	10,194,165	\$1,553			16,620,490	184,374
Total.....	32,814,045	177,765	11,103,863	84,795			43,917,908	262,560
TRAPS.								
Coho, or silver.....	673,278	8,977	539,508	3,597	59,580	\$397	1,272,366	12,971
Dog, or chum.....	2,699,160	5,061			811,648	1,015	3,510,808	6,076
Humpback, or pink.....	14,515,760	36,289	14,960	28	60	1	14,530,780	36,318
King, or spring.....	112,354	3,371	981,904	2,232	68,112	155	1,162,370	5,758
Red, or sockeye.....	5,362,896	71,505	10,762,775	86,102	2,540,055	20,320	18,665,726	177,927
Total.....	23,363,448	125,203	12,299,147	91,959	3,479,455	21,888	39,142,050	239,050

CATCH, BY SPECIES AND APPARATUS, IN THE SALMON FISHERIES OF ALASKA IN 1909—Continued.

Apparatus and species.	Southeast Alaska.		Central Alaska.		Western Alaska.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
GILL NETS.								
Coho, or silver.....	473,070	\$6,308			428,358	\$6,010	901,428	\$12,318
Dog, or chum.....	72,328	136			2,770,720	3,554	2,843,048	3,690
Humpback, or pink.	509,688	1,274			127,244	796	636,932	2,070
King, or spring.....	1,510,498	45,315	397,298	\$902	2,835,646	10,781	4,743,442	56,998
Red, or sockeye.....	2,391,990	38,272	2,439,920	19,519	75,669,360	605,355	80,501,270	663,146
Total.....	4,957,574	91,305	2,837,218	20,421	81,831,328	626,496	89,626,120	738,222
LINES.								
Coho, or silver.....	48,000	640					48,000	640
King, or spring.....	2,961,332	88,840					2,961,332	88,840
Steelhead trout.....	11,650	400					11,650	400
Total.....	3,020,982	89,880					3,020,982	89,880
SPEARS.								
Red, or sockeye....	227,000	3,632					227,000	3,632
TOTAL.								
Coho, or silver.....	2,185,410	29,139	853,056	5,687	487,938	6,407	3,526,404	41,233
Dog, or chum.....	5,873,680	11,014			3,582,368	4,569	9,456,048	15,583
Humpback, or pink.	37,313,468	93,283	525,156	985	127,304	797	37,965,928	95,065
King, or spring.....	4,590,630	137,719	1,465,156	3,329	2,903,758	10,936	8,959,544	151,984
Red, or sockeye.....	14,408,211	216,230	23,396,860	187,174	78,209,415	625,675	116,014,486	1,029,079
Steelhead trout.....	11,650	400					11,650	400
Grand total..	64,383,049	487,785	26,240,228	197,175	85,310,783	648,384	175,934,060	1,333,344

Products canned.—The total canned pack amounted to 2,403,669 pound and half-pound cases, valued at \$9,438,152. More than two-thirds of the pack was composed of red salmon. Three canneries were not operated, which very materially reduced the size of the pack.

OUTPUT OF SALMON FROM THE CANNERIES IN ALASKA IN 1909, BY SPECIES AND SIZE OF CANS.^a

Products.	Southeast Alaska.		Central Alaska.		Western Alaska.		Total.	
	Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
Coho, or silver:								
1-pound flat.....	1,206	\$5,543					1,206	\$5,543
1-pound tall.....	38,714	155,431	10,275	\$43,155	6,361	\$26,900	55,350	225,486
Total.....	39,920	160,974	10,275	43,155	6,361	26,900	56,556	231,029
Dog, or chum:								
1-pound tall.....	83,001	186,454			37,711	87,656	120,712	274,110
Humpback, or pink:								
1-pound tall.....	455,999	1,092,389	5,581	13,394	3,293	9,056	464,873	1,114,839
King, or spring:								
1-pound tall.....	857	3,598	16,913	74,418	30,264	129,608	48,034	207,624
Red, or sockeye:								
½-pound flat.....	14,898	58,535			1,487	5,353	16,385	63,888
1-pound flat.....	80,200	209,962	2,936	15,539	2,057	11,108	85,193	236,609
1-pound tall.....	185,444	825,926	355,349	1,625,371	1,071,123	4,858,756	1,611,916	7,310,053
Total.....	280,542	1,094,423	358,285	1,640,910	1,074,667	4,875,217	1,713,494	7,610,550
Grand total....	860,319	2,537,838	391,054	1,771,877	1,152,296	5,128,437	2,403,669	9,438,152

^a All 1-pound cases contain forty-eight 1-pound cans; the ½-pound cases contain forty-eight ½-pound cans. Reduced to a common basis of cases containing forty-eight 1-pound cans the pack is 2,395,477½ cases.

Miscellaneous products.—The total miscellaneous products prepared amounted to 9,473,005 pounds, valued at \$374,324. Owing to the low prices prevailing for pickled salmon, the pack of such very materially declined. Restrictive regulations in regard to the pickling of salmon bellies also aided in reducing the pack. The mild-cured pack shows a gratifying increase over 1908.

MISCELLANEOUS SECONDARY SALMON PRODUCTS PREPARED IN ALASKA IN 1909.

Products.	Southeast Alaska.		Central Alaska.		Western Alaska.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Frozen:								
Coho, or silver.....	35,721	\$1,072	35,721	\$1,072
Dog, or chum.....	77,882	1,558	77,882	1,558
Steelhead trout.....	9,450	473	9,450	473
Total.....	123,053	3,103	123,053	3,103
Mild-cured:								
King, or spring.....	1,833,600	149,300	1,833,600	149,300
Pickled:								
Coho, or silver.....	40,400	1,405	17,800	\$810	5,400	\$270	63,600	2,485
Coho bellies.....	311,400	9,405	227,750	3,843	227,750	3,843
Dog, or chum.....	3,000	90	4,000	100	7,000	190
Humpback.....	11,200	224	311,400	9,405
Humpback backs.....	123,480	6,896	46,000	500	11,200	224
Humpback bellies.....	6,200	248	82,000	3,550	169,480	7,396
King, or spring.....	7,000	175	88,200	3,798
King bellies.....	437,800	17,319	4,863,700	149,979	7,000	175
Red, or sockeye.....	783,600	13,902	5,301,500	167,298
Redbellies.....	783,600	13,902
Total.....	502,680	18,443	1,512,950	36,374	4,955,100	153,899	6,970,730	208,716
Dry-salted and dried:								
Coho backs.....	14,500	549	14,500	549
Dog.....	71,600	1,038	71,600	1,038
Humpback backs.....	50,000	500	1,500	45	51,500	545
King.....	800	45	800	45
Redbacks.....	83,000	2,302	83,000	2,302
Total.....	122,400	1,583	99,000	2,896	221,400	4,479
Smoked:								
Coho backs.....	4,000	400	4,000	400
Dog.....	585	43	585	43
Redbacks.....	28,300	1,580	12,000	1,200	40,300	2,780
Total.....	585	43	32,300	1,980	12,000	1,200	44,885	3,223
Fertilizer.....	159,224	2,287	159,224	2,287
Oil.....	120,113	3,216	^a 120,113	3,216
Grand total.....	2,862,202	177,975	1,644,250	41,250	4,967,100	155,099	9,473,005	374,324

^a Represents 16,015 gallons.

As the fisheries of Alaska are carried on almost wholly in innumerable bays, straits, and sounds, but little being done in the rivers, it does not seem desirable to show them by waters, owing to the amount of space required for the tables.

BRITISH COLUMBIA.

The canned salmon pack of British Columbia was the only branch of the salmon industry of the Province which could be shown by species. Owing to the quadrennially heavy run occurring in the Fraser River in 1909, the pack of British Columbia is quite large. The pack is shown by water areas.

PACK OF CANNED SALMON IN BRITISH COLUMBIA, CANADA, IN 1909.

Species.	Fraser River.		Skeena River.		Rivers Inlet.		Nass River.	
	Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
Coho, or silver:								
½-pound flat.....	710	\$1,988	1,158	\$3,242	264	\$739
1-pound flat.....	5,735	27,528	176	845
1-pound tall.....	15,459	64,928	11,671	49,034	1,092	4,586	6,818	\$28,636
Total.....	21,904	94,444	12,829	52,276	1,532	6,170	6,818	28,636
Dogs, or chums:								
1-pound tall.....	725	1,740	12,000	28,800
Humpback, or pink:								
1-pound flat.....	227	624	40	110
1-pound tall.....	1,053	2,527	16,080	38,640	3,589	8,614
Total.....	1,280	3,151	16,120	38,750	3,589	8,614
King, or spring:								
½-pound flat.....	304	1,216	56	224
1-pound flat.....	1,167	7,032	47	282
½-pound tall.....	176	516
1-pound tall.....	173	934	12,025	64,935	388	1,095	2,309	12,469
1-pound oval.....	444	2,886
Total.....	1,516	8,482	12,469	67,821	739	2,593	2,365	12,693
Sockeye, or red:								
½-pound flat.....	309,634	1,238,536	72,838	291,352	51,520	206,080	11,162	44,648
1-pound flat.....	243,697	1,462,182	19,789	118,734	28,750	172,500	2,070	12,420
½-pound tall.....	2,600	8,580	10,280	33,924
1-pound tall.....	126,597	683,624	30,393	164,122	29,377	158,636	20,189	109,021
½-pound oval.....	17,650	75,013
1-pound oval.....	406	2,639
1-pound squats.....	8,312	49,872
Total.....	705,890	3,509,227	125,620	582,788	119,927	571,140	33,827	168,728
Grand total.....	731,315	3,617,044	181,038	770,435	122,198	579,903	46,599	218,671

PACK OF CANNED SALMON IN BRITISH COLUMBIA, CANADA, IN 1909—Continued.

Species.	Northern miscellaneous waters.		Vancouver Island.		Total.	
	Cases.	Value.	Cases.	Value.	Cases.	Value.
Coho, or silver:						
½-pound flat.....					2,132	\$5,969
1-pound flat.....					5,911	28,373
1-pound tall.....	13,071	\$54,898	13,409	\$56,318	61,520	258,400
Total.....	13,071	54,898	13,409	56,318	69,563	292,742
Dogs, or chums:						
1-pound tall.....	1,568	3,763	2,280	5,472	16,573	39,775
Humpback, or pink:						
1-pound flat.....			2,000	5,500	2,267	6,234
1-pound tall.....	3,000	7,200	4,000	9,600	27,722	66,581
Total.....	3,000	7,200	4,000	15,100	29,989	72,815
King, or spring:						
½-pound flat.....					360	1,440
1-pound flat.....					1,214	7,314
½-pound tall.....					176	516
1-pound tall.....	2,218	11,977	500	2,700	17,613	94,110
1-pound oval.....					444	2,886
Total.....	2,218	11,977	500	2,700	19,807	106,266
Sockeye, or red:						
½-pound flat.....	18,806	75,224	19,800	79,200	483,760	1,935,040
1-pound flat.....			20,400	122,400	314,706	1,888,236
½-pound tall.....					12,880	42,504
1-pound tall.....	29,694	160,348	41,643	224,872	277,893	1,500,623
½-pound oval.....					17,650	75,013
1-pound oval.....					406	2,639
1-pound squats.....					8,312	49,872
Total.....	48,500	235,572	81,843	426,472	1,115,607	5,493,927
Grand total.....	68,357	313,410	102,032	506,062	1,251,539	6,005,525

^a All pound cases contain forty-eight 1-pound cans; the ½-pound cases contain forty-eight ½-pound cans. Reduced to a common basis of cases containing forty-eight 1-pound cans the pack is 993,060 cases.

VIII. STATISTICAL DATA FOR OTHER YEARS.

CANNING INDUSTRY OF PACIFIC COAST FROM 1864 TO 1910.

From the beginning of the canning of salmon on this coast it has been the most important branch of the industry, and the table below shows in condensed form the number of cases packed in each year on the Pacific coast of North America from the beginning of the industry in 1864 to 1910.

As British Columbia is a Province of the Dominion of Canada it does not come strictly within the scope of this report, but in order to show the pack of canned salmon on the North American shores of the Pacific Ocean, which would be incomplete without that of the Province, it has been included also.

PACK OF CANNED SALMON ON THE PACIFIC COAST, BY YEARS AND WATERS.

Year.	Puget Sound.	Grays Harbor.	Willapa Harbor.	Columbia River.	Coastal streams of Oregon.	Smith River, Cal.
	<i>Cases.</i>	<i>Cases.</i>	<i>Cases.</i>	<i>Cases.</i>	<i>Cases.</i>	<i>Cases.</i>
1866.....				4,000		
1867.....				18,000		
1868.....				28,000		
1869.....				100,000		
1870.....				150,000		
1871.....				200,000		
1872.....				250,000		
1873.....				250,000		
1874.....				350,000		
1875.....				375,000		
1876.....				450,000		
1877.....	5,500			380,000	7,804	
1878.....	238	5,420		400,000	16,634	4,277
1879.....	1,300			480,000	8,571	
1880.....	5,100			530,000	7,772	7,500
1881.....	8,500			550,000	12,320	
1882.....	7,900			541,300	19,186	
1883.....	1,500			629,400	16,156	
1884.....	5,500			620,000	12,376	
1885.....	12,000			553,800	9,310	
1886.....	17,000			448,500	49,147	
1887.....	22,000			356,000	73,996	
1888.....	21,975	37,000	22,500	372,477	92,863	2,347
1889.....	11,674			309,885	98,800	
1890.....	8,000			435,774	47,009	
1891.....	20,529	500	8,000	398,953	24,500	
1892.....	26,426	16,500	14,500	487,338	83,600	
1893.....	89,774	22,000	16,195	415,876	52,778	
1894.....	95,400	21,400	15,100	490,100	54,815	1,500
1895.....	179,968	11,449	22,600	634,696	77,878	2,500
1896.....	195,664	21,274	24,941	481,697	87,300	
1897.....	494,026	13,300	29,600	552,721	60,158	
1898.....	400,200	12,100	21,420	487,944	75,679	
1899.....	919,611	24,240	21,314	332,774	82,041	
1900.....	460,450	30,800	26,300	358,772	12,237	
1901.....	1,380,590	41,500	34,000	390,183	58,618	
1902.....	581,659	31,500	39,492	317,143	44,236	
1903.....	478,488		5,890	339,577	54,861	
1904.....	291,488	27,559	26,400	395,104	98,874	
1905.....	1,018,641	22,050	14,950	397,273	89,055	
1906.....	430,602	22,000	14,140	394,898	107,332	
1907.....	698,080	14,000	13,382	324,171	79,712	
1908.....	448,765	14,000	20,457	253,341	52,478	
1909.....	1,632,949	19,787	12,024	274,087	58,169	
1910.....	567,883	51,130	14,508	391,415	103,617	
Total.....	10,548,380	459,509	418,013	16,960,199	1,829,942	19,374

PACK OF CANNED SALMON ON THE PACIFIC COAST, BY YEARS AND WATERS—Con

Year.	Klamath River, Cal.	Eel River, Cal.	Sacramento River.	Alaska.	British Columbia.	Total.
	<i>Cases.</i>	<i>Cases.</i>	<i>Cases.</i>	<i>Cases.</i>	<i>Cases.</i>	<i>Cases.^a</i>
1864.....			2,000			2,000
1865.....			2,000			2,000
1866.....						4,000
1867.....						18,000
1868.....						28,000
1869.....						100,000
1870.....						150,000
1871.....						200,000
1872.....						250,000
1873.....						250,000
1874.....			2,500			352,500
1875.....			3,000			378,000
1876.....			10,000		7,247	467,247
1877.....		8,500	21,500		58,387	481,691
1878.....		10,500	34,017	8,159	89,946	629,191
1879.....			13,855	12,530	61,093	577,349
1880.....		6,250	62,000	6,539	61,849	687,010
1881.....			181,200	8,977	169,576	930,573
1882.....			200,000	21,745	240,461	1,030,592
1883.....			123,000	48,337	163,438	981,831
1884.....			81,450	64,886	123,706	907,918
1885.....			90,000	83,415	108,517	857,042
1886.....			39,300	142,065	152,964	848,976
1887.....			36,500	206,677	204,083	899,256
1888.....	4,400		68,075	412,115	184,040	1,217,792
1889.....			57,300	719,196	417,211	1,614,066
1890.....			25,065	682,591	411,257	1,609,696
1891.....			10,353	801,400	314,511	1,578,746
1892.....			2,281	474,717	248,721	1,354,083
1893.....	1,600		23,336	643,654	610,202	1,876,915
1894.....	1,700		28,463	686,440	492,232	1,887,150
1895.....	1,600		25,185	626,530	587,092	2,169,848
1896.....			13,387	966,707	617,782	2,408,812
1897.....			38,543	909,078	1,027,183	3,124,609
1898.....			29,731	965,097	492,551	2,484,722
1899.....	1,600		32,580	1,078,146	765,519	3,257,825
1900.....			39,304	1,548,139	606,540	3,091,542
1901.....			17,500	2,016,804	1,247,212	5,186,407
1902.....	2,500		14,043	2,536,824	627,161	4,194,558
1903.....			8,200	2,246,210	473,847	3,607,073
1904.....	3,400		14,407	1,953,756	465,894	3,276,882
1905.....			2,780	1,894,516	1,167,822	4,607,087
1906.....				2,219,044	629,460	3,817,776
1907.....				2,169,873	547,459	3,522,506
1908.....				2,606,973	566,303	3,962,317
1909.....	5,633			2,395,477	993,060	5,393,670
1910.....	8,016	6,000		2,413,054	760,830	4,316,453
Total.....	30,449	31,250	1,352,855	33,569,671	15,695,756	80,593,711

^a Reduced to a common basis of forty-eight 1-pound cans to the case.

CANNING INDUSTRY, BY SPECIES AND WATERS.

The tables below show separately, by waters and as far as possible by species, the salmon canned on the Pacific coast from the beginning of the industry until 1910. It is only within recent years that the published statistics have shown the pack of the different species separately. In the early years of canning, the chinook, or quinnat, salmon was used exclusively, the other species not being utilized until the chinook had begun to decrease in abundance, or a demand had arisen for a cheaper product. There is a very great difference in the selling value of the highest and lowest grades, and it is necessary to have complete statistical data now in order intelligently to comprehend the trend of the industry. While every effort has been made to make these tables complete, there are, unfortunately, some gaps which it was found impossible to fill.

PACK OF CANNED SALMON ON PUGET SOUND FROM 1877 TO 1910.

Year.	Number of canneries.	Chinook.		Blueback.		Silver.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.
1877.....	1					5,000	
1878.....	1					238	
1879.....	1					1,300	\$5,090
1880.....	1						
1881.....	1						
1882.....	1						
1883.....	1						
1884.....	1						
1885.....							
1886.....							
1887.....							
1888.....	4						
1889.....	2	240	\$1,200			7,480	37,400
1890.....	1	1,000	5,000			3,000	15,000
1891.....	2	382	2,101	5,538	\$24,921	5,869	19,368
1892.....	2	86	473	2,954	11,816	7,206	24,500
1893.....	3	1,200	6,480	47,852	103,371	11,812	59,060
1894.....	3			41,781	188,014	22,418	89,672
1895.....	7	1,542	7,325	65,143	273,108	50,865	154,218
1896.....	11	13,495	67,475	72,979	350,299	82,640	264,448
1897.....	12	9,500	39,045	312,048	1,248,192	91,900	282,133
1898.....	18	11,200	50,624	252,000	1,058,400	98,600	335,240
1899.....	19	24,364	103,180	499,646	2,368,354	111,387	418,176
1900.....	19	22,350	134,100	229,800	1,149,000	128,200	512,800
1901.....							
1902.....	21	30,049	150,245	372,301	2,047,655	85,817	429,085
1903.....	22	14,500	72,500	167,211	1,003,260	103,450	413,800
1904.....	13	14,441	69,352	109,264	653,871	118,127	447,851
1905.....	24	1,804	9,922	825,453	4,952,718	79,335	337,174
1906.....	16	8,139	48,834	178,748	1,251,236	94,497	472,485
1907.....	14	1,814	16,326	93,122	698,416	119,472	476,288
1908.....	11	95,210	666,470	170,951	1,196,657	128,922	644,922
1909.....	24	13,019	72,604	1,097,904	6,183,300	143,133	630,446
1910.....	15	10,064	60,324	248,014	1,673,095	162,755	895,155

PACK OF CANNED SALMON ON PUGET SOUND FROM 1877 TO 1910—Continued.

Year.	Number of canneries.	Dog.		Humpback.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.
1877.	1			500		5,500	
1878.	1					238	
1879.	1					1,300	\$5,690
1880.	1					5,100	
1881.	1					8,500	
1882.	1					7,900	
1883.	1					1,500	
1884.	1					5,500	
1885.						12,000	
1886.						17,000	
1887.						22,000	
1888.	4					21,975	120,356
1889.	2	1,145	\$3,435	2,809	\$7,584	11,674	49,619
1890.	1	4,000	12,000			8,000	32,000
1891.	2	3,093	10,825	5,647	15,246	20,529	72,461
1892.	2	16,180	56,630			26,426	93,419
1893.	3	11,380	31,295	17,530	47,331	89,774	247,537
1894.	3	22,152	60,918	9,049	24,432	95,400	363,036
1895.	7	38,785	94,741	23,633	62,556	179,968	591,948
1896.	11	26,550	73,013			195,664	755,235
1897.	12	23,310	64,103	57,268	171,804	494,026	1,805,277
1898.	18	38,400	105,600			400,200	1,549,864
1899.	19	31,481	86,427	252,733	734,241	919,611	3,710,358
1900.	19	89,100	245,025			469,540	1,940,925
1901.						1,380,590	
1902.	21	93,492	467,460			581,659	3,094,445
1903.	22	12,001	30,002	181,326	407,984	478,488	1,927,546
1904.	13	49,656	124,254			291,488	1,295,328
1905.	24	41,057	102,643	70,992	212,976	1,018,641	5,615,433
1906.	16	149,218	708,781			430,602	2,481,336
1907.	14	50,249	150,847	433,423	1,300,269	698,080	2,642,146
1908.	11	47,607	142,821	6,075	18,225	448,765	2,669,095
1909.	24	53,688	128,916	370,993	902,342	1,632,949	7,917,608
1910.	15	146,942	514,297	108	388	567,883	3,143,256

PACK OF CANNED SALMON ON GRAYS HARBOR FROM 1878 TO 1910.

Year.	Number of canneries.	Chinook.		Silver.		Dog or chum.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
1878.	1							5,420	\$29,268
1879.	1								
1880.									
1881.									
1882.									
1883.									
1884.									
1885.									
1886.									
1887.									
1888.	4							37,000	\$212,750
1889.									
1890.									
1891.	1			500	\$1,500			500	1,500
1892.	1	4,500	\$15,390	9,000	30,780	3,000	\$9,415	16,500	55,585
1893.	1	4,500	22,500	12,000	48,000	5,500	14,850	22,000	85,350
1894.	1	12,300	61,500	4,100	16,400	5,000	13,500	21,400	91,400
1895.	1	56	202	8,876	28,403	2,517	6,922	11,449	35,527
1896.	2	7,816	36,806	9,278	29,689	4,180	11,495	21,274	57,990
1897.	1	3,100	11,741	8,300	23,481	1,900	5,000	13,300	40,222
1898.	2	5,100	23,052	4,800	16,320	2,200	6,050	12,100	45,422
1899.	1	5,000	21,250	15,740	59,025	3,500	8,750	24,240	89,025
1900.	2	6,700	33,500	12,900	51,600	11,200	30,800	30,800	115,900
1901.								41,500	
1902.	1	4,000	20,000	10,000	45,000	17,500	70,000	31,500	135,000
1903.									
1904.	2	4,339	20,163	14,904	51,854	8,316	21,022	27,559	93,039
1905.	2	2,050	9,225	13,000	52,000	7,000	18,200	22,050	79,425
1906.	2	2,500	10,000	11,500	43,900	8,000	21,500	22,000	75,400
1907.	1	1,000	7,000	9,500	47,500	3,500	11,500	14,000	66,000
1908.	1	1,000	7,000	9,500	47,500	3,500	11,500	14,000	66,000
1909.	1	5,721	20,819	9,019	38,146	5,047	11,608	a 19,787	70,573
1910.	3	15,495	90,718	21,768	108,840	13,867	48,534	b 51,130	248,092

a Also 1,649 cases, valued at \$9,051, with sockeyes brought from Puget Sound.

b Also 4,350 cases of "Quinault," or sockeye salmon.

PACK OF CANNED SALMON ON WILLAPA HARBOR FROM 1887 TO 1910.

Year.	Number of canneries.	Chinook or Black.		Silver.		Dog.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
1887.....	4								
1888.....	3							22,500	\$129,375
1889.....									
1890.....									
1891.....	1			8,000	\$24,000			8,000	24,000
1892.....	1	3,000	\$10,260	9,000	30,780	2,500	\$7,745	14,500	48,785
1893.....	1	1,700	9,180	7,895	31,580	6,600	18,150	16,195	58,910
1894.....	1	2,700	14,580	5,600	22,400	6,800	18,700	15,100	55,680
1895.....	2	4,636	23,180	13,047	41,150	4,917	13,222	22,600	77,552
1896.....	2	4,551	22,755	11,940	38,208	8,450	21,238	24,941	82,201
1897.....	1	8,100	33,291	14,600	44,822	6,900	18,975	29,600	97,088
1898.....	2	5,865	26,510	9,809	33,351	5,746	15,802	21,420	75,663
1899.....	3	5,650	25,425	10,675	40,031	4,989	13,720	21,314	79,176
1900.....	3	6,700	33,500	12,400	49,600	7,200	19,800	26,300	102,900
1901.....								34,000	
1902.....	2	5,836	29,180	9,128	41,076	24,528	97,112	39,492	167,368
1903.....	1	2,300	13,800	2,390	10,755	1,200	3,300	5,890	27,855
1904.....	2	3,000	12,000	7,400	28,440	16,000	38,700	26,400	79,140
1905.....	2	4,650	20,925	4,300	17,200	6,000	15,000	14,950	53,125
1906.....	2	4,000	16,000	5,340	21,360	5,100	13,260	14,440	50,620
1907.....	2	3,530	15,354	9,228	36,682	624	2,496	13,382	54,532
1908.....	2	4,017	20,585	5,923	23,692	10,517	36,809	20,457	81,086
1909.....	1	1,455	5,809	4,822	17,359	5,747	13,163	12,024	36,391
1910.....	1	2,923	15,077	5,096	25,480	3,489	22,711	14,508	63,268

PACK OF CANNED SALMON ON THE COLUMBIA RIVER FROM THE INCEPTION OF THE INDUSTRY TO 1910.

Year.	Num-ber of can-ner-ies.	Chinook.		Blueback.		Silversides.		Dog or chum.		Steelhead trout.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
1866													864,000
1867													288,000
1868													392,000
1869													1,350,000
1870													1,800,000
1871													2,100,000
1872													2,325,000
1873													2,550,000
1874													2,625,000
1875													2,550,000
1876													2,475,000
1877													2,052,000
1878													2,300,000
1879	30												2,640,000
1880	29												2,650,000
1881													2,475,000
1882													2,600,000
1883													2,475,000
1884													2,640,000
1885													2,475,000
1886													2,640,000
1887													2,475,000
1888	28												2,640,000
1889	21	296,697	\$1,600,182	17,797	\$101,051					25,391	\$108,587	309,885	1,809,850
1890	1890	335,004	1,946,087	57,345	290,069					42,825	171,300	435,774	2,407,456
1891	1891	353,907	2,038,566	15,482	281,242					29,564	118,156	398,953	2,440,964
1892	24	344,207	1,996,388	66,547	372,909					72,348	288,892	487,338	2,679,069
1893	24	288,773	1,559,374	30,459	152,295	29,107	\$6,933	2,311	\$6,933	62,522	260,904	415,876	2,985,354
1894	24	351,106	1,895,976	49,814	224,430	42,758	171,032			52,422	209,688	490,100	2,801,126
1895	24	444,909	2,428,658	18,015	86,523	99,601	329,683	22,493	62,591	49,678	203,542	634,696	3,110,897
1896	24	370,943	1,840,511	16,983	81,518	44,108	141,145			49,663	198,652	481,697	2,361,826
1897	22	432,753	1,804,221	12,972	51,888	60,850	197,762			46,146	165,440	552,721	2,219,311
1898	23	329,566	1,490,394	66,670	300,015	65,431	222,465			26,277	60,352	487,944	2,073,226
1899	17	255,824	1,458,175	26,968	134,723	29,608	112,055	11,379	33,836	11,994	39,186	332,774	1,777,975
1900	16	262,392	1,821,268	13,162	92,184	44,925	202,163	17,696	63,706	20,597	102,985	358,772	2,282,296
1901	14	270,580	1,428,743	17,037	86,465	10,522	44,732	10,401	41,004	8,593	42,965	317,143	1,644,509
1902	16	301,762	1,610,614	8,383	42,807	12,181	49,869	10,000	37,500	7,251	36,255	339,577	1,776,105
1903	20	320,378	1,944,690	12,911	78,048	31,254	118,357	20,693	52,691	9,868	48,892	395,104	2,242,678

1905.....	327,106	1,962,636	7,768	46,608	26,826	114,011	25,751	65,206	9,822	\$49,110	397,273	2,237,571
1906.....	311,334	1,868,007	7,816	54,712	41,446	124,338	27,802	69,505	6,500	32,500	394,898	2,149,062
1907.....	258,433	5,504	31,757	22,556	5,921	324,171	1,763,490
1908.....	210,006	8,581	31,432	16,884	10,736	253,341	1,380,708
1909.....	102,131	1,203,546	^a 27,908	214,561	42,178	185,070	24,542	57,115	17,382	99,796	674,087	1,700,088
1910.....	244,285	1,882,137	6,234	34,287	68,922	363,688	66,538	232,883	5,436	31,203	391,415	2,544,198
Total.....	16,960,199	94,792,931

^a Of these, 2,846 cases, valued at \$23,203, were packed with sockeyes brought from Puget Sound.

^b 55 cases of humpbacks, valued at \$132, were also packed with humpbacks brought from Puget Sound.

PACK OF CANNED SALMON ON THE NEHALEM RIVER, OREG., FROM 1887 TO 1910.

Year.	Number of canneries.	Chinook.		Silver.		Dog.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
1887	1							5,000	\$30,000
1888									
1889								6,000	32,000
1890								9,000	45,500
1891	1							3,500	14,000
1892	1			10,000	\$40,000			10,000	40,000
1893	1	1,092	\$6,768	5,031	20,124			6,723	26,892
1894	1	1,627	6,508	4,566	19,464			6,493	25,972
1895	1	1,752	7,008	5,152	16,486			6,904	23,494
1896	1	2,828	8,484	5,218	15,654			8,046	24,138
1897	2	3,384	10,152	8,366	25,098			11,750	35,250
1898	1	3,808	9,891	5,700	19,380			9,508	29,271
1899	1	1,384	5,536	7,405	26,658	1,288	\$3,864	10,077	36,058
1900	1								
1901	1	268	1,139	3,273	13,092	2,669	7,206	6,210	21,437
1902	1	271	1,431	3,169	13,468	2,570	10,280	6,010	25,179
1903	1	686	3,670	4,615	19,614			5,301	
1904	1	500	2,500	5,000	20,000	6,000	12,000	11,500	34,500
1905		2,700	16,200	2,900	12,325	6,000	15,000	11,600	43,525
1906	1	3,987	23,922	4,976	14,928	2,057	5,143	11,020	42,993
1907	1	4,000		6,600		2,000		12,600	
1908	1	5,000		6,100		2,016			
1909	1	1,985	10,542	4,554	20,253	909	2,091	7,448	32,886
1910	1	3,500		5,400		1,500		10,400	

PACK OF CANNED SALMON ON TILLAMOOK BAY, OREG., FROM 1886 TO 1910.

Year.	Number of canneries.	Chinook.		Silver.		Dog.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
1886	2							37,000	
1887	2							21,000	\$115,500
1888	2							14,633	84,140
1889								9,500	52,250
1890								14,009	79,049
1891	1								
1892	1			18,000	\$72,000			18,000	72,000
1893	1	497	\$1,988	4,000	16,000	6,919	\$17,297	11,416	35,285
1894	1	700	2,800	7,763	31,052	700	1,750	9,163	35,602
1895	1			6,514	20,845	7,001	19,253	13,515	40,098
1896	1	2,200	6,600	4,860	14,580			7,060	21,150
1897	1	2,000	6,000	9,000	27,000			11,000	33,000
1898	1	5,000	13,000	10,342	35,162			15,342	48,162
1899	1	2,180	8,720	3,889	14,036	5,121	15,363	11,190	38,119
1900	1								
1901	1	848	4,240	2,133	9,598	3,901	10,728	6,882	24,566
1902	1	215	1,135	2,287	9,720	4,093	16,372	6,595	27,227
1903	1			2,727	11,590	2,620	10,480	5,347	22,070
1904	1			4,400	17,600	6,500	13,000	10,900	30,600
1905	1	1,100	6,600	1,700	7,650	8,800	22,000	11,600	36,250
1906	1	1,870	11,220	2,364	7,092	1,270	3,175	5,504	21,487
1907	1	2,000		3,410		2,314		7,724	
1908	1	2,300		6,000		4,000			
1909	1	2,615	15,663	5,029	21,809	3,712	8,538	11,356	46,010
1910	1	2,900		4,500		2,000		9,400	

PACK OF CANNED SALMON ON NESTUCCA RIVER, OREG., FROM 1887 TO 1910.

Year.	Number of canneries.	Chinook.		Silver.		Dog.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
1887.....	1							4,300	\$23,650
1888.....	1							5,000	28,750
1889.....	1							6,700	36,850
1890.....									
1891.....	1								
1892.....									
1893.....									
1894.....									
1895.....									
1896.....									
1897.....									
1898.....									
1899.....	1	1,109	\$4,436	3,034	\$10,922	513	\$1,539	4,656	16,897
1900.....	1								
1901.....	1	279	1,116	3,553	13,323	396	1,089	4,228	15,528
1902.....	1								
1903.....									
1904.....									
1905.....	1	3,000	18,000	1,000	4,250	400	1,000	4,400	23,250
1906.....	1	2,622	15,732	2,468	7,404	165	413	5,255	23,549
1907.....	1	2,100		3,540		150		5,790	
1908.....	1	2,000		3,000		100			
1909.....									
1910.....	1	2,000		3,300		140		5,440	

PACK OF CANNED SALMON ON SILETZ RIVER, OREG., FROM 1896 TO 1910.

Year.	Number of canneries.	Chinook.		Silver.		Dog.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
1896.....	1	2,500	\$7,500	1,900	\$5,700			4,400	\$13,200
1897.....	1	3,510	10,530	5,015	15,045			8,525	25,575
1898.....	1	3,200	8,360	4,330	14,722			7,530	23,082
1899.....	1	2,200	9,900	2,319	8,696	200	\$550	4,719	19,146
1900.....	1								
1901.....	1	876	4,380	3,740	16,830	360	1,260	4,976	22,470
1902.....	1	600	3,168	1,917	8,147	500	2,000	3,017	13,315
1903.....									
1904.....	1	1,000	5,000	3,300	13,200	1,000	2,000	5,300	20,200
1905.....	1	1,500	9,000	1,700	7,225	900	2,250	4,100	18,475
1906.....	1	2,635	15,810	3,192	9,576	167	418	5,994	25,804
1907.....	1	2,333		4,300		200		6,833	
1908.....	1	2,100		4,700		300		7,100	
1909.....									
1910.....	1	2,200		4,600		250		7,050	

PACK OF CANNED SALMON ON YAQUINA BAY AND RIVER, OREG., FROM 1887 TO 1910.

Year.	Number of canneries.	Chinook.		Silver.		Dog.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
1887.....	2								
1888.....	3							5,088	\$29,256
1889.....								5,000	27,500
1890.....									
1891.....	1								
1892.....									
1893.....									
1894.....									
1895.....									
1896.....	1	1,714	\$5,142	615	\$1,845			2,329	6,987
1897.....									
1898.....	1	170	442	1,530	5,202			1,700	5,644
1899.....	2	316	1,422	3,234	12,127	1,300	\$3,575	4,850	17,124
1900.....	1								
1901.....	1	96	480	2,848	12,816	549	1,647	3,493	14,943
1902.....									
1903.....	1			1,238	5,202	315	787	1,553	6,049
1904.....	1	50	200	2,600	8,840	450	1,080	3,100	10,120
1905.....	1	200	1,200	2,050	8,613	62	155	2,312	9,968
1906.....	1	500	3,000	3,100	9,300	60	150	3,660	12,450
1907.....	1	834		1,000		49		1,883	
1908.....	1			4,000				4,000	
1909.....	1			1,139	4,556	33	76	1,172	4,632
1910.....	1			2,669	13,345			2,669	13,345

PACK OF CANNED SALMON ON ALSEA RIVER AND BAY, OREG., FROM 1886 TO 1910.

Year.	Number of canneries.	Chinook.		Silver.		Dog.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
1886.....	1								
1887.....	2							11,180	\$64,285
1888.....	3							9,620	55,315
1889.....								10,000	55,000
1890.....									
1891.....	1								
1892.....	1			3,600	\$14,400			3,600	14,400
1893.....	1	1,260	\$6,300	3,240	12,960			4,500	19,260
1894.....	1	440	2,200	4,160	16,640			4,600	18,840
1895.....	1	1,700	6,375	3,280	11,808			4,980	18,183
1896.....	1	3,500	10,500	3,400	10,200			6,900	20,700
1897.....	1	1,800	5,400	3,200	9,600			5,000	15,000
1898.....	1	4,296	11,170	2,170	7,378			6,466	18,548
1899.....	1	2,150	9,138	5,010	19,038			7,160	28,176
1900.....	1								
1901.....	1	695	3,475	4,629	18,790	891	\$3,118	6,215	25,383
1902.....	1	701	3,702	4,530	19,253	670	2,680	5,901	25,635
1903.....	1	1,031	5,516	4,242	18,029	44	88	5,317	23,633
1904.....	1	1,000	5,000	6,500	26,000	300	600	7,800	31,600
1905.....	1	2,500	15,000	1,800	7,650	700	1,750	5,000	24,400
1906.....	1	3,702	22,212	3,843	11,529			7,545	33,741
1907.....	1	800		5,100		350		6,250	
1908.....	1	1,200		6,000		400		7,600	
1909.....	1	1,119	6,714	5,486	24,027	80	184	6,685	30,925
1910.....	1	2,500		5,900		100		8,500	

PACK OF CANNED SALMON ON THE SIUSLAW RIVER, OREG., FROM 1878 TO 1910.

Year.	Number of canneries.	Chinook.		Silver.		Dog.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
1878.....	2							10,300	\$55,620
1879.....	2								
1880.....									
1881.....									
1882.....									
1883.....									
1884.....									
1885.....									
1886.....									
1887.....									
1888.....	1							11,960	68,770
1889.....	1							12,000	66,000
1890.....									
1891.....	2								
1892.....	2			18,000	\$72,000			18,000	72,000
1893.....	2	1,471	\$7,355	11,830	47,320			13,301	54,675
1894.....	2	1,871	9,355	14,987	59,948			16,858	69,303
1895.....	2	1,637	6,139	10,465	35,274			12,102	41,413
1896.....	1	2,700	8,100	9,000	27,000			11,700	35,100
1897.....	1	1,100	3,300	3,900	11,700			5,000	15,000
1898.....	1	850	2,210	10,000	34,000			10,850	36,210
1899.....	1	1,162	4,648	7,323	26,363	115	\$345	8,600	31,356
1900.....	2								
1901.....	1	1,735	8,675	7,488	29,952			9,223	38,627
1902.....	1	1,288	6,800	4,320	18,260			5,608	25,060
1903.....	1	1,519	8,127	6,842	29,079			8,361	37,206
1904.....	1	500	2,500	6,500	26,000			7,000	28,500
1905.....	1								
1906.....	2	4,500	27,000	15,000	45,000	1,500	3,750	21,000	75,750
1907.....	1			15,773				15,773	
1908.....	1			8,600				8,600	
1909.....	2	632	3,792	7,436	32,956			8,068	36,748
1910.....	2	856		12,800		8,502		22,158	

PACK OF CANNED SALMON ON THE UMPQUA RIVER, OREG., FROM 1878 TO 1910.

Year.	Number of canneries.	Chinook.		Silver.		Dog.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.	Cases.	Value.
1878.....	2							8,100	\$43,740
1879.....	2								
1880.....									
1881.....									
1882.....									
1883.....									
1884.....	2								
1885.....	1								
1886.....	1							4,000	22,000
1887.....	1							9,000	51,750
1888.....	1							12,000	66,000
1889.....	1								
1890.....									
1891.....	1			10,000	\$40,000			10,000	40,000
1892.....	1			3,204	12,816			4,013	16,861
1893.....	1	809	\$4,045	6,875	27,500			7,110	28,675
1894.....	1	235	1,175	7,697	28,863			8,689	32,583
1895.....	1	992	3,720	8,000	24,000			9,300	27,900
1896.....	1	1,300	3,900						
1897.....									
1898.....									
1899.....	2	925	3,860	7,576	27,006	115	\$345	8,616	31,211
1900.....	2								
1901.....									
1902.....									
1903.....	1	23	123	6,733	28,615			6,756	28,738
1904.....	1	500	2,500	9,500	38,000	500	1,000	10,500	41,500
1905.....	1	6,100	36,600	10,500	44,625			16,600	81,225
1906.....	1	1,143	6,858	5,613	16,839			6,756	23,697
1907.....									
1908.....									
1909.....	1	500	3,000	7,753	31,012			8,253	34,012
1910.....	1	2,000		11,000				13,000	

PACK OF CANNED SALMON ON COOS BAY AND RIVER, OREG., FROM 1887 TO 1910.

Year.	Number of canneries.	Chinook.		Silver.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.
1887.....	2					11,300	\$62,150
1888.....	1					5,500	31,625
1889.....	1					7,000	38,500
1890.....							
1891.....	2						
1892.....							
1893.....	1			3,125	\$12,500	3,125	12,500
1894.....	1	163	\$815	8,428	33,712	8,591	34,527
1895.....	1	5,110	19,163	2,332	8,934	7,442	28,097
1896.....	1	13,000	39,000	2,000	6,000	15,000	45,000
1897.....	1	6,200	18,600	2,200	6,600	8,400	25,200
1898.....	2	3,142	8,169	7,180	24,412	10,322	32,581
1899.....	2	1,273	5,092	5,174	18,626	6,447	23,718
1900.....	2						
1901.....	1	1,215	6,075	4,082	16,328	5,297	22,403
1902.....	1	412	2,175	2,640	11,220	3,052	13,395
1903.....							
1904.....	1	2,033	7,725	7,200	24,480	9,233	32,205
1905.....							
1906.....	1	2,043	12,258	1,755	5,265	3,798	17,523
1907.....							
1908.....							
1909.....	1	275	1,475	3,959	17,927	4,234	19,402
1910.....	1	500		5,500		6,000	

PACK OF CANNED SALMON ON THE COQUILLE RIVER, OREG., FROM 1883 TO 1910.

Year.	Number of canneries.	Chinook.		Silver.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.
1883.....	1						
1884.....							
1885.....							
1886.....	2						
1887.....	3						
1888.....	2					11,000	\$63,250
1889.....						8,600	47,300
1890.....							
1891.....	1						
1892.....	1			5,000	\$20,000	5,000	20,000
1893.....	1			6,500	26,000	6,500	26,000
1894.....	α 1			2,000	8,000	2,000	8,000
1895.....	2	760	\$2,887	8,724	32,615	9,484	35,502
1896.....	2	1,225	3,675	7,800	23,400	9,025	27,075
1897.....							
1898.....	2	541	1,407	7,485	25,499	8,026	26,906
1899.....	2	950	3,800	7,550	28,500	8,500	32,300
1900.....	1	2,636	13,180	9,601	38,404	12,237	51,584
1901.....	1	133	665	5,096	20,384	5,229	21,049
1902.....	1	286	1,510	5,877	24,927	6,163	26,437
1903.....	1	331	1,771	8,685	36,911	9,016	38,682
1904.....	2	600	2,400	13,686	54,744	14,286	57,144
1905.....	2	2,100	12,600	11,343	48,208	13,443	60,808
1906.....	2	821	4,926	17,979	53,937	18,800	58,863
1907.....	2	306		13,220		13,526	
1908.....	2			19,174		19,174	
1909.....	2	250	1,255	9,818	42,687	10,068	43,942
1910.....	2	420		16,637		17,057	

α Burned.

PACK OF CANNED SALMON ON ROGUE RIVER, OREG., FROM 1877 TO 1910.

Year.	Number of canneries.	Chinook.		Silver.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.
1877	1					7,804	
1878	1					8,534	
1879	1					8,571	
1880	1					7,772	
1881	1					12,320	
1882	1					19,186	
1883	1					16,156	
1884	1					12,376	
1885	1					9,310	
1886	1					12,147	
1887	1					17,216	
1888	1					21,062	\$121,107
1889	1					22,000	132,000
1890	1					24,000	120,000
1891	1					21,000	105,000
1892	1	10,000	\$59,000	9,000	\$36,000	19,000	95,000
1893	^a 1	3,200	16,000			3,200	16,000
1894	(b)						
1895	1	10,377	41,508	4,385	15,347	14,762	56,855
1896	1	15,000	75,000	3,000	9,000	18,000	84,000
1897	1	15,355	61,420	3,653	10,959	19,008	72,379
1898	1	12,964	51,550	501	1,303	13,465	52,853
1899	1	5,481	30,145	1,745	6,980	7,226	37,125
1900	1						
1901	1	2,681	13,405	4,184	17,736	6,865	31,141
1902	1	3,799	20,058	4,091	17,387	7,890	37,445
1903	1	8,418	45,036	4,792	20,366	13,210	65,402
1904	1	16,000	64,000	3,255	11,392	19,255	75,392
1905	1	18,500	111,000	1,500	6,375	20,000	117,375
1906	1	12,000	72,000	6,000	18,000	18,000	90,000
1907	1	7,537		1,796		9,333	
1908	1	4,354		2,650		6,004	
1909	1	186	1,300	699	2,977	885	4,277
1910	1	232		2,711		1,943	

^a Burned down during season.

^b Not operated.

PACK OF CANNED SALMON ON SMITH RIVER, CAL., IN SPECIFIED YEARS.

Years.	Number of canneries.	Chinook salmon.		Silver salmon.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.
1878	1	4,277	\$23,096			4,277	\$23,096
1880	1	7,500				7,500	
1888	1	2,347	14,082			2,347	14,082
1893	1	1,500		500		2,000	
1894	1	1,500		500		2,000	
1895	1	2,250	9,990			2,250	9,990

PACK OF CANNED SALMON ON KLAMATH RIVER, CAL., IN SPECIFIED YEARS.

Year.	Number of canneries.	Chinook.		Silver.		Total.	
		Cases.	Value.	Cases.	Value.	Cases.	Value.
1888	1	4,400	\$26,400			4,400	\$26,400
1893	1	1,600				1,600	
1894	1	1,700				1,700	
1895	1	1,200	5,321	400	\$1,500	1,600	6,821
1899	1	1,600	8,800			1,600	8,800
1902	1	2,500				2,500	
1904	1	3,400	18,360			3,400	18,360
1909	1	5,633	28,315			5,633	28,315
1910	1	8,016				8,016	

PACK OF CANNED SALMON ON EEL RIVER, CAL., IN SPECIFIED YEARS.

Year.	Number of canneries.	Chinooks.	
		Cases.	Value.
1877.....	1	8,500	\$51,000
1878.....	1	10,500	56,700
1880.....	1	6,250
1910.....	1	6,000

PACK OF CANNED SALMON ON THE SACRAMENTO RIVER, FROM 1864 TO 1905.

Year.	Number of canneries.	Cases packed. ^a	Value.	Year.	Number of canneries.	Cases packed. ^a	Value.
1864.....	1	2,000	1886.....	39,300
1865.....	1	2,000	1887.....	36,500
1866.....	1888.....	6	68,075	\$423,750
1867.....	1889.....	3	57,300
1868.....	1890.....	25,065
1869.....	1891.....	10,353
1870.....	1892.....	2,281
1871.....	1893.....	23,336
1872.....	1894.....	28,463
1873.....	1895.....	3	25,185	111,821
1874.....	2,500	1896.....	13,387
1875.....	3,000	1897.....	38,543
1876.....	2	10,000	1898.....	29,731
1877.....	21,500	1899.....	32,580	150,688
1878.....	6	34,017	\$183,692	1900.....	39,304
1879.....	4	13,855	59,577	1901.....	17,500
1880.....	9	62,000	1902.....	14,043
1881.....	181,200	1903.....	8,200
1882.....	19	200,000	1904.....	2	14,407	66,936
1883.....	21	123,000	1905.....	1	2,780
1884.....	81,450	Total.....	1,352,855
1885.....	6	90,000				

^a All were quinnat or chinook salmon.

PACK OF CANNED SALMON IN ALASKA, BY DISTRICTS, FROM THE INCEPTION OF THE INDUSTRY.

Year.	Southeast Alaska.		Central Alaska.		Western Alaska.		Total.	
	Canneries.	Pack.	Canneries.	Pack.	Canneries.	Pack.	Canneries.	Pack.
1878.....	2	<i>Cases.</i> 8,159	<i>Cases.</i>	<i>Cases.</i>	2	8,159
1879.....	2	12,530	2	12,530
1880.....	1	6,539	1	6,539
1881.....	1	8,977	1	8,977
1882.....	1	11,501	2	10,244	3	21,745
1883.....	4	20,040	2	28,297	6	48,337
1884.....	4	22,189	2	42,297	1	^a 400	7	64,886
1885.....	3	16,728	2	52,687	1	14,000	6	83,415
1886.....	4	18,660	2	74,583	3	48,822	9	142,065
1887.....	5	31,462	2	102,515	3	72,700	10	206,677
1888.....	6	81,128	6	241,101	4	89,886	16	412,115
1889.....	12	141,760	21	461,451	4	115,985	37	719,196
1890.....	12	142,901	19	421,300	4	118,390	35	682,591
1891.....	11	156,615	14	511,367	5	133,418	30	801,400
1892.....	7	115,722	6	295,496	2	63,499	15	474,717
1893.....	8	136,053	11	399,815	3	107,786	22	643,654
1894.....	7	142,544	10	435,052	4	108,844	21	686,440
1895.....	7	148,476	10	327,919	6	150,135	23	626,530
1896.....	9	262,381	12	485,990	8	218,336	29	966,707
1897.....	9	271,867	13	382,899	7	254,312	29	909,078
1898.....	9	251,385	14	395,009	7	318,703	30	965,097
1899.....	9	310,219	14	356,095	9	411,832	32	1,078,146
1900.....	16	456,639	14	492,223	12	599,277	42	1,548,139

^a Experimental pack.

PACK OF CANNED SALMON IN ALASKA, BY DISTRICTS, FROM THE INCEPTION OF THE INDUSTRY—Continued.

Year.	Southeast Alaska.		Central Alaska.		Western Alaska.		Total.	
	Can-neries.	Pack.	Can-neries.	Pack.	Can-neries.	Pack.	Can-neries.	Pack.
1901.....	21	<i>Cases.</i> 735, 449	13	562, 142	21	719, 213	55	2, 016, 804
1902.....	26	906, 676	12	583, 690	26	1, 046, 458	64	2, 536, 824
1903.....	21	642, 305	12	417, 175	27	1, 186, 730	60	2, 246, 210
1904.....	12	569, 093	11	499, 485	32	885, 268	55	1, 953, 756
1905.....	13	433, 607	9	371, 755	25	1, 089, 154	47	1, 894, 516
1906.....	20	767, 285	8	473, 024	19	978, 735	47	2, 219, 044
1907.....	22	887, 503	8	522, 836	18	759, 534	48	2, 169, 873
1908.....	23	1, 011, 648	8	425, 721	19	1, 169, 604	50	2, 606, 973
1909.....	19	852, 870	8	391, 054	18	1, 151, 553	45	2, 395, 477
1910.....	23	1, 066, 399	10	432, 517	19	914, 138	52	2, 413, 054
Total.....	10, 647, 220	10, 195, 739	12, 726, 712	33, 569, 671

PACK OF CANNED SALMON IN ALASKA FROM 1898 TO 1910, BY SPECIES.

Year.	Coho, or silver.		Dog, or chum.		Humpback, or pink.	
	Cases.	Value.	Cases.	Value.	Cases.	Value.
1898.....	54, 711	5, 184	109, 399
1899.....	39, 402	1, 931	149, 159
1900.....	50, 984	30, 012	232, 022
1901.....	65, 509	47, 464	541, 427
1902.....	82, 723	159, 849	549, 602
1903.....	120, 506	35, 052	355, 799
1904.....	85, 741	21, 178	299, 333
1905.....	67, 394	\$215, 875	41, 972	\$113, 056	168, 597	\$498, 194
1906.....	109, 141	382, 109	254, 812	730, 235	348, 297	1, 046, 951
1907.....	85, 190	337, 384	184, 173	547, 757	561, 973	1, 799, 280
1908.....	68, 827	274, 089	218, 513	554, 197	644, 133	1, 733, 379
1909.....	56, 556	231, 029	120, 712	274, 110	464, 873	1, 114, 839
1910.....	114, 026	559, 666	254, 218	773, 409	554, 322	1, 764, 055

Year.	King, or spring.		Red, or sockeye.		Total.	
	Cases.	Value.	Cases.	Value.	Cases.	Value.
1898.....	12, 862	782, 941	965, 097
1899.....	23, 400	864, 254	1, 078, 146
1900.....	37, 715	1, 197, 406	1, 548, 139
1901.....	43, 069	1, 319, 335	2, 016, 804
1902.....	59, 104	1, 685, 546	2, 536, 824
1903.....	47, 609	1, 687, 244	2, 246, 210
1904.....	41, 956	1, 505, 548	1, 953, 756
1905.....	42, 125	\$141, 999	1, 574, 428	\$5, 335, 547	1, 894, 516	\$6, 304, 671
1906.....	30, 834	116, 222	1, 475, 961	5, 620, 875	2, 219, 044	7, 896, 392
1907.....	43, 424	181, 718	1, 295, 113	5, 915, 227	2, 169, 873	8, 781, 366
1908.....	23, 730	99, 867	1, 651, 770	7, 524, 251	2, 606, 973	10, 185, 783
1909.....	48, 034	207, 624	1, 705, 302	7, 610, 550	2, 395, 477	9, 438, 152
1910.....	40, 221	214, 802	1, 450, 267	7, 774, 390	2, 413, 054	11, 086, 322

PACK OF CANNED SALMON IN BRITISH COLUMBIA SINCE THE INCEPTION OF THE INDUSTRY, BY WATERS.

Year.	Number of canneries.	Fraser River.	Skeena River.	Rivers inlet.	Nass River.	Vancouver Island.	Northern miscellaneous waters.	Total.
		<i>Cases.</i>	<i>Cases.</i>	<i>Cases.</i>	<i>Cases.</i>	<i>Cases.</i>	<i>Cases.</i>	<i>Cases.</i>
1876.....	2	7,247						7,247
1877.....	5	55,387	3,000					58,387
1878.....	8	81,446	8,500					89,946
1879.....	9	50,490	10,603					61,093
1880.....	9	42,155	19,694					61,849
1881.....	11	142,516	21,560			5,500		169,576
1882.....	16	199,204	24,522	5,635	6,500	4,600		240,461
1883.....	20	105,701	31,157	10,780	9,400	6,400		163,438
1884.....	14	34,037	53,786	20,383	8,500	7,000		123,706
1885.....	9	89,617	12,900			6,000		108,517
1886.....	16	99,177	37,587	15,000		1,200		152,964
1887.....	20	130,088	58,592	11,203		4,200		204,083
1888.....	21	76,616	70,106	20,000	12,318	5,000		184,040
1889.....	28	310,122	58,405	21,722	19,800	7,162		417,211
1890.....	33	244,352	91,645	33,500	24,700	11,060	6,000	411,257
1891.....	38	177,989	77,057	36,500	11,058	4,300	8,057	314,511
1892.....	36	98,491	90,750	14,955	26,100	4,300	14,125	248,721
1893.....	44	474,237	59,021	35,416	15,680	8,098	17,750	610,202
1894.....	42	363,566	61,005	40,161	20,000		7,500	492,232
1895.....	49	432,920	69,356	58,575	20,541	3,300	3,000	587,692
1896.....	56	375,344	97,863	107,473	14,649	7,903	14,550	617,782
1897.....	65	879,776	61,310	40,090	20,000	13,807	12,200	1,027,183
1898.....	67	264,225	80,102	105,362	20,000	12,539	10,323	492,551
1899.....	68	527,396	112,562	76,428	19,442	12,150	17,541	765,519
1900.....	69	331,371	135,424	74,196	20,200	17,102	28,247	606,540
1901.....	78	998,913	125,845	66,794	15,004	11,005	29,651	1,247,212
1902.....	69	327,197	155,936	70,298	23,212	16,432	34,086	627,161
1903.....	61	237,162	98,688	69,389	18,094	12,360	38,154	473,847
1904.....	51	128,903	154,869	94,292	29,587	14,888	43,355	465,894
1905.....	64	846,998	114,085	83,122	32,725	50,975	39,917	1,167,822
1906.....	59	226,744	162,420	122,878	32,534	40,511	44,343	629,460
1907.....	42	163,116	159,255	94,064	31,832	76,616	22,576	547,459
1908.....	50	89,184	209,177	75,090	46,908	83,918	62,026	566,303
1909.....		567,230	142,740	91,014	40,990	58,954	92,132	993,060
1910.....		223,148	222,035	129,398	39,720	53,964	92,565	760,830
Total.....		9,402,095	2,891,557	1,623,718	579,494	560,794	638,098	15,695,756

PICKLING INDUSTRY.

The salmon-pickling industry was so overshadowed by its giant brother, the canning industry, that statistical data, except for Alaska, were found in extremely fragmentary shape, and only that portion is shown relating to Alaska from the time of annexation to 1909.

PACK OF SALTED SALMON IN ALASKA, 1868 TO 1909.

Year.	Salmon.		Salmon bellies.		Dry-salted salmon.	
	Barrels.	Value.	Barrels.	Value.	Pounds.	Value.
1868.....	2,000	\$16,000				
1869.....	1,700	13,600				
1870.....	1,800	14,400				
1871.....	700	6,300				
1872.....	1,000	9,000				
1873.....	900	7,200				
1874.....	1,400	11,200				
1875.....	1,200	9,600				
1876.....	1,800	14,400				
1877.....	1,950	15,700				
1878.....	2,100	16,800				
1879.....	3,500	28,000				
1880.....	3,700	29,600	300	\$3,300		
1881.....	1,760	15,840				
1882.....	5,890	53,010				

PACK OF SALTED SALMON IN ALASKA, 1868 TO 1909—Continued.

Year.	Salmon.		Salmon bellies.		Dry-salted salmon.	
	Barrels.	Value.	Barrels.	Value.	Pounds.	Value.
1883.....	7,251	\$65,259				
1884.....	6,106	54,954				
1885.....	3,230	29,070				
1886.....	4,861	43,749				
1887.....	3,978	35,802				
1888.....	9,500	85,500				
1889.....	6,457	58,013				
1890.....	18,039	162,351				
1891.....	8,913	71,304				
1892.....	17,374	140,057	53	\$815		
1893.....	24,005	120,083				
1894.....	32,011	176,060				
1895.....	14,234	85,404				
1896.....	9,314	65,198	150	1,200		
1897.....	15,848	110,936	2,846	28,460		
1898.....	22,670	181,360	580	5,800		
1899.....	22,382	167,865	235	2,350		
1900.....	31,852	238,890	2,353	23,530	511,400	\$10,228
1901.....	24,477	171,339	652	3,816		
1902.....	30,384	212,688	328	2,952		
1903.....	27,921	223,368	3,667	32,973	300,000	5,500
1904.....	13,674	89,209	208	1,950	966,812	16,180
1905.....	19,071	143,811	1,360	11,355	7,280,234	115,643
1906.....	17,283	126,194	1,338	13,644	1,107,680	16,969
1907.....	22,307	203,127	2,965	37,422	107,580	1,505
1908.....	31,472	266,713	7,600	85,994	20,800	416
1909.....	28,443	183,400	1,970	25,358	71,600	1,038
1910.....	12,779	111,634	1,626	19,007	22,178	554
Total.....	517,236	3,883,988	28,231	299,926	10,388,284	168,033

MILD CURING INDUSTRY.

The beginning of this industry on the Pacific coast is of comparatively recent date, and the following table is complete, with the possible exception of a few tierces, which may not have been reported for the coastal rivers of Oregon:

NUMBER OF TIERCES OF MILD-CURED SALMON PACKED ON THE PACIFIC COAST FROM 1897 TO 1910.^a

Year.	Alaska.	British Columbia.	Puget Sound, Wash.	Grays Harbor, Wash.	Willapa Harbor, Wash.	Columbia River, (both sides).	Coastal rivers, Ore.	Eel River, Cal.	Sacramento River, Cal.	Monterey Bay, Cal.	Total.
1897.....						400					400
1898.....	70					700					770
1899.....	130			375		1,250					1,755
1900.....						1,275		950			2,225
1901.....	67		600			3,000		3,100			6,767
1902.....	67		425			4,213	188	2,325		504	7,222
1903.....	8		824			6,725		3,600		354	11,511
1904.....	34		1,250			9,088		200	4,719	248	15,539
1905.....	189	1,175	3,000			9,805	415	2,979		310	17,873
1906.....	1,126	957				8,000	740	175	2,177	510	13,685
1907.....	1,657	1,993	2,060	20	100	6,070	740	140	4,102	582	17,464
1908.....	1,378	1,060				4,960			3,243	252	10,893
1909.....	2,292	1,560	2,109	75	29	5,540	560	80	5,111	911	18,267
1910.....	3,357	1,638	2,435	67		7,922	1,398		5,516	75	22,408
Total.....	10,375	8,383	12,703	537	129	68,948	4,041	595	37,822	3,746	147,279

^a The net weight of fish in a tierce is about 800 pounds. King, chinook, or spring salmon were used almost exclusively. From most places the data are complete from the time of the inception of the industry, but from a few minor places the data are somewhat fragmentary.

IX. TRADE WITH OUTLYING POSSESSIONS.

As a result of the war with Spain the United States in 1898 acquired possession of Porto Rico, Guam, and the Philippine Islands, while in the same year Hawaii became a part of this country at its own request, and in 1900 two islands of the Samoan group were acquired by a partition agreement with Great Britain and Germany. The trade with the Philippine Islands is shown to date in the tables of exports and imports to foreign countries, but the trade with the other possessions has been eliminated from these tables and shown separately ever since their annexation to the United States.

HAWAII.

The islands constituting this Territory, owing to their reciprocity treaty with this country for a number of years before annexation, purchased their supplies of salmon from the United States almost exclusively. In recent years the Territory has imported the following quantities of salmon from the mainland:

Year.	Canned.		All other, fresh or cured.
	Pounds.	Value.	
1907.....	1,126,217	\$89,286	<i>Value.</i> \$64,232
1908.....	965,029	89,025	67,143
1909.....	1,440,410	121,716	73,848
1910.....	1,381,398	113,526	72,194

PORTO RICO.

Of recent years, the following shipments of domestic salmon have been made to this island:

Year.	Canned.		All other, fresh or cured.
	Pounds.	Value.	
1907.....	604,627	\$53,916	<i>Value.</i> \$2,893
1908.....	512,038	48,195	1,428
1909.....	381,171	34,777	3,810
1910.....	511,055	43,494	6,243

GUAM.

Since annexation, this country and Japan have been competing for the trade of this island, which, in earlier years, Japan controlled quite largely. During the last two years, however, the United States has secured the advantage. The following table shows the extent of the trade, which is made up almost entirely of salted or pickled salmon:

Year and country.	Pickled salmon.		Fresh salmon.	
	Pounds.	Value.	Pounds.	Value.
1905.				
United States.....	1,415	\$71		
Japan.....	16,526	1,221		
1907.				
United States.....	13,604	1,086		
Japan.....	19,862	1,601		
1908.				
United States.....	7,406	623	900	\$92
Japan.....	6,130	465		
1909.				
United States.....	10,779	740		
Japan.....	4,295	344		
1910.				
United States.....				
Japan.....				

TUTUILA, SAMOA.

The customs statistics lump the imports of fish under one general heading, thus making it impossible to show separately the imports of salmon.

X. FOREIGN TRADE IN SALMON.

As we do not consume all of the salmon produced by our fisheries, it is necessary to find a foreign market for the surplus each season, but as canned salmon has become one of the staples of the world, there is not much difficulty in this respect, especially since our only competitors are Canada and Japan. The latter has not yet become much of a factor in the canned-salmon market, though she will as her fishing operations are extended. There is more competition in the pickled, fresh, and frozen markets, several European and Asiatic countries being large producers of these goods, as is Canada also, for a considerable proportion of which she is compelled to find an outside market.

EXPORTS OF CANNED SALMON.

From the beginning of the industry a considerable proportion of the salmon canned has been exported, especially of the higher grades. In Europe the chief customer is Great Britain, taking about nine-tenths of all sent to European ports. Great Britain does not, however, consume this quantity, for a considerable part of her importations are reexported. On the North American Continent and adjacent islands the best customers are Mexico, Panama, and the British West Indies, in the order named. In South America, Peru, Argentina, and British Guiana were the leading markets in 1910. In 1908 Chile imported 4,196,060 pounds; in 1909 the importations dropped to 97,993 pounds, but increased in 1910 to 1,556,629 pounds. In Asia, Hongkong and China import canned salmon, although neither buys great quantities. The islands of the Pacific and Indian Oceans are large consumers. British Australasia took 5,474,818 pounds, valued at \$551,312, in 1910, and other good customers were the British East Indies and British, French, and German Oceania. In Africa the British and Portuguese possessions are the largest importers.

The movements of these products are naturally often influenced favorably or adversely as the tariffs of the various countries in which they are marketed are raised or lowered.

The following table shows the yearly exports of domestic canned salmon and the countries to which exported for the period from 1900 to 1910, inclusive:

EXPORTS, BY COUNTRIES, OF DOMESTIC CANNED SALMON, 1900 TO 1910.

Countries.	1900		1901		1902	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
North America:						
Dominion of Canada—						
Nova Scotia, New Brunswick, etc.					10	\$1
Quebec, Ontario, Manitoba, etc.	24,137	\$2,514	101	\$10	22,442	2,493
British Columbia	382,811	33,454	1,725,251	223,230	1,866,272	159,682
Newfoundland and Labrador					810	73
Miquelon, Langley, etc.	240	20				
Mexico	162,785	14,806	160,425	14,967	387,905	31,041
Central American States—						
British Honduras	16,488	1,604	19,331	2,054	23,467	2,370
Costa Rica	70,458	6,114	69,135	6,768	70,036	5,954
Guatemala	2,666	277	11,361	1,151	15,325	1,324
Honduras	7,193	677	7,681	776	4,924	498
Nicaragua	26,647	2,684	21,543	2,256	17,125	1,635
Salvador	550	60	550	55	1,828	161
Bermuda	59,672	6,158	63,786	7,398	76,456	7,768
West Indies—						
British	259,249	25,651	315,209	33,635	242,999	24,191
Danish	9,085	873	8,612	929	14,526	1,390
Dutch	13,303	1,610	16,591	1,944	13,112	1,506
French	432	45	1,084	127	960	96
Haiti	468	44	505	65	920	88
Santo Domingo	2,764	297	1,899	192	1,531	140
Cuba	8,406	786	20,407	1,883	20,196	1,618
Porto Rico	4,394	390				
South America:						
Argentina	104,367	8,822	127,751	10,916	88,622	7,816
Bolivia			240	37	15,110	1,147
Brazil	637,638	76,152	207,033	23,506	87,800	8,350
Chile	647,328	61,800	645,323	64,059	384,766	28,529
Colombia	92,868	9,075	97,163	9,975	86,046	7,451
Ecuador	50,387	5,631	98,587	10,387	24,937	1,868
Guiana—						
British	168,718	16,197	136,192	14,807	146,502	14,604
Dutch	43,096	3,553	61,334	6,542	92,971	8,718
French	3,240	299	2,248	261	8,316	850
Peru	75,621	7,392	124,823	12,526	313,476	24,444
Uruguay	2,837	285	9,408	933	1,016	104
Venezuela	42,125	3,712	66,911	6,913	42,436	4,026
Europe:						
Austria-Hungary	2,208	309			250	25
Azores, and Madeira Islands	48	7	950	92		
Belgium	31,118	3,186	5,800	600	336	39
Denmark	24,492	2,455	3,168	326	860	92
France	22,544	2,130	61,790	6,565	23,956	1,889
Germany	16,110	1,431	77,921	7,567	10,905	1,068
Italy	120	10	2,496	244		
Malta, Gozo, etc.			141	21		
Netherlands	3,048	299	288	30	4,800	400
Portugal	19,776	1,779			336	35
Russia, on Baltic and White Seas					8,400	932
Spain			1,536	151	675	67
Sweden and Norway	1,168	112	720	70	72	8
Switzerland	24	3				
United Kingdom	18,820,453	1,870,004	31,722,853	3,219,196	30,632,961	2,620,729
Asia and Oceania:						
Aden	216	22				
Chinese Empire	40,960	4,255	149,295	15,263	117,043	8,716
China—Russian			20,634	2,058	9,460	772
Hongkong	63,210	6,488	78,960	8,056	551,860	40,261
Japan	11,560	1,200	285,036	28,990	14,578	1,220
Korea			1,105	115	2,208	179
Russia, Asiatic			1,495	145	6,572	521
Turkey in Asia			144	16		
East Indies—						
British	538,180	55,976	312,805	31,528	733,685	56,912
Dutch			3,960	400	161,940	12,093

EXPORTS, BY COUNTRIES, OF DOMESTIC CANNED SALMON, 1900 TO 1910—Continued.

Countries.	1900		1901		1902	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Asia and Oceania—Continued.						
British Australasia.....	2,804,004	\$283,110	3,442,085	\$343,540	7,131,641	\$599,671
British Oceania.....					151,998	10,555
French Oceania.....	103,940	10,732	118,355	12,026	142,570	11,355
German Oceania.....			8,480	874	12,900	997
Guam ^a	480	50				
Hawaii ^b	860,682	84,808				
Philippine Islands.....	1,160	120	39,316	3,925	718,876	46,712
Tonga, Samoa, and all other.....	112,380	11,646	73,040	7,168		
Tutuila ^c					21,176	1,451
Africa:						
British Africa.....	632,012	57,387	816,433	79,063	2,581,088	219,233
Canary Islands.....			656	66		
French Africa.....	4,320	421	4,080	415	200	21
Liberia.....	312	30				
Portuguese Africa.....	47,812	4,696	35,384	3,459	52,726	4,931
All other Africa.....					6,200	582
Total.....	27,082,370	2,693,648	41,289,500	4,230,271	47,173,114	3,991,402
RECAPITULATION.						
Europe.....	18,941,109	1,881,725	31,877,663	3,234,862	30,683,551	2,625,284
North America.....	1,051,808	98,064	2,443,561	297,440	2,780,844	242,029
South America.....	1,868,225	192,918	1,577,013	160,862	1,291,998	107,907
Asia.....	654,126	67,941	853,434	86,571	1,597,346	120,674
Oceania.....	3,882,646	390,466	3,681,276	367,533	8,179,161	670,741
Africa.....	684,456	62,534	856,553	83,003	2,640,214	224,767

Countries.	1903		1904		1905	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
North America:						
Dominion of Canada.....					290,850	\$21,121
Nova Scotia, New Brunswick, etc.....			49	\$4		
Quebec, Ontario, Manitoba, etc.....	43,107	\$5,171	153,697	9,558		
British Columbia.....	3,246,082	287,212	1,086,370	95,021		
Newfoundland and Labrador.....					240	25
Mexico.....	356,951	26,787	538,949	38,691	493,371	40,597
Central American States—						
British Honduras.....	24,187	2,316	28,044	2,534	28,959	2,534
Costa Rica.....	36,806	3,072	58,828	4,668	93,580	8,179
Guatemala.....	3,527	295	15,732	1,131	20,498	1,583
Honduras.....	7,455	716	12,428	1,090	14,434	1,221
Nicaragua.....	20,089	1,771	28,159	2,394	42,103	3,146
Panama ^d			18,466	1,671	112,320	9,211
Salvador.....	3,360	252	4,304	326	2,396	184
Bermuda.....	64,264	6,792	36,022	3,778	33,821	3,634
West Indies—						
British.....	418,636	38,434	409,219	37,389	366,747	34,262
Danish.....	9,647	903	7,442	752	9,474	965
Dutch.....	22,981	2,480	17,878	1,999	13,051	1,419
French.....	892	92	984	86	660	64
Haiti.....	2,496	238	2,115	228	1,611	164
Santo Domingo.....	3,290	335	7,660	719	4,855	452
Cuba.....	21,636	1,789	24,677	2,324	36,903	3,373
South America:						
Argentina.....	72,445	6,808	66,275	6,612	120,586	11,263
Bolivia.....	384	40	672	80	170	17
Brazil.....	88,740	8,481	114,033	11,742	188,342	17,908
Chile.....	1,044,490	59,354	1,218,266	72,205	821,171	56,160
Colombia.....	149,272	11,194	118,269	10,104	81,239	7,491
Ecuador.....	45,126	3,115	59,266	4,041	121,894	7,941
Guiana—						
British.....	172,300	16,829	112,360	11,226	135,424	13,617
Dutch.....	52,138	4,959	78,464	8,280	45,231	4,797
French.....	18,752	1,805	11,169	1,307	11,684	1,228
Peru.....	89,440	7,309	214,982	15,530	151,832	11,369
Uruguay.....	2,140	185	2,246	225	3,250	325
Venezuela.....	20,987	1,839	59,857	5,981	28,005	2,825

^a Guam was annexed to the United States in 1898.^b Hawaii was annexed to the United States in 1898.^c Tutuila was acquired in 1898.^d Panama separated from Colombia in 1903.

EXPORTS, BY COUNTRIES, OF DOMESTIC CANNED SALMON, 1900 TO 1910—Continued.

Countries.	1903		1904		1905	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Europe:						
Austria-Hungary.....	400	\$25	384	\$36		
Azores, and Madeira Islands.....			48	5	384	\$41
Belgium.....	788	73	480	53	9,760	1,019
Denmark.....	80	8	100	8		
France.....	2,400	260	4,800	600	21,995	2,262
Germany.....	32,268	2,470	18,790	1,747	1,210	122
Italy.....	1,120	114	5,232	556	5,760	465
Netherlands.....	1,072	124	4,072	414	3,250	349
Norway ^a	96	10	1,440	150		
Spain.....	3,108	316	1,400	140	2,700	249
Sweden ^a			70	7	96	10
Switzerland.....	240	24				
United Kingdom.....	35,369,196	3,121,774	33,555,080	3,505,102	21,026,108	1,872,992
Asia and Oceania:						
Aden.....					2,520	180
Chinese Empire.....	166,522	13,602	218,142	18,770	249,386	17,587
China—Russian.....	53,368	5,111	40,000	3,932		
Hongkong.....	814,008	56,225	160,367	11,870	518,423	36,635
Japan.....	13,536	1,015	11,817,343	841,461	2,437,484	162,524
Korea.....	2,152	179	3,888	292	2,572	186
Russia, Asiatic.....	48	4	482	41		
Siam.....					384	31
East Indies—						
British.....	473,740	39,367	636,320	44,669	673,897	55,599
French.....					720	69
Dutch.....	235,680	19,256	119,216	9,018	109,476	7,893
All other Asia.....	240	24	10	1		
British Australasia.....	4,268,652	360,720	3,136,728	290,307	4,075,094	389,518
British Oceania.....	36,018	2,290	28,670	1,941	42,624	3,645
French Oceania.....	153,696	12,179	185,848	15,305	133,204	11,414
German Oceania.....	451,824	26,614	340,464	19,326	324,888	20,651
Philippine Islands.....	601,324	42,702	206,896	14,970	681,636	42,700
Africa:						
British Africa.....	1,454,226	127,921	794,758	77,911	1,259,269	121,120
Canary Islands.....	144	15			900	90
French Africa.....	2,220	207	3,200	320	4,800	460
Liberia.....	384	41	140	14	140	14
Portuguese Africa.....	167,964	17,043	137,640	13,906	200,826	20,365
Turkey in Africa—Egypt.....			388	30	2,448	204
All other Africa.....	5,200	506				
Total.....	50,353,334	4,350,791	55,924,278	5,224,598	35,066,555	3,035,469
RECAPITULATION.						
Europe.....	35,410,768	3,125,197	33,591,896	3,508,818	21,071,263	1,877,509
North America.....	4,285,406	378,655	2,446,023	204,363	1,565,773	132,134
South America.....	1,756,214	121,918	2,055,859	147,333	1,708,828	134,941
Asia.....	1,759,294	134,783	12,995,768	930,054	3,994,862	280,704
Oceania.....	5,511,514	444,505	3,898,606	341,849	5,257,446	467,928
Africa.....	1,630,138	145,733	936,126	92,181	1,468,383	142,253

Countries.	1906		1907		1908	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
North America:						
Dominion of Canada.....	236,664	\$14,814	793,247	\$65,356	7,320	\$587
Mexico.....	699,002	56,747	877,989	73,582	1,068,824	94,278
Central American States—						
British Honduras.....	43,155	3,639	36,020	3,214	32,632	3,080
Costa Rica.....	106,879	8,968	148,157	12,260	138,421	12,260
Guatemala.....	26,925	1,989	31,242	2,535	29,777	2,319
Honduras.....	15,148	1,319	23,508	2,048	33,955	3,202
Nicaragua.....	39,949	3,022	41,106	3,335	27,721	2,302
Panama ^b	308,624	25,965	443,687	38,642	487,079	46,883
Salvador.....	2,880	197	4,092	331	5,854	467
Bermuda.....	24,679	2,406	29,139	2,711	25,183	2,579
West Indies—						
British.....	471,814	43,368	515,664	46,510	687,620	64,275
Danish.....	9,713	1,011	13,336	1,340	15,604	1,658
Dutch.....	11,643	1,230	24,275	2,428	21,368	2,234
French.....	200	20	100	9	96	11
Haiti.....	2,953	291	91	91	864	85
Santo Domingo.....	11,688	1,137	9,278	891	13,887	1,371
Cuba.....	57,441	5,823	60,904	5,855	57,970	5,288

^a Sweden and Norway separated in 1905.^b Panama separated from Colombia in 1903.

EXPORTS, BY COUNTRIES, OF DOMESTIC CANNED SALMON, 1900 TO 1910—Continued.

Countries.	1906		1907		1908	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
South America:						
Argentina.....	200,206	\$20,339	262,667	\$25,801	394,306	\$30,759
Bolivia.....	1,720	181	18,951	1,577	11,762	1,217
Brazil.....	188,278	18,975	150,592	14,880	146,826	14,055
Chile.....	4,462,147	154,396	4,168,876	286,229	4,196,060	295,194
Colombia.....	51,987	4,667	41,964	3,850	51,786	4,880
Ecuador.....	80,876	5,855	203,930	15,599	174,920	12,486
Guiana—						
British.....	120,016	12,391	116,120	12,202	140,514	16,014
Dutch.....	65,654	6,246	66,530	6,494	59,390	6,053
French.....	12,650	1,305	17,950	1,829	23,218	2,599
Peru.....	269,858	20,342	551,160	40,431	316,701	22,229
Uruguay.....	10,436	1,075	16,124	1,546	17,934	1,693
Venezuela.....	35,775	3,280	44,826	4,336	37,583	3,564
Europe:						
Austria-Hungary.....	1,260	135	1,220	112		
Azores, and Madeira Is- lands.....			883	89		
Belgium.....	500	60				
Denmark.....	40,200	4,112				
France.....	29,980	3,000			10,575	961
Germany.....	4,896	420	9,150	976	45,977	4,572
Italy.....	4,920	413	10,230	861		
Malta, Gozo, etc.....	420	36				
Netherlands.....	8,280	959	11,098	850		
Norway ^a	40,200	3,981			17,670	1,860
Portugal.....					7,577	731
Spain.....	1,930	193	3,208	303	27,900	2,735
Sweden ^a	10,000	1,050			10,500	1,000
United Kingdom.....	31,918,816	2,739,284	7,720,991	788,245	13,200,887	1,193,516
Asia and Oceania:						
Aden.....	480	50				
Chinese Empire.....	32,189	2,321	59,110	4,386	23,126	2,154
Hongkong.....	105,581	7,632	122,482	9,959	144,624	13,367
Japan.....	9,051	713	22,881	1,775	2,472	269
Korea.....	1,632	128	1,500	129	1,156	126
Russia, Asiatic.....	1,440	102	770	84	582	65
Siam.....			1,440	90	3,264	282
Turkey in Asia.....	750	90			290	30
East Indies—						
British.....	477,234	38,263	1,043,618	75,001	702,169	59,254
French.....	16,262	1,162			720	75
Dutch.....	134,796	9,692	167,590	13,940	126,168	11,286
British Australasia.....	5,230,076	426,814	5,451,378	462,648	3,654,756	330,029
British Oceania.....	11,952	923	40,080	2,958	14,660	1,278
French Oceania.....	125,998	10,274	137,472	11,494	185,608	15,732
German Oceania.....	214,920	14,503	156,939	11,267	105,696	8,345
Philippine Islands.....	757,400	56,743	933,288	63,838	1,171,834	84,533
Africa:						
British Africa.....	1,029,787	87,881	504,848	47,748	454,892	43,883
Canary Islands.....	782	76	144	17		
French Africa.....	144	14			48	6
German Africa.....			600	60		
Liberia.....					5,079	482
Portuguese Africa.....	161,178	16,001	104,837	10,307	83,640	8,325
Turkey in Africa—Egypt.....	2,400	200				
Total.....	45,944,414	3,847,943	25,218,105	2,183,049	28,226,045	2,438,518
RECAPITULATION.						
Europe.....	32,061,402	2,753,643	7,756,780	791,436	13,321,086	1,205,375
North America.....	2,069,357	171,946	3,052,658	261,138	2,654,175	242,879
South America.....	3,499,160	249,052	5,659,690	414,774	5,571,000	410,743
Asia.....	779,415	60,173	1,419,391	105,364	1,004,571	86,908
Oceania.....	6,340,346	509,257	6,719,157	552,205	5,131,554	439,917
Africa.....	1,194,291	103,872	610,429	58,132	543,659	52,696

^a Sweden and Norway separated in 1905.

EXPORTS, BY COUNTRIES, OF DOMESTIC CANNED SALMON, 1900 TO 1910—Continued.

Countries.	1900		1910	
	Pounds.	Value.	Pounds.	Value.
North America:				
Dominion of Canada.....	229,934	\$21,773	99,022	\$7,570
Mexico.....	756,052	58,124	697,217	50,782
Central American States—				
British Honduras.....	35,195	3,261	28,310	2,606
Costa Rica.....	118,266	9,828	157,946	12,237
Guatemala.....	13,957	1,117	16,821	1,361
Honduras.....	14,112	1,179	16,240	1,361
Nicaragua.....	21,534	1,656	28,116	2,066
Panama ^a	528,228	50,940	482,717	45,404
Salvador.....	9,184	754	5,498	423
Bermuda.....	23,774	2,461	26,484	2,383
West Indies—				
British.....	358,114	36,644	548,561	53,939
Danish.....	14,848	1,568	14,655	1,512
Dutch.....	16,621	1,883	9,838	1,160
French.....	564	69	196	18
Haiti.....	2,184	203	2,038	185
Santo Domingo.....	13,258	1,306	22,120	2,058
Cuba.....	53,580	5,277	68,737	6,486
South America:				
Argentina.....	259,192	17,030	229,461	15,690
Bolivia.....	6,184	647	33,502	2,941
Brazil.....	176,150	17,109	267,354	28,241
Chile.....	97,993	6,918	1,556,629	92,259
Colombia.....	58,518	5,767	114,274	9,494
Ecuador.....	139,808	10,952	272,411	16,487
Guiana—				
British.....	255,039	25,981	222,398	22,133
Dutch.....	100,259	9,906	57,509	6,297
French.....	22,816	2,164	17,724	1,784
Peru.....	295,885	22,640	367,676	24,817
Uruguay.....	15,140	1,330	11,730	1,167
Venezuela.....	34,618	3,058	43,144	4,887
Europe:				
Azores, and Madeira Islands.....			100	12
Denmark.....	192	18		
France.....			1,878	223
Germany.....	17,096	1,757	424	51
Italy.....	5,148	500		
Netherlands.....	11,612	1,017	9,744	1,020
Russia on Baltic and White Seas.....	2,050	205	11,580	1,210
Spain.....	3,160	311	5,100	506
Sweden ^b	20,000	1,940		
United Kingdom.....	22,969,218	2,201,446	44,737,072	4,709,160
Asia and Oceania:				
Chinese Empire.....	53,448	4,887	28,522	2,688
China—British leased territory.....			3,120	345
Hongkong.....	103,448	9,707	121,558	12,234
Japan.....	15,078	1,245	3,716	352
Korea.....	2,652	266	2,016	220
Russia, Asiatic.....	5,380	394		
Siam.....	14,880	1,025	1,008	93
East Indies—				
British.....	989,592	85,094	1,246,751	101,619
French.....	528	56		
Dutch.....	201,696	16,908	189,604	15,920
All other Asia.....			480	45
British Australasia.....	5,704,960	590,094	5,474,818	551,312
British Oceania.....	109,936	7,437	66,826	5,160
French Oceania.....	162,336	14,570	241,200	22,889
German Oceania.....	279,792	18,311	360,576	22,554
Philippine Islands.....	1,126,470	74,792	5,425,404	396,604
Africa:				
British Africa.....	484,196	48,220	357,051	37,707
Canary Islands.....	510	51		
German Africa.....	350	36	910	92
Portuguese Africa.....	162,314	14,604	151,470	14,674
Turkey in Africa—Egypt.....			1,440	120
Total.....	36,117,109	3,416,436	63,860,696	6,314,258
RECAPITULATION.				
Europe.....	23,028,476	2,207,194	44,765,898	4,712,182
North America.....	2,209,405	193,043	2,224,516	191,551
South America.....	1,461,662	123,502	3,193,812	226,197
Asia.....	1,386,702	119,582	1,596,775	133,516
Oceania.....	7,383,494	705,204	11,568,824	998,219
Africa.....	647,370	62,911	510,871	52,593

^a Panama separated from Colombia in 1903.^b Sweden and Norway separated in 1898.

The table following shows for the past 11 years the customs districts from which the canned salmon was exported. Up to 1910 about two-thirds of the total exports have gone from the port of San Francisco, while about one-fifth of the total passed through the port of Puget Sound, Wash. In 1910, however, the exports from Puget Sound exceeded those from San Francisco. The only other port through which any considerable quantity is shipped is New York City. It is usual now to load the salmon on steamers and sailing vessels at San Francisco and the Puget Sound cities to go direct to Europe.

EXPORTS, BY CUSTOMS DISTRICTS, OF CANNED SALMON, 1900 TO 1910.

Customs districts from which exported.	1900		1901		1902	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Atlantic ports:						
Baltimore, Md.....	648	\$65	334,580	\$33,053	324	\$34
Bangor, Me.....					10	1
Boston and Charlestown, Mass.....	222,770	20,488	192,676	27,372	172,110	20,224
New York, N. Y.....	3,485,326	340,538	7,960,104	847,294	4,365,074	407,009
Philadelphia, Pa.....	110,500	9,100	77,000	9,050		
Savannah, Ga.....	1,012	81	582	72	480	60
St. Johns, Fla.....					75	7
Norfolk and Portsmouth, Va.....			269,380	30,888		
Charleston, S. C.....	400	30				
Gulf ports:						
Key West, Fla.....			400	43		
Mobile, Ala.....	10,536	958	7,340	816	11,032	1,055
New Orleans, La.....	28,332	2,472	47,685	4,567	39,084	3,910
Mexican border ports:						
Arizona.....	6,253	706	18,104	1,869	23,879	2,350
Brazos de Santiago, Tex.....	168	21	816	115	300	29
Paso del Norte, Tex.....	23,843	2,134	1,220	98	164,167	13,119
Pacific ports:						
Alaska.....	289	38	4,859	291	3,636	558
Hawaii.....					48	4
Puget Sound, Wash.....	1,477,232	144,059	2,271,306	282,441	9,864,259	872,912
San Diego, Cal.....	3,094	220	3,574	293	6,202	487
San Francisco, Cal.....	21,611,030	2,164,904	30,014,055	2,983,982	32,327,572	2,654,020
Willamette, Oreg.....	76,800	5,320	43,318	3,517	155,500	11,250
Northern border and Lake ports:						
Detroit, Mich.....			26,200	2,700		
Minnesota, Minn.....			101	10		
Vermont, Vt.....	120	12				
Duluth, Minn.....	24,000	2,500	16,200	1,800	39,312	4,368
Memphremagog, Vt.....	17	2			50	5
Total.....	27,082,370	2,693,648	41,289,500	4,230,271	47,173,114	3,991,402
RECAPITULATION.						
Atlantic ports.....	3,820,656	370,302	8,834,322	947,729	4,538,073	427,335
Gulf ports.....	38,868	3,430	55,425	5,426	50,116	4,965
Mexican border ports.....	30,264	2,861	20,140	2,082	188,346	15,498
Pacific ports.....	23,168,445	2,314,541	32,337,112	3,270,524	42,357,217	3,539,231
Northern border and Lake ports.....	24,137	2,514	42,501	4,510	39,362	4,373

EXPORTS, BY CUSTOMS DISTRICTS, OF CANNED SALMON, 1900 TO 1910—Continued.

Customs districts from which exported.	1903		1904		1905	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Atlantic ports:						
Baltimore, Md.....	840	\$92	490	\$50	576	\$62
Bangor, Me.....			121	9	294	26
Boston and Charlestown, Mass.....	104,750	12,266	2,400	215		
New York, N. Y.....	5,627,654	599,393	2,129,523	214,016	2,683,775	266,599
Philadelphia, Pa.....	540	54	587	42	8,858	576
Providence, R. I.....	685	63				
Gulf ports:						
Key West, Fla.....			1,500	125	460	23
Mobile, Ala.....	9,612	824	9,203	811	7,102	561
New Orleans, La.....	44,404	4,261	61,909	5,503	89,999	7,841
Tampa, Fla.....			180	16		
Mexican border ports:						
Arizona.....	26,988	2,803	7,568	745	20,845	1,878
Brazos de Santiago, Tex.....			96	7		
Paso del Norte, Tex.....	103,375	8,938	347,218	23,401	262,014	20,687
Saluria, Tex.....			366	30	6,580	583
Pacific ports:						
Alaska.....			153,600	9,550	4,848	557
Hawaii.....			48	7	148	15
Puget Sound, Wash.....	16,527,456	1,549,319	19,766,003	1,655,666	4,444,562	326,485
San Diego, Cal.....	5,897	421	5,678	422	3,594	259
San Francisco, Cal.....	27,448,182	2,138,019	33,212,614	3,303,292	27,498,325	2,406,422
Willamette, Oreg.....	409,444	29,142	224,549	10,628	5,775	531
Oregon, Oreg.....	400	25				
Northern border and Lake ports:						
Detroit, Mich.....			580	58		
North and South Dakota.....			20	2		
Superior, Mich.....					28,800	2,364
Vermont, Vt.....	74	7	25	3		
Duluth, Minn.....	43,033	5,164				
Total.....	50,353,334	4,350,791	55,924,278	5,224,598	35,066,555	3,035,469
RECAPITULATION.						
Atlantic ports.....	5,734,469	611,868	2,133,121	214,332	2,693,503	267,263
Gulf ports.....	54,016	5,085	72,792	6,455	97,561	8,425
Mexican border ports.....	130,363	11,741	355,248	24,183	289,439	23,148
Pacific ports.....	44,391,379	3,716,926	53,362,492	4,979,565	31,957,232	2,734,269
Northern border and Lake ports.....	43,107	5,171	625	63	28,800	2,364
Customs districts from which exported.						
Atlantic ports:						
Baltimore, Md.....	196	\$21	156	\$28	301	\$37
New York, N. Y.....	3,275,875	318,128	2,313,335	227,646	2,332,392	226,850
Philadelphia, Pa.....	1,400	159	722	67	720	71
Portland and Falmouth, Me.....	100	13				
St. Johns, Fla.....			322	38	1,250	155
Gulf ports:						
Galveston, Tex.....	60	8	40,213	3,216	292	23
Key West, Fla.....	890	94	312	25	190	18
Mobile, Ala.....	38,267	3,031	11,675	992	10,823	1,051
New Orleans, La.....	88,014	7,775	112,850	10,217	194,711	18,144
Sabine, Tex.....					104	9
Tampa, Fla.....	24	2				
Mexican border ports:						
Arizona.....	45,883	4,128	34,479	3,268	43,035	3,856
Corpus Christi, Tex.....					30,930	2,775
Paso del Norte, Tex.....	387,568	30,336	513,202	42,548	626,837	56,147
Saluria, Tex.....	21,962	1,666	22,662	1,960	22,887	2,341
Pacific ports:						
Alaska.....			305,294	33,315	790	99
Hawaii.....					144	14
Los Angeles, Cal.....	840	53				
Puget Sound, Wash.....	17,286,930	1,499,819	9,340,000	845,982	6,351,440	528,558
San Diego, Cal.....	4,228	331	8,456	661	6,994	567
San Francisco, Cal.....	24,613,868	1,969,214	12,502,876	1,012,199	18,601,705	1,597,735
Willamette, Oreg.....	540	55	3,723	241	100	22

EXPORTS, BY CUSTOMS DISTRICTS, OF CANNED SALMON, 1900 TO 1910—Continued.

Customs districts from which exported.	1906		1907		1908	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
Northern border and Lake ports:						
Huron, Mich.....	177,734	\$13,107	7,000	\$570		
Minnesota, Minn.....			48	5		
Oswegatchie, N. Y.....			780	71	400	\$46
Vermont, Vt.....	35	3				
Total.....	45,944,414	3,847,943	25,218,105	2,183,049	28,226,045	2,438,518
RECAPITULATION.						
Atlantic ports.....	3,277,571	318,321	2,314,535	227,779	2,334,663	227,113
Gulf ports.....	127,255	10,910	165,050	14,450	206,120	19,245
Mexican border ports.....	455,413	36,130	570,343	47,776	723,689	65,119
Pacific ports.....	41,906,406	3,469,472	22,160,349	1,892,398	24,961,173	2,126,995
Northern border and Lake ports.....	177,769	13,110	7,828	646	400	46
Customs districts from which exported.						
	1909		1910			
	Pounds.	Value.	Pounds.	Value.		
Atlantic ports:						
Baltimore, Md.....	192	\$22	36		\$3	
Bangor, Me.....	216	25				
Boston and Charlestown, Mass.....	162,024	16,837	3,000		280	
New York, N. Y.....	3,848,870	390,266	2,999,480		305,732	
Philadelphia, Pa.....	405	44	700		89	
Norfolk and Portsmouth, Va.....	32,100	2,739				
Perth Amboy, N. J.....			214		18	
Gulf ports:						
Galveston, Tex.....	876	88	155		12	
Key West, Fla.....	40	4	340		27	
Mobile, Ala.....	13,565	1,247	14,018		1,322	
New Orleans, La.....	92,537	7,615	103,980		8,187	
Tampa, Fla.....			66		6	
Mexican border ports:						
Arizona.....	27,735	2,733	54,425		4,612	
Brazos de Santiago, Tex.....	138	13	641		64	
Corpus Christi, Tex.....	26,220	2,450	27,365		2,414	
Paso del Norte, Tex.....	150,636	14,850	125,169		11,560	
Saluria, Tex.....	14,399	1,528	47,117		2,853	
Pacific ports:						
Alaska.....	66,020	6,263				
Los Angeles, Cal.....	13,370	934	9,229		820	
Puget Sound, Wash.....	7,858,552	716,370	32,406,617		3,331,174	
San Diego, Cal.....	5,546	460	6,355		583	
San Francisco, Cal.....	23,761,656	2,247,957	28,027,911		2,641,608	
Willamette, Oreg.....			78		11	
Northern border and Lake ports:						
Detroit, Mich.....	42,000	3,990				
North and South Dakota.....	12	1				
Duluth, Minn.....			33,200		2,800	
Montana and Idaho.....			600		83	
Total.....	36,117,109	3,416,436	63,860,696		6,314,258	
RECAPITULATION.						
Atlantic ports.....		4,043,807		409,933		306,122
Gulf ports.....		107,018		8,954		9,554
Mexican border ports.....		219,128		21,574		21,503
Pacific ports.....		31,705,144		2,971,984		5,974,196
Northern border and Lake ports.....		42,012		3,991		2,883

EXPORTS OF FRESH AND CURED SALMON.

The following table shows, by countries, the value of the exports of fresh and cured salmon for the period 1900 to 1910, inclusive. As with the canned salmon, the greater part of these exports go to European countries, Germany taking by far the largest quantity. A small portion of this is salmon caught in eastern waters.

EXPORTS, BY COUNTRIES RECEIVING, OF DOMESTIC PICKLED, FRESH, ETC., SALMON, 1900 TO 1910.

Exported to—	1900	1901	1902	1903	1904	1905
North America:						
Bermuda.....	\$88	\$14	\$11	\$21		\$246
British Honduras.....	7	9		22	\$120	94
Dominion of Canada—						
Nova Scotia, New Brunswick, etc.....					418	3
Quebec, Ontario, Manitoba, etc.....	1,516	2,555	1,051	6,083	3,572	7,499
British Columbia.....	80,652	53,922	125,916	53,592	25,913	10,299
Central American States—						
Costa Rica.....	220	703	218	178	340	192
Guatemala.....			27	11	1	208
Honduras.....		5		1	2	26
Nicaragua.....	53	26	40	78	40	75
Panama.....					167	315
Salvador.....		22		7		2
Mexico.....	1,330	664	1,925	1,397	1,266	1,136
West Indies—						
British.....	943	939	2,348	5,150	3,867	4,999
Cuba.....	429	376	273	114	194	162
Danish.....	12	31	38	84	13	67
Dutch.....	195	167	293	177	197	238
French.....	126	122	315	199	273	100
Haiti.....	181	191	164	54	11	124
Porto Rico.....	1,214					
Santo Domingo.....	998	670	85	57	14	26
South America:						
Argentina.....					143	1,641
Bolivia.....			1,200			
Brazil.....	172	38	419	385	227	1,160
Chile.....	142			70	164	
Colombia.....	416	223	657	441	17	
Ecuador.....			65			15
Guiana—						
British.....	30	82	30	262	60	161
Dutch.....	400	226	286	11	766	176
French.....	420	290	134	434	251	65
Peru.....	26		27	62	194	112
Venezuela.....	96	42	245	25		108
Europe:						
Azores, and Madeira Islands.....	3				123	85
Belgium.....		1,062	88		4,750	
Denmark.....	378	15,285	16,904	653	2,315	22,952
France.....	180	300			57	
Germany.....	300,291	320,369	470,657	741,634	1,061,944	1,666,787
Greece.....						158
Italy.....						100
Malta, Gozo, etc.....	475	55	280	28		
Netherlands.....	50	184	3,023	4,127	3,105	300
Norway.....				12,765	12,295	7,896
Russia in Europe.....	300					2,574
Spain.....						56
Sweden and Norway ^a	7	5,595	5,685			
Sweden.....					1,838	17,776
United Kingdom.....	38,959	1,528		990	8,523	29,355
Asia:						
Chinese Empire.....		400	25	9	54	201
China—Russian.....				15		
East Indies—						
British.....		121	71	30	115	135
Dutch.....					275	
Hongkong.....	507		519	1,840	462	4,797
Japan.....	2,807	14,516	25,228	3,499	476	25,037
Russia—Asiatic.....	10					
Oceania:						
British Australasia.....	39,867	618	33,785	31,503	25,208	21,595
All other British Oceania.....			346	29	27	22
French Oceania.....	1,958	1,729	1,325	1,877	1,838	2,299
German Oceania.....			13	948	977	861
Guam.....	57	3,420				
Hawaii.....	58,870					

^aSweden and Norway separated in 1898.

EXPORTS, BY COUNTRIES RECEIVING, OF DOMESTIC PICKLED, FRESH, ETC., SALMON,
 1900 to 1910—Continued.

Exported to—	1900	1901	1902	1903	1904	1905
Oceania—Continued.						
Philippine Islands.....			\$384	\$478	\$13	\$308
Tonga, Samoa, and all other.....	\$636	\$215				
Tutuila.....			10			
Africa:						
British Africa—						
West.....			304			
South.....	170	24	21	12	859	114
French Africa.....	85					
Liberia.....					5	
Total.....	535,276	426,738	694,435	869,352	1,163,489	1,832,655
RECAPITULATION.						
North America.....	87,964	60,416	132,704	67,225	36,408	25,809
South America.....	1,702	901	3,063	1,690	1,822	3,438
Europe.....	340,643	344,368	496,637	760,197	1,094,950	1,748,039
Asia.....	3,324	15,037	25,843	5,393	1,382	30,170
Oceania.....	101,388	5,982	35,863	34,835	28,063	25,085
Africa.....	255	24	325	12	864	114

Exported to—	1906	1907	1908	1909	1910
North America:					
Bermuda.....	\$173	\$20	\$23	\$68	\$630
British Honduras.....	14		1,036		
Dominion of Canada—Nova Scotia, New Brunswick, etc.....	32,925	18,785	16,964	21,973	23,559
Central American States—					
Costa Rica.....	46	213	189	217	197
Guatemala.....	40		902	18	62
Honduras.....	39	92	2,451		
Nicaragua.....	39	27	1,317	31	11
Panama.....	380	2,211	1,878	175	775
Mexico.....	1,231	528	460	199	555
West Indies—					
British.....	1,646	208	975	4,890	3,067
Cuba.....	128	371	104	121	97
Danish.....	30	108	39	165	42
Dutch.....	94	93		49	78
French.....		16	19	14	19
Haiti.....	97	277	678	335	283
Santo Domingo.....	100	255	228	128	313
South America:					
Argentina.....	85	500			
Brazil.....	308			120	3,029
Chile.....	15	20	56		
Colombia.....	105	67	90	22	167
Ecuador.....		391		290	
Guiana—					
British.....	218	5	48	76	823
Dutch.....	287	133	130	271	217
French.....	57	36	75	21	695
Peru.....	1,317	1,163	118	555	
Venezuela.....	208	36			311
Uruguay.....				10	
Europe:					
Azores, and Madeira Islands.....		95			
Belgium.....	114			410	
Denmark.....	36,623	108,269	90,015	81,195	83,580
France.....		150		250	415
Germany.....	1,670,366	1,601,166	1,422,846	1,038,530	1,223,595
Italy.....	137				
Netherlands.....	793	264	2,947		
Norway.....	9,303	11,390	22,104	22,917	45,885
Portugal.....		1,650			
Russia in Europe.....		140		14,735	5,260
Spain.....		55		289	
Sweden.....	32,554	23,469	21,540	23,670	42,725
United Kingdom.....	26,196	48,237	28,083	43,952	66,555
Asia:					
Chinese Empire.....	3,391	293	170	41	89
East Indies—					
British.....					
Dutch.....	63		66	18	60
Hongkong.....	1,339	687	13	809	41
Japan.....	88,068	18,395	3,592	2,772	90
Korea.....		3			3
Russia—Asiatic.....		6	121		
Turkey in Asia.....					55

EXPORTS, BY COUNTRIES RECEIVING, OF DOMESTIC PICKLED, FRESH, ETC., SALMON,
1900 TO 1910—Continued.

Exported to—	1906	1907	1908	1909	1910
Oceania:					
British Australasia.....	\$15,169	\$23,186	\$26,591	\$25,466	\$22,826
All other British Oceania.....	21		11		89
French Oceania.....	2,154	2,136	1,792	1,528	1,886
German Oceania.....	749	1,112	373	1,229	1,189
Philippine Islands.....	821	12,287		712	2,089
Africa:					
British Africa—South.....	20				1,268
Liberia.....	40				
Portuguese Africa.....			198		
Spanish Africa.....				289	
Total.....	1,927,464	1,878,743	1,648,044	1,288,560	1,532,640
RECAPITULATION.					
North America.....	36,943	23,204	27,263	28,383	29,688
South America.....	2,600	2,351	517	1,365	5,242
Europe.....	1,776,086	1,794,885	1,587,535	1,225,948	1,468,015
Asia.....	92,861	19,384	3,962	3,640	348
Oceania.....	18,914	38,721	28,767	28,935	28,079
Africa.....	60	198		289	1,268

The exports of domestic fresh and cured salmon from 1900 to 1910, inclusive, are shown below, by customs districts. The greater part of the shipments pass through the New York City customs district:

EXPORTS, BY CUSTOMS DISTRICTS, OF DOMESTIC PICKLED, FRESH, ETC., SALMON,
1900 TO 1910.

Customs districts from which exported.	1900	1901	1902	1903	1904	1905
Atlantic ports:						
Baltimore, Md.....			\$158			\$8
Bangor, Me.....						3
Belfast, Me.....	\$12	\$17	12	\$19	87	
Boston and Charlestown, Mass.....	16		34	52	418	
New York, N. Y.....	346,853	330,805	503,219	766,128	1,102,542	1,757,742
Philadelphia, Pa.....	10			1,151	7	
Portland and Falmouth, Me.....	11	68	46	47	60	79
Savannah, Ga.....	22					
Gulf ports:						
Mobile, Ala.....				30	8	96
New Orleans, La.....		5	143		116	63
Mexican border ports:						
Arizona.....	18	85	416	115		14
Brazos de Santiago, Tex.....				19	4	
Corpus Christi, Tex.....	414	13		30	208	
Paso del Norte, Tex.....	760	67	13		80	206
Saluria, Tex.....		370	1,428	1,063	868	777
Pacific ports:						
Alaska.....	2,377	12,422	293	4,375	1,003	1,184
Oregon, Oreg.....		17,500				
Puget Sound, Wash.....	80,493	55,727	150,906	58,278	29,212	36,145
San Diego, Cal.....	108	19	20	34	73	4
San Francisco, Cal.....	102,666	7,030	36,958	36,331	25,851	27,939
Willamette, Oreg.....					28	1,500
Northern border and Lake ports:						
Champlain, N. Y.....	234	1,464	449	1,542	1,183	2,142
Detroit, Mich.....		742	24		1,393	4,445
Genesee, N. Y.....					26	
Huron, Mich.....	456	121	225	55		
Memphremagog, Vt.....			6	7	24	
Montana and Idaho.....	2	6				6
North and South Dakota.....	523	162	95	36	378	247
Superior, Mich.....						33
Vermont, Vt.....	301	115	20	40		22
Total.....	535,276	426,738	694,435	869,352	1,163,489	1,832,655
RECAPITULATION.						
Atlantic ports.....	346,924	330,890	503,439	767,397	1,103,034	1,757,832
Gulf ports.....		5	143	30	124	159
Mexican border ports.....	1,192	535	1,857	1,227	1,160	997
Pacific ports.....	185,644	92,698	188,177	99,018	56,167	66,772
Northern border and Lake ports.....	1,516	2,610	819	1,680	3,004	6,895

EXPORTS, BY CUSTOMS DISTRICTS, OF DOMESTIC PICKLED, FRESH, ETC., SALMON,
1900 TO 1910—Continued.

Customs districts from which exported.	1906	1907	1908	1909	1910
Atlantic ports:					
Baltimore, Md.....	\$11			\$31	
Bangor, Me.....			\$7	58	
Belfast, Me.....	15	\$8		11	\$12
New York, N. Y.....	1,781,330	1,786,105	1,590,757	1,230,436	1,479,625
Philadelphia, Pa.....	105				
Portland and Falmouth, Me.....	15	11,298	14	6	19
Gulf ports:					
Mobile, Ala.....	14		128		
New Orleans, La.....		276	7,098	49	74
Mexican border ports:					
Arizona.....	700	134	13	25	
Brazos de Santiago, Tex.....					5
Paso del Norte, Tex.....	8	290	154		
Saluria, Tex.....	80				197
Pacific ports:					
Alaska.....	44,436	451	803	1,091	212
Puget Sound, Wash.....	63,626	44,492	14,370	11,677	22,666
San Diego, Cal.....	44		28	4	12
San Francisco, Cal.....	31,500	28,984	29,112	37,305	27,628
Willamette, Oreg.....				743	3
Hawaii.....				14	
Northern border and Lake ports:					
Buffalo Creek, N. Y.....				3,069	
Cape Vincent, N. Y.....		92			
Champlain, N. Y.....	992	4,333	1,359	2,079	598
Detroit, Mich.....	3,954	1,972	1,667		
Duluth, Minn.....					68
Huron, Mich.....	428		284	891	
Memphremagog, Vt.....					20
Minnesota, Minn.....	40	52	798	59	
Montana and Idaho.....	69	92	45	154	82
North and South Dakota.....	36	3	20		
Vermont, Vt.....	61	161	1,387	858	1,419
Total.....	1,927,464	1,878,743	1,648,044	1,288,560	1,532,640
RECAPITULATION.					
Atlantic ports.....	1,781,476	1,797,411	1,590,778	1,230,542	1,479,656
Gulf ports.....	14	276	7,226	49	74
Mexican border ports.....	788	424	167	25	202
Pacific ports.....	139,606	73,927	44,313	50,834	50,521
Northern border and Lake ports.....	5,580	6,705	5,560	7,110	2,187

IMPORTS OF FRESH SALMON.

For some years it was the custom of the canneries on Puget Sound, when fish were scarce on the American side and abundant on the Canadian side, to import fresh salmon to fill out the domestic supply, and the Canadian canneries would do the same when the conditions were reversed. In 1904 the Canadian Government prohibited the export of fresh salmon to Puget Sound for packing purposes, and in 1910 an effort was made to have Congress retaliate by enacting a similar law for this side of the line, but the bill failed of passage. The reciprocity agreement with Canada now before Congress provides for the free entry of fresh fish and would permit the canneries of either country to import salmon as they wished. This agreement, if adopted, will undoubtedly be of considerable importance to the Puget Sound canneries in securing full packs in certain poor years.

The table below shows the yearly imports of fresh salmon from British Columbia:

IMPORTS OF FRESH SALMON FROM BRITISH COLUMBIA, CANADA, FOR A SERIES OF YEARS.

Year.	Pounds.	Value.	Year.	Pounds.	Value.	Year.	Pounds.	Value.
1890.....	4,660	\$241	1897.....	93,454	\$2,681	1904.....	40,610	\$1,025
1891.....	4,950	170	1898.....	11,580	278	1905.....	1,015	35
1892.....	6,288	301	1899.....	58,002	4,101	1906.....	3,457,738	64,408
1893.....	64,811	3,639	1900.....	19,404	855	1907.....	113,224	4,131
1894.....	3,872	219	1901.....	27,072	2,050	1908.....	8,880	795
1895.....	14,000	1,403	1902.....	22,353	739	1909.....	41,073	2,346
1896.....	11,799	419	1903.....	6,860	343	1910.....	198,251	10,116

IMPORTS OF CURED SALMON.

Below are shown the imports into this country of foreign-cured salmon, the product of the Pacific salmon fisheries, from 1886 to 1909, inclusive.

IMPORTS OF FOREIGN PICKLED PACIFIC SALMON, 1886 TO 1909.

Year.	British Columbia.		Japan.		Hongkong.		Russia, Asiatic.		Total.	
	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.	Pounds.	Value.
1886.....	5,600	\$224							5,600	\$224
1887.....	200	4							200	4
1888.....	86,000	4,031							86,000	4,031
1889.....	18,200	800							18,200	800
1890.....	600	36							600	36
1891.....	200	5							200	5
1892.....										
1893.....	5,478	291							5,478	291
1894.....	149,410	17,592			1,200	\$29	11,875	\$298	162,485	17,919
1895.....	6,550	250			600	13			7,150	263
1896.....	6,530	474							6,530	474
1897.....	6,890	156							6,890	156
1898.....	4,145	188			30	2	9,870	206	14,045	456
1899.....	15,875	1,554							a 16,032	a 1,560
1900.....	162,558	11,061	600	\$41					163,158	11,102
1901.....	165,243	11,225							165,243	11,225
1902.....	175,411	13,794	606	28					176,017	13,822
1903.....	161,549	11,756	360	18					161,909	11,774
1904.....	282,210	23,319	1,400	52					283,610	23,371
1905.....	282,027	25,584	3,015	133					285,042	25,717
1906.....	35,475	1,730	5,510	175					40,985	1,905
1907.....	6,393	322	680	31					7,073	353
1908.....	13,230	631	4,185	174					17,415	805
1909.....	30,710	1,523	3,537	148					34,247	1,671
1910.....	111,645	5,505								

a Includes 157 pounds, valued at \$6, from China.

XI. SALMON CULTURE.

CALIFORNIA.

HISTORY.

The first fish-cultural station on the Pacific coast was located on McCloud River, a stream of the Sierra Nevada Mountains emptying into Pitt River, a tributary to the Sacramento, 323 miles nearly due north of San Francisco. The site on the west bank of the river, about 3 miles above the mouth, was chosen after investigation of a number of places on the Sacramento, by Mr. Livingston Stone, one of America's pioneer fish culturists, and the station was named Baird, in honor of the then Commissioner of Fisheries, Prof. Spencer F. Baird. Although the season had nearly passed when the station was sufficiently advanced to handle eggs, 50,000 eggs were secured, and while 20,000 were lost, owing to the excessive heat, the remaining 30,000 were shipped east, all of which were eventually lost but 7,000 fry, which were planted in the Susquehanna River, in Pennsylvania.

The main object of the hatchery the first few years was to secure eggs to ship to the East for the purpose of introducing Pacific salmon in the waters in that section. The Commission early made an agreement with the State of California, however, under which the latter at first paid part of the expense, and the Commission hatched and planted a portion of the take in the McCloud River. Later, part of the eggs were turned over to the State, which hatched and planted the salmon in local waters.

In 1881 the station buildings were washed away in a freshet, but were immediately rebuilt. From 1884 to 1887, both inclusive, all operations were suspended.

In 1889 a hatchery was established at Fort Gaston, on the Army reservation in the Hoopa Indian Reservation in Humboldt County, but it was not put into operation until 1890. As the reservation was abolished on July 1, 1892, the Commission took complete charge of the plant, and in 1893 established a tributary station on Redwood Creek. The same year Korbels station was established about one-half mile above Korbels, on Mad River, in Humboldt County. Owing to the lack of money this station was closed in the fiscal year 1896, but was reopened during the fiscal year 1897.

That same year the Commission erected, on ground owned by the State, a hatchery at Battle Creek, in Tehama County, and also took charge of and operated the hatchery erected at this place by the State fish commission the previous year. Under the terms of an

agreement the Commission was to deliver to the State as many eyed spawn as the latter could hatch at Sisson, its own station.

Owing to their inaccessibility, the Fort Gaston hatchery and its substations were abandoned in 1898. The same year an experimental station was established at Olema, Bear Valley, in Marin County, whence eggs were transferred from Baird station, hatched out here, and planted in Olema Creek in order to see if they could not be domesticated here, where they had not been found previously.

During the fiscal year 1902 a substation was established on Mill Creek, a stream which has its source in the foothills of the Sierra Mountains, in the northeastern part of Tehama County, and empties into the Sacramento River from the east about a mile above the town of Tehama. The eggs are retained here until eyed and then shipped to other hatcheries.

As stated above, the State aided the work of the United States Fish Commission in a financial way and also by hatching and distributing the eggs turned over to its care. In 1885 the State legislature passed a bill authorizing the establishment of a hatchery of its own, and the same year such a station was built upon Hat Creek about $2\frac{1}{2}$ miles above its junction with Pitt River, a tributary of the Sacramento River. As the work of the first few seasons developed that the location was unsuitable, the hatchery was removed in 1888 to Sisson, in Siskiyou County. The work of this hatchery was to handle the eggs turned over to it by the United States Fish Commission.

In 1895 another hatchery was built by the State near the mouth of Battle Creek, a tributary of the Sacramento River. In 1896 and 1897 this hatchery was operated jointly by the State and the United States Fish Commission while awaiting the appropriation of money by the Commission to purchase it from the State.

In the fall of 1897 a hatchery was established by the State on Price Creek, a tributary of Eel River, in Humboldt County, and in 1902 this hatchery made the first plant in the State of steelhead trout fry.

Santa Cruz County has had a hatchery at Brookdale for a number of years.

OUTPUT.

The following tables show separately the quantity of eggs, fry, etc., distributed by the United States Fish Commission and the State since the inception of the work. The large quantity of eggs shown by the Commission represents largely the eggs supplied to the State, which hatched and distributed them, and eggs sent to other States and to foreign countries.

OUTPUT OF HATCHERIES OWNED BY THE UNITED STATES BUREAU OF FISHERIES.

Year ending June 30 ^a —	Chinook.		Silver fry.	Steelhead trout.		Total.	
	Eggs.	Fry.		Eggs.	Fry.	Eggs.	Fry.
1872.....	30,000					30,000	
1873.....	1,400,000					1,400,000	
1874.....	4,155,000	850,000				4,155,000	850,000
1875.....	6,250,000	1,750,000				6,250,000	1,750,000
1876.....	5,065,000	1,500,000				5,065,000	1,500,000
1877.....	4,983,000	2,000,000				4,983,000	2,000,000
1878.....	7,810,000	2,500,000				7,810,000	2,500,000
1879.....	4,250,000	2,300,000				4,250,000	2,300,000
1880.....	3,800,000	2,000,000				3,800,000	2,000,000
1881.....	4,300,000	3,100,000				4,300,000	3,100,000
1882.....		3,991,750					3,991,750
1883.....		776,125					776,125
1889 ^b	3,450,000	1,500,000				3,450,000	1,500,000
1890.....	1,554,000	84,000				1,554,000	84,000
1891.....	2,988,000	777,000				2,988,000	777,000
1892.....	2,902,000	315,500				2,902,000	315,500
1893.....	3,530,000	1,190,100				3,530,000	1,190,100
1894.....	7,500,000	438,500	280,000	75,000	308,500	7,575,000	1,027,000
1895.....	3,676,000	500,000	c1,250,000		d1,184,500	3,676,000	2,934,500
1896.....	6,170,800	715,700		175,000	107,808	6,345,800	823,508
1897.....	18,232,590	3,056,701	298,137	50,000	257,000	18,282,590	3,611,838
1898.....	30,605,000	15,643,300		60,000	650,000	30,665,000	16,293,300
1899.....	27,665,000	3,275,110				27,665,000	3,275,110
1900.....	2,925,000	3,533,950				2,925,000	3,533,950
1901.....	3,934,036	889,570				3,934,036	889,570
1902.....	17,580,410	2,115,560				17,580,410	2,115,560
1903.....	11,275,777	1,618,066				11,275,777	1,618,066
1904.....	64,598,354	2,350,130				64,598,354	2,350,130
1905.....	96,025,765	7,561,380				96,025,765	7,561,380
1906.....	107,905,945	e3,496,405				107,905,945	3,496,405
1907.....	73,376,315	2,512,250				73,376,315	2,512,250
1908.....	64,990,550	4,780,855				64,990,550	4,780,855
1909.....	32,278,265	3,590,078				32,278,265	3,590,078
1910.....	30,539,467	2,286,257				30,539,467	2,286,257
Total.....	655,746,274	82,998,287	1,828,137	360,000	2,507,808	656,106,274	87,334,232

^a The calendar year was used up to 1889.^b The hatchery was closed from 1884 to 1888.^c Includes 560,000 fingerlings, yearlings, or adults.^d Includes 332,000 fingerlings, yearlings, or adults.^e Includes 138 fingerlings, yearlings, or adults.

OUTPUT OF HATCHERIES OWNED BY THE STATE OF CALIFORNIA.

Year.	Chinook.		Steel- head fry.	Total.	
	Eggs.	Fry. ^a		Eggs.	Fry.
1873.....			520,000		520,000
1874.....			850,000		850,000
1875.....	b250,000	2,250,000		250,000	2,250,000
1876.....		2,000,000			2,000,000
1877.....		2,200,000			2,200,000
1878.....		2,500,000			2,500,000
1879.....		2,300,000			2,300,000
1880.....		2,225,000			2,225,000
1881.....		2,420,000			2,420,000
1882.....		3,991,750			3,991,750
1884.....		600,000			600,000
1886.....		150,000			150,000
1887.....		200,000			200,000
1888.....		1,290,000			1,290,000
1889.....		2,168,000			2,168,000
1890.....		1,320,000			1,320,000
1891.....		2,798,000			2,798,000
1892.....		2,651,000			2,651,000
1893.....		3,941,650			3,941,650

^a The greater part of the output of chinook fry was from eggs supplied by the United States Bureau of Fisheries hatcheries in California.^b All were lost.

OUTPUT OF HATCHERIES OWNED BY THE STATE OF CALIFORNIA—Continued.

Year.	Chinook.		Steel-head fry.	Total.	
	Eggs.	Fry.		Eggs.	Fry.
1894.....		7,776,400			7,776,400
1895.....		3,435,000			3,435,000
1896.....		15,283,183			15,283,183
1897.....		18,123,000			18,123,000
1898.....		31,476,388			31,476,388
1899.....		21,234,000			21,234,000
1900.....		2,536,000			2,536,000
1901.....		3,239,000			3,239,000
1902.....		16,852,040	301,000		17,153,040
1903.....		20,040,487	120,000		20,160,487
1904.....		63,632,000	90,000		63,722,000
1905.....		87,000,000	108,000		87,108,000
1906.....		105,815,920	243,000		106,058,920
1907.....		71,267,000	352,000		71,619,000
1908.....		60,619,000	170,000		60,789,000
1909.....		28,000,000	517,000		28,517,000
1910.....		28,469,745	667,800		29,137,545
Total.....	250,000	621,174,563	2,568,800	250,000	623,743,363

DISTRIBUTION.

The following table shows, by streams and species, the distribution in California of the eggs, fry, etc., from the hatcheries of the United States Fish Commission and the State. This far from represents the work of the hatcheries, as large quantities of eggs were sent to other States and foreign countries.

DISTRIBUTION OF SALMON EGGS, FRY, ETC., IN THE WATERS OF CALIFORNIA.

Year.	Klamath River and tributaries.				Redwood Creek and tributaries.			
	Chinook.		Silver.		Chinook.	Silver.		Steel-head.
	Fry.	Year-lings.	Fry.	Adults and year-lings.	Fry.	Fry.	Adults and year-lings.	Fry.
1890.....	90,000							
1891.....	30,000				25,000			
1892.....	147,600	25,000			142,500			
1893.....	487,200				170,000			
1895.....			300,000	160,000		140,000	400,000	
1896.....					65,700			107,808
1897.....					280,250	124,750		202,000
1898.....	16,000				1,260,000			650,000
1903.....	40,000							
Total.....	810,800	25,000	300,000	160,000	1,943,450	264,750	400,000	959,808

DISTRIBUTION OF SALMON EGGS, FRY, ETC., IN THE WATERS OF CALIFORNIA—CON.

Year.	Mad River and North Fork.			Eel River.		Russian River.	Skaggs Springs.	Marin County creeks.
	Chinook.	Silver.	Steel-head.	Chinook.	Steel-head.	Chinook.	Chinook.	Chinook.
	Fry.	Fry.	Fry.	Fry.	Fry.	Fry.	Fry.	Fry.
1881.....						15,000	15,000	
1894.....		280,000	308,500					
1895.....		470,000						
1897.....	145,365	173,387	60,000					635,000
1898.....				7,857,388				1,970,000
1899.....				8,202,000				900,000
1900.....				885,000				
1902.....				2,069,500	301,000			
1903.....				5,257,947	120,000			
1904.....				5,200,000	90,000			
1905.....				8,100,000				
1906.....				9,265,920	243,000			
1907.....				7,570,000	352,000	25,000		25,000
1908.....				6,154,000				
1909.....				5,500,000	349,000			
Total.....	145,365	923,387	368,500	66,061,755	1,455,000	40,000	15,000	3,530,000

Year.	Sacramento River and tributaries.				San Francisco Bay streams	San Gregorio River.	Pescadero Creek.	Monterey Bay and tributaries.
	Chinook.			Steel-head.	Chinook.	Chinook.	Chinook.	Chinook.
	Eggs.	Fry.	Year-lings, finger-ling, and adults.	Fry.	Fry.	Fry.	Fry.	Fry.
1873.....	20,000	520,000						
1874.....		850,000						
1875.....	250,000	2,000,000						
1876.....		2,000,000						
1877.....		2,200,000						
1878.....		2,500,000						
1879.....		2,300,000						
1880.....		2,225,000						
1881.....		2,300,500			20,000	15,000	15,000	30,000
1882.....	80,300	3,991,750						
1884.....		600,000						
1886.....		150,000						
1887.....		200,000						
1888.....		1,290,000						
1889.....		3,668,000						
1890.....		1,404,000						
1891.....		3,520,000						
1892.....		2,676,500						
1893.....		4,474,750						
1894.....		8,214,800		45,000				
1895.....		3,935,000						
1896.....		15,683,183	250,000					
1897.....		19,264,086						
1898.....		33,998,300						
1899.....	85,200	16,307,110						
1900.....		5,184,950						
1901.....		4,128,570						
1902.....		16,898,100						
1903.....		16,359,605						
1904.....		60,782,130						
1905.....		94,561,380		108,000				
1906.....		100,038,552						900,000
1907.....		66,209,250		135,000				1,200,000
1908.....		59,245,855		170,000				800,000
1909.....		26,090,000		168,000				
Total.....	435,500	585,771,472	250,000	626,000	20,000	15,000	15,000	2,930,000

a All were lost.

DISTRIBUTION OF SALMON EGGS, FRY, ETC., IN THE WATERS OF CALIFORNIA—CON.

Year.	Monterey Bay and tributaries.		Truckee River.	Total.					
	Silver.	Steel-head.	Chinook.	Chinook.			Silver.		Steel-head.
	Fry.	Fry.	Fry.	Eggs.	Fry.	Yearlings, fingerlings, and adults.	Fry.	Adults and yearlings.	Fry.
1873.....				20,000	520,000				
1874.....					850,000				
1875.....			250,000	250,000	2,250,000				
1876.....					2,000,000				
1877.....					2,200,000				
1878.....					2,500,000				
1879.....					2,300,000				
1880.....					2,225,000				
1881.....			10,000		2,420,500				
1882.....				80,300	3,991,750				
1884.....					600,000				
1886.....					150,000				
1887.....					200,000				
1888.....					1,790,000				
1889.....					3,668,000				
1890.....					1,494,000				
1891.....					3,575,000				
1892.....					2,966,600	25,000			
1893.....					5,131,950				
1894.....					8,214,900		280,000		353,500
1895.....					3,935,000		910,000	560,000	
1896.....					15,748,883	250,000			107,808
1897.....					20,324,701		298,137		262,000
1898.....					45,101,688				650,000
1899.....				85,200	25,409,110				
1900.....					6,072,950				
1901.....					4,128,570				
1902.....					18,967,600				301,000
1903.....					5,297,947				120,000
1904.....					65,982,130				90,000
1905.....					102,661,380				108,000
1906.....					110,204,472				243,000
1907.....	80,000				75,029,250		80,000		487,000
1908.....	80,000				66,199,855		80,000		170,000
1909.....	42,000	1,200			31,590,000		42,000		518,200
Total.....	202,000	1,200	260,000	435,500	645,201,236	275,000	1,690,137	560,000	3,410,508

OREGON. .

HATCHERIES ON COASTAL STREAMS.

Rogue River.—In 1877 Mr. R. D. Hume, who had been packing salmon on this river for some years, erected a hatchery at Ellensburg. In 1888 the Oregon Legislature appropriated a sum of money for the enlargement and support of this hatchery, Mr. Hume to retain complete control. As the location is on tidewater it is necessary to catch the parent fish and hold them until they are ready to spawn, and in order to do this Mr. Hume had an excavation 32 by 62 feet and 11 feet deep made in the bank of the river. This was lined with concrete 1 foot thick, which, when filled with water, made a pond 30 by 60 feet and 10 feet deep. Over the entire pond he constructed a building which could be closed up so as virtually to

exclude the light. It is supposed that retaining the fish in a dark place aids in keeping them in good physical condition until ready to spawn. The death of Mr. Hume in 1908 may lead to the abandonment of this hatchery, unless the State or Government takes it over.

In 1897 Mr. Hume built and equipped a hatchery on the upper Rogue River at the mouth of Elk Creek, about 26 miles from the town of Central Point, in Jackson County, and, in pursuance of an understanding with the United States Fish Commission, the latter operated then and still continues to operate this plant.

In 1900 the Government established an auxiliary station for the collection of steelhead trout eggs on Elk Creek, about 10 miles above the main station. In 1905 a substation was operated at Grants Pass, while during the fiscal year 1908 substations were operated at Findley Eddy, on the Rogue River, Illinois River, and Applegate Creek, tributaries of the Rogue.

Many of the eggs gathered at the upper Rogue River stations were shipped to Mr. Hume's hatchery, on the lower river, and there hatched out and planted.

Coquille River.—The State formerly had a hatchery on this river, but it was abandoned during the winter of 1902-3. In the winter of 1904-5 a substation was established on one of the tributaries of the Coquille River, about 6 miles from the South Coos River hatchery, and was used in hatching eggs brought to it from the latter place.

Coos River.—A hatchery was built by the State in 1900 on the South Coos River, about 20 miles from the town of Marshfield.

Umpqua River.—In 1900 the State built a hatchery on the north fork of the Umpqua River, near the town of Glide and about 24 miles east of Roseburg. In 1901 a station was established farther up the north fork, at the mouth of Steamboat Creek. After working here two years the station was moved a couple of miles farther up the stream. In 1907 work was resumed again at the original station near Glide, as winter freshets had seriously damaged the upper station.

Siuslaw River.—In 1893 the State erected a hatchery on Knowles Creek, a tributary of the Siuslaw River, about 20 miles above the mouth of the river. It was turned over to the United States Fish Commission to operate, but no fish came up to the hatchery because the fishermen lower down stretched their nets entirely across the river.

In 1897 and 1898 the United States Fish Commission operated a hatchery owned by a Mr. McGuire and located close to Mapleton, about 2 miles below the head of tidewater.

In 1902 the State established an experimental station at the Bailey place, near Meadow post office. In 1907 a permanent station was established by the State on Land Creek fork of the Siuslaw River.

Alsea River.—In 1902 the State established a station on the Willis Vidito place, near the town of Alsea. In 1907 an experimental station was established on this river at the mouth of Rock Creek, about 14 miles above the head of tidewater.

Yaquina River.—In 1902 the State established a hatching station on the Big Elk River, a tributary of Yaquina River, about 3 miles above its confluence with the main river. This station was made permanent the next year.

Tillamook Bay.—In 1902 the State established a station on Wilson River, a tributary of Tillamook Bay, and about 8 miles above tide water. In 1906 the station was removed to the Trask River, a tributary of Tillamook Bay.

DISTRIBUTION.

The following table shows the distribution of fry in the coastal streams of the State by the Government and the State.

DISTRIBUTION OF SALMON FRY IN THE COASTAL STREAMS OF OREGON.

Year ending June 30—	Tillamook Bay and tributaries.			Yaquina River.			Alsea River.	
	Chinook.	Silverside.	Steel-head.	Chinook.	Silverside.	Steel-head.	Chinook.	Silverside.
	Fry.	Fry.	Fry.	Fry.	Fry.	Fry.	Fry.	Fry.
1898.....	19,994							
1901.....				213,500				
1903.....	251,875			557,700			67,750	
1904.....	799,300			3,144,380	985,220			
1905.....				1,407,470	3,009,075	780,500	1,000,000	1,000,000
1906.....				816,608	4,178,000	1,033,150	806,938	1,785,351
1907.....	312,700	2,648,000		1,919,508	1,955,793	376,245		
1908.....	2,124,000	1,629,000		2,193,043	909,855		199,700	812,300
1909.....		4,896,000	509,690	485,500	1,006,309			
1910.....	624,800	3,506,990	2,309,770	324,038	28,815			
Total.....	4,132,669	12,679,990	2,879,460	11,061,747	12,073,067	2,189,895	2,074,388	3,597,651

Year ending June 30—	Siuslaw River.			Umpqua River.	Coos Bay and tributaries.		
	Chinook.	Silverside.	Steel-head.	Chinook.	Chinook.	Silverside.	Steel-head.
	Fry.	Fry.	Fry.	Fry.	Fry.	Fry.	Fry.
1897.....	180,000						
1898.....	440,275						
1899.....	2,700,000						
1901.....	213,500			730,000	235,000		
1902.....	112,000	214,800		1,136,000	2,416,350		
1903.....	389,239			1,596,213			
1904.....	822,567			1,399,860	4,079,274		
1905.....	435,162	311,900		2,654,925	3,877,172		
1906.....	1,826,531	1,296,732	397,355	4,903,700	2,744,000		
1907.....	608,949	1,030,486		4,685,900	4,014,400		
1908.....	729,130	1,127,293		2,378,853	3,000,000		
1909.....	191,267	1,092,540	98,243	4,093,848	2,084,500	1,032,000	222,000
1910.....	273,352	25,289		5,686,273	1,683,738		
Total.....	8,921,972	5,099,040	495,598	29,265,572	24,134,434	1,032,000	222,000

DISTRIBUTION OF SALMON FRY IN THE COASTAL STREAMS OF OREGON—Continued.

Year ending June 30—	Coquille River.		Rogue River and tributaries.			
	Chinook.	Silverside.	Chinook.		Silverside.	Steelhead.
	Fry.	Fry.	Fry.	Yearlings, fingerlings, and adults.	Fry.	Fry.
1877.....			50,000			
1898.....			1,910,045			
1900.....			2,156,945			
1901.....	235,000		2,967,058		128,000	65,850
1902.....			4,750,763		424,530	20,250
1903.....	3,084,577		3,480,300		680,800	
1904.....	1,000,000		9,023,428			8,073
1905.....	2,210,000		4,758,653		1,250,432	531,000
1906.....	2,978,700		47,500	75,000		12,625
1907.....	2,840,000		5,880,290		1,375,000	105,300
1908.....	2,450,000	226,600	6,597,027	170,051	158,000	937,680
1909.....		1,185,800	771,710		643,000	878,847
1910.....			1,430,292			89,850
Total.....	14,798,277	1,412,400	43,824,011	245,051	4,659,762	2,649,475

Year ending June 30—	Total.			
	Chinook.		Silverside.	Steelhead.
	Fry.	Yearlings, fingerlings, and adults.	Fry.	Fry.
1877.....	50,000			
1897.....	180,000			
1898.....	2,370,314			
1899.....	2,700,000			
1900.....	2,156,945			
1901.....	4,594,058		128,000	65,850
1902.....	8,415,113		639,330	20,250
1903.....	9,427,654		680,800	
1904.....	20,268,809		985,220	8,073
1905.....	16,343,382		5,571,407	1,311,500
1906.....	14,123,977	75,000	7,260,083	1,443,130
1907.....	20,261,747		7,009,279	481,545
1908.....	19,671,753	170,051	4,863,048	937,680
1909.....	7,626,825		9,855,649	1,768,780
1910.....	10,022,493		3,561,094	2,399,620
Total.....	138,213,070	245,051	40,553,910	8,436,428

The following tables show the total output of the hatcheries in Oregon owned by the United States Bureau of Fisheries and the State of Oregon:

OUTPUT OF HATCHERIES OWNED BY THE UNITED STATES BUREAU OF FISHERIES.

Year ending June 30—	Chinook.			Silver.		
	Eggs.	Fry.	Fingerlings, yearlings, and adults.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
1889.....		4,500,000				
1890.....	1,000,000	2,776,475				
1891.....	700,000	4,901,525				
1892.....		1,332,400				
1893.....		4,100,000				
1894.....		213,000				
1895.....	23,000					
1896.....		^a 2,832,150	^b 557,150			
1897.....		4,922,634				
1898.....		16,915,512				
1899.....	27,000	4,300,200				
1900.....	1,800,000	4,126,367			146,824	
1901.....	1,100,000	1,669,857	1,668		128,000	
1902.....	1,866,000	11,587,061			424,530	
1903.....	4,884,400	5,453,860	250	680,800		
1904.....	3,113,000	15,270,675				
1905.....	30,000	9,822,636			1,250,432	
1906.....	28,200	2,454,371	122,980			300
1907.....	1,661,390	8,542,104				
1908.....	2,045,000	7,844,827	627,856		158,000	57,932
1909.....	3,531,000	5,021,655	2,763		1,799,915	
1910.....	3,953,992	4,220,197	225			
Total.....	25,762,982	122,807,506	1,312,892	680,800	3,907,701	58,232

Year ending June 30—	Steelhead trout.			Total.		
	Eggs.	Fry.	Fingerlings, yearlings, and adults.	Eggs.	Fry.	Fingerlings, yearlings, and adults.
1889.....					4,500,000	
1890.....				1,000,000	2,776,475	
1891.....				700,000	4,901,525	
1892.....					1,332,400	
1893.....					4,100,000	
1894.....					213,000	
1895.....				23,000		
1896.....					2,832,150	557,150
1897.....					4,922,634	
1898.....					16,915,512	
1899.....	159,000	12,125		180,000	4,312,325	
1900.....	415,000	99,000		2,215,000	4,372,191	
1901.....	246,000	65,850	25,000	1,346,000	1,863,707	26,668
1902.....	481,000	20,250		2,347,000	12,031,841	
1903.....	400,000	262,700	62,033	5,965,200	5,716,560	62,283
1904.....		23,205	11,090	3,113,000	15,293,880	11,090
1905.....	50,000	534,000		80,000	11,607,068	
1906.....	10,000	1,294,485	40,383	38,200	3,748,856	163,663
1907.....	50,000	105,300		1,711,390	8,647,404	
1908.....	263,725	952,680		2,308,725	8,955,507	685,788
1909.....	51,468	1,374,308		3,582,468	8,195,878	2,763
1910.....		2,074,188		3,953,992	6,294,385	225
Total.....	2,126,193	6,818,091	138,506	28,569,975	133,533,298	1,509,630

^a All but 17,000 of these were from eggs received from the California stations.

^b All raised from eggs received from the California stations.

OUTPUT OF HATCHERIES OWNED BY THE STATE OF OREGON.

Year.	Chinook fry.	Silverside fry.	Steelhead trout fry.	Total.
1877.....	50,000			50,000
1878.....	79,620			79,620
1879.....	1,876,500			1,876,500
1880.....	1,834,290			1,834,290
1881.....	2,554,290			2,554,290
1888.....	1,300,000			1,300,000
1889.....	4,500,000			4,500,000
1890.....	990,000			990,000
1891.....	^a 792,000			792,000
1895.....	2,500,000			2,500,000
1896.....	2,500,000			2,500,000
1899.....	2,700,000			2,700,000
1900.....	2,500,000		200,000	2,700,000
1901.....	7,562,000		245,000	7,807,000
1902.....	11,220,550	7,957,000	256,327	19,433,877
1903.....	18,502,072	3,288,600	300,850	22,091,522
1904.....	^b 48,730,791	3,974,185	143,849	52,848,825
1905.....	16,393,249	^c 5,509,085	1,495,735	23,398,069
1906.....	^c 27,404,596	7,503,655	1,859,696	36,767,947
1907.....	^d 25,156,732	6,446,628	376,245	31,979,605
1908.....	^e 21,209,394	5,359,709		26,569,103
1909.....	^f 20,108,990	9,212,649	1,403,129	30,724,768
1910.....	^g 24,169,365	3,631,827	2,364,120	30,165,312
Total.....	244,634,439	52,883,338	8,644,951	306,162,728

^a Eggs from which hatched obtained from United States Bureau of Fisheries.

^b 6,826,540 eggs were obtained from United States Bureau of Fisheries.

^c 7,714,000 eggs were obtained from United States Bureau of Fisheries.

^d 3,550,000 eggs were obtained from United States Bureau of Fisheries.

^e 3,020,000 eggs were obtained from United States Bureau of Fisheries.

^f 6,581,000 eggs were obtained from United States Bureau of Fisheries.

^g 6,465,300 eggs were obtained from United States Bureau of Fisheries.

COLUMBIA RIVER AND TRIBUTARIES.

The first fish-cultural work upon the Columbia River and in Oregon was at Clackamas, on the Clackamas River, a tributary of the Willamette River, which empties into the Columbia River about 180 miles from its mouth.

This hatchery was built in 1876 by the Oregon & Washington Fish Propagating Co., which operated it until 1880. In 1887 the State provided for and there was appointed a State fish commission. Almost the first work of the commission was to spend \$12,000 appropriated by the legislature to put in repair and operate this hatchery. On July 1, 1888, it was informally turned over to the United States Commission of Fish and Fisheries, which paid over the purchase price, took formal possession in the following winter, and has operated it ever since, with the exception of several years when the building of dams stopped the progress of salmon to the hatchery. During this period a temporary station for the collection of eggs was established on Sandy River, about 15 miles away, and on Salmon River, a tributary of Sandy River, both tributaries of the Columbia River. Some eggs were also brought in from the California hatcheries and hatched at the Clackamas station. In 1901 the hatchery was moved about 4 miles down the river and has since been operated as both a rearing and a collecting station. In 1901 the State established

another hatchery on the Clackamas River about 30 miles below the main station and between the North and South Forks. In 1904 all were turned over to the United States. In 1907 an experimental station for the collection of eggs of the early variety of chinook salmon was established by the State of Oregon on the Clackamas River below the Portland Railway, Light & Power Co.'s dam at Cazadero, but this is now operated by the United States Bureau of Fisheries.

In 1889 the State established a hatchery in the cannery of Mr. F. M. Warren, at Warrendale, in Multnomah County, on the Columbia River, which was operated in that year and in 1890.

In 1895 some of the Oregon salmon packers combined and organized the Columbia River Packers' Propagating Co., which established a hatchery on the upper Clackamas River at the junction of the Warm Springs and the Clackamas and operated it in 1895 and 1896. The Government operated it in 1897 and 1898, after which it was turned over to the State and moved to the opposite side of the river.

In 1898 the collection of steelhead trout eggs was first undertaken on the northwest coast by the State of Oregon on Salmon River, a tributary of the Columbia River, and met with fair success. In March, 1899, the Government sent a party to the falls of the Willamette River, near Oregon City, to collect steelhead eggs, and also operated for this purpose at its substation on the Salmon River, but the latter effort met with failure, as the rack was washed away. This station was turned over to the State on June 15, 1899.

In 1901 the State of Oregon did some experimental work at Swan Falls, on Snake River, the boundary for a considerable distance between Oregon and Idaho. During the winter and early spring of 1902 the State also worked Tucannon River, which is a tributary of Snake River, for steelhead, but met with poor success. Snake River was worked again in 1902 at the foot of Morton Island, which is situated 2 miles above Ontario, in Malheur County. Title to the necessary property was secured from the War Department in 1903 and permanent buildings were erected.

In 1901 the State of Oregon established an experimental hatchery in Wallowa County, on the Grande Ronde River, at the mouth of a small tributary called the Wenaha River, which enters the main stream about 50 miles from its mouth. A permanent station was established in the canyon about $1\frac{1}{2}$ miles below the Wallowa bridge on the Wallowa River, a tributary of the Grande Ronde River, in 1903.

In 1902 the State of Oregon erected a permanent plant on Salmon River at its junction with Boulder Creek.

In the same year the State established an experimental station on the McKenzie River, a tributary of the Willamette River, about

one-half mile above Vida post office. This experimental work was resumed in 1905 at a point 2 miles below Gate Creek. The hatchery was permanently established at a spot about 30 miles from Eugene and near the town of Leaburg a year or two later.

In 1906 an experimental station was established by the State on Breitenbush Creek a short distance above its junction with the Santiam River, a tributary of the Willamette River, but the plant was destroyed very shortly after its establishment, by a forest fire. An experimental station was reestablished here in 1909, but a heavy freshet raised the river so high that the penned fish escaped around the rack.

In 1909 the State of Oregon built at Bonneville, on Tanner Creek, a tributary of the Columbia River, a large central hatchery capable of handling 60,000,000 eggs, it being the intention of the State to hatch at this plant the eggs collected at other stations.

The first entrance of Washington (then a Territory) into fish-cultural operations was in 1879, when the State fish commissioner paid the Oregon & Washington Fish Propagating Co., which was operating the hatchery on the Clackamas River, \$2,000 for salmon fry deposited in that river. In 1893 the State legislature established a hatchery fund which was to be supplied by licenses from certain lines of the fishery business. In 1895 its first hatchery in the Columbia River Basin was built on the Kalama River, about 4 miles distant from its junction with the Columbia, and in Cowlitz County. Another station for the collection and eyeing of eggs was established on the Chinook River, a small stream which empties into Baker Bay near the mouth of the Columbia.

During the fiscal year 1897 the United States Fish Commission established a station on Little White Salmon River, a stream which empties into the Columbia, on the Washington side, about 14 miles above the Cascades. During the fiscal year 1901 an auxiliary station was operated on Big White Salmon River, while fishing was carried on in Eagle and Tanner Creeks, in Oregon, the eggs obtained from these creeks being brought to the Little White Salmon hatchery.

In 1899 the State of Washington built and operated hatcheries on the Wenatchee River, a tributary of the Columbia River, about $1\frac{1}{2}$ miles from Chiwaukum station on the Great Northern Railway, and on Wind River, a tributary of the Columbia, about 1 mile from the junction.

In 1900 Washington State hatcheries were established in the Columbia River basin as follows: White River hatchery, which was built on Coos Creek, which empties into a tributary of the White River, the location being about $2\frac{1}{2}$ miles from where the Green River joins the White River; Methow River hatchery, built on the Methow River at the point where it is joined by the Twisp, about

22 miles from the Columbia River; Colville River hatchery, built on the north bank of Colville River, about $1\frac{1}{2}$ miles from its mouth, and about 1 mile from Kettle Falls; Klickitat River hatchery, located on the east bank of the Klickitat River, about 6 miles from its mouth; and one on the Little Spokane River, about 10 miles from its mouth and about 9 miles north of the city of Spokane. The Klickitat River hatchery never was operated, while most of the others were operated intermittently.

In 1906 a hatchery was established by the State of Washington on the Lewis River, some distance above the town of Woodland.

The following table shows the plants of salmon and steelhead trout in the Columbia River and its tributaries by the Bureau of Fisheries and the States of Oregon and Washington:

TABLE SHOWING THE PLANTS OF SALMON FRY IN THE COLUMBIA RIVER BASIN SINCE 1877.

Year ending June 30—	Columbia River and tributaries.			Total.
	Chinook fry.	Silverside fry.	Steelhead trout fry.	
1877.....	300,000			300,000
1878.....	79,620			79,620
1879.....	3,076,500			3,076,500
1880.....	1,834,290			1,834,290
1881.....	2,554,290			2,554,290
1888.....	1,300,000			1,300,000
1889.....	4,500,000			4,500,000
1890.....	3,756,475			3,756,475
1891.....	5,694,000			5,694,000
1892.....	1,332,400			1,332,400
1893.....	4,100,000			4,100,000
1894.....	213,000			213,000
1895.....	a 2,523,000			2,523,000
1896.....	b 10,389,300			10,389,300
1897.....	10,641,394			10,641,394
1898.....	26,212,074			26,212,074
1899.....	19,979,241		8,625	19,987,866
1900.....	22,510,869	7,175,824	299,000	29,985,693
1901.....	c 24,978,978	5,559,750	245,000	30,783,728
1902.....	44,328,085	17,545,724	256,327	62,130,136
1903.....	40,174,313	8,721,720	d 600,583	49,496,616
1904.....	71,694,587	8,422,085	e 158,981	80,275,653
1905.....	17,107,217	1,354,610	e 768,235	19,230,062
1906.....	f 36,372,785	g 828,872	h 1,769,494	38,971,151
1907.....	23,171,235	2,657,349	26,640	25,855,224
1908.....	i 34,852,008	1,705,543	15,000	36,572,551
1909.....	+ j 33,098,943	2,439,415	k 1,058,657	36,597,015
1910.....	l 37,744,002	3,374,733	m 2,063,688	43,182,423
Total.....	484,518,606	59,785,625	7,270,230	551,574,461

a Includes 23,000 eggs.

b Includes 557,150 yearlings, fingerlings, or adults.

c Includes 1,668 yearlings, fingerlings, or adults.

d Includes 37,033 yearlings, fingerlings, or adults.

e Includes 50,000 eggs.

f Includes 48,200 eggs and 47,980 yearlings, fingerlings, or adults.

g Includes 300 yearlings, fingerlings, or adults.

h Includes 24,383 yearlings, fingerlings, or adults, and 58,000 eggs.

i Includes 1,995,746 yearlings, fingerlings, or adults.

j Includes 16,949 yearlings, fingerlings, or adults.

k Includes 50,000 eggs.

l Includes 225 yearlings, fingerlings, or adults.

m Includes 25,000 eggs.

WASHINGTON.

Willapa River.—In 1899 Washington established a hatchery on Trap Creek, a tributary of the Willapa River, situated about 200 yards from the creek's mouth.

Chehalis River.—The construction of a hatchery on the Chehalis River, about 4 miles above the city of Montesano, was begun by the State in October, 1897, but owing to bad weather and extreme high water was not completed until late in 1898. The hatchery was a failure until 1902 when a fair season was had, as was again true in 1903. It was not operated in 1904. Since the State began taking eggs from the Satsop River, a tributary of the Chehalis, it has been possible to fill the hatchery each season.

Puget Sound and tributaries.—In 1896 the State established a hatchery on Baker Lake, which is the head of Baker River, a tributary of the Skagit River, and this was the first establishment for the hatching of sockeye salmon. In July, 1899, it was sold to the United States Fish Commission. In 1901 steelhead trout eggs were collected on Phinney Creek, about 5 miles from the town of Birdsvew, and some 30 miles from Baker Lake. In 1901 an auxiliary station was opened at Birdsvew, on Skagit River, and steelhead trout eggs were collected on Phinney and Grandy Creeks and brought to Baker Lake to be hatched.

In 1898 a private hatchery (the necessary money being raised by subscription among the residents of Fairhaven, now Bellingham, and vicinity) was built near Lake Samish, a few miles from Fairhaven.

In 1899 a hatchery was built by the State on Kendall Creek, a tributary of the Nooksack River, about 300 yards from same, and about 2 miles from the railway station of Kendall. Except in 1903, this hatchery has since been operated continuously. An eyeing station was built in 1907 on the south fork of the Nooksack River, about 1 mile from Acme.

In the same year the State built a hatchery on the Skokomish River, about 4 miles from its mouth. An eyeing station was also erected on the north fork of the same river. The main station was not operated in 1904 and only on a small scale in 1903 and 1905.

The State in 1899 built a hatchery on Friday Creek, a tributary of the Samish River, situated about 1 mile from the mouth of the creek.

The following State hatcheries were first operated in 1900. Snohomish hatchery, built on the west bank of the Skykomish River, a few miles from its mouth; Nisqually River hatchery, built on Muck Creek, about one-half mile from the Nisqually River, and about 4 miles from the town of Roy, in Pierce County; and the Stillaguamish hatchery, located on the Stillaguamish River, about 4 miles from the

town of Arlington, in Snohomish County. The latter has since been moved to Jim Creek, a tributary of the south branch of the Stillaguamish River.

The Startup hatchery, located near Startup, on the Skykomish River, was formerly used as a collecting station for the Snohomish hatchery. It is still used for this purpose, but also retains and hatches a considerable quantity of spawn. The station is about 4 miles from the Snohomish hatchery.

In 1900 the State established a fisheries experimental station at Keyport Landing, on the east arm of Port Orchard Bay, with Pearson as the nearest post office. The work of the station is devoted to salmon and oysters.

The State established a hatchery on the Dungeness River, about 7 miles from the town of Dungeness, in Clallam County, in 1901. In 1906 it constructed a hatchery on a small tributary of the Skagit River, between Hamilton and Lyman. The station built on Sauk River, a tributary of the Skagit, has been operated only occasionally since the Skagit hatchery was built.

The United States Bureau of Fisheries has now (1911) under construction hatcheries on the Duckabush and Quilcene Rivers in Hoods Canal.

The following tables show the total output of the salmon hatcheries in the State of Washington owned by the United States Bureau of Fisheries and the hatcheries owned by the State itself:

OUTPUT OF THE SALMON HATCHERIES IN WASHINGTON OWNED BY THE UNITED STATES BUREAU OF FISHERIES.

Year ending June 30—	Chinook.			Sockeye, or blueback.			Silver.	
	Eggs.	Fry.	Finger- lings, yearlings, and adults.	Eggs.	Fry.	Finger- lings, yearlings, and adults.	Eggs.	Fry.
1897.....		1,848,760						
1898.....		7,391,886						
1899.....	4,926,000	1,791,056						
1900.....	2,686,000	6,626,947			10,683,000			
1901.....	6,581,000	5,427,680			3,834,453			174,041
1902.....		15,637,687			3,371,000			
1903.....		16,774,030			3,731,789			81,812
1904.....	7,506,000	17,386,183			3,855,000			3,984,645
1905.....		4,236,276			7,819,281	10,000	107,000	8,071,081
1906.....	7,714,000	14,846,905		880,000	3,285,130	9,500	239,180	6,445,574
1907.....	3,550,000	6,512,738			4,224,255		760,000	3,636,952
1908.....	1,485,000	12,372,503	1,537,941	75,000	8,514,305		296,000	13,262,714
1909.....	3,050,000	11,565,553	14,186	100,000	5,430,626		272,000	7,661,110
1910.....	3,813,250	9,175,610			4,554,825		275,000	10,888,025
Total..	41,311,250	131,593,814	1,552,127	1,055,000	59,303,664	19,500	1,949,180	54,205,954

OUTPUT OF THE SALMON HATCHERIES IN WASHINGTON OWNED BY THE UNITED STATES
BUREAU OF FISHERIES—Continued.

Year ending June 30—	Humpback.		Steelhead trout.			Total.		
	Eggs.	Fry.	Eggs.	Fry.	Finger- lings, yearlings, and adults.	Eggs.	Fry.	Finger- lings, yearlings, and adults.
1897.....							1,848,760	
1898.....							7,391,886	
1899.....						4,926,000	1,791,056	
1900.....				26,000		2,686,000	17,335,947	
1901.....						6,581,000	9,436,174	
1902.....				110,000			19,118,687	
1903.....			80,000	440,000	223,815	80,000	21,027,631	223,815
1904.....		176,597	255,000	70,000		7,761,000	25,472,425	
1905.....			414,400	3,205		521,400	20,129,843	
1906.....	2,000	969,990	348,000	540,000		9,183,180	26,087,599	9,500
1907.....			200,000	941,505		4,510,000	15,315,450	
1908.....	502,000	6,764,762	224,000	136,916		2,582,000	41,051,200	1,537,941
1909.....			220,000	717,691		3,642,000	25,374,980	14,186
1910.....		1,368,000	300,000	1,437,038		4,388,250	27,423,498	
Total.....	504,000	9,279,349	2,041,400	4,422,355	223,815	46,860,830	258,805,136	1,795,442

OUTPUT OF THE SALMON HATCHERIES OWNED BY THE STATE OF WASHINGTON.

Year ending June 30—	Chinook fry.	Dog fry.	Hump- back fry.	Silverside, or coho, fry.	Sockeye, or blue- back, fry.	Steelhead trout fry.	Total.
1896.....	4,500,000						4,500,000
1897.....	4,050,000						9,550,000
1898.....	4,275,000				5,500,000		9,675,000
1899.....	8,595,000			189,000			8,784,000
1900.....	12,251,600	10,301,760		13,778,280		1,736,560	38,068,200
1901.....	12,275,400	16,478,280		19,747,894		1,398,476	49,900,050
1902.....	14,766,822	9,937,390		32,964,593		2,481,371	60,150,176
1903.....	14,283,499	9,937,390		28,659,079		3,134,076	56,014,044
1904.....	13,261,184			15,725,196		3,868,866	33,150,446
1905.....	7,101,180		295,200	12,226,294		2,433,635	21,761,109
1906.....	10,943,550	3,268,800		28,906,380		2,769,784	45,888,514
1907.....	8,897,670	6,120,000		28,668,600		3,575,943	47,262,213
1908.....	18,647,600	4,342,350	2,655,900	29,273,202		4,578,075	59,497,127
1909.....	17,440,950	8,218,000		24,543,200		4,080,450	54,282,600
1910.....	21,168,350	8,607,500	519,600	30,894,100		4,855,000	66,044,550
Total.....	172,457,805	77,211,470	3,470,700	265,575,818	10,900,000	34,912,236	564,528,029

NOTE.—As the printed reports of the State in many instances report as the output the number of eggs gathered, it has been necessary in such cases to make an arbitrary reduction from these figures, in order to allow for the loss in the egg stage.

The following table shows the plantings made in waters of Washington other than the Columbia River by the United States Bureau of Fisheries and the State of Washington:

PLANTS OF SALMON FRY IN THE WATERS OF WASHINGTON OTHER THAN THE COLUMBIA RIVER.

Year ending June 30—	Puget Sound and tributaries.					
	Chinook.	Sockeye.	Silver, or coho.	Hump-back.	Dog.	Steelhead.
1897.....		5,500,000				
1898.....		5,400,000				
1899.....	7,470,000		189,000			
1900.....		10,683,000	6,749,280		10,301,760	1,572,560
1901.....	300,000	3,834,453	14,360,185		16,478,280	1,398,476
1902.....	2,141,322	3,371,000	23,161,069		9,937,390	2,591,371
1903.....	2,113,850	3,731,789	21,507,771		9,937,390	a 3,326,091
1904.....	1,865,933	3,855,000	14,071,845	471,797		b 3,518,476
1905.....	2,590,738		16,441,375			b 1,329,940
1906.....	4,819,290	c 3,582,630	d 29,770,414	969,990	1,800,000	e 3,177,174
1907.....	3,907,598		26,960,552	4,224,255	5,230,000	3,964,308
1908.....	8,356,709	8,514,305	37,613,466	9,420,662	2,278,350	4,566,491
1909.....	9,647,288	5,430,626	28,622,310		6,048,000	f 4,499,141
1910.....	11,681,060	4,554,825	36,837,125	1,887,600	7,748,500	6,292,338
Total.....	54,893,788	58,457,628	256,284,392	16,974,304	69,749,670	36,236,366

Year ending June 30—	Chehalis River.			Willapa River.		
	Chinook.	Silver, or coho.	Dog.	Chinook.	Silver, or coho.	Steelhead.
1899.....	1,215,000					190,000
1900.....	2,355,300			881,000		653,400
1901.....	1,909,800					2,163,019
1903.....					1,800,000	500,000
1904.....	900,000			819,504	204,876	420,390
1905.....				630,000	1,800,000	288,000
1906.....		2,563,380	1,468,800	529,650	2,160,000	171,550
1907.....		2,250,000	960,000	393,660	2,250,000	526,500
1908.....	163,000	3,275,000	2,064,000	678,600	654,500	148,500
1909.....	148,000	1,800,000	1,757,000	322,200	504,000	399,000
1910.....	403,000	1,577,000	859,000	455,200	64,000	
Total.....	7,094,100	11,465,380	7,048,800	7,526,233	9,437,376	2,643,940

Year ending June 30—	Total by species.						Grand total.
	Chinook.	Sockeye.	Silver, or coho.	Hump-back.	Dog.	Steelhead.	
1878.....	g 3,000						3,000
1897.....		5,500,000					5,500,000
1898.....		5,400,000					5,400,000
1899.....	8,685,000		189,000				8,874,000
1900.....	3,236,300	10,683,000	6,749,280		10,301,760	1,762,560	32,732,900
1901.....	2,863,200	3,834,453	14,360,185		16,478,280	1,398,476	38,934,594
1902.....	2,141,322	3,371,000	23,161,069		9,937,390	2,591,371	41,202,152
1903.....	4,276,869	3,731,789	23,307,771		9,937,390	3,826,091	45,079,910
1904.....	3,585,437	3,855,000	14,276,721	471,797		3,938,866	26,127,821
1905.....	3,220,738		18,241,375			1,617,940	23,080,053
1906.....	5,348,940	3,582,630	34,493,794	969,990	3,268,800	3,348,724	51,012,878
1907.....	4,301,258		31,460,552	4,224,255	6,120,000	4,490,808	50,586,873
1908.....	9,198,309	8,514,305	41,542,966	9,420,662	4,342,350	4,714,991	77,733,583
1909.....	10,117,488	5,430,626	30,926,310		7,805,000	4,898,141	59,177,565
1910.....	12,539,260	4,554,825	38,478,125	1,887,600	8,607,500	6,292,338	72,359,648
Total.....	69,517,121	58,457,628	277,187,148	16,974,304	76,798,470	38,880,306	537,814,977

a Of these, 218,200 were yearlings, fingerlings, or adults.

b Of these, 14,400 were eggs.

c Of these, 9,500 were yearlings, fingerlings, or adults.

d Of these, 14,840 were yearlings, fingerlings, or adults.

e Of these, 15,000 were yearlings, fingerlings, or adults.

f Includes 100,000 eggs.

g These were brought from the Clackamas (Oregon) station and planted in some unnamed lake.

BRITISH COLUMBIA.

Fraser River.—The first hatchery established by the Dominion of Canada on the Pacific coast was erected in 1884 at what is now Bon Accord, a point on the lower river some 4 miles above New Westminster, and on the opposite shore. The next built was in 1901 on Granite Creek, Shuswap Lake, which discharges into the Fraser through the South Thompson River, the lake being about 280 miles from New Westminster. In 1904 another hatchery was established on Harrison Lake on the Lillooet River, first large tributary of the Fraser on the north side; also one about 4 miles east of the lower extremities of Pemberton Meadows, at the junction of Owl Creek and the Birkenhead River, 4 miles above its confluence with the eastern branch of the Lillooet River, which in turn discharges into Lillooet Lake. In 1907 a hatchery was built on Stuart Lake, near the headwaters of the Fraser.

The Province of British Columbia owns Seton Lake Hatchery, which was established in 1903 on Lake Creek, on the north side, about half a mile from the outlet of Seton Lake, and it has been operated continuously ever since. Seton Lake is a part of the Fraser River chain and is some 300 miles above the mouth of the river. Lake Creek, the outlet of Seton Lake, empties into the Cayoosh Creek, a tributary of the Fraser, 45 miles north of the latter's junction with the Thompson, and 1 mile south of the town of Lillooet.

Nimkish River.—In 1902 Mr. S. A. Spencer, of the Alert Bay cannery (now belonging to the British Columbia Packers' Association), in return for certain special fishery privileges granted by the Dominion, established a hatchery on this river, which is located on the northeast shore of Vancouver Island. The hatchery was burned down in 1903, but was immediately rebuilt. Since its establishment it has been operated by the Dominion.

Rivers Inlet.—A hatchery was established by the Dominion on McTavish Creek, one of the tributaries of Oweekayno Lake, about 20 miles up Rivers Inlet, in 1905, and has been operated ever since.

Skeena River.—In 1902 the Dominion established a hatchery on Lakelse Lake, in the Skeena River basin, about 65 miles up the river from Port Essington. In 1907 another was constructed on Babine Lake, the source of the Skeena River.

The following table shows the plantings made in the waters of British Columbia from the Dominion and provincial hatcheries:

PLANTS OF SALMON FRY MADE IN THE WATERS OF BRITISH COLUMBIA.

Year.	Fraser River.						
	Dog.	Coho.	Spring, or king.	Hump- back.	Sockeye.	Steel- head trout.	Total.
1885.....					1,800,000		1,800,000
1886.....					2,625,000		2,625,000
1887.....					4,414,000		4,414,000
1888.....					5,807,000		5,807,000
1889.....					4,419,000		4,419,000
1890.....					6,640,000		6,640,000
1891.....					3,603,800		3,603,800
1892.....					6,000,000		6,000,000
1893.....					5,674,000		5,674,000
1894.....					6,300,000		6,300,000
1895.....					6,390,000		6,390,000
1896.....					10,393,000		10,393,000
1897.....					5,928,000		5,928,000
1898.....					5,850,000		5,850,000
1899.....					4,742,000		4,742,000
1900.....					6,200,000		6,200,000
1901.....					[No fish.]		
1902.....		90,000			15,808,000	75,000	15,973,000
1903.....	75,000	1,750,000	22,000		12,521,000		14,368,000
1904.....		210,000		50,000	13,729,200	12,000	14,001,200
1905.....		5,576,100	4,381,400		9,244,300		19,201,800
1906.....		4,774,000	1,791,500		100,479,000	4,000	107,048,500
1907.....		3,219,200	1,814,900		36,965,900		42,000,000
1908.....		5,890,000	2,815,000	22,500,000	51,855,200		83,060,200
1909.....		7,375,400	5,772,400		41,909,500		55,057,300
1910.....		450,000	6,300,000		105,312,500		112,062,500
Total.....	75,000	29,334,700	22,897,200	22,550,000	474,610,400	91,000	549,558,300

Year.	Skeena River.	Rivers Inlet.			Nimkish River.
	Sockeye.	Sockeye.	Spring, or king.	Total.	Sockeye.
1903.....	3,450,000				1,636,000
1904.....	4,000,000				2,496,000
1905.....	3,767,900				2,850,000
1906.....	3,784,450	8,000,000		8,000,000	4,873,400
1907.....	4,125,750	8,440,000		8,440,000	4,870,000
1908.....	8,946,950	8,594,000	4,706,000	13,300,000	4,800,000
1909.....	11,882,400	13,300,000		13,300,000	4,500,000
1910.....	^a 11,521,700	12,750,000		12,750,000	5,055,000
Total.....	51,479,150	51,084,000	4,706,000	55,790,000	31,080,400

^a Includes 80,000 coho fry.

PLANTS OF SALMON FRY MADE IN THE WATERS OF BRITISH COLUMBIA—CON.

Year.	Total by species.						Grand total.
	Dog.	Coho.	Spring, or king.	Hump-back.	Sockeye.	Steel-head trout.	
1885.....					1,800,000		1,800,000
1886.....					2,625,000		2,625,000
1887.....					4,414,000		4,414,000
1888.....					5,807,000		5,807,000
1889.....					4,419,000		4,419,000
1890.....					6,640,000		6,640,000
1891.....					3,603,800		3,603,800
1892.....					6,000,000		6,000,000
1893.....					5,674,000		5,674,000
1894.....					6,300,000		6,300,000
1895.....					6,390,000		6,390,000
1896.....					10,393,000		10,393,000
1897.....					5,928,000		5,928,000
1898.....					5,850,000		5,850,000
1899.....					4,742,000		4,742,000
1900.....					6,200,000		6,200,000
1902.....		90,000			15,808,000	75,000	15,973,000
1903.....	75,000	1,750,000	22,000		17,607,000		19,454,000
1904.....		210,000		50,000	20,225,200	12,000	20,497,200
1905.....		5,576,100	4,381,400		15,862,200		25,819,700
1906.....		4,774,000	1,791,500		117,136,850	4,000	123,706,350
1907.....		3,219,200	1,814,900		54,401,650		59,435,750
1908.....		5,890,000	7,521,000	22,500,000	74,196,150		110,107,150
1909.....		7,375,400	5,772,400		71,591,900		84,739,700
1910.....		450,000	6,300,000		134,639,200		141,389,200
Total.....	75,000	29,334,700	27,603,200	22,550,000	308,253,950	91,000	687,907,850

ALASKA.

In 1891 several of the canneries operating at Karluk, on Kodiak Island, combined forces and built a hatchery on the lagoon at that place. As the cannery men were at swords' points in regard to their fishing rights on the spit, in 1892 the hatchery was closed. In May, 1896, the Alaska Packers' Association broke ground for a hatchery at the eastern end of the lagoon, near the outlet of Karluk River, a short distance from where the hatchery was located in 1891, and has operated this plant ever since.

In 1892 Capt. John C. Callbreath, manager of the Point Ellis cannery, on Kuiu Island, operated a small hatchery on the left bank of Kutlakoo stream. It was a very primitive place, and an exceptionally high tide destroyed the whole plant in September. It was never rebuilt.

Capt. Callbreath, however, after seeing to the operation of the hatchery, had returned to Wrangell during the summer, where his attention was again attracted to hatchery work, and in the fall of 1892 he built a small hatchery on Jadjeska stream, Etolin Island, about 200 yards from its mouth. The stream is about one-half mile in length and is the outlet of a small lake. Finding the location unsuitable Capt. Callbreath removed the hatchery in 1893 to the northern side of the lake, about three-eighths of a mile from the head of the outlet, where it still stands. The owner's intention was to build up a stream which had a small natural run of red salmon until it had a large run,

with the hope that the Government would then give him the exclusive right to take these fish from the stream for commercial purposes. The experiment was kept up until the end of the season of 1905, when Capt. Callbreath's failing eyesight compelled the cessation of the actual hatching. Since then a man has been stationed on the stream during the run of spawning fish for the purpose of lifting them over the dam, so that they could reach the spawning beds at the head of the lake. The owner's expectation of a big run as a result of hatching operations was never realized.

In 1896 the Baranof Packing Company, which operated a cannery on Redfish Bay, on the western coast of Baranof Island, built a small hatchery on the lake at the head of Redfish stream. The following winter was so cold that not only the flume, but the whole cataract, froze solid, and as the hatchery was thus left without water the eggs were put into the lake and left to their fate and the hatchery closed down permanently.

In 1897 the North Pacific Trading & Packing Company, at Klawak, Prince of Wales Island, established a hatchery near the head of Klawak stream, close to Klawak Lake. In 1898 the plant was moved to the mouth of a small stream entering the lake about halfway up the western shore. This hatchery has been operated continuously ever since. In 1909 the North Alaska Salmon Co. acquired a half interest in it.

The Pacific Steam Whaling Company in 1898 erected a small hatchery on Hetta Lake, on the west side of Prince of Wales Island, which was operated until the close of the hatching season of 1903-4, when the Pacific Packing & Navigation Company, successor to the original owner, went into the hands of a receiver. In 1907 it was reopened by the Northwestern Fisheries Company, which had acquired the interests of the old company, and has been operated each season since.

Up to 1900 the work of hatching salmon was entirely voluntary on the part of the packers. On May 2 of that year the following regulation was promulgated at the Treasury Department, which at that time had control of the Alaska salmon-inspection service:

7. Each person, company, or corporation taking salmon in Alaskan waters shall establish and conduct, at or near the fisheries operated by him or them, a suitable artificial propagating plant or hatchery; and shall produce yearly and place in the natural spawning waters of each fishery so operated red salmon fry in such numbers as shall be equal to at least four times the number of mature fish taken from the said fisheries, by or for him or them, during the preceding fishing season. The management and operation of such hatcheries shall be subject to such rules and regulations as may hereafter be prescribed by the Secretary of the Treasury. They shall be open for inspection by the authorized official of this department; annual reports shall be made, giving full particulars of the number of male and female salmon stripped, the number of eggs treated, the number and percentage of fish hatched, and all other conditions of interest; and there shall be made a sworn yearly statement of the number of fry planted and the exact location where said planting was done.

On January 24, 1902, this regulation was amended so as to require the planting of "red salmon fry in such numbers as shall be equal to at least ten times the number of salmon of all varieties taken from the said fisheries."

Although the regulation was mandatory, but few of the packers obeyed it, some because no suitable place was to be found within a reasonable distance of their plants, others because the establishment and operation of such a hatchery would cost more than their returns from the industry justified, and others because of lack of knowledge required in hatchery work. The greater number of them absolutely ignored it, and as a result those who conformed to the regulation were placed under a heavy financial handicap. The injustice of this arrangement was patent on its face, and in 1906, when a comprehensive revision of the law was made by Congress, provision was made for reimbursing in the future those cannery men who operated salmon hatcheries. The section covering this point reads as follows:

SEC. 2. That the catch and pack of salmon made in Alaska by the owners of private salmon hatcheries operated in Alaska shall be exempt from all license fees and taxation of every nature at the rate of ten cases of canned salmon to every one thousand red or king salmon fry liberated, upon the following conditions:

That the Secretary of Commerce and Labor may from time to time, and on the application of the hatchery owner shall, within a reasonable time thereafter, cause such private hatcheries to be inspected for the purpose of determining the character of their operations, efficiency, and productiveness, and if he approve the same shall cause notice of such approval to be filed in the office of the clerk or deputy clerk of the United States district court of the division of the District of Alaska wherein any such hatchery is located, and shall also notify the owners of such hatchery of the action taken by him. The owner, agent, officer, or superintendent of any hatchery the effectiveness and productiveness of which has been approved as above provided shall, between the thirtieth day of June and the thirty-first day of December of each year, make proof of the number of salmon fry liberated during the twelve months immediately preceding the thirtieth day of June, by a written statement under oath. Such proof shall be filed in the office of the clerk or deputy clerk of the United States district court of the division of the District of Alaska wherein such hatchery is located, and when so filed shall entitle the respective hatchery owners to the exemption as herein provided; and a false oath as to the number of salmon fry liberated shall be deemed perjury and subject the offender to all the pains and penalties thereof. Duplicates of such statements shall also be filed with the Secretary of Commerce and Labor.

It shall be the duty of such clerk or deputy clerk in whose office the approval and proof heretofore provided for are filed to forthwith issue to the hatchery owner, causing such proofs to be filed, certificates which shall not be transferable and of such denominations as said owner may request (no certificate to cover fewer than one thousand fry), covering in the aggregate the number of fry so proved to have been liberated; and such certificates may be used at any time by the person, company, corporation, or association to whom issued for the payment pro tanto of any license fees or taxes upon or against or on account of any catch or pack of salmon made by them in Alaska; and it shall be the duty of all public officials charged with the duty of collecting or receiving such license fees or taxes to accept such certificates in lieu of money in payment of all license fees or taxes upon or against the pack of canned salmon at the ratio of one thousand fry for each ten cases of salmon. No hatchery

owner shall obtain the rebates from the output of any hatchery to which he might otherwise be entitled under this act unless the efficiency of said hatchery has first been approved by the Secretary of Commerce and Labor in the manner herein provided for.

In 1901 the Pacific Steam Whaling Company established two small hatcheries—one on Nagel stream, which enters the northern side of Quadra Lake, on the mainland of southeast Alaska, and one on a stream entering Freshwater Lake Bay, Chatham Strait. Both were closed down in 1904 when the company failed. In 1908 the Northwestern Fisheries Company, which had acquired the Quadra plant, removed it to a small stream entering the head of the lake and has operated it ever since.

In 1901 the Alaska Packers' Association erected a hatchery on Heckman Lake, the third of a series of lakes on Naha stream, Revilaigedo Island, and about 8 miles from Loring, where the association has a cannery. This is without question the largest and costliest salmon hatchery in the world, having a capacity of 110,000,000 eggs, and the association is entitled to great credit for the public spirit it has shown and the work it has done, entirely without remuneration until 1906, in building and operating not only this hatchery but also the one at Karluk.

The Union Packing Company, at Kell Bay, on Kuiu Island, and Mr. F. C. Barnes, at Lake Bay, on Prince of Wales Island, in 1902 built and operated small hatcheries, both of which were abandoned after one season's work.

Up to 1905 the work of hatching salmon in Alaska was confined to the salmon cannery men. In that year, however, the United States Bureau of Fisheries erected a hatchery on Yes Lake, which empties through a short stream into Yes Bay, on Cleveland Peninsula. In 1907 the bureau constructed another hatchery, on Afognak Lake, near Litnik Bay, Afognak Island.

The following tables show the eggs gathered and the fry planted from the government and privately owned hatcheries in Alaska:

OUTPUT OF THE SALMON HATCHERIES IN ALASKA OWNED BY THE UNITED STATES BUREAU OF FISHERIES, 1906 TO 1910.

Year ending June 30—	Yes Lake hatchery.						Afognak hatchery.			
	Red, or sockeye.		Coho, or silver.		Steelhead trout.		Red, or sockeye.		Humpback.	
	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.
1906....	7,031,480	6,638,550
1907....	58,210,000	54,610,800	182,000	143,500
1908....	65,550,000	61,369,000
1909....	50,000,000	48,653,000	17,000	9,900	46,380,000	39,325,870	12,000	10,000
1910....	72,000,000	69,879,600	76,020,000	71,647,170	499,400	363,740
Total.	252,791,480	241,150,950	17,000	9,900	182,000	143,500	122,400,000	110,973,040	511,400	373,740

OUTPUT OF THE SALMON HATCHERIES IN ALASKA OWNED BY THE UNITED STATES BUREAU OF FISHERIES, 1906 TO 1910—Continued.

Year ending June 30—	Total by species.							Grand total.		
	Red, or sockeye.		Coho, or silver.		Humpback.		Steelhead trout.			
	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.
1906.....	7,031,480	6,638,550							7,031,480	6,638,550
1907.....	58,210,000	54,610,800					182,000	143,500	58,392,000	54,754,300
1908.....	65,550,000	61,369,000							65,550,000	61,369,000
1909.....	96,380,000	87,978,870	17,000	9,900	12,000	10,000			96,409,000	87,998,770
1910.....	148,020,000	141,526,770			499,400	363,740			148,519,400	141,890,510
Total.....	375,191,480	352,123,990	17,000	9,900	511,400	373,740	182,000	143,500	375,901,880	352,651,130

OUTPUT OF PRIVATE SALMON HATCHERIES OF ALASKA, 1893 TO 1910.

NOTE.—Unless otherwise stated in footnotes, all of the fry liberated were red salmon.

Year ended June 30—	Callbreath's hatchery.		Karluk hatchery.		Klawak hatchery.	
	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.
1893.....	900,000	600,000				
1894.....	3,000,000	2,204,000				
1895.....	6,300,000	5,291,000				
1896.....	6,200,000	5,475,000				
1897.....	4,400,000	4,390,000	3,236,000	2,556,440		
1898.....	3,400,000	2,526,000	8,454,000	6,340,000	2,023,000	800,000
1899.....	3,000,000	2,050,000	4,491,000	3,369,000	3,600,000	3,000,000
1900.....	3,400,000	2,335,000	10,496,900	7,872,000	3,600,000	a 1,000,000
1901.....	(b)		19,334,000	15,566,800	(c)	
1902.....	6,000,000	5,500,000	32,800,000	28,700,000	3,500,000	2,800,000
1903.....	6,000,000	5,000,000	23,400,000	17,555,000	3,500,000	1,500,000
1904.....	6,000,000	5,000,000	28,113,000	22,000,000	3,000,000	1,700,000
1905.....	6,050,000	5,250,000	45,500,000	33,670,000	2,800,000	2,000,000
1906.....	7,700,000	6,500,000	36,933,000	28,236,412	2,800,000	2,300,000
1907.....	(d)	(d)	38,679,200	36,846,000	3,600,000	1,187,000
1908.....	(e)	(e)	47,808,200	43,655,000	3,500,000	2,776,000
1909.....	(e)	(e)	40,320,000	37,105,000	3,500,000	3,200,000
1910.....	(e)	(e)	43,228,000	40,620,000	5,800,000	5,300,000
Total.....	f 63,350,000	52,121,000	284,793,300	324,091,652	41,223,000	27,563,000

Year ended June 30—	Hetta hatchery.		Quadra Bay hatchery.		Freshwater Bay hatchery.	
	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.
1893.....						
1894.....						
1895.....						
1896.....						
1897.....						
1898.....						
1899.....	2,800,000	2,600,000				
1900.....	2,000,000	1,500,000				
1901.....	1,800,000	a 500,000				
1902.....	2,500,000	1,700,000	4,500,000	3,500,000	1,500,000	1,000,000
1903.....	4,800,000	4,000,000	5,500,000	4,000,000	(b)	(b)
1904.....	5,127,500	3,750,000	600,000	c 400,000	(d)	(d)
1905.....	(g)	(g)	(g)	(g)	(g)	(g)
1906.....	(g)	(g)	(g)	(g)	(g)	(g)
1907.....	(g)	(g)	(g)	(g)	(g)	(g)
1908.....	8,000,000	6,125,000	(g)	(g)	(g)	(g)
1909.....	8,400,000	8,134,000	3,325,000	3,025,750	(g)	(g)
1910.....	10,313,000	9,000,000	10,863,000	9,850,000	(g)	(g)
Total.....	45,710,500	37,309,000	24,788,000	20,775,750	1,500,000	1,000,000

a Many eggs frozen.

b No run of fish.

c Hatchery was not used, the eggs being hatched out in the lake.

d No report.

e Fish coming in to spawn were lifted over the dam.

f A considerable proportion of these are coho eggs.

g Not operated.

OUTPUT OF PRIVATE SALMON HATCHERIES OF ALASKA, 1893 TO 1910—Continued.


Year ended June 30—	Fortmann hatchery.		Kell Bay hatchery.		Total.	
	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.	Eggs taken.	Fry liberated.
1893.....					900,000	600,000
1894.....					3,000,000	2,204,000
1895.....					6,300,000	5,291,090
1896.....					6,200,000	5,475,000
1897.....					8,636,000	6,946,440
1898.....					13,877,000	9,666,000
1899.....					13,891,000	11,019,000
1900.....					19,496,900	12,707,000
1901.....					21,134,000	16,066,800
1902.....	11,460,000	10,300,000			62,260,000	53,500,000
1903.....	40,050,000	29,005,000	2,500,000	2,000,000	85,750,000	63,060,000
1904.....	22,203,000	13,780,000	(a)	(a)	65,043,500	46,630,006
1905.....	65,010,000	63,181,000	(a)	(a)	119,360,000	104,101,000
1906.....	68,715,000	67,643,000	(a)	(a)	116,148,000	104,679,412
1907.....	105,450,000	80,973,000	(a)	(a)	147,729,200	119,006,000
1908.....	^b 41,280,000	33,920,000	(a)	(a)	100,588,200	86,476,000
1909.....	24,465,000	22,785,000	(a)	(a)	80,010,000	74,249,750
1910.....	53,340,000	50,725,000	(a)	(a)	125,544,000	115,495,000
Total.....	431,973,000	372,312,000	2,500,000	2,000,000	995,867,800	837,172,402


^a Not operated.^b Includes 30,000 coho eggs taken and 27,000 fry liberated.


















SMITHSONIAN INSTITUTION LIBRARIES



3 9088 01017 6378